ERRATUM

Erratum to: An Up-regulation of IRF-1 After a Spinal Cord Injury: Implications for Neuronal Apoptosis

Jian Zhao¹ · Chen Chen² · Jian-Ru Xiao¹ · Hai-Feng Wei¹ · Xu-hui Zhou¹ · Xing-Xing Mao² · Wei-dong Zhang² · Rong Qian² · Xin-lei Chen² · Ming-qing He⁴ · Xiao-Wei Yu^{3,5} · Jian Zhao¹

Published online: 21 March 2016
© Springer Science+Business Media New York 2016

Erratum to: J Mol Neurosci (2015) 57:595–604
DOI 10.1007/s12031-015-0642-2

The original version of this article unfortunately contained a missing reference information. This should be cited in the text, as presented below:

Seung-Won Park et al., JPET 320:1001–1012, 2007 (cited below) showed that IRF-1 mRNA is up-regulated in spinal cord

injury in rats, and this increase is blocked by thiozolidinedione derivatives.

References

Park S-W, Yi J-H, Miranpuri G, Satriotomo I, Bowen K, Resnick DK, Vemuganti R (2007) Thiazolidinedione class of peroxisome proliferator-activated receptor γ agonists prevents neuronal damage, motor dysfunction, myelin loss, neuropathic pain, and inflammation after spinal cord injury in adult rats. JPET 320:1001–1012

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12031-015-0642-2>.

✉ Xiao-Wei Yu
njh306@163.com

✉ Jian Zhao
drzhaojian123@163.com

¹ Department of Orthopedics, Shanghai Changzheng Hospital, Shanghai 200000, China

² Department of Orthopedics, Affiliated Hospital of Nantong University, Nantong, Jiangsu Province 226001, China

³ Department of Neurology, Affiliated Hospital of Nantong University Nantong Second People's Hospital, Nantong, Jiangsu Province 226001, China

⁴ Department of Geriatrics, Affiliated Hospital of Nantong University, Nantong, Jiangsu Province 226001, China

⁵ Department of Neurology, Affiliated Hospital of Nantong University Nantong Second People's Hospital, Nantong, Jiangsu Province 226001, China