CASE REPORT

Extra Nodal Rosai–Dorfman Disease (Sinus Histiocytosis with Massive Lymphadenopathy) Presenting as Asymmetric Bilateral Optic Atrophy

An Atypical Ocular Presentation

Eesha Shukla¹ · Anjali Nicholson¹ · Anamika Agrawal¹ · Darshana Rathod¹

Received: 6 April 2016 / Accepted: 11 April 2016 / Published online: 18 April 2016
© Springer Science+Business Media New York 2016

Abstract Rosai–Dorfman disease (sinus histiocytosis with massive lymphadenopathy, SHML) is a rare, non-hereditary, benign histiocytic proliferative disorder, presenting as painless bilateral cervical lymphadenopathy, with systemic symptoms. Extra nodal manifestations have been reported in 28–43 % cases with rare ocular involvement. We report a case of a 57 year old female presenting with gradual progressive decrease of vision OU since 8 months associated with epistaxis. Fundus examination revealed established optic atrophy in right eye with features of chronic papilloedema in left eye suggestive of compressive lesion. CT of brain, paranasal sinuses confirmed the presence of homogenously enhancing mass in left ethmoid sinus, left sphenoid sinus extending into suprasellar region. The biopsy of this mass revealed extra nodal SHML with tissue sections being S100 and CD68 positive with emperipolesis noted. Here we describe this atypical ocular presentation of extra nodal SHML to highlight that this rare disease can manifest as an aggressive sight threatening entity, even in older age group.

Keywords Extra nodal Rosai Dorfman disease · Optic atrophy · S100 · CD68 · Sinus histiocytosis with massive lymphadenopathy

Introduction

Rosai–Dorfman disease (RDD) is a rare histiocytic disorder initially described as a separate entity in 1969 by Rosai and Dorfman under the term sinus histiocytosis with massive lymphadenopathy (SHML) [1]. Rosai–Dorfman disease is a benign systemic histioproliferative disease characterized clinically by massive bilateral cervical lymphadenopathy, fever, leukocytosis, raised erythrocyte sedimentation rate, and pathologically, by hyperplastic lymph node sinuses containing large histiocytosis with intact phagocytosed lymphocytes (emperipolesis) [2].

The causes of RDD are not fully understood, and treatment strategies can be different according to severity or vital organ involvement. Extra nodal involvement by RDD has been documented in 43 % of cases with the most frequent sites being skin, soft tissue, upper respiratory tract, multifocal bone, eye, and retro-orbital tissue with lymphadenopathy or as an isolated initial manifestation.

Extra nodal involvement has been reported in 28–43 % of the cases with the skin as the most commonly involved site. Ocular involvement is relatively rare (8.5 %), and mostly manifested as lymphoproliferation in the soft tissues of the orbit [3].

Case Report

A 57 year old female came referred to the ophthalmic OPD from ENT services in view of gradual progressive loss of vision since 8–9 months. The patient had an episode of epistaxis 2–3 months back. There was history of associated floaters and gradual diminution of vision in both eyes, painless and progressive since 8–9 months. Left eye cataract surgery was done 3 years back with poor post

✉ Eesha Shukla
eesha3388@gmail.com

¹ Department of Ophthalmology, T.N.M.C & B.Y.L Nair Charitable Hospital, Mumbai, India

operative vision. There was no perception of light in either of the eye. Right eye pupil was sluggishly reacting to light while the left eye pupil was circular, mid dilated, and not reacting to light. On slit lamp examination, the right eye showed immature cataract and the left eye was pseudophakic. A pendular nystagmus was noted on examination. Intra ocular pressure measured with applanation was 20 mm of hg in the right eye and 14 mm hg in the left eye. Gonioscopy showed all angles to be open in both eyes. Dilated 78D examination showed neurological cupping in both eyes (left more than right, with optic disc pallor being more than the cupping). Dilated ophthalmoscopy examination confirmed the diagnosis of bilateral optic atrophy (Fig. 1a, b).

Patient underwent further investigations and neuroimaging. CT scan of the brain, orbit and paranasal sinuses revealed an ill defined enhancing soft tissue mass; in left nasal cavity extending into the left maxillary sinus, ethmoidal, frontal and sphenoid sinus (3×4 cms) with bone destruction of wall of the sinuses. The same lesion showed continuity in suprasellar region (3.4×3.8 cms). Bilateral bony orbit and optic nerves were normal (Fig. 2a, b).

Nasal mass biopsy was advised by ENT services and the tissue sent for histopathology evaluation. Characteristic features consistent with the diagnosis of Extra Nodal Rosai Dorfman Disease were found on immunohistochemistry.

Fig. 1 **a** Right eye fundus photo. **b** Left eye Fundus Photo

Marked histiocytosis was present. Hematoxylin and eosin stain showed emperipolesis (lymphophagocytosis) (Fig. 3). Histiocytes were positive for S100 and CD68, which are histiocyte markers.

The salient features of this case were that the patient presented late for ophthalmic evaluation, owing to social reasons. Usually Rosai Dorfman Disease is self limiting, not requiring treatment, however in our case it caused sinus wall destruction with compressive optic atrophy. Size, extent, and location of tumor was such that it was in a precarious location.

Discussion

RDD is characterized by abnormal proliferation of histiocytes which are tissue macrophages belonging to the monocyte–macrophage lineage. Usually histiocytes are present in the lymph node sinuses i.e. the sub capsular region of a lymph node. Hence the term sinus histiocytosis, which usually presents with massive lymphadenopathy [4, 5].

The current working hypothesis on pathogenesis of RDD is recruitment of bone marrow monocytes from peripheral blood into lymph nodal sinuses or extranodal sites and their transformation into the immunophenotypically distinct RDD histiocytes which demonstrate emperipolesis. Release of cytokines like TNF alpha from these cells is responsible for the genesis of fever and other systemic symptoms.

Amongst extra nodal manifestations of RDD, eye and adnexal involvement is seen in 11 % of patients and CNS involvement, as in our case is seen in only 7 % (Table 1).

Ocular involvement is mostly seen in the form of orbital involvement, subconjunctival nodules, sub retinal exudates, or anterior uveitis [6].

Immunohistochemistry markers and H&E staining are diagnostic investigations. They demonstrate histiocytes of RDD which are S100-positive. CD68, a marker for phagocytic histiocytes is also positive in histiocytes of RDD as is Alpha 1 antichymotrypsin. Emperipolesis (lymphophagocytosis) literally means intact cell within the cytoplasm of another cell. It is a distinguishing feature of the disease.

Surgical excision, radiotherapy, and chemotherapy along with systemic steroids are the available treatment modalities.

The ideal treatment for Rosai–Dorfman disease has not been established, owing to the rarity of the disease. Only about 50 % of patients with Rosai–Dorfman disease need some form of treatment [7].

Radiotherapy and antimetabolite treatment have been considered in a few cases but the literature review does not

Fig. 2 **a** Axial and coronal CT Brain, **b** 3D CT reconstruction of lesion

suggest any conclusive role of these treatment modalities. No randomized control trials have been done as the disease in itself is rare. A large literature review by Pulsoni et al.

[5] concluded that clinical observation without treatment is advisable when possible. In the presence of vital organ compression and/or extra nodal localization with important

Fig. 3 Histopathology (H&E stain showing Emperipolesis) (white arrow)

Table 1 Frequency of distribution of clinical features of RDD

Anatomic location	Frequency (%)	Clinical manifestation
Lymph nodes	87	Massive painless lymphadenopathy, unilateral or bilateral, most commonly affecting cervicallymph nodes
Skin and soft tissue	16	Cutaneous lesions (maculopapular rash, xantomatous lesion, reddish or bluish mass), or subcutaneous nodules
Nasal cavity and paranasal sinuses	16	Mucosal thickening, polyp, infiltrative paranasal sinus mass
Eye and adnexa	11	Orbital or eyelid mass
Bone	11	Osteolytic lesions of axial or appendicular skeleton with or without sclerosis
Salivary gland	7	Bilateral parotid or submandibular gland mass
CNS	7	Intracranial, epidural, dural or spinal mass

clinical signs, surgical debulking may be necessary. Radiotherapy has shown limited efficacy, while chemotherapy is in general ineffective [8, 9].

In conclusion, Rosai–Dorfman disease can present, even in an elderly age group, with features suggestive of compressive optic atrophy. This case highlights an unusual manifestation of a rare disease, diagnosed on ophthalmic

evaluation. Usually considered benign, this disease can also manifest as an aggressive sight threatening entity. Liaising with other specialties is essential for a complete diagnosis and favorable patient outcomes.

Acknowledgments This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Authors' Contribution All authors have contributed equally to the varied aspects of this case report such as data collection, patient treatment, literature review and final editing of the manuscript.

Compliance with Ethical Standards

Conflict of interest None.

Ethical Standard This work presented above complies with the guidelines for human studies and animal welfare regulations. The subject has given their informed consent and as this was a case report, the institute's ethics committee has allowed the case report without a study protocol. No animal experimentation was done during the course of this work. IRB/Ethics Committee had ruled that approval was not required for the study.

References

1. Rosai J, Dorfman RF. Sinus histiocytosis with massive lymphadenopathy. A newly recognized benign clinicopathological entity. *Arch Pathol.* 1969;87:63–70.
2. Vemuganti GK, Naik MN, Honavar SG. Rosai dorfman disease of the orbit. *J Hematol Oncol.* 2008;1:7.
3. Kala C, Agarwal A, Kala S. Extranodal manifestation of Rosai–Dorfman disease with bilateral ocular involvement. *J Cytol.* 2011;28(3):131–3.
4. Huang YC, Tan HY, Jung SM, Chuang WY, Chuang CC, Hsu PW, et al. Spinal epidural Rosai–Dorfman disease preceding by relapsing uveitis: a case report with literature review. *Spinal Cord.* 2007;45(9):641–4.
5. Pulsoni A, Anghel G, Falcucci P, Matera R, Pescarmona E, Ribersani M, et al. Treatment of sinus histiocytosis with massive lymphadenopathy (Rosai–Dorfman disease): report of a case and literature review. *Am J Hematol.* 2002;69(1):67–71.
6. Payne JF, Srivastava SK, Wells JR, Grossniklaus HE. Rosai–Dorfman disease simulating nodular scleritis and panuveitis. *Arch Ophthalmol.* 2011;129(4):512–6.
7. Iyer VK, Handa KK, Sharma MC. Variable extent of emperipolesis in the evolution of Rosai Dorfman disease: diagnostic and pathogenetic implications. *J Cytol.* 2009;26(3):111–6.
8. Sarwal R, Tu E, Mendelblatt FI, Sugar J, Gross SA, Pulido JS, et al. Atypical ocular presentations of Rosai–Dorfman disease. *OculImmunolInflamm.* 2008;16(1):9–15.
9. Symss NP, Cugati G, Vasudevan MC, Ramamurthi R, Pande A. Intracranial Rosai Dorfman disease: report of three cases and literature review. *Asian J Neurosurg.* 2010;5(2):19–30.