Published in final edited form as:

Psychiatry Res. 2016 August 30; 242: 365–374. doi:10.1016/j.psychres.2016.04.115.

Examination of trait impulsivity on the response to a brief mindfulness intervention among college student drinkers

Christine Vinci^{a,*}, MacKenzie Peltier^b, Krystal Waldo^b, Jessica Kinsaul^c, Sonia Shah^b, Scott F. Coffey^d, and Amy L. Copeland^b

^aRice University, Psychology Department, 6100 Main St., Houston, TX 77005, United States

^bLouisiana State University, Department of Psychology, 236 Audubon Hall, Baton Rouge, LA 70803, United States

^cDurham Veteran Affairs Medical Center, Mental Health Service Line, Um 508 Fulton St, Durham, NC 27705, United States

^dUniversity of Mississippi Medical Center, Department of Psychiatry and Human Behavior, 2500 North State St., Jackson, MS 39216, United States

Abstract

Mindfulness-based strategies show promise for targeting the construct of impulsivity and associated variables among problematic alcohol users. This study examined the moderating role of intervention (mindfulness vs relaxation vs control) on trait impulsivity and three outcomes examined post-intervention (negative affect, positive affect, and urge to drink) among 207 college students with levels of at-risk drinking. Moderation analyses revealed that the relationship between baseline impulsivity and the primary outcomes significantly differed for participants who underwent the mindfulness versus relaxation interventions. Notably, simple slope analyses revealed that negative urgency was positively associated with urge to drink following the mindfulness intervention. Among participants who underwent the relaxation intervention, analysis of simple slopes revealed that negative urgency was negatively associated with urge to drink, while positive urgency was positively associated with positive affect following the relaxation intervention. Findings suggest that level (low vs high) and subscale of impulsivity matter with regard to how a participant will respond to a mindfulness versus relaxation intervention.

Keywords

Impulsivity; Mindfulness; Alcohol; Affect; College students

1. Introduction

Among college students, 39% report binge drinking (five or more alcoholic drinks on one occasion) and 12.7% indicate heavy drinking (five or more drinks on one occasion on five days or more over the past month; Substance Abuse and Mental Health Services

*Correspondence to: Rice University, Department of Psychology – MS 25, 6100 Main Street, Houston, TX 77005, United States., christine.vinci@rice.edu, cvinci28@gmail.com (C. Vinci).

Administration, 2014). College students engage in more problematic drinking behaviors than their non-college aged peers (Substance Abuse and Mental Health Services Administration, 2014; Wechsler et al., 2002), and several negative consequences have been associated with problematic drinking among college students such as impaired driving, interpersonal violence, academic impairment, and suicidal ideation and attempts (Perkins, 2002). Previous research has indicated that certain traits, including impulsivity, may impact problematic drinking among this population (Labrie et al., 2014).

Impulsivity is considered a multi-faceted construct and has been defined according to five subscales – (1) Negative Urgency (an individual's likelihood of acting impulsively when experiencing negative affect), (2) Lack of Premeditation (not thinking about the consequences of an action before engaging in that act), (3) Lack of Perseverance (lacking the tendency to focus on a boring or difficult task), (4) Sensation Seeking (engagement in activities that are exciting and that may or may not be dangerous), and (5) Positive Urgency (an individual's likelihood of acting impulsively when experiencing positive affect; Cyders and Smith 2007; Whiteside and Lynam, 2001). These facets of impulsivity have demonstrated differential relationships with alcohol-use outcomes. For instance, among college students, Sensation Seeking is related to increased frequency of alcohol use (Cyders et al., 2009; Labrie et al., 2014), whereas Positive and Negative Urgency have been linked to increased quantity of alcohol use (Cyders et al., 2009; Labrie et al., 2014) and negative consequences from drinking (Cyders et al., 2009, 2007). The relationship between urge to drink and trait impulsivity appears less clear, as some research has indicated that trait impulsivity does not impact alcohol craving (impulsivity measured via the Barratt Impulsiveness Scale [BIS]; Papachristou et al., 2012), whereas other research does indicate a relationship between increased impulsivity and craving (impulsivity also measured by the BIS; Papachristou et al., 2013). While differences in findings are likely due to the type of sample examined in these studies, the literature could benefit from additional research on the relationship between impulsivity and urge to drink.

Impulsivity is also related to certain mood states, including increased negative affect/depression in the general population (Corruble et al., 2003; Cyders and Coskunpinar, 2011) and among college students (Emmons and Diener, 1986; Langhinrichsen-Rohling et al., 2004). One study found that the relationship between depressive symptoms and alcohol use was moderated by impulsivity (specifically Negative Urgency and Sensation Seeking), such that the association between depressive symptoms and alcohol use was strengthened as impulsivity increased (Cyders and Coskunpinar, 2011). The association of positive affect and impulsivity, on the other hand, is more complicated. Some research has found that positive affect is a protective factor, and is linked to increased self-control among college students (Isen and Reeve, 2005; Ramezani and Gholtash, 2015). However, when looking at the impact of positive affect on specific impulsive behaviors related to alcohol use, positive affect has been associated with increased alcohol use and the experience of negative consequences related to drinking among college students (Del Boca et al., 2004). Furthermore, when considering the role of drinking motives and impulsivity, it has been shown that the facet of Positive Urgency (and not the other facets) was related to problematic drinking, but only among those individuals who drink to enhance their mood (as opposed to those who do not drink for this reason; Cyders et al., 2007). Thus, while Positive

Urgency may be a specific facet of impulsivity associated with problematic drinking, this relationship may be unique among individuals endorsing specific motives for drinking.

1.1. Mindfulness and substance use

Mindfulness has been defined as, “paying attention in a particular way: on purpose, in the present moment, and non-judgmentally” (Kabat-Zinn, 1994, p. 4). Increases in state mindfulness, defined as one’s ability to be mindful in a given moment (Lau et al., 2006), has been linked to increases in trait mindfulness, defined as one’s natural tendency to be mindful (Baer et al., 2006a; Kiken et al., 2015). While trait mindfulness captures an individual’s general tendency to be mindful (Baer et al., 2006a), mindfulness-based interventions often aim to change some behavior through increasing mindfulness (usually measured through changes in state and/or trait mindfulness). These interventions typically consist of about 8 group treatment sessions, with a specific topic for each session (e.g., awareness of triggers, how to be mindful in daily life and in high-risk situations, approaching thoughts from a decentered perspective). Additionally, participants are (1) guided through formal meditation each session, and (2) asked to practice meditation at home and incorporate the skills learned in each session into day-to-day life (Bowen et al., 2010).

Mindfulness-based interventions to treat problematic substance use behaviors have been well-supported, as results have indicated decreases in craving (Chen et al., 2010; Chiesa and Serretti, 2014; Garland et al., 2014), substance use behavior (Bowen et al., 2006; Bowen et al., 2014; Brewer et al., 2011; Chiesa and Serretti, 2014; Witkiewitz et al., 2014), and substance-related consequences (Bowen et al., 2006; Witkiewitz et al., 2014) post-treatment. While fewer studies have examined brief, laboratory-based mindfulness interventions, Bowen and Marlatt (2009) demonstrated that a brief, 11-min, intervention decreased smoking behavior in college student cigarette smokers.

Mindfulness practices are rooted in taking a nonjudgmental stance towards difficult experiences (e.g., unpleasant affect, negative thoughts; Kabat-Zinn, 1994) and have been found to decrease levels of negative affect (Arch and Craske, 2006; Jain et al., 2007; Ortner et al., 2007; Tang et al., 2007; Vinci et al., 2014) and increase positive affect (Davidson et al., 2003; Jain et al., 2007; Tang et al., 2007). Even a very brief, 15-min mindfulness-based breathing induction assisted individuals in managing negative affect when compared to two control groups (Arch and Craske, 2006). Performance on attentional tasks has improved following mindfulness interventions (e.g., Black et al., 2011; Ortner et al., 2007; Slagter et al., 2007; Tang et al., 2007). For example, Ortner et al. (2007) found that college students who received a 7-week course in mindfulness meditation demonstrated decreases in interference (on a cognitive task) when presented with affective images, as opposed to those in the relaxation group. The authors concluded that the ability to disengage from the emotional content allowed participants to perform better on the cognitive interference task, and thus potentially increase attention control.

1.2. Potential mechanisms of impulsivity and mindfulness

The management of attention and emotions are common mechanisms to both impulsivity and mindfulness (Ainslie and Haslam, 1992; Teper et al., 2013). Regarding impulsivity,

having the ability to attend to a boring or challenging task should result in increased perseverance (as captured by the Lack of Perseverance facet) and acting mindfully when experiencing certain mood states, as opposed to rashly, should result in the management of certain moods (as captured by the Negative and Positive Urgency facets; Cyders et al., 2007; Whiteside and Lynam, 2001). Together, these constructs suggest that managing emotion and attention are central aspects of impulsivity. Similarly, trait mindfulness is linked to both the regulation of emotion/affect and the control of attention that are hypothesized to be associated with impulsivity (Arch and Craske, 2006; Black et al., 2011; Davidson et al., 2003; Jain et al., 2007; Ortner et al., 2007; Slagter et al., 2007; Tang et al., 2007; Vinci et al., 2014). In fact, a recent theoretical model has proposed that mindfulness improves attention through two related processes – increased awareness of affect and acceptance of experiences (Teper et al., 2013).

1.3. Direct relationship between impulsivity and mindfulness

While previous work indicates that impulsivity and mindfulness may function through both emotion regulation and attentional control, research attempting to extrapolate the relationship between mindfulness and impulsivity has only recently been directly examined. Both the constructs of mindfulness and impulsivity share an emphasis on present moment focus. However, the decisions and subsequent consequences following engagement in mindful versus impulsive processes vary greatly (Murphy and MacKillop, 2011). Research has shown that mindfulness and impulsivity are generally inversely correlated; individuals reporting increased mindfulness are less impulsive and vice versa (Murphy and MacKillop, 2011; Peters et al., 2011).

Murphy and MacKillop (2011) examined the relationship between mindfulness (measured by the Five Factor Mindfulness Questionnaire [FFMQ]) and impulsivity (measured by the Urgency-Premeditation-Perseverance-Sensation Seeking-Positive [UPPS-P] Impulsivity Scale) in a sample of college student drinkers, in order to better understand the relationship between the multi-faceted constructs of both mindfulness and impulsivity in this sample. Their findings indicated that (1) Positive and Negative Urgency were negatively and strongly associated with Acting with Awareness, Nonreactivity, and Nonjudgment, but only weakly associated with Describe, (2) Lack of Premeditation was moderately and negatively associated with Describe and Nonreactivity, (3) Sensation Seeking was positively correlated with Observing, and (4) Lack of Perseverance had a strong and negative association with Describe and Acting with Awareness, but a weak association with Nonjudging and Nonreactivity. Similar results were found by Peters et al. (2011), who examined these same constructs among college students while also controlling for negative affect and general distress. The strongest associations (all negative) were found for Negative Urgency and both Acting with Awareness and Nonjudging, in addition to Lack of Perseverance and Acting with Awareness. Given this study controlled for negative affect and distress, these significant relationships suggests that affect does not completely account for the relationship between mindfulness and impulsivity, but instead demonstrates an important consideration for future research (Peters et al., 2011). It is possible that other factors may account for the mindfulness/impulsivity relationship. For example, “reactivity” is associated with certain aspects of impulsivity (e.g., positive and negative urgency; Cyders and Smith 2007;

Whiteside and Lynam, 2001), whereas “nonreactivity” is a facet of mindfulness (Baer et al., 2006b). Thus, for impulsive versus mindful individuals, reactivity level may be implicated in one’s attentional ability and emotional experience.

1.4. Impulsivity and mindfulness as related to substance use

Consistent with previous literature, Murphy and MacKillop (2011) also discovered that both alcohol use and problematic alcohol use were positively associated with facets of impulsivity. Furthermore, individuals endorsing problematic alcohol use were less likely to endorse certain facets of mindfulness (specifically Acting with Awareness, Nonreactivity, and Nonjudgment), indicating that they do not engage in these mindfulness practices. The authors postulate that while mindfulness and impulsivity are similar in some aspects regarding their relationships to alcohol use (i.e., negative associations generally exist with both regarding problematic alcohol use), they are still distinct constructs when examined individually (i.e., variation exists regarding which facets of mindfulness versus facets of impulsivity are associated alcohol use behaviors; Murphy and MacKillop, 2011).

1.5. Summary

Problematic alcohol use among college students is high and associated with several negative consequences. Both impulsivity and mindfulness have been associated not only with alcohol use and urge to drink, but also with the management of negative and positive affect. Additionally, two mechanisms common to impulsivity and mindfulness are the management of emotion and attention. Prior research examining the role of mindfulness on impulsivity and alcohol use through self-report measures has been promising, as mindfulness not only appears to be related to constructs relevant to both impulsivity and alcohol use as noted above, but these constructs may be particularly relevant to college student drinkers (given the association between impulsivity and problematic drinking in this population). Based on prior research, a logical next step would be to determine whether among at-risk college student drinkers, trait impulsivity would impact how an individual responds to a mindfulness intervention.

1.6. The current study

The present study aimed to determine whether receiving a particular type of brief intervention (mindfulness, relaxation, or control) would moderate the relationship between baseline level of trait impulsivity (each subscale of the UPPS-P was examined individually) and response to the intervention in a sample of at-risk college student drinkers. Specific outcome variables of interest included positive affect, negative affect, and urge to drink, which were measured pre- and post-intervention (see Fig. 1). We first hypothesized that the mindfulness group would significantly moderate the relationship between impulsivity and the outcome measures, such that those in the mindfulness group would have decreased negative affect and urge and increased positive affect, when compared to those in the relaxation and control groups. Second, given the lack of research in this area, we were unsure exactly how level of impulsivity (e.g., low vs high) would impact negative affect, positive affect, and urge to drink for those in the mindfulness group. As such, understanding the role of mindfulness interventions, when directly matched to other comparison

interventions, on impulsivity and associated variables among college students will provide important information on how to best intervene with this population.

2. Method

2.1. Participants

Data presented in the current paper were collected from a larger study ($n = 207$) examining the impact of a mood induction procedure following a brief mindfulness intervention on affect and urge to drink in at-risk college student drinkers (data from all 207 participants are used in the current analyses; see Vinci et al. (2014) for details of full study). The primary outcome study presented findings on the differential effects of each intervention type on the stated outcomes following a mood induction procedure, and did not examine how trait impulsivity interacted with intervention type.

Participants were screened for at-risk drinking via a secure, online system and then invited to attend the experimental portion of the study if eligible. Eligibility criteria included having a score of six or greater on the Alcohol Use Disorders Identification Test (AUDIT; a score of 6–8 indicates hazardous drinking in college students; Adewuya, 2005; Aertgeerts et al., 2000; Devos-Comby and Lange, 2008; Kokotailo et al., 2004; Menses-Gaya et al., 2009) and being elevated on at least one of two subscales of the Drinking Motives Questionnaire-Revised (DMQ-R; Cooper, 1994). Specifically, participants needed to be elevated on either the Coping or Enhancement motive (or both) on the DMQ-R to be eligible, in order to recruit those individuals who endorse drinking as a way to cope with emotions (a goal for the larger study). Thus, individuals were eligible if they met both the AUDIT and DMQ-R screening criteria outlined here. Eligible participants attended the experimental portion of the study one to two weeks following screening.

2.2. Self-report measures

2.2.1. Demographic questionnaire—Participants completed a demographic questionnaire developed by the experimenters assessing areas such as age, race, and gender.

2.2.2. Alcohol use disorders identification test—(AUDIT; Babor et al., 2001). The AUDIT is a 10-item self-report measure that assesses hazardous alcohol use on a 0–4 Likert scale. Research examining the AUDIT has found that among college students, a cut-off score of six and above best identifies hazardous drinking when examining sensitivity and specificity outcomes (Adewuya, 2005; Aertgeerts et al., 2000; Devos-Comby and Lange, 2008; Kokotailo et al., 2004; Menses-Gaya et al., 2009); thus, this cut-off was used in the present study to recruit at-risk drinkers. Internal consistency for the present study was adequate ($r = 0.64$)

2.2.3. Drinking motives questionnaire-revised—(DMQ-R; Cooper, 1994). The DMQ-R is 20-item self-report measure used to assess four drinking motives: Enhancement, Coping, Social Affiliative, and Social Conformity. The present study included only those participants primarily endorsing drinking for Enhancement or Coping motives, given these motives reflect those individuals who drink to manage mood. Specifically, out of the four

subscales, participants had to have the highest scores on either the Enhancement or Coping subscales. Internal consistency results for each subscale were as follows: Enhancement: ($r = 0.84$), Coping ($r = 0.83$), Social Affiliative ($r = 0.62$), and Social Conformity ($r = 0.84$).

2.2.4. The five-facet mindfulness questionnaire—(FFMQ; Baer et al., 2006b). The FFMQ is a 39-item measure assessing five facets of dispositional mindfulness on a 5-point Likert scale. This questionnaire was developed through factor analysis of previous mindfulness rating scales assessing mindfulness. Five factors emerged and constitute the FFMQ: Observing, Describing, Acting with Awareness, Nonjudging, and Non-reactivity. The current study resulted in the following internal consistency results: Observing ($r = 0.70$), Describing ($r = 0.91$), Acting with Awareness ($r = 0.87$), Nonjudging ($r = 0.87$), and Non-reactivity ($r = 0.71$). In the current study, this measure was used to assess participants' general degree of trait mindfulness pre-intervention (i.e., their general level of mindfulness in their daily lives).

2.2.5. Urgency-premeditation-perseverance-sensation seeking-positive impulsivity scale—(UPPS – P; Cyders and Smith 2007; Cyders et al., 2007; Whiteside and Lynam, 2001). The UPPS – P is a 59-item self-report measure of impulsivity, with each question being rated on a 4-point Likert scale. The UPPS – P was developed based on the Five Factor Model of personality, with questions being combined from several other measures of impulsivity. The five subscales of the UPPS-P include: Negative Urgency, (lack of) Premeditation, (lack of) Perseverance, Sensation Seeking, and Positive Urgency. Many items on the measure are reversed scored and elevations on any subscale indicate higher levels of impulsivity. The current study yielded internal reliability estimates for the following subscales: Negative Urgency ($r = 0.87$), Premeditation ($r = 0.85$), Perseverance ($r = 0.84$), Sensation Seeking ($r = 0.86$), and Positive Urgency ($r = 0.94$). The UPPS – P was used in the present study to examine participants' general degree of impulsivity pre-intervention.

2.2.6. Toronto mindfulness scale—(TMS; Lau et al., 2006). The TMS is a 13-item self-report measure of state mindfulness. Participants respond to each item using a 5-point Likert scale. The TMS has two subscales: Curiosity (approaching the present moment with a sense of curiosity) and Decentering (observing feelings and thoughts, while keeping distance from them). Previous research on this measure has indicated that the TMS has good internal consistency ($r = 0.95$) and post-treatment predictive validity regarding stress and psychological symptoms (Lau et al., 2006). In the present study, the Curiosity subscale demonstrated an internal consistency of 0.85 and the Decentering subscale 0.65. This measure was used in the present analyses to determine state mindfulness pre- and post-intervention. Thus, while the FFMQ assesses participants' general degree of mindfulness, the TMS allows for changes in mindfulness to be examined from moment-to-moment, and therefore represents a theoretically different measure of mindfulness. Specifically, the TMS asks participants about "what they just experienced," whereas the FFMQ asks participants about how mindful they are in their day-to-day lives (and not specifically about the previous moment).

2.2.7. Positive and negative affective schedule—(PANAS; Watson et al., 1988). The PANAS is a 20-item self-report measure that assesses an individual's negative and positive affect at a given point in time (for the current study, participants were asked to respond to questions according to how they feel "right now, at the present moment"). Twenty different emotions are listed, and individuals rate items on a 5-point Likert scale (possible range of responses: 10–50 for each subscale). The current study yielded the following internal consistency estimates – positive affect: $r = 0.84$; negative affect: $r = 0.70$. The present study utilized the PANAS to examine positive and negative affect at pre-and post-intervention.

2.2.8. Urge to drink—Participants' urge to drink was assessed via a 10-point Likert scale, stating "Please rate your urge to drink at this moment by circling a number on the scale below." Participants indicated their response from 1 (*absolutely no urge*) to 10 (*very strong urge*). Single-item measures have been found to be both reliable and valid in assessing urge to drink (Monti et al., 2000). Urge to drink was assessed at both pre- and post-intervention.

2.3. Brief interventions

2.3.1. Mindfulness intervention—Participants in the mindfulness intervention underwent a 10-min guided meditation (listened to on a cassette tape) instructing them to focus on the present moment, while noting their breath and any other physical sensations that may be occurring (e.g., touch, taste, sound). They were asked to do this with an attitude of acceptance and nonjudgment. A tape recording utilized by Adams et al. (2013) was used in the present study (for a written transcript of these instructions, please see Vinci et al. (2014)). Results from both Adams et al. (2013) and Vinci et al. (2014) demonstrated that participants who were guided through the mindfulness meditation significantly increased in state levels of mindfulness when compared to the groups that did not listen to the tape. This mindfulness meditation was primarily adapted of selections from Kabat-Zinn (1994, 2002); the five facets of mindfulness (Baer et al., 2006a) were also integrated throughout the recording.

2.3.2. Relaxation intervention—Participants in this group underwent a guided, 10-min relaxation intervention (also listened to on cassette tape), which was based on passive, progressive muscle relaxation (Feldman et al., 2010). This active comparison intervention was chosen in order to determine if the mindfulness intervention provided skills above and beyond just increased relaxation. See Vinci et al. (2014) for both more detail regarding this intervention and a written transcript of the instructions provided to participants.

2.3.3. Control intervention—Participants in the control group completed a word search puzzle for 10 min. This intervention was chosen to control for the passage of time and to have participants engaged in a focused task (as opposed to allowing their minds to wander).

2.4. Procedure

All study procedures were approved by the university's Institutional Review Board. College student drinkers were recruited for the study through the psychology department's experiment website and provided course credit for participating. Participants were also recruited via flyers on the campus; these individuals did not receive any incentive for

participating. Data for the screening phase was collected and stored through a secure online survey engine. Participants signed up for the study and indicated their consent to participate. They then completed the following questionnaires to determine eligibility: demographic form, the AUDIT, and the DMQ-R. Participants interested in the second phase of the study were asked to provide their email address in order to be contacted for scheduling if they were eligible (given most participants completed the study for course credit, some expressed disinterest due to already receiving all of their credit points). Eligible participants were then contacted and those interested in participating were scheduled within one to two weeks to complete the experimental portion of the study.

The second phase was conducted between 3:00 pm and 8:00 pm Monday through Friday. Sessions were conducted individually (and not in groups) for all participants. Participants completed the FFMQ and UPPS – P upon arrival. Participants were randomly assigned to groups and then completed baseline measures of the PANAS, TMS, and single-item urge question. They then underwent their respective intervention, followed by immediately completing the PANAS, TMS, and single-item urge question post-intervention. The approximate amount of time that lapsed from the completion of the baseline measures to the completion of the post-intervention measures was 15–20 min. The remainder of the study is not presented here, as it is not relevant to the primary hypotheses (Vinci et al., 2014).

3. Results

A total of 1831 participants completed the initial screening phase, and 394 were eligible for the experimental portion of the study. Though all 394 participants were invited to attend, 207 chose to do so (96.1% participated for course credit). Sixty-seven participants completed the mindfulness intervention, 74 were in the relaxation group, and 66 in the control group. The average age of the sample was 20.13 ($SD = 1.89$) and consisted of 76.3% women. Participants identified as 85.5% Caucasian, 6.3% African American, and 8.3% Other.

Participants had an average score of 10.03 ($SD = 4.28$) on the AUDIT (indicating that participants were engaging in levels of hazardous drinking; Babor et al., 2001). Intervention groups did not differ on age, AUDIT score, FFMQ subscales, UPPS – P subscales, or any of the outcome variables at baseline. Chi square analyses revealed no significant differences between the proportions of males and females in the intervention groups. Following the interventions, the following means and standard deviations were found for the outcome variables of interest: Negative Affect ($M = 12$, $SD = 3.08$), Positive Affect ($M = 24.48$, $SD = 7.56$), Urge ($M = 2.42$, $SD = 1.87$).

3.1. Initial bivariate correlations

Given our interest in further understanding the relationship between impulsivity and mindfulness among at-risk drinkers, correlational analyses for the entire sample on UPPS – P subscales, FFMQ subscales, TMS (at baseline) scores, and AUDIT scores were conducted (see Table 1). Associations were in the expected directions for several variables. Exceptions to this included: state mindfulness was positively associated with Sensation Seeking and Positive Urgency; Sensation Seeking was positively associated with the trait mindfulness

subscales of Observe and Nonreactivity; and state mindfulness was negatively associated with the trait mindfulness subscales of Acting with Awareness and Nonjudging.

3.2. Primary analyses

As this is a secondary data analysis, it is important to note that the primary outcome study did find that immediately following the interventions, participants in the mindfulness group decreased significantly in negative affect (when compared to the control group), participants in the relaxation group decreased in positive affect (when compared to the mindfulness and control groups), and that there were no significant changes in urge (Vinci et al., 2014). Additionally, the mindfulness group reported significant increases in state mindfulness from pre- to post-mindfulness intervention, and were also higher in state mindfulness than both the relaxation and control groups post-intervention (Vinci et al., 2014), indicating that the mindfulness intervention was effective at increasing state mindfulness.

To examine our hypothesis that intervention type (specifically the mindfulness group) would moderate the relationship between baseline trait impulsivity (Negative Urgency, Premeditation, Perseverance, Sensation Seeking, and Positive Urgency) and our outcome variables (Negative Affect, Positive Affect, and Urge), a series of hierarchical multiple regression analyses were conducted. Covariates were entered into Step 1 and included the baseline measurement of the outcome variable (e.g., baseline negative affect was entered when predicting negative affect post-intervention to control for the baseline value), gender (as previous research has indicated that college student males drink more alcohol than college student females; Berkowitz and Perkins, 1986; Nolen-Hoeksema, 2004; O'Malley and Johnston, 2002), and total AUDIT score (to control for level of problematic drinking). Step 2 included the main effects of group and impulsivity subscale score. Step 3 included the product of group and impulsivity subscale score. For this interaction term, the mindfulness group was coded as the reference group.

Results from the hierarchical multiple regression analyses indicated that intervention type significantly moderated the relationship in four models (see Table 2 for results these models). First, the relaxation group (when compared to the mindfulness group) moderated the relationship between Sensation Seeking and Negative Affect [$F(8,198) = 10.15, p = 0.001$]. Specifically, for every one point increase on Sensation Seeking, participants in the relaxation group decreased 1.54 points in Negative Affect, when compared to those in the mindfulness group. To probe the interaction, examination of the simple slopes was conducted and revealed that neither slope was significant from zero (mindfulness: $b = 0.36, p = 0.52$; relaxation: $b = -1.18, p = 0.34$). Thus, while the effect of Sensation Seeking was significantly different for participants in the mindfulness versus relaxation groups regarding their ratings of Negative Affect, the degree to which Sensation Seeking matters within each group was not significant.

Second, the relaxation group (when compared to the mindfulness group) moderated the relationship between Negative Urgency and Urge [$F(8,195) = 93.48, p = 0.001$]. Specifically, for every one point increase on Negative Urgency, participants in the relaxation group decreased 0.82 points in Urge, when compared to the mindfulness group. Examination of the simple slopes revealed a significant association between Negative Urgency and Urge for

participants in both the mindfulness and relaxation groups. For participants in the mindfulness group, having low trait Negative Urgency was associated with low reports of Urge, whereas participants with high trait Negative Urgency reported high Urge ($b = 0.4$, $t = 2$, $p = 0.046$). The opposite pattern was observed for participants in the relaxation group, such that for those with low trait Negative Urgency, Urge was high; for those with high trait Negative Urgency, Urge was low ($b = -0.42$, $t = -2.1$, $p = 0.04$). See Fig. 2.

Third, the relaxation group (when compared to the mindfulness group) moderated the relationship between Positive Urgency and Positive Affect [$F(8,198) = 24.26$, $p = 0.001$]. Specifically, for every one point increase on Positive Urgency, participants in the relaxation group increased 3.74 points in Positive Affect, when compared to the mindfulness group. Examination of the simple slopes revealed a significant effect for the relaxation group, such that for participants low in trait Positive Urgency, Positive Affect was low; for participants with high trait Positive Urgency, Positive Affect was high ($b = 2.95$, $t = 2.68$, $p = 0.01$). See Fig. 3.

Fourth, the control group (when compared to the mindfulness group) moderated the relationship between (lack of) Perseverance and Positive Affect [$F(8,198) = 23.29$, $p = 0.001$]. Specifically, for every one point increase on (lack of) Perseverance, participants in the control group decreased 3.78 points in Positive Affect, when compared to the mindfulness group. Examination of the simple slopes revealed no significant findings (mindfulness: $b = 1.67$, $p = 0.20$; relaxation: -2.11 , $p = 0.13$).

4. Discussion

The current study examined the moderating role of a brief mindfulness intervention (vs relaxation and control) on the relationship between trait impulsivity and negative affect, positive affect, and urge to drink. First, and somewhat unexpectedly, three of the four significant moderation analyses revealed that for participants who underwent the mindfulness intervention (compared to the relaxation group), higher levels of certain facets of impulsivity were associated with increased negative affect and urge, and decreased positive affect post-mindfulness intervention. Second, results of simple slope analyses revealed that the level of impulsivity interacted with intervention type, such that changes in affect and urge differed depending on whether participants were low versus high on certain subscales of impulsivity.

To expand upon the significant interactions, we found that the relationship between certain subscales of impulsivity (Sensation Seeking, Negative Urgency, and Positive Urgency) and outcome (Negative Affect, Urge, and Positive Affect, respectively) was moderated by type of intervention; in particular, higher levels of these facets of trait impulsivity were associated with higher levels of negative affect and urge, and lower levels of positive affect among those receiving mindfulness intervention compared to those receiving relaxation intervention. These findings indicate that for individuals elevated on Sensation Seeking, Negative Urgency, and Positive Urgency, a brief mindfulness intervention may increase negative affect and urge to drink and decrease positive affect. However, the fourth significant interaction revealed that for those who underwent the mindfulness intervention (when

compared to the control group) decreases in the impulsivity facet of (lack of) Perseverance was associated with increased Positive Affect post-mindfulness intervention. Thus, for individuals with elevated levels of Perseverance, a brief mindfulness intervention may increase positive affect.

Second, analysis of the simple slopes revealed that for participants with low levels of Negative Urgency, undergoing a mindfulness intervention resulted in a low urge to drink. However, for participants with high levels of Negative Urgency the opposite effect was found, such that the mindfulness intervention was associated with high urge to drink. These findings suggest that how low/high an individual is on Negative Urgency will impact their urge to drink following a brief mindfulness intervention, with those low in Negative Urgency responding more favorably to the intervention. For participants with low levels of Negative and Positive Urgency, the relaxation intervention was associated with high levels of urge to drink and low levels of positive affect, respectively. On the other hand, those with high Negative and Positive Urgency reported low levels of urge and high levels of positive affect, respectively. These results suggest that for those with elevated Negative and Positive Urgency, a brief relaxation intervention will impact urge to drink and positive affect, with those elevated on these facets responding more favorably to this intervention. Overall, it appears that the level (low versus high) and specific type of impulsivity subscale matters with regard to how a participant will respond to a mindfulness versus relaxation intervention.

While previous research has indicated an inverse relationship between the various subscales of impulsivity and trait mindfulness (Murphy and MacKillop, 2011; Peters et al., 2011), this is the first study to examine the relationship between the subscales of impulsivity and response to a mindfulness intervention among problematic alcohol drinkers. Results from the current study were intriguing, in that the majority of the significant interactions and simple slope analyses revealed that for participants with higher trait impulsivity who underwent the mindfulness intervention, these individuals had increased negative affect and urge to drink and decreased positive affect. However, these findings could make sense in light of the strength model of self-regulation (Baumeister et al., 2007; Hagger et al., 2009; Muraven and Baumeister, 2000). This model posits that individuals have a limited capacity to regulate certain states (e.g., affect, hunger), and when the ability to self-regulate has been depleted, individuals may have difficulty continuing to self-regulate on other tasks. It is possible that undergoing a brief mindfulness intervention might be more depleting than experiencing the relaxation intervention. For instance, upon examining the moderating role of group (brief mindfulness intervention versus control group) on pain tolerance, Evans and colleagues found that for participants in the control group (and not in the mindfulness group), higher heart rate variability, a measure of self-regulatory capacity, was associated with increased pain tolerance (Evans et al., 2014). The authors posited that such a brief mindfulness intervention may have increased effort and depleted self-regulatory capacity. While the current study did not examine self-regulatory capacity, it is possible that for individuals with high levels of impulsivity, the mindfulness intervention reduced self-regulatory capacity (whereas this perhaps did not happen for those in the relaxation group). If self-regulatory capacity was depleted, the mindfulness intervention would be less effective on decreasing negative affect and urge and increasing positive affect than the relaxation intervention. Possible reasons that such a mindfulness intervention could deplete self-

regulatory capacity would be that participants only had a brief period of time to learn and practice mindfulness, and that once practiced, had to apply these newly learned skills to a negative affect situation. Future research could examine this hypothesis by including a measure of self-regulatory capacity when examining the role of mindfulness (both brief mindfulness and lengthier interventions) on impulsivity.

Given the lack of research on impulsivity and mindfulness among at-risk drinkers, a brief discussion of the bivariate correlations is warranted. Most of the associations were in the expected directions and consistent with previous findings (Murphy and MacKillop, 2011; Peters et al., 2011). However, positive associations were found between the impulsivity subscales of Sensation Seeking and Negative Urgency and state mindfulness. Prior research on the relationship between mindfulness and various measures of impulsivity, to our knowledge, have only utilized measures of trait mindfulness (Baer et al., 2006a; Lakey et al., 2009; Lattimore et al., 2011; Murphy and MacKillop, 2011; Peters et al., 2011; Williams and Grishman, 2012; Wupperman et al., 2009), thus making the current study's findings on the association between the UPPS – P subscales and TMS important. When considering that the TMS measures state levels of mindfulness and the FFMQ examines dispositional mindfulness, it might be the case that the two questionnaires vary considerably when related to impulsivity. Our results also indicated that associations between the FFMQ and TMS subscales varied, as some were positive and others negative. This finding is less surprising, given the theoretical differences between state and trait mindfulness (i.e., state mindfulness measures momentary shifts in mindfulness, whereas trait mindfulness captures general levels of mindfulness in daily living; Baer et al., 2006b; Lau et al., 2006), as well as evidence that different mindfulness measures are often assessing different constructs (Grossman, 2011; Grossman and Van Dam, 2011).

Clinical implications of these findings suggest that practitioners providing mindfulness and relaxation interventions to at-risk college student drinkers may want to consider their patients' level of impulsivity and subsequent response to these interventions. Monitoring patient response and assessing level of impulsivity at the outset of therapy would be useful, as individuals with elevations on certain aspects of impulsivity may be more/less likely to experience certain affective states and urges to drink following these two interventions. For example, for individuals reporting low levels of negative urgency, a mindfulness intervention appears to have a positive effect on urge. However, the opposite is true for individuals with high scores on the negative urgency subscale, suggesting that a relaxation intervention may be more appropriate. It is possible that with the increased practice of mindfulness skills, individuals with high negative urgency may benefit in a similar way to those with low negative urgency when undergoing a mindfulness intervention. However, additional research examining these constructs is needed before such recommendations can be made.

Regarding future research in this area, replication of these findings is a necessary first step. Little research has been conducted on how impulsivity is related to mindfulness among at-risk college student drinkers, and to our knowledge, no research has examined the role of a brief mindfulness intervention on these constructs, aside from the current study. Several moderation analyses were not significant in the current study (e.g., intervention type did not moderate the relationship between Lack of Premeditation and any of the outcome variables).

Replication is needed to determine whether the non-significant relationships found in this study extend to other samples before meaningful conclusions can be drawn from the lack of significance found here. Research on more extensive interventions (e.g., longer than 10 min, multiple sessions) is also suggested, as it is possible that with increased practice, individuals with high impulsivity may respond differently to the mindfulness intervention over time. Consideration of how impulsivity may relate to outcomes following a mindfulness intervention with other populations could be beneficial (e.g., young adults engaging in at-risk drinking who are not in college, individuals seeking treatment for alcohol use problems through an outpatient clinic). For instance, prior work has shown that when only the mindfulness portion of DBT was provided to those with borderline personality disorder, level of impulsivity decreased (Soler et al., 2012). Other mental health problems that involve impulsivity as a critical mechanism of the disorder may want to explore the differential impact of mindfulness-based versus relaxation-based interventions, as well as variations in length of intervention on outcomes; such populations may include those with eating disorders (Rosval et al., 2006), anger management problems (Horesh et al., 1997), and suicidality (Horesh et al., 1997).

Limitations of the present study should be noted. First, the sample consisted of at-risk college student drinkers (consisting primarily of women and Caucasians), thus we do not know how these results would generalize to other populations. However, the average AUDIT score of 10.03 indicates that the sample consisted of at-risk alcohol drinkers. Second, the correlations presented in Table 1 should be interpreted with caution, as while they indicate many significant relationships, the strength of these relationships is relatively weak. Third, the present study only examined the immediate change in affect and urge; we do not know whether these changes would maintain over time or if they would vary over time. Fourth, while we utilized a self-report measure to gather level of impulsivity, future studies should incorporate behaviorally-based measures of impulsivity. And fifth, some have argued that many questionnaires assessing mindfulness likely do not accurately capture an individual's level of mindfulness due to inherent difficulties in assessing one's own level of mindfulness, problems with defining mindfulness through questionnaires, and the simplistic nature of questionnaire items (Grossman and Van Dam, 2011; Purser and Milillo, 2014). Additionally, self-report measures often include items that are very similar to the mindfulness instructions, which may not truly represent whether an individual has become more mindful (Grossman and Van Dam, 2011). Future research may want to consider additional assessment methods aside from self-report measures of mindfulness (e.g., interviews).

In conclusion, the current study presents interesting findings on the role of a brief mindfulness intervention on the relationship between aspects of trait impulsivity and negative affect, positive affect, and urge to drink in a sample of at-risk college student drinkers. These results are applicable to both future research and in clinical settings where brief mindfulness and relaxation exercises are conducted. Nonetheless, future work examining the role of mindfulness on impulsivity, affect, and urge is needed, in order to replicate these findings. Ultimately, determining the effects of lengthier interventions, the impact of such interventions over time, and ways in which to best examine changes in mindfulness and relevant constructs are suggested. While this study warrants such research efforts in problematic drinkers, examining the effects of mindfulness-based interventions in

other populations with increased impulsivity, as well as impulsive, non-problematic drinkers are recommended. Lastly, practical considerations of utilizing brief mindfulness and relaxation exercises within a therapeutic context for at-risk drinkers should be considered in light of the findings.

Acknowledgments

Funding statement

This work was supported the National Institute on Minority Health and Health Disparities (K99MD010468). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

- Adams CE, Benitez L, Kinsaul J, McVay M, Barbry A, Thibodeaux A, Copeland AL. Effects of brief mindfulness instructions on reactions to body image stimuli among female smokers: an experimental study. *Nicotine Tob Res.* 2013; 15(2):376–384. <http://dx.doi.org/10.1093/ntr/nts133>. [PubMed: 22987786]
- Adeyuya AO. Validation of the alcohol use disorders identification test (AUDIT) as a screening tool for alcohol-related problems among Nigerian university students. *Alcohol Alcohol.* 2005; 40:575–577. <http://dx.doi.org/10.1093/alcalc/agh197>. [PubMed: 16115823]
- Aertgeerts B, Buntinx F, Bande-Knops J, Vandermeulen C, Roelants M, Ansoms S, Fevery J. The value of CAGE, CUGE, and AUDIT in screening for alcohol abuse and dependence among college freshman. *Alcohol: Clin Exp Res.* 2000; 24:53–57. [PubMed: 10656193]
- Ainslie, G.; Haslam, N. Self-Control. In: Loewenstein, G.; Elster, J., editors. *Choice Over Time*. Russell Sage Foundation; New York, NY: 1992. p. 177-209.
- Arch JJ, Craske MG. Mechanisms of mindfulness: emotion regulation following a focused breathing induction. *Behav Res Ther.* 2006; 44(12):1849–1858. <http://dx.doi.org/10.1016/j.brat.2005.12.007>. [PubMed: 16460668]
- Babor, TF.; Higgins-Biddle, JC.; Saunders, JB.; Monteiro, MG. AUDIT: The Alcohol Use Disorders Identification Test: Guidelines for Use in Primary Care. World Health Organization; Geneva, Switzerland: 2001.
- Baer RA, Fischer S, Huss DB. Mindfulness and acceptance in the treatment of disordered eating. *J Ration Emot Cogn Behav Ther.* 2006b; 23:281–300. <http://dx.doi.org/10.1007/s10942-005-0015-9>.
- Baer RA, Smith GT, Hopkins J, Krietemeyer J, Toney L. Using self-report assessment methods to explore facets of mindfulness. *Assessment.* 2006a; 13:27–45. <http://dx.doi.org/10.1177/1073191105283504>. [PubMed: 16443717]
- Baumeister RF, Vohs KD, Tice DM. The strength model of self-control. *Curr Dir Psychol Sci.* 2007; 16:351–355.
- Berkowitz AD, Perkins HW. Problem drinking among college students: a review of recent research. *J Am Coll Health.* 1986; 35:21–28. [PubMed: 3771936]
- Black DS, Semple RJ, Pokhrel P, Grenard JL. Component processes of executive function – mindfulness, self-control, and working memory – and their relationships with mental and behavioral health. *Mindfulness.* 2011; 2:179–185. <http://dx.doi.org/10.1007/s12671-011-0057-2>. [PubMed: 22013495]
- Bowen, S.; Chawla, N.; Marlatt, GA. Mindfulness-Based Relapse Prevention for Substance Use Disorders: A Clinician's Guide. Guilford Press; New York, NY: 2010.
- Bowen S, Marlatt A. Surfing the urge: brief mindfulness-based intervention for college student smokers. *Psychol Addict Behav.* 2009; 23:666–671. <http://dx.doi.org/10.1037/a0017127>. [PubMed: 20025372]
- Bowen S, Witkiewitz K, Clifasefi S, Grow J, Chawla N, Hsu S, Larimer ME. Relative long-term efficacy of mindfulness-based relapse prevention, standard relapse prevention and treatment as usual for substance use disorders. *JAMA Psychiatry.* 2014; 71:547–565. [PubMed: 24647726]

- Bowen S, Wikiewitz K, Dillworth TM, Chawla N, Simpson TL, Ostafin BD, Marlatt GA. Mindfulness meditation and substance use in an incarcerated population. *Psychol Addict Behav*. 2006; 20:343–347. <http://dx.doi.org/10.1037//0893>. [PubMed: 16938074]
- Brewer JA, Mallik S, Babuscio TA, Nich C, Johnson HE, Deleone CM, Rounsaville BJ. Mindfulness training for smoking cessation: Results from a randomized controlled trial. *Drug Alcohol Depend*. 2011; 119:72–80. <http://dx.doi.org/10.1016/j.drugalcdep.2011.05.027>. [PubMed: 21723049]
- Chen KW, Comerford A, Shinnick P, Ziedonis DM. Introducing Qigong. Meditation into residential addiction treatment: a pilot study where gender makes a difference. *J Altern Complement Med*. 2010; 16:875–882. <http://dx.doi.org/10.1089/acm.2009.0443>.
- Chiesa A, Serretti A. Are mindfulness-based interventions effective for substance use disorders? A systematic review of the evidence. *Subst Use Misuse*. 2014; 49:492–521. <http://dx.doi.org/10.3109/10826084.2013.770027>. [PubMed: 23461667]
- Cooper ML. Motivations for alcohol use among adolescents: development and validation of a four-factor model. *Psychol Assess*. 1994; 6:117–128. 1040-3590/94.
- Corruble E, Benyamina A, Bayle F, Falissard B, Hardy P. Understanding impulsivity in severe depression? A psychometrical contribution. *Prog Neuro-Psychopharmacol Biol Psychiatry*. 2003; 27:829–833. [http://dx.doi.org/10.1016/S0278-5846\(03\)00115-5](http://dx.doi.org/10.1016/S0278-5846(03)00115-5).
- Cyders MA, Coskunpinar A. Depression, impulsivity, and health-related disability: a moderated mediation analysis. *J Res Personal*. 2011; 45:679–682. <http://dx.doi.org/10.1016/j.jrp.2011.08.005>.
- Cyders MA, Flory K, Rainer S, Smith GT. The role of personality dispositions to risky behavior in predicting first year college drinking. *Addiction*. 2009; 104:193–202. <http://dx.doi.org/10.1111/j.1360-0443.2008.02434.x>. [PubMed: 19149813]
- Cyders MA, Smith GT. Mood-based rash action and its components: positive and negative urgency. *Personal Individ Differ*. 2007; 43(4):839–850. <http://dx.doi.org/10.1016/j.paid.2007.02.008>.
- Cyders MA, Smith GT, Spillane NS, Fischer S, Annus AM, Peterson C. Integration of impulsivity and positive mood to predict risky behavior: development and validation of a measure of positive urgency. *Psychol Assess*. 2007; 19:107–118. <http://dx.doi.org/10.1037/1040-3590.19.1.107>. [PubMed: 17371126]
- Davidson RJ, Kabat-Zinn J, Schumacher J, Rosenkranz M, Muller D, Santorelli SF, Sheridan JF. Alterations in brain and immune function produced by mindfulness meditation. *Psychosom Med*. 2003; 65(4):564–570. <http://dx.doi.org/10.1097/01.PSY.0000077505.67574.E3>. [PubMed: 12883106]
- Del Boca FK, Darkes J, Greenbaum PE, Goldman MS. Up close and personal: temporal variability in the drinking of individual students during their first year. *J Consult Clin Psychol*. 2004; 72:155–164. <http://dx.doi.org/10.1037/0022-006X.72.2.155>. [PubMed: 15065951]
- Devos-Comby L, Lange JE. Standardized measures of alcohol-related problems: a review of their use among college students. *Psychol Addict Behav*. 2008; 22:349–361. <http://dx.doi.org/10.1037/0893-164X.22.3.349>. [PubMed: 18778128]
- Emmons RA, Diener E. Influence of impulsivity and sociability on subjective well-being. *J Personal Soc Psychol*. 1986; 50:1211–1215.
- Evans DR, Eisenlohr-Moul TA, Button DF, Baer RA, Segerstrom SC. Self-regulatory deficits associated with unpracticed mindfulness strategies for coping with acute pain. *J Appl Soc Psychol*. 2014; 44:23–30. <http://dx.doi.org/10.1111/jasp.12196>. [PubMed: 25843972]
- Feldman G, Greeson J, Senville J. Differential effects of mindful breathing, progressive muscle relaxation, and loving-kindness meditation on decentering and negative reaction to repetitive thoughts. *Behav Res Ther*. 2010; 48:1002–1011. <http://dx.doi.org/10.1016/j.brat.2010.06.006>. [PubMed: 20633873]
- Garland EL, Manusov EG, Froeliger B, Kelly A, Williams JM, Howard MO. Mindfulness-oriented recovery enhancement for chronic pain and prescription opioid misuse: results from an early-stage randomized controlled trial. *J Consult Clin Psychol*. 2014; 82:448–459. <http://dx.doi.org/10.1037/a0035798>. [PubMed: 24491075]
- Grossman P. Comment: defining mindfulness by how poorly I think I pay attention during everyday awareness and other intractable problems for psychology's (re)invention of mindfulness: comment

- on Brown et al. Psychol Assess. 2011; 23:1034–1040. <http://dx.doi.org/10.1037/a0022713>. [PubMed: 22122674]
- Grossman P, Van Dam NT. Mindfulness, by any other name....: Trials and tribulations of sati in western psychology and science. Contemp Buddh: Interdiscip J. 2011; 12:219–239. <http://dx.doi.org/10.1080/14639947.2011.564841>.
- Hagger MS, Wood C, Stiff C, Chatzisarantis NLD. The strength model of self-regulation failure and health-related behavior. Health Psychol Rev. 2009; 3:208–238. <http://dx.doi.org/10.1080/17437190903414387>.
- Horesh N, Rolnick T, Iancu I, Dannon P, Lepkifker E, Apter A, Kotler M. Anger, impulsivity, and suicide risk. Psychother Psychosom. 1997; 66:92–96. <http://dx.doi.org/10.1159/000289115>. [PubMed: 9097337]
- Isen AM, Reeve J. The influence of positive affect on intrinsic and extrinsic motivation: facilitating enjoyment of play, responsible work behavior, and self-control. Motiv Emot. 2005; 29:297–325. <http://dx.doi.org/10.1007/s11031-006-9019-8>.
- Jain S, Shapiro SL, Swanick S, Roesch SC, Mills PJ, Bell I, Schwartz GR. A randomized controlled trial of mindfulness meditation versus relaxation training: Effects on distress, positive states of mind, rumination, and distraction. Ann Behav Med. 2007; 33(1):11–21. http://dx.doi.org/10.1207/s15324796abm3301_2. [PubMed: 17291166]
- Kabat-Zinn, J. Wherever You Go, There You Are: Mindfulness Meditation in Everyday Life. Hyperion; New York, NY: 1994.
- Kabat-Zinn, J. On Guided Mindfulness Meditation [compact disc]. Lexington, MA: Stress Reduction CDs and Tapes; 2002. Sitting Meditations.
- Kiken LG, Garland EL, Bluth K, Palsson OS, Gaylord SA. From a state to a trait: trajectories of state mindfulness in meditation during intervention predict changes in trait mindfulness. Personal Individ Differ. 2015; 81:41–46. <http://dx.doi.org/10.1016/j.paid.2014.12.044>.
- Kokotailo PK, Egan J, Gangnon R, Brown D, Mundt M, Fleming M. Validity of the alcohol use disorders identification test in college students. Alcohol: Clin Exp Res. 2004; 28(6):914–920. <http://dx.doi.org/10.1097/01.ALC.0000128239.87611.F5>. [PubMed: 15201634]
- Labrie JW, Kenney SR, Napper LE, Miller K. Impulsivity and alcohol-related risk among college students: Examining urgency, sensation seeking and the moderating influence of beliefs about alcohol's role in the college experience. Addict Behav. 2014; 39:1–15. <http://dx.doi.org/10.1016/j.addbeh.2013.09.018>. [PubMed: 24119712]
- Lakey CE, Campbell WK, Brown KW, Goodie AS. Dispositional mindfulness as a predictor of the severity of gambling outcomes. Personal Individ Differ. 2009; 43:1698–1710. <http://dx.doi.org/10.1016/j.paid.2007.05.007>.
- Langhinrichsen-Rohling J, Arata C, Bowers D, O'Brien N, Morgan A. Suicidal behavior, negative affect, gender, and self-reported delinquency in college students. Suicide Life Threat Behav. 2004; 34:255–266. [PubMed: 15385180]
- Lattimore P, Fisher N, Malinowski P. A cross-sectional investigation of trait disinhibition and its association with mindfulness and impulsivity. Appetite. 2011; 56:241–248. <http://dx.doi.org/10.1016/j.appet.2010.12.007>. [PubMed: 21146571]
- Lau MA, Bishop SR, Segal ZV, Buis T, Anderson ND, Carlson L, Devins G. The Toronto mindfulness scale: development and validation. J Clin Psychol. 2006; 62(12):1445–1467. <http://dx.doi.org/10.1002/jclp.20326>. [PubMed: 17019673]
- Menses-Gaya C, Zuardi AW, Loureiro SR, Crippa JAS. Alcohol use disorders identification test (AUDIT): An updated systematic review of psychometric properties. Psychol Neurosci. 2009; 2:83–97. <http://dx.doi.org/10.3922/j.psns.2009.1.12>.
- Monti PM, Rohsenow DJ, Hutchison KE. Toward bridging the gap between biological, psychobiological, and psychosocial models of alcohol craving. Addiction. 2000; 95:229–236. [PubMed: 10723851]
- Muraven M, Baumeister RF. Self-regulation and depletion of limited resources: does self-control resemble a muscle? Psychol Bull. 2000; 126:247–259. <http://dx.doi.org/10.1037/0033-2909.126.2.247>. [PubMed: 10748642]

- Murphy C, MacKillop J. Living in the here and now: Interrelationships between impulsivity, mindfulness, and alcohol misuse. *Psychopharmacology*. 2011; 219:527–536. <http://dx.doi.org/10.1007/s00213-011-2573-0>. [PubMed: 22169883]
- Nolen-Hoeksema S. Gender differences in risk factors and consequences for alcohol use and problems. *Clin Psychol Rev*. 2004; 24:981–1010. <http://dx.doi.org/10.1016/j.cpr.2004.08.003>. [PubMed: 15533281]
- O'Malley PM, Johnston LD. Epidemiology of alcohol and other drug use among American college students. *J Stud Alcohol*. 2002; 14:23–29.
- Ortner CM, Kilner SJ, Zelazo P. Mindfulness meditation and reduced emotional interference on a cognitive task. *Motiv Emot*. 2007; 31(4):271–283. <http://dx.doi.org/10.1007/s11031-007-9076-7>.
- Papachristou H, Nederkoorn C, Havermans R, Bongers P, Beunen S, Jansen A. Higher levels of trait impulsiveness and a less effective response inhibition are linked to more intense cue-elicited craving for alcohol in alcohol-dependent patients. *Psychopharmacology*. 2013; 228:641–649. <http://dx.doi.org/10.1007/s00213-013-3063-3>. [PubMed: 23508556]
- Papachristou H, Nederkoorn C, Havermans R, van der Horst M, Jansen A. Can't stop the craving: the effect of impulsivity on cue-elicited craving for alcohol in heavy and light social drinkers. *Psychopharmacology*. 2012; 219:511–518. <http://dx.doi.org/10.1007/s00213-011-2240-5>. [PubMed: 21384105]
- Perkins HW. Surveying the damage: a review of research on consequences of alcohol misuse in college populations. *J Stud Alcohol*. 2002; 14:91–139.
- Peters JR, Erisman SM, Upton BT, Baer RA, Roemer L. A preliminary investigation of the relationships between dispositional mindfulness and impulsivity. *Mindfulness*. 2011; 2:228–235. <http://dx.doi.org/10.1007/s12671-011-0065-2>.
- Purser, RE.; Milillo, J. Mindfulness revisited: a Buddhist-based conceptualization. *J Manag Inq*. 2014. 2014. <http://dx.doi.org/10.1177/1056492614532315> (e-Pub)
- Ramezani SG, Gholtash A. The relationship between happiness, self-control, and locus of control. *Int J Educ Psychol Res*. 2015; 1:100–104.
- Rosval L, Steiger H, Bruce K, Israel M, Richardson J, Aubut M. Impulsivity in women with eating disorders: Problems of response inhibition, planning, or attention? *Int J Eat Disord*. 2006; 39:590–593. <http://dx.doi.org/10.1002/eat>. [PubMed: 16826575]
- Slagter HA, Lutz A, Greischar LL, Francis AD, Nieuwenhuis S, Davis JM, Davidson RJ. Mental training affects distribution of limited brain resources. *Public Library Sci Biol*. 2007; 5(6) <http://dx.doi.org/10.1371/journal.pbio.0050138>.
- Soler J, Valdepérez A, Feliu-Soler A, Pascual JC, Portella MJ, Martín-Blanco A, Pérez V. Effects of the dialectical behavioral therapy-mindfulness module on attention in patients with borderline personality disorder. *Behav Res Therapy*. 2012; 50:150–157. <http://dx.doi.org/10.1016/j.brat.2011.12.002>.
- Substance Abuse and Mental Health Services Administration. Results from the 2013 National Survey on Drug Use and Health: Summary of National Findings. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2014. NSDUH Series H-48, HHS Publication no. (SMA) 14-4863. NSDUH Series H-41, HHS Publication No. (SMA) 11-4658
- Tang Y, Ma Y, Wang J, Fan Y, Feng S, Lu Q, Posner M. Short-term meditation training improves attention and self-regulation. *Proc Natl Acad Sci*. 2007; 104:17152–17156. DOI: 10.1073/pnas.0707678104 [PubMed: 17940025]
- Teper R, Segal ZV, Inzlicht M. Inside the mindful mind: how mindfulness enhances emotion regulation through improvements in executive control. *Curr Dir Psychol Sci*. 2013; 22:449–454. doi: 10.1177-0963721413495869.
- Vinci C, Peltier MR, Shah S, Kinsaul J, Waldo K, McVay M, Copeland AL. Effects of a brief mindfulness intervention on negative affect and urge to drink among college student drinkers. *Behav Res Ther*. 2014; 59:82–93. <http://dx.doi.org/10.1016/j.brat.2014.05.012>. [PubMed: 24972492]
- Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: The PANAS scales. *J Personality Soc Psychol*. 1988; 54:1063–1070.

- Wechsler H, Lee JE, Kuo M, Seibring M, Nelson TF, Lee H. Trends in college binge drinking during a period of increased prevention efforts. *J Am Coll Health*. 2002; 50(5):203–217. [PubMed: 11990979]
- Whiteside SP, Lynam DR. The Five Factor Model and impulsivity: using a structural model of personality to understand impulsivity. *Personal Individ Differ*. 2001; 30:669–689. 0191-8869/01.
- Williams AD, Grishman JR. Impulsivity, emotion regulation, and mindful attentional focus in compulsive buying. *Cogn Ther Res*. 2012; 36:451–457. <http://dx.doi.org/10.1007/s10608-011-9384-9>.
- Witkiewitz K, Warner K, Sully B, Barricks A, Stauffer C, Steckler G, Luoma J. Randomized trial comparing mindfulness based relapse prevention with relapse prevention for women offenders at a residential addiction treatment center. *Subst Use Misuse*. 2014; 49:536–546. [PubMed: 24611849]
- Wupperman P, Neumann CS, Whitman JB, Axelrod SR. The role of mindfulness in borderline personality disorder features. *J Nerv Ment Dis*. 2009; 197(10):766–771. <http://dx.doi.org/10.1097/NMD.0b013e3181b97343>. [PubMed: 19829206]

Fig. 1.
Tested moderation model.

Fig. 2.
Significant interaction of intervention type and Negative Urgency on Urge.

Fig. 3.
Significant interaction of intervention type and Positive Urgency on Positive Affect.

Table 1

Bivariate correlations of UPPS – P, TMS, FFMQ, and AUDIT scores for entire sample (n = 207) at baseline.

	Neg Urg	Premed	Persev	SS	Pos Urg	TMS Cur	TMS Dec	Observe	Describe	AA	Nonjudging	Nonreactivity
Neg Urg	–	–	–	–	–	–	–	–	–	–	–	–
Premed	0.40**	–	–	–	–	–	–	–	–	–	–	–
Persev	0.31**	0.38**	–	–	–	–	–	–	–	–	–	–
SS	0.12	0.35**	0.01	–	–	–	–	–	–	–	–	–
Pos Urg	0.57**	0.37**	0.37**	0.31**	–	–	–	–	–	–	–	–
TMS Cur	0.16*	–0.02	–0.03	0.14*	0.21**	–	–	–	–	–	–	–
TMS Dec	0.09	0.05	0.01	0.27**	0.22**	0.46**	–	–	–	–	–	–
Observe	–0.07	–0.08	–0.03	0.19**	0.03	0.24**	0.19**	–	–	–	–	–
Describe	–0.24**	–0.11	–0.17*	–0.01	–0.18**	0.03	0.14	0.16*	–	–	–	–
AA	–0.32**	–0.21**	–0.41**	–0.07	–0.34**	–0.17	–0.21**	–0.01	0.24**	–	–	–
Nonjudging	–0.41**	–0.03	–0.05	–0.07	–0.25**	–0.22**	–0.13	–0.12	0.13	0.21**	–	–
Nonreactivity	–0.31**	–0.15*	–0.13	0.18**	–0.07	0.01	0.22**	0.19**	0.23**	0.15*	0.07	–
AUDIT	0.30**	0.33**	0.30**	0.16*	0.34**	0.01	0.10	0.04	–0.13	–0.20**	–0.14*	–0.12

Note. Neg Urg = UPPTS-P Negative Urgency; Premed = UPPTS-P Premeditation; Persev = UPPTS-P Perseverance; SS = UPPTS-P Sensation Seeking; Pos Urg = UPPTS-P Positive Urgency; AA = Acting with Awareness; TMS = Toronto Mindfulness Scale; Cur = Curiosity; Dec = Decentering; AUDIT = Alcohol Use Disorders Identification Test.

**
p<0.01.

*
p<0.05.

Table 2

Significant hierarchical regression models.

Relaxation group moderates the relationship between sensation seeking and negative affect when compared to the mindfulness group								
Baseline negative affect	Gender	AUDIT	Control group ^a	Relaxation group ^b	Sensation seeking	Sensation seeking × control group ^a	Sensation seeking × relaxation group ^b	
B	0.34	-0.39	0.15	1.83	0.36	-0.16	-1.54	
SE B	0.05	0.45	0.05	2.31	0.56	0.77	0.76	
β	0.40***	-0.05	0.21**	0.28	0.07	-0.08	-0.71*	
Adjusted R ²				0.62				
R ²				0.02				
Relaxation group moderates the relationship between negative urgency and urge when compared to the mindfulness group								
Baseline Urge	Gender	AUDIT	Control group ^a	Relaxation group ^b	Negative urgency	Negative urgency × control group ^a	Negative urgency × relaxation group ^b	
B	0.77	-0.25	0.47	1.84	0.40	0.01	-0.82	
SE B	0.04	0.15	0.66	0.68	0.21	0.27	0.28	
β	0.81***	-0.06	0.12	0.47**	0.12	0.01	-0.54**	
Adjusted R ²			0.79					
R ²			0.01**					
Relaxation group moderates the relationship between positive urgency and positive affect when compared to the mindfulness group								
Baseline positive affect	Gender	AUDIT	Control group ^a	Relaxation group ^b	Positive urgency	Positive urgency × control group ^a	Positive urgency × relaxation group ^b	
B	0.71	-0.42	-0.08	2.20	-0.79	0.10	3.74	
SE B	0.06	0.96	0.10	3.21	1.22	1.63	1.63	
β	0.60***	-0.02	-0.05	0.14	-0.06	0.01	0.49*	
Adjusted R ²				0.48				
R ²				0.02*				
Control group moderates the relationship between perseverance and positive affect when compared to the mindfulness group								
Baseline positive affect	Gender	AUDIT	Control group ^a	Relaxation group ^b	Perseverance	Perseverance × control group ^a	Perseverance × relaxation group ^b	
B	0.72	-0.06	4.03	9.38	1.67	-3.78	-1.15	

Control group moderates the relationship between perseverance and positive affect when compared to the mindfulness group

	Baseline positive affect	Gender	AUDIT	Control group ^a	Relaxation group ^b	Perseverance	Perseverance × control group ^a	Perseverance × relaxation group ^b
SE B	0.06	0.92	0.10	3.58	3.65	1.30	1.84	1.83
β	0.61 ***	-0.01	0.01	0.58 *	-0.08	0.11	-0.46 *	-0.15
Adjusted R ²					0.46			
R ²					0.01			

Note:

AUDIT = Alcohol Use Disorders Identification Test.

B = unstandardized coefficient; SE = standard error; β = standardized coefficient.

* $p < 0.05$.
** $p < 0.01$.
*** $p < 0.001$.

^a dummy coded as: 0 = mindfulness and relaxation groups; 1 = control group.

^b dummy coded as: 0 = mindfulness and control groups; 1 = relaxation group.