bmj.com news roundup

Full versions of these stories are available at: bmj.com/content/vol329/issue7460/#NEWS_ROUNDUP

Childhood exposure to smoke may increase risk of back pain in later life

Exposure to secondhand smoke during childhood may increase the risk of developing back and neck problems in later life, says a new study. The research showed that adults who were exposed to tobacco smoke as children were more likely to take long term sick leave because of spinal pain.

One explanation, say the researchers, could be the effects of the smoke on the developing spine. "In several studies, smoking has been associated with the occurrence of spinal pain, mostly low back pain, but also neck pain and prolapsed cervical intervertebral discs," says the report in the *European Journal of Public Health* (2004;14:296-300).

The study, which was carried out at the University of Oslo and supported by the Norwegian Ministry of Health and Social Affairs, looked at a cohort of nurses' aides—the assistants who provide practical care to patients—to see whether exposure to environmental tobacco smoke at home during childhood led to a greater likelihood of long term sick leave.

The results showed that the aides who had been exposed to environmental tobacco smoke during childhood were more likely than aides who had not been exposed to smoke to take sick leave exceeding 14 days for neck pain (odds ratio 1.34 (95% confidence interval 1.04 to 1.73)), high back pain (1.49 (1.07 to 2.06)), and any illness (1.23 (1.07 to 1.42)), after demographic and family characteristics, former smoking status, current smoking status, physical leisure time activities, nature of work before neck injury, and affective symptoms were adjusted for. The difference in the number of aides who took leave exceeding 14 days for low back pain was not significant. Aides who had been exposed to environmental tobacco smoke were also more likely to take sick leave exceeding eight weeks (odds ratio 1.29 (1.08 to 1.55)).

Roger Dobson *Abergavenny*

Life expectancy in Great Britain rises—but later years are still spent in poor health

People are living longer but the extra years are not lived in good health, new data published by the Office for National Statistics show.

Life expectancy in Great Britain increased between 1981 and 2001. Females born in 2001 can now expect to live an average of 80.4 years (versus 76.8 years in 1981) and males an average of 75.7 years (versus 70.9 years). Although women have also in the past had a longer life expectancy than men, the new data show that life expectancy is increasing at a faster rate for males than for females.

Healthy life expectancy was quantified and defined as the expected years of life in good or fairly good health. In 2001 healthy life expectancy at birth was 67.0 years for males and 68.8 years for females. Although these values are slightly higher than for people born in 1981, healthy life expectancy has not been rising as fast as life expectancy.

People may be living longer, but the extra years are largely lived in poor health (defined as the difference between life expectancy and healthy life expectancy). In 1981, the expected time lived in poor health for males was 6.5 years; by 2001 this had risen to 8.7 years.

Women are even worse off, according to the data. Those born in 1981 could have expected to live in poor health for 10.1 years, but this rose to 11.6 years for those born in 2001.

Karen Hébert *BMJ*

See www.statistics.gov.uk

Bush signs law to protect US from bioterrorism

President George Bush last week signed into law Project BioShield to protect Americans against biological, chemical, radiological, or nuclear attack. He did this a day before the National Commission on Terrorist Attacks upon the United States delivered its final report. The commission found missed opportunities to intercept one or more of the hijackers who attacked the World Trade Center and the Pentagon in 2001.

The new law will introduce several important changes. It will earmark \$5.6bn (£3bn; €4.6bn) over 10 years for countermeasures against anthrax; smallpox; and other chemical, biological, radiological, and nuclear agents. Research on and development of medical countermeasures by the National Institutes of Health will be accelerated, and a national network of eight new regional Centers of Excellence for Biodefense will be set up. The Food and Drug Administration will be able to make promising treatments speedily available in emergency situations.

The Departments of Health

and Human Services, Homeland Security, and Agriculture will extend the surveillance of human, plant, and animal health and of disease by using an early warning system, Biowatch.

The strategic national stockpile of drugs and medical supplies will be increased to five times its size compared with 2001, including more antibiotics for treating exposure to anthrax, enough smallpox vaccine for every person in the United States, and more treatments for radiation poisoning and exposure to chemicals and pathogens.

Fred Charatan *Florida*

Combination anticoagulation therapy doubles risk of bleeding in patients at high risk of stroke

Patients at high risk of stroke should not be treated with combination anticoagulation therapy using aspirin and clopidogrel—known to be beneficial for those with cardiovascular disease—because it doubles the risk of bleeding, concludes new research from the University of Essen, Germany (*Lancet* 2004;364:331-7).

Previous research showed that clopidogrel was better than aspirin for patients who had recently had an ischaemic stroke, heart attack, or symptomatic peripheral arterial disease. Its benefits were greater in some high risk subgroups of patients, such as those with diabetes or previous cardiac surgery (*Lancet* 1996;348:1329-39).

The current findings are based on a randomised, double blind, placebo controlled trial, which included 7599 high risk patients from 507 stroke units and neurology departments in 28 countries.

The patients were already taking clopidogrel for either an ischaemic stroke or a transient ischaemic attack that they had had within the preceding three months. They also had at least