*Interactive case report***Abnormal liver function found after an unplanned consultation: case presentation**

James Heathcote

Mrs Prior is a 40 year old housewife whose husband frequently attends the surgery because of his diabetes. They are a close couple, and she takes the lead in managing his condition. In the past three years she has consulted her general practitioner just once on her own account—for dyspepsia. During one of her husband's consultations (and quite out of character) she mentions three problems of her own: a seven day history of intermittent "cystitis," for which she had taken over the counter sodium citrate; generalised itching; and a rash that looks to her general practitioner like seborrhoeic dermatitis.

In the limited time available (this is, after all, her husband's consultation) the general practitioner ascertains that her weight is steady and she is otherwise well, before examining her quickly to rule out any obvious skin disease to account for the itching. He finds no

Questions

- 1 What do these results suggest?
- 2 If you were her general practitioner, what would you do now and with what urgency?
- 3 What would you tell Mrs Prior?

Please respond on bmj.com

South View Lodge,
South View,
Bromley, Kent
BR1 3DR

James Heathcote
general practitioner

[james.heathcote@
gp-G84001.nhs.uk](mailto:james.heathcote@gp-G84001.nhs.uk)

BMJ 2004;329:273

Results of biochemical analysis and full blood count

	Result
Serum sodium (mmol/l)	141
Serum potassium (mmol/l)	4.2
Serum creatinine (μ mol/l)	44*
Serum total protein (g/l)	73
Serum albumin (g/l)	43
Alkaline phosphatase (U/l)	893*
γ -glutamyl transferase (U/l)	1227*
Aspartate aminotransferase (U/l)	103*
Total bilirubin (μ mol/l)	119*
Plasma glucose (nmol/l)	4.7
White blood cells ($\times 10^9$ /l)	10.2
Haemoglobin (g/l)	143
Mean corpuscular volume (fl)	101.8*
Mean corpuscular haemoglobin (pg/cell)	34.4*
Platelet count ($\times 10^9$ /l)	273

* Abnormal result.

hepatomegaly and no jaundice. He then orders a laboratory midstream urine culture, full blood count, electrolytes, plasma glucose, and liver function tests and prescribes a short course of cefalexin for the cystitis and a tube of cream containing hydrocortisone (1%) and miconazole (2%) for the rash.

Two days later, the urine result confirms an *Escherichia coli* infection with a high white blood cell count and culture of $> 100\,000$ organisms/ml. It is sensitive to cefalexin. Eight days after the initial consultation, Mrs Prior goes for her blood tests. The results come back to the general practitioner the next day (table). Mrs Prior has not yet made a follow up appointment.

Competing interests: None declared.

We welcome contributions of interactive case reports. Cases should raise interesting clinical, investigative, diagnostic, and management issues but not be so rare that they appeal to only a minority of readers. Full details of criteria are available at: bmj.com/cgi/content/full/3267/7389/564/DC1

This is the first of a three part case report where we invite readers to take part in considering the diagnosis and management of a case using the rapid response feature on bmj.com. Next week we will report the case progression and in four weeks' time we will report the outcome and summarise the responses

bmjlearning.com

None of the above?

Which of the following cities are furthest from London?

- A: Plymouth
- B: Cardiff
- C: York
- D: Paris
- E: None of the above.

"None of the above" often puzzles doctors when answering multiple choice questions.¹ It can work if all the other options are clearly incorrect: it is then the only correct answer. However, it causes confusion when the options are not absolutely true or false. You then have to weigh in your mind whether one of the specific options is more correct than the infinite number of other options that are encapsulated in "None of the above."¹ Sometimes, as in the above question, it makes no sense at all.

BMJ Learning's online learning resources can help you with your appraisal and revalidation. We ask questions throughout the modules, but we avoid imprecise and ambiguous questions. Our new learning module on influenza outlines how to diagnose and treat patients with this condition. It also brings you up to date with recent advances such as the role of neuraminidase inhibitors and amantadine in treatment. It states the circumstances in which patients should receive these drugs for treatment and for post-exposure prophylaxis.

To find out more, go to our website, bmjlearning.com.

Kieran Walsh *editorial registrar, BMJ Learning*
(bmjlearning@bmjgroup.com)

1 Case SM, Swanson DB. *Constructing written test questions for the basic and clinical sciences*. 3rd ed. Philadelphia PA: National Board of Medical Examiners, 2001.