CORRESPONDENCE

Inflammatory Bowel Disease in Pediatric Patients—Characteristics of Newly Diagnosed Patients From the CEDATA-GPGE Registry

by Dr. med. Stephan Buderus, Dr. med. Dietmar Scholz, Prof. Dr. med. Rolf Behrens, Dr. med. Martin Classen, Dr. med. Jan de Laffolie, Prof. Dr. med. Klaus-Michael Keller, Prof. Dr. med. Klaus-Peter Zimmer, Prof. Dr. med. Sibylle Koletzko in issue 8/2015

Poor Data Quality

Congratulations to the authors for their presentation of patient characteristics and related healthcare data (1). Unfortunately, their article is flawed by a few errors which can result in wrong conclusions being drawn (2). This applies in particular to Figure 3 where the completeness of the initial diagnostic evaluation is depicted. The authors describe an improvement in the guideline-compliant diagnostic assessment during the observation period (2004 to 2014). However, 24% of cases were excluded for incomplete data. It remains unclear how cases of missing documentation were distinguished from those with missing diagnostic evaluation. It would be important to know whether documentation quality has evolved over the observation period and at what time improvement initiatives were implemented, if any.

The analysis is primarily based on the reported data available in 2010 (935 cases); following the reporting gap between 2011 and 2013, only 23 newly diagnosed cases seem to have been added, despite the postulated rise in incidence. It appears to be plausible that a selection of cases of very committed reporting colleagues contrasted the poor documentation quality of less enthusiastic contributors to this project. The usability of the data should have been critically discussed against the background of lower reporting activity. Furthermore, the confidence intervals of the 2013/2014 data are so wide that the low case numbers alone make it impossible to ascertain the increase. In addition, limiting the y-axis scale to 100% halves the current confidence intervals, creating the false impression of high precision.

Thus, it would appear that the now presented data only allow for a very limited evaluation of care quality, if any. Likewise, due to insufficient data quality, other information, for example about phenotyping and medication, should be viewed critically. The authors overestimate the value of the registry in this regard. In any case, the issue has not been discussed in appropriate detail.

DOI: 10.3238/arztebl.2015.0544a

REFERENCES

1. Buderus S, Scholz D, Behrens R, et al. for the CEDATA-GPGE study group: Inflammatory bowel disease in pediatric patients—characteristics of newly diagnosed patients from the CEDATA-GPGE registry. *Dtsch Arztebl Int* 2015; 112: 121–7.
2. Drummond GB, Vowler SL: Show the data, don't conceal the data. *Adv Physiol Educ* 2011; 35: 130–2.

Prof. Dr. med. Antje Timmer

Carl von Ossietzky Universität Oldenburg
Abteilung Epidemiologie und Biometrie
Fakultät VI, Department für Versorgungsforschung
antje.timmer@uni-oldenburg.de

Conflict of interest statement

Prof. Timmer coordinated this registry between 2008 and 2010 and wrote a publication based on the registry data.

In Reply:

We like to thank Prof. Timmer for her critical evaluation focused on the limitations of large multicenter registries. Registries reflect the everyday and real-world healthcare situation of a large number of patients. However, along with this advantage comes the disadvantage of not being able to conduct and/or document every examination and variable with the same level of accuracy that is achievable within the tight set-up of controlled studies. Prof. Timmer herself made a very valuable contribution to the improvement of data quality by structuring plausibility checks and creating detailed SOPs, among others, during her time at the CEDATA registry between 2008 and 2010. Under her leadership, the registry data on the latency to diagnosis and associated parameters were published (reference 12 in the article). The new online version represents a further step to improve data quality. At the time of data analysis (end of 2013), only 23 new patients had been included. These data were only included into our current analysis on the request of the reviewers, and we reported these numbers transparently. The confidence interval was computed using the Clopper-Pearson method; thus it is accurately presented and the top part has not been cut off. Even without the patients included in 2013, an improvement of the complete diagnostic assessment was noted between 2004 and 2010. For phenotyping, analysis was limited to data of patients for whom the required endoscopic and imaging information was available. The validity of our results is supported by the fact that they are in line with the results of the European registry (references 18, 25). With regard to medication, we agree with Prof. Timmer that there are significant limitations in this respect. We are trying to close the gap in documentation using the online reporting version and hope that, in doing so, we will be able to get closer to our goal of improving healthcare quality.

DOI: 10.3238/arztebl.2015.0544b

REFERENCES

1. Buderus S, Scholz D, Behrens R, et al. for the CEDATA-GPGE study group: Inflammatory bowel disease in pediatric patients—characteristics of newly diagnosed patients from the CEDATA-GPGE registry. *Dtsch Arztebl Int* 2015; 112: 121–7.

On behalf of the authors:

Dr. med. Stephan Buderus

GFO-Kliniken Bonn
stephan.buderus@marien-hospital-bonn.de

Conflict of interest statement

Dr. Buderus has received consultancy fees from MSD. He has received fees for preparing scientific continuing professional development events from AbbVie, MSD, and Nestlé. He has received reimbursement of conference fees from the Falk Foundation.