COMMENT

Response to “Consumption of wild-harvested meat from New Zealand feral animals provides a unique opportunity to study the health effects of lead exposure in hunters” by Buenz et al.

Niels Kanstrup, Vernon G. Thomas,
Oliver Krone, Carl Gremse

Published online: 25 June 2016

Comment to: Buenz, E., G. Parry, and M.M. Peacey. 2016. Consumption of wild-harvested meat from New Zealand feral animals provides a unique opportunity to study the health effects of lead exposure in hunters. *Ambio*. doi:[10.1007/s13280-016-0798-1](https://doi.org/10.1007/s13280-016-0798-1).

While acknowledging endorsement of our paper from Buenz et al. (2016), we believe that further study of relations between elevated human blood lead levels and lead in game meat is not required to validate the transition to non-lead bullets. The essence of our article (Kanstrup et al. 2016) was that Denmark could extend its existing bans on possession and use of lead gunshot and fishing weights to lead-core rifle ammunition, and thus complete the transition. Denmark faces several binding legal obligations to do so. We presented evidence for defined risks from bullet-derived lead: human health concerns from eating game meat, and the health of scavenging wildlife. Because Denmark has not monitored these areas, our evidence was indirect.

Sport hunting is international, as are the types of lead-core ammunition used, and behaviours of such bullets in animals. Thus, international research can support progressive changes in government policy on environmental lead reduction, especially given the same sources and routes of exposure.

Levels of lead in blood and game meat were correlated for US and Norwegian hunters (Iqbal et al. 2009; Meltzer et al. 2013; Knutsen et al. 2015), and confirmed by isotope analyses (Tsuji et al. 2008). High levels of bullet-derived lead exist in wild game meat (Dobrowolska and Melosik 2008; Morales et al. 2011; Fachehoun et al. 2015) that often exceed the European Commission threshold of 0.2 mg Pb/Kg.

Our paper emphasized the risks of lead exposure to wild scavengers from ingested lead fragments in shot animals, an aspect not addressed by Buenz et al. (2016). The scientific evidence for this relationship is compelling, especially for avian scavengers of the Baltic-European region (Helander et al. 2009; Krone et al. 2009; Nadjafzadeh et al. 2013).

Given the consistent conclusions from the above-cited research, further monitoring is not likely to alter our understanding of the risks to human and avian health. However, the study proposed by Buenz et al. (2016) could facilitate awareness at the local level and stimulate a regulated transition to non-lead hunting ammunition in New Zealand.

REFERENCES

- Buenz, E., G. Parry, and M.M. Peacey. 2016. Consumption of wild-harvested meat from New Zealand feral animals provides a unique opportunity to study the health effects of lead exposure in hunters. *Ambio*. doi:[10.1007/s13280-016-0798-1](https://doi.org/10.1007/s13280-016-0798-1).
- Dobrowolska, A., and M. Melosik. 2008. Bullet-derived lead in tissues of the wild boar (*Sus scrofa*) and red deer (*Cervus elaphus*). *European Journal of Wildlife Research* 54: 231–235. doi:[10.1007/s10344-007-0134-y](https://doi.org/10.1007/s10344-007-0134-y).
- Fachehoun, R.C., B. Lévesque, P. Dumas, A. St-Louis, M. Dubé, and P. Ayotte. 2015. Lead exposure through consumption of big game meat in Quebec, Canada: Risk assessment and perception. *Food Additives and Contaminants: Part A* 32: 1501–1511. doi:[10.1080/19440049.2015.1071921](https://doi.org/10.1080/19440049.2015.1071921).
- Helander, B., J. Axelsson, H. Borg, K. Holm, and A. Bignert. 2009. Ingestion of lead from ammunition and lead concentrations in white-tailed sea eagles (*Haliaeetus albicilla*) in Sweden. *Science of the Total Environment* 407: 5555–5563. doi:[10.1016/j.scitotenv.2009.07.027](https://doi.org/10.1016/j.scitotenv.2009.07.027).

- Iqbal, S., W. Blumenthal, C. Kennedy, F.Y. Yip, S. Pickard, W.D. Flanders, K. Loring, K. Kruger, et al. 2009. Hunting with lead: association between blood lead levels and wild game consumption. *Environmental Research* 109: 952–959. doi:[10.1016/j.envres.2009.08.007](https://doi.org/10.1016/j.envres.2009.08.007).
- Kanstrup, N., V.G. Thomas, O. Krone, and C. Gremse. 2016. The transition to non-lead rifle ammunition in Denmark: National obligations and policy considerations. *Ambio*. doi:[10.1007/s13280-016-0780-y](https://doi.org/10.1007/s13280-016-0780-y).
- Knutsen, H.K., A.L. Brantsæter, J. Alexander, and H.M. Meltzer. 2015. Associations between consumption of large game animals and blood lead levels in humans in Europe: the Norwegian experience. In *Proceedings of the Oxford Lead Symposium. Lead Ammunition: understanding and minimising the risks to human and environmental health*, ed. R.J. Delahay and C.J. Spray, 44–50. Oxford, UK: The Edward Grey Institute.
- Krone, O., N. Kennntner, A. Trinogga, N. Nadjafzadeh, F. Scholz, J. Sulawa, K. Totschek, P. Schuck-Wersig, et al. 2009. Lead poisoning in white-tailed sea eagles: Causes and approaches to solutions in Germany. In *Ingestion of lead from spent ammunition: Implications for wildlife and humans*, ed. R.T. Watson, M. Fuller, M. Pokras, and W.G. Hunt, 289–301, Boise, ID: The Peregrine Fund. doi:[10.4080/ilsa.2009.0207](https://doi.org/10.4080/ilsa.2009.0207).
- Meltzer, H.M., H. Dahl, A.L. Brantsæter, B.E. Birgisdottir, H.K. Knutsen, A. Bernhoft, B. Oftedal, U.S. Lande, et al. 2013. Consumption of lead-shot cervid meat and blood lead concentrations in a group of adult Norwegians. *Environmental Research* 127: 29–39. doi:[10.1016/j.envres.2013.08.007](https://doi.org/10.1016/j.envres.2013.08.007).
- Morales, J.S.S., R.M. Rojas, F. Pérez-Rodríguez, A.A. Casas, and M.A.A. López. 2011. Risk assessment of the lead intake by consumption of red deer and wild boar meat in Southern Spain. *Food Additives & Contaminants* 28: 1021–1033. doi:[10.1080/19440049.2011.583282](https://doi.org/10.1080/19440049.2011.583282).
- Nadjafzadeh, M., H. Hofer, and O. Krone. 2013. The link between feeding ecology and lead poisoning in White-Tailed Eagles. *Journal of Wildlife Management* 77: 48–57. doi:[10.1002/jwmg.440](https://doi.org/10.1002/jwmg.440).
- Tsuji, L.J.S., B.C. Wainman, I.D. Martin, C. Sutherland, J.-P. Weber, P. Dumas, and E. Nieboer. 2008. Lead shot contribution to blood lead of First Nations people: the use of lead isotopes to identify the source of exposure. *Science of the Total Environment* 405: 180–185. doi:[10.1016/j.scitotenv.2008.06.048](https://doi.org/10.1016/j.scitotenv.2008.06.048).
- Niels Kanstrup**
Address: Danish Academy of Hunting, Skrejrupvej 31, 8410 Rønne, Denmark.
e-mail: nk@danskjagtakademi.dk
- Vernon G. Thomas** (✉)
Address: Department of Integrative Biology, University of Guelph, Guelph, ON N1G 2W1, Canada.
e-mail: vthomas@uoguelph.ca
- Oliver Krone**
Address: Department of Wildlife Diseases, Leibniz Institute for Zoo and Wildlife Research, P.O. 601103, 10252 Berlin, Germany.
e-mail: krone@izw-berlin.de
- Carl Gremse**
Address: Wildlife Department, Eberswalde-University, Schicklerstrasse 5, 16225 Eberswalde, Germany.
e-mail: carl.gremse@hnee.de