Published in final edited form as:

Neurosci Lett. 2016 August 26; 629: 15–18. doi:10.1016/j.neulet.2016.06.037.

***Tsc1* haploinsufficiency is sufficient to increase dendritic patterning and Filamin A levels**

Longbo Zhang^{1,2,*}, Tianxiang Huang^{1,2,*}, and Angélique Bordey²

¹Department of Neurosurgery, Xiangya Hospital, Central South University, 85 Xiangya Street, Changsha, 410008, China

²Departments of Neurosurgery, and Cellular & Molecular Physiology, Yale University School of Medicine, 333 Cedar Street, New Haven, CT 06520-8082, USA

Abstract

Most individuals with tuberous sclerosis complex (TSC) are born with a mutant allele of either *TSC1* or *TSC2* and a mosaic of psychological and cognitive defects. *Tsc1* loss of heterozygosity contributes to severe dendritic abnormalities that are rescued by normalizing the levels of the actin-cross linking protein, Filamin A (FLNA). However, it is unclear whether dendrites and FLNA levels are abnormal in an heterozygote *Tsc1* condition. Here, we examined dendritic morphology and FLNA levels in the olfactory bulb of *Tsc1* wild type and heterozygote mice. Using *in vivo* neonatal electroporation to label newborn neurons followed by sholl analysis, we found that *Tsc1* haploinsufficiency is associated with increased dendritic complexity and total dendritic length as well as increased FLNA levels. Since reducing FLNA levels has been shown to decrease *Tsc1*^{+/-} dendritic complexity, these data suggest that increased FLNA levels in *Tsc1*^{+/-} mice contribute to abnormal dendritic patterning in the *Tsc1* heterozygote condition of individuals with TSC.

Keywords

dendrite; mTOR; tuberous sclerosis complex; Filamin A; circuit; network; neurogenesis

1. Introduction

TSC is an autosomal dominant monogenetic disorder observed in 1/6,000 individuals characterized by discrete lesions in diverse tissues, including the skin, heart, kidney, lung, and brain [1]). Most patients are born with at least one detectable mutation in *TSC1* or *TSC2* and are thus heterozygous for one of these genes [2;3]. Very often, there are subsequent

Address for correspondence: Angélique Bordey, Ph.D., Department of Neurosurgery, Yale University School of Medicine, 333 Cedar Street, FMB 422, New Haven, CT 06520-8082, Phone: 203-737-2515, Fax: 203-737-2159, angelique.bordey@yale.edu.
*co-first author

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

inactivating mutations of the other functional allele during development leading to the formation of discrete lesions in several organs, including the brain [4–7].

The cortical lesions, called cortical tubers, are responsible for seizures occurring in >80% of TSC individuals [1]. However, TSC individuals display a mosaic of neuropsychological problems (*e.g.* autistic traits) and about half of them present with cognitive impairments (*i.e.* mental retardation). Although epilepsy can lead to both cognitive and psychiatric deficits, it is thought that heterozygosity for *TSC1* and *TSC2* is sufficient to account for some of these deficits. Indeed, *Tsc1*^{+/-} and *Tsc2*^{+/-} mice as well as mice with a heterozygote *Tsc2* dominant negative allele (*Tsc2*^{RG}) display cognitive and social deficits despite the absence of brain lesions or seizures ([8] and for review see [9]). These deficits likely result from a collection of cellular and molecular alterations in heterozygote mice. At the molecular level, mTORC1 and ERK activities have been shown to be increased in *Tsc1*^{+/-} and *Tsc2*^{RG} mice, respectively [10;11]. At the cellular level, retinogeniculate projections have been reported to show abnormalities in targeting in *Tsc1*^{+/-} mice [12]. Alterations in dendritic patterning have not been observed in hippocampal neurons of *Tsc1*^{+/-} mice in vivo [13]. This is somehow surprising considering that dendritic defects are often observed in disorders associated with cognitive or psychiatric deficits [14–17] and are a classical outcome of increased mTORC1 or ERK activity (see [9;18] for additional references). Because, the defects in axonal patterning were modest, it is possible that a modest defect in dendritic patterning may not be detectable by analyzing the number of first and second dendrites as performed in the published study using *Tsc1*^{+/-} mice [13].

Here, we thus set out to examine the dendritic complexity of neurons in *Tsc1*^{+/-} mice. To efficiently label dendrites, we used mice crossed with a tdTomato reporter line, in which expression of a Cre recombinase (Cre)-encoding plasmid leads to tdTomato expression. We electroporated a Cre plasmid in newborn neurons of the subventricular zone that migrate and become interneurons in the olfactory bulb (OB) [19]. We found that *Tsc1*^{+/-} neurons display a modest but significant increase in dendritic complexity and total length compared to wild type *Tsc1*^{+/+} neurons. Intriguingly, a recent study reported that decreasing the level of an actin-cross-linking protein filamin A (FLNA) [20] in conditional *Tsc1*^{+/-} neurons of the OB decreased their dendritic complexity [18]. The recent study also reported increased levels of FLNA in conditional *Tsc1*^{-/-} (null) neurons compared to *Tsc1*^{+/-} neurons and in TSC patients, but FLNA levels in *Tsc1*^{+/-} versus wild type *Tsc1* mice were not examined. In light of the data on dendrites in *Tsc1*^{+/-} mice reported here, we examined FLNA levels in these mice and found that there were increased compared to those in *Tsc1*^{+/+} mice. In addition, correlation analysis between TSC1 and FLNA levels in *Tsc1* mice suggest a gene-dosage relationship between *Tsc1* and FLNA levels.

2. Materials and Methods

Animals

Research protocols were approved by the Yale University Institutional Animal Care and Use Committee. Experiments were performed on the following lines of transgenic mice of either gender: *Tsc1*^{fl/+} (fl, floxed, obtained from Jackson Labs), *Tsc1*^{+/+}, and *Tsc1*^{+/-} mice (obtained from the National Cancer Institute). All these lines were crossed with Rosa26R-

STOP-tdTomato reporter mice (Jackson labs). Tail or toe samples were taken and were subjected to DNA isolation, PCR amplification using previously published primers [21] and amplicons separated by standard electrophoresis methods.

Neonatal electroporation and vectors

Postnatal electroporation was performed as previously described [19;22;23]. Plasmids (2–3 µg/µl) were diluted in PBS containing 0.1% fast green as a tracer. 0.5–1 µl of plasmid solution was injected into the lateral ventricles of neonatal pups using a pulled glass pipette (diameter <50 µm). 5 square-pulses of 50 ms-duration with 950 ms-intervals at 100 V were applied using the ECM 830 square wave pulse generator (BTX) and tweezer-type electrodes (model 520, BTX) placed on the heads of P0-P1 pups. The electrodes were positioned to direct current rostrally in the dorso-lateral SVZ. Vectors included pCAG-tdTomato [24] and pCAG-Cre (addgene, C. Cepko).

Morphometric analysis

Images of tdTomato⁺ basal dendrites were acquired in coronal sections using a Fluoview 1000 confocal microscope and traced with simple neurite tracer software (FIJI, GNU GPL v3, an ImageJ plugin, URL: http://fiji.sc/Simple_Neurite_Tracer). We analyzed the basal dendrites of granule cells in a specific layer for the following technical reason; the basal dendrite can be fully imaged and traced without confusion from a single labeled granule cell. Due to its length, the apical dendrite is often cut or partially cut, and this could lead to erroneous data. In addition, in the external plexiform layer where apical dendrites branch and terminate, it is difficult to distinguish whether the dendrite come from one cell or several cells. Sholl analyses were carried out using the number of intersections in 10 µm-increment concentric circles as a measure of morphological complexity. Z-stacks from 3 different square fields of view were taken from 3 different olfactory bulb (OB) sections. Analysis was performed blindly from at least 3 animals per condition.

Olfactory bulb lysate and western blot

Mice were anesthetized with pentobarbital (50 mg/kg) prior to olfactory bulb (OB) dissection. Western blotting protocol is detailed in [18]. Samples were homogenized in RIPA buffer, 1x Halt Protease and phosphatase inhibitor cocktail (Thermo Fisher Scientific), and 8 U/ml DNase. Samples were boiled in 2× Laemmli's buffer. 10 or 20 µg protein/sample was loaded into a 4–15% polyacrylamide gel (Bio-Rad Mini Protean TGX gel). Proteins were transferred to PVDF and blocked in 5% milk or 5% BSA. Quantifications were performed using NIH Image J software. Primary antibodies included: FLNA (molecular weight: 281 kDa, Abcam, ab51217, 1:5,000), TSC1 (predicted band size: 130 kDa, observed at 150–160 kDa as reported for the company site, Abcam, ab32936, 1:5,000), and ERK1/2 (molecular weight of ERK1: 44 kDa, Santa Cruz, sc-94, 1:250,000). We used the Thermo Scientific BenchMark™ Pre-Stained Protein Ladder containing a mixture of 10 proteins spanning 6–180 kDa (Catalog number: 10748010, invitogen).

Statistics

Analysis was performed on N indicating the number of animals in each condition or n indicating the number of replicates. Data were presented in GraphPad Prism 6. Statistical significance was determined using unpaired Student t-test or one way ANOVA with $p < 0.05$ or two way ANOVA with *post hoc* Bonferroni's test for comparison of dendritic crossing. Data are presented as mean $\pm$ standard error of the mean (SEM).

3. Results

To investigate whether *Tsc1* haploinsufficiency would lead to alterations in dendritogenesis *in vivo*, we employed neonatal electroporation to fluorescently label developing OB neurons in *Tsc1*^{+/+}, *Tsc1*^{+/-}, and *Tsc1*^{fl/+} mice crossed with Rosa26R-STOP-tdTomato reporter mice (Fig. 1A). We electroporated pCAG-Cre to excise the floxed *Tsc1* allele and the STOP cassette leading to tdTomato expression at P0 into neural progenitor cells of the subventricular zone. During division, electroporated neural progenitors pass their exogenous plasmids on to neuroblasts, which migrate and integrate into the OB [19]. By 2 weeks, these neuroblasts differentiate into GABAergic interneurons with both apical and basal dendrites. We assessed the complexity and length of the basal dendrites of tdTomato⁺ neurons *in vivo* at 14 days post-electroporation by sholl analysis in coronal sections from littermate mice (Fig. 1B). Sholl analysis revealed that the dendrites of *Tsc1*^{+/-} neurons in both constitutive and conditional *Tsc1*^{+/-} mice displayed a significant increase in their dendritic complexity (2-way ANOVA with *post hoc* Bonferroni's test, $n=27-30$ neurons, $N=3$ mice each, Fig. 1C) and total dendritic length (One way ANOVA, $P=0.0184$, $F=4.194$, Fig. 1D).

Considering that decreasing FLNA levels in conditional *Tsc1*^{+/-} neurons decreased their dendritic complexity and total length [18], we examined whether FLNA levels were increased in *Tsc1*^{+/-} versus *Tsc1*^{+/+} mice at P14. We performed immunoblotting for FLNA, TSC1 and the loading control ERK1/2, which is not altered in *Tsc1* mice of different genotypes [10]. We found that FLNA levels were significantly increased in the OB as a result of *Tsc1* haploinsufficiency ($N=7$ per condition, $P=0.0045$, unpaired t-test, Fig. 2A and B). In addition, correlation analysis between FLNA and TSC1 levels revealed that the levels of TSC1 and FLNA were inversely correlated ($R^2=0.53$, $P=0.003$, $N=14$ from *Tsc1*^{+/+} and *Tsc1*^{+/-} mice, Fig. 2C).

4. Discussion

Our data suggest that *Tsc1* haploinsufficiency as reported in TSC individuals is associated with increased dendritic patterning and increased levels of FLNA. We have previously reported that decreasing FLNA levels in heterozygote neurons was sufficient to reduce dendritic complexity [18], but it was unclear whether dendrites were altered in the heterozygote condition or whether FLNA was just required for dendritic maintenance. Together with the published findings, our data suggest that increased FLNA levels in *TSC1* heterozygote condition contributes to altered dendritic complexity.

FLNA is an actin-cross linking protein with multiple binding partners and a spectrum of functions inside cells, including neurons [20;25;26]. As such the mechanism of FLNA action

on dendritic development remains unclear and is outside the scope of this study. Nevertheless, our finding has several important implications. First, it was also recently reported that increased FLNA levels can alter spine morphology [27] in addition to dendrites as shown here. A change in dendritic morphology, including spine morphology is a hallmark of several disorders associated with cognitive and deficits [15;28]. In addition, *Tsc1*^{+/-} mice have both cognitive and social deficits [8;13]. Our findings thus beg the question as to whether increased FLNA levels could contribute to cognitive and psychiatric deficits in TSC. Second, our findings are not limited to TSC. Indeed, we recently reported that increased FLNA levels are increased as a result of increased MEK/ERK1/2 signaling in *Tsc1* null condition [18]. Several neurodevelopmental disorders are considered to be part of a RAS/MAPK syndrome associated with increased MEK/ERK activity. It would thus be intriguing to examine whether FLNA levels are increased in these disorders associated with neurocognitive deficits. As part of this non-neurocognitive group is cancer which is associated with altered FLNA levels [29]. Another condition leading to increased FLNA levels is hypoxia, in which increased FLNA contributed to altered spine morphology.

Finding a correlation between FLNA and TSC1 levels suggests that TSC individuals will have differences in their FLNA levels due to allele-specific variations in *Tsc1* mRNA expression [30] that could contribute to a spectrum of dendritic dysmorphogenesis and cognitive or psychiatric disparities. In addition, a copy number variation of *FLNA* is found in the genome of patients suffering from autism disorder [31]. Whether this could contribute to patterning abnormalities remain to be examined.

All together these findings emphasize the need to further examine FLNA contribution to neuronal morphology and patterning as well as network activity and behavior in several disorders associated with alterations in RAS/MAPK and AKT/TSC/mTOR signaling.

Acknowledgments

This work was supported by grants from NIH-NINDS R01NS093704 (AB), training grant from the China Scholarship Council (LZ and TH), and Young Investigator Grant of Xiangya Hospital, CSU (2015Q02, LZ).

References

1. Crino PB, Nathanson KL, Henske EP. The tuberous sclerosis complex. *N Engl J Med*. 2006; 355:1345–1356. [PubMed: 17005952]
2. van Slegtenhorst M, de Hoogt R, Hermans C, Nellist M, Janssen B, Verhoef S, Lindhout D, van den OA, Halley D, Young J, Burley M, Jeremiah S, Woodward K, Nahmias J, Fox M, Ekong R, Osborne J, Wolfe J, Povey S, Snell RG, Cheadle JP, Jones AC, Tachataki M, Ravine D, Sampson JR, Reeve MP, Richardson P, Wilmer F, Munro C, Hawkins TL, Sepp T, Ali JB, Ward S, Green AJ, Yates JR, Kwiatkowska J, Henske EP, Short MP, Haines JH, Jozwiak S, Kwiatkowski DJ. Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34. *Science*. 1997; 277:805–808. [PubMed: 9242607]
3. Kwiatkowski DJ, Short MP. Tuberous sclerosis. *Arch Dermatol*. 1994; 130:348–354. [PubMed: 8129414]
4. Green AJ, Smith M, Yates JR. Loss of heterozygosity on chromosome 16p13.3 in hamartomas from tuberous sclerosis patients. *Nat Genet*. 1994; 6:193–196. [PubMed: 8162074]
5. Sepp T, Yates JR, Green AJ. Loss of heterozygosity in tuberous sclerosis hamartomas. *J Med Genet*. 1996; 33:962–964. [PubMed: 8950679]

6. Henske EP, Scheithauer BW, Short MP, Wollmann R, Nahmias J, Hornigold N, van Slegtenhorst M, Welsh CT, Kwiatkowski DJ. Allelic loss is frequent in tuberous sclerosis kidney lesions but rare in brain lesions. *Am J Hum Genet.* 1996; 59:400–406. [PubMed: 8755927]
7. Crino PB, Aronica E, Baltuch G, Nathanson KL. Biallelic TSC gene inactivation in tuberous sclerosis complex. *Neurology.* 2010; 74:1716–1723. [PubMed: 20498439]
8. Sato A, Kasai S, Kobayashi T, Takamatsu Y, Hino O, Ikeda K, Mizuguchi M. Rapamycin reverses impaired social interaction in mouse models of tuberous sclerosis complex. *Nat Commun.* 2012; 3:1292. [PubMed: 23250422]
9. Feliciano DM, Hartman NW, Lin TV, Bartley C, Kubera C, Hsieh L, Lafourcade C, O'Keefe RA, Bordey A. A circuitry and biochemical basis for tuberous sclerosis symptoms: from epilepsy to neurocognitive deficits. *Int J Dev Neurosci.* 2013;667–678. [PubMed: 23485365]
10. Bartley CM, O'Keefe RA, Bordey A. FMRP S499 is phosphorylated independent of mTORC1-S6K1 activity. *PLoS ONE.* 2014; 9:e96956. [PubMed: 24806451]
11. Chevere-Torres I, Maki JM, Santini E, Klann E. Impaired social interactions and motor learning skills in tuberous sclerosis complex model mice expressing a dominant/negative form of tuberin. *Neurobiol Dis.* 2012; 45:156–164. [PubMed: 21827857]
12. Nie D, Di NA, Han JM, Baharanyi H, Kramvis I, Huynh T, Dabora S, Codeluppi S, Pandolfi PP, Pasquale EB, Sahin M. Tsc2-Rheb signaling regulates EphA-mediated axon guidance. *Nat Neurosci.* 2010; 13:163–172. [PubMed: 20062052]
13. Goorden SM, van Woerden GM, van der WL, Cheadle JP, Elgersma Y. Cognitive deficits in Tsc1+/- mice in the absence of cerebral lesions and seizures. *Ann Neurol.* 2007; 62:648–655. [PubMed: 18067135]
14. Jan YN, Jan LY. Branching out: mechanisms of dendritic arborization. *Nat Rev Neurosci.* 2010; 11:316–328. [PubMed: 20404840]
15. Kaufmann WE, Moser HW. Dendritic anomalies in disorders associated with mental retardation. *Cereb Cortex.* 2000; 10:981–991. [PubMed: 11007549]
16. Dierssen M, Ramakers GJ. Dendritic pathology in mental retardation: from molecular genetics to neurobiology. *Genes Brain Behav.* 2006; 5(Suppl 2):48–60. [PubMed: 16681800]
17. Nestor MW, Hoffman DA. Aberrant dendritic excitability: a common pathophysiology in CNS disorders affecting memory? *Mol Neurobiol.* 2012; 45:478–487. [PubMed: 22528602]
18. Zhang L, Bartley CM, Gong X, Hsieh LS, Lin TV, Feliciano DM, Bordey A. MEK-ERK1/2-Dependent FLNA Overexpression Promotes Abnormal Dendritic Patterning in Tuberous Sclerosis Independent of mTOR. *Neuron.* 2014; 84:78–91. [PubMed: 25277454]
19. Lacar B, Young SZ, Platel JC, Bordey A. Imaging and recording subventricular zone progenitor cells in live tissue of postnatal mice. *Front Neurosci.* 2010; 4
20. Nakamura F, Stossel TP, Hartwig JH. The filamins: organizers of cell structure and function. *Cell Adh Migr.* 2011; 5:160–169. [PubMed: 21169733]
21. Feliciano DM, Su T, Lopez J, Platel JC, Bordey A. Single-cell Tsc1 knockout during corticogenesis generates tuber-like lesions and reduces seizure threshold in mice. *J Clin Invest.* 2011; 121:1596–1607. [PubMed: 21403402]
22. Feliciano DM, Lafourcade CA, Bordey A. Neonatal subventricular zone electroporation. *J Vis Exp.* 2013
23. Platel JC, Dave KA, Gordon V, Lacar B, Rubio ME, Bordey A. NMDA receptors activated by subventricular zone astrocytic glutamate are critical for neuroblast survival prior to entering a synaptic network. *Neuron.* 2010; 65:859–872. [PubMed: 20346761]
24. Pathania M, Torres-Reveron J, Yan L, Kimura T, Lin TV, Gordon V, Teng ZQ, Zhao X, Fulga TA, Van VD, Bordey A. miR-132 enhances dendritic morphogenesis, spine density, synaptic integration, and survival of newborn olfactory bulb neurons. *PLoS ONE.* 2012; 7:e38174. [PubMed: 22693596]
25. Feng Y, Walsh CA. The many faces of filamin: a versatile molecular scaffold for cell motility and signalling. *Nat Cell Biol.* 2004; 6:1034–1038. [PubMed: 15516996]
26. Sarkisian MR, Bartley CM, Rakic P. Trouble making the first move: interpreting arrested neuronal migration in the cerebral cortex. *Trends Neurosci.* 2008; 31:54–61. [PubMed: 18201775]

27. Segura I, Lange C, Knevels E, Moskalyuk A, Pulizzi R, Eelen G, Chaze T, Tudor C, Boulegue C, Holt M, Daelemans D, Matondo M, Ghesquiere B, Giugliano M, Ruiz de AC, Dewerchin M, Carmeliet P. The Oxygen Sensor PHD2 Controls Dendritic Spines and Synapses via Modification of Filamin A. *Cell Rep.* 2016; 14:2653–2667. [PubMed: 26972007]
28. Penzes P, Cahill ME, Jones KA, VanLeeuwen JE, Woolfrey KM. Dendritic spine pathology in neuropsychiatric disorders. *Nat Neurosci.* 2011; 14:285–293. [PubMed: 21346746]
29. Savoy RM, Ghosh PM. The dual role of filamin A in cancer: can't live with (too much of) it can't live without it. *Endocr Relat Cancer.* 2013; 20:R341–R356. [PubMed: 24108109]
30. Jentarra GM, Rice SG, Olfers S, Saffen D, Narayanan V. Evidence for population variation in TSC1 and TSC2 gene expression. *BMC Med Genet.* 2011; 12:29. [PubMed: 21345208]
31. Gilman SR, Chang J, Xu B, Bawa TS, Gogos JA, Karayiorgou M, Vitkup D. Diverse types of genetic variation converge on functional gene networks involved in schizophrenia. *Nat Neurosci.* 2012; 15:1723–1728. [PubMed: 23143521]

Highlights

- *Tsc1* heterozygote neurons display increased dendritic arborization.
- *Tsc1* heterozygosity leads to increased filamin A levels.
- Filamin A and TSC1 levels are correlated.

Figure 1. Constitutive and conditional *Tsc1*^{+/-} neurons display increased dendritic patterning compared to *Tsc1*^{+/+} neurons

(A) Diagram illustrating electroporation in P0 neonates. Most transfected newborn neurons reach their final location in the olfactory bulb (OB) by P14. (B) Image and reconstruction of the basal dendrite of an OB granule cell expressing tdTomato with superimposed concentric circles for sholl analysis. (C) # crossings as a function of distance for the basal dendrites of tdTomato+ neurons at P14 in *Tsc1*^{+/+} (black), *Tsc1*^{+/-} (violet) and *Tsc1*^{fl/+} (green) mice. Neurons were electroporated with pCAG-Cre leading to *Tsc1* heterozygosity in *Tsc1*^{fl/+} mice and tdTomato expression due to crossing with R26R-Stop-tdTomato mice. *: P<0.05, **: P<0.01, purple for comparisons between *Tsc1* wild type and heterozygote mice, and green between wild type and *Tsc1*^{fl/+} mice using two way ANOVA followed by Bonferroni multiple comparisons. (D) Total dendritic length corresponding to the conditions in C.

Figure 2. FLNA levels are increased in *Tsc1*^{+/-} mice and correlate with TSC1 levels
(A) Immunoblots for FLNA, TSC1, and ERK1/2 from P14 OB of *Tsc1*^{+/+} (black) and *Tsc1*^{+/-} (grey). **(B)** Normalized % changes for FLNA/ERK. **(C)** Plot of FLNA versus TSC1 levels.