Published in final edited form as:

Adv Cancer Res. 2015 ; 127: 49–121. doi:10.1016/bs.acr.2015.04.006.

Examination of Epigenetic and other Molecular Factors Associated with *mda-9/Syntenin* Dysregulation in Cancer Through Integrated Analyses of Public Genomic Datasets

Manny D. Bacolod^{1,2}, Swadesh K. Das^{1,2,3}, Upneet K. Sokhi¹, Steven Bradley⁴, David A. Fenstermacher^{3,5}, Maurizio Pellecchia⁶, Luni Emdad^{1,2,3}, Devanand Sarkar^{1,2,3}, and Paul B. Fisher^{1,2,3,*}

¹Department of Human and Molecular Genetics, Virginia Commonwealth University, School of Medicine, Richmond, Virginia 23298

²VCU Institute of Molecular Medicine, Virginia Commonwealth University, School of Medicine, Richmond, Virginia 23298

³VCU Massey Cancer Center, Virginia Commonwealth University, School of Medicine, Richmond, Virginia 23298

⁴VCU Bioinformatics Program, Virginia Commonwealth University, School of Medicine, Richmond, Virginia 23298

⁵Department of Biostatistics, Virginia Commonwealth University, School of Medicine, Richmond, Virginia 23298

⁶Sanford-Burnham Medical Research Institute, La Jolla, California 92037

Abstract

mda-9/Syntenin (melanoma differentiation-associated gene 9) is a PDZ domain-containing, cancer invasion-related protein. In this study, we employed multiple integrated bioinformatic approaches to identify the probable epigenetic factors, molecular pathways, and functionalities associated with *mda-9* dysregulation during cancer progression. Analyses of publicly available genomic data (e.g., expression, copy number, methylation) from TCGA, GEO, ENCODE, and Human Protein Atlas projects led to the following observations: a) *mda-9* expression correlates with both copy number and methylation level of an intronic CpG site (cg17197774) located downstream of the CpG island, b) cg17197774 methylation is a likely prognostic marker in glioma, c) Among 22 cancer types, melanoma exhibits the highest *mda-9* level, and lowest level of methylation at cg17197774, d) cg17197774 hypomethylation is also associated with histone modifications (at the *mda-9* locus) indicative of more active transcription, e) Using Gene Set Enrichment Analysis (GSEA), and the **VIGOR (Virtual Gene Over-expression or Repression)** analytical scheme, we were able to predict *mda-9*'s association with extracellular matrix organization (e.g., MMPs, collagen, integrins), IGFBP2 and NF- κ B signaling pathways, phospholipid metabolism, cytokines (e.g., interleukins), CTLA-4, and components of complement cascade pathways. Indeed, previous

*Correspondence: Dr. Paul B. Fisher, Professor and Chairman, Department of Human and Molecular Genetics, Director, VCU Institute of Molecular Medicine, Virginia Commonwealth University, School of Medicine, 1101 East Marshall Street, Sanger Hall Building, Room 11-015, Richmond, VA 23298-0033, Tel: 804-628-3506 (Office), Fax: 804-827-1124, pbfisher@vcu.edu.

publications have shown that many of the aforementioned genes and pathways are associated with *mda-9*'s functionality.

Keywords

MDA-9; Syntenin; SDCBP; TCGA; GEO; ENCODE; epigenetics; glioma; melanoma; VIGOR

1. Background on *mda-9/Syntenin*

mda-9/Syntenin (*Melanoma differentiation-associated gene-9*), also known as *SDCBP* (*Syndecan binding protein*), is a human gene transcribed in the plus strand of the 8q12.1 region. The gene consists of 9 exons (including the UTR regions), and has 5 transcript variants coding for 3 protein isoforms. Isoforms 1, 2 and 3, code for proteins with 298, 292 and 297 amino acids, respectively. As indicated in the UCSC genome Browser (genome.ucsc.edu), there is a possible 10th exon located 3' of the 5' UTR exon, which may result in an even longer isoform (318 aa). The prominent features of the protein (isoform 1 is 33 KDa) are its two PDZ domains (PDZ1 and PDZ2). PDZ domains, due to their repertoire of possible interactions (C-terminal peptide recognition, interactions with internal peptide ligands, PDZ–PDZ interactions, PDZ–phospholipid interactions) (Ivarsson, 2012) are known to mediate a wide array of signaling pathways and cellular functions. Among the molecules the protein interacts with are, phosphoinositides, IL-5 receptor α (IL5RA), proTGF α , syndecan, eIF5A, Schwannomin and CD6, [see (Beekman and Coffey, 2008; Sarkar *et al.*, 2008) for reviews]. A short list of cellular processes involving MDA-9 are: axon outgrowth (Tomoda *et al.*, 2004), chemotaxis (Sala-Valdes *et al.*, 2012), HIV-entry (Gordon-Alonso *et al.*, 2012), development of neuronal membrane architecture (Hirbec *et al.*, 2005), protein-cell surface localization (Fernandez-Larrea *et al.*, 1999), cell adhesion (Zimmermann *et al.*, 2001) and pro-metastatic and pro-angiogenic activities (Boukerche *et al.*, 2005; Das *et al.*, 2013).

mda-9, as numerous studies have already shown, also plays an important role in cancer progression, particularly during the invasion/metastasis stage [reviewed in (Kegelman *et al.*, 2015)]. The earliest report describing this gene in our laboratory dates back to 1996 (Lin *et al.*, 1998) when it was first cloned using a subtraction hybridization approach developed in our laboratory (Jiang *et al.*, 1995) using melanoma cells induced to terminally differentiate by treatment with interferon β (IFN- β) and the anti-leukemic compound mezerein (Fisher *et al.*, 1985). Subsequent studies have demonstrated that MDA-9 is a positive regulator of metastasis in melanoma partially attributed to its interaction with c-Src, which eventually leads to the activation of the transcription factor NF- κ B (Boukerche *et al.*, 2010; Boukerche *et al.*, 2008; Das *et al.*, 2012). These changes induce an increase in the transcription of matrix metalloproteinases (MMPs), necessary for the degradation of extracellular matrix during invasion (Das *et al.*, 2012). MDA-9's interaction with c-Src can also lead to transcriptional activation of insulin growth factor binding protein 2 (IGFBP2), which can promote angiogenesis in melanoma (Das *et al.*, 2013). As our group recently reported, *mda-9*'s invasion-promoting property is also evident in glioma, in which its over-expression led to activation of c-Src, p38 MAPK and NF- κ B, and eventually the elevated expression of

MMP2 and secretion of interleukin-8 (IL-8) (Kegelman *et al.*, 2014). *mda-9*-driven proliferation via p38 MAPK activation and MMP2 upregulation was similarly demonstrated in hepatocellular carcinoma (Liu *et al.*, 2014) and small cell lung cancer (Kim *et al.*, 2014) cells. *mda-9* is also over-expressed in metastatic (Qian *et al.*, 2013), as well as ER-negative (Koo *et al.*, 2002) breast cancer. *mda-9*'s regulation of cell migration was also demonstrated in colorectal cancer (Lee *et al.*, 2011), a cancer type in which poor clinical outcome is associated with elevated expression of this gene (Hao *et al.*, 2010). Another recent finding from our group is *mda-9*'s regulation of urothelial cell proliferation through its modulation of EGFR signaling (Dasgupta *et al.*, 2013).

Despite the wealth of *mda-9*-related knowledge listed above, there is still much to learn regarding *mda-9* and its involvement in cancer progression. One area that has not yet been investigated is how genetic and epigenetic factors contribute to its elevated expression during cancer progression. For any gene, two factors that can lead to elevated expression would be: a gain or amplification of copy number, and a reduced methylation at the appropriate CpG site on its promoter region. These are, by no means the only factors which can influence the transcription of a gene. Activation of transcription factors (such as NF- κ B, shown in previous experiments) and the upstream signaling pathways are also necessary for the gene's transcriptional activation. The activities of DNA methyltransferases (DNMTs), which initially methylate the CpG sites, may also factor in the gene's dysregulation. The elucidation of genetic and epigenetic factors associated with *mda-9*'s transformation into a cancer- or metastasis-promoting state may now be conducted *in-silico*. This is due to the availability of comprehensive and publicly available genomic datasets (such as genome-wide expression, methylation and copy number data) for various types of cancer. Integrated analyses of these datasets may help define the effects of *mda-9* copy number and CpG methylation on its transcription. Analysis of expression datasets may also lead to the identification of various genes and molecular pathways associated with *mda-9* dysregulation, which can then be validated experimentally.

2. Publicly available cancer genomic datasets

In the past decade, cancer-related data generated using various genome-wide molecular profiling tools have been made publicly available. Currently, there are tens of thousands of cancer genomic datasets available to the public for analysis. Two very extensive repositories are NCBI's Gene Expression Omnibus (GEO) (<http://www.ncbi.nlm.nih.gov/geo/>) and EMBL's Array Express (www.ebi.ac.uk/arrayexpress/). A huge proportion of these datasets are genome-wide expression profiles (but also include genome-wide methylation and copy number data) of cancer tissue samples, cell lines and mouse models. However, what is perhaps the most comprehensive and organized cancer genomic repository is The Cancer Genome Atlas (TCGA) (<https://tcga-data.nci.nih.gov/tcga/>). Launched in 2005 (Kaiser, 2005), TCGA has now examined the genome-wide expression (mRNA, miRNA, exon, limited protein), copy number variations, methylation status and mutations in more than 20 adult cancer types (a total of more than 6000 tissue samples). The primary advantages of TCGA datasets are: a) each patient sample is accompanied by very comprehensive clinico-pathological data (e.g., follow up survival records, TNM staging, treatment records), b) a huge portion of the samples have integrated molecular profiles (i.e., same sample being

profiled for expression, copy number, sequence, methylation), c) many of the tumor samples have matched normals and d) the data are generated using the latest and widely considered standards in molecular profiling technology. These include Illumina HiSeq 2000 for expression profiling, Illumina Infinium 450k BeadChip for methylation analysis, Affymetrix SNP 6 array for copy number analysis, and various Next Gen Sequencing platforms for mutational profiling. Also available to the public is the Human Protein Atlas Database (<http://www.proteinatlas.org/>) (Uhlen *et al.*, 2010). This database (although not as quantitative as the mRNA levels reported using RNASeq or array chips) is a genome-wide immunohistochemical staining analyses portal of a large number of human tissues, cancers and cell lines. Another data repository which is very useful for cancer research is ENCODE (Encyclopedia of DNA Elements) (<https://genome.ucsc.edu/ENCODE/>) (2004), a project which aims to build a comprehensive list of functional elements in the human genome, and further our understanding of gene regulations. In ENCODE, numerous experimental tools, such as ChIP Seq technology, were employed to examine how proteins (e.g., transcription factors, histones) recognize and bind to genomic sequences such as promoter regions.

3. Specific datasets examined for *mda-9* analysis and the analytical approaches employed

3.1. Publicly available genomic datasets

This review is a foray into *mda-9/Syntenin*'s genetic and epigenetic regulation through a careful and rigorous examination of various public genomic datasets (See Figure 1 for the scheme). Most datasets originated from TCGA, and some were downloaded from NCBI-GEO (Described in Table 1). Specifically, we analyzed the Illumina HiSeq 2000 (expression), Illumina Infinium 450k BeadChip (CpG Methylation), and Affymetrix SNP 6-derived GISTIC2 copy number datasets for TCGA Glioma (combined Glioblastoma Multiforme and Lower Grade Glioma; GBM and LGG, respectively), Skin Cutaneous Melanoma (SKCM), Liver Hepatocellular Cancer (LIHC), Prostate Adenocarcinoma (PRAD), Colon Adenocarcinoma (COAD) and Kidney Renal Papillary Cell Carcinoma (KIRP). In addition, we also examined the TCGA Pan Cancer (PANCAN) dataset, which is the merging of all the 22 unique TCGA expression datasets (Chang *et al.*, 2013). At level 3, The TCGA datasets coming from TCGA Genome Characterization (and Data Coordination) Centers consist of single text files representing genome-wide expression, copy number or methylation values for individual samples (see: <https://tcga-data.nci.nih.gov/tcga/tcgaDataType.jsp>). These datasets were then merged, further processed (e.g., normalization for expression data), and made available for download by the UCSC Cancer Genomics Brower team (<https://genome-cancer.ucsc.edu/>) (Goldman *et al.*, 2013; Zhu *et al.*, 2009). For the purposes of this report, the processed datasets along with relevant clinical information were downloaded for further analyses. Also analyzed were tissue expression datasets from GEO: GSE4290 (glioma) (Sun *et al.*, 2006), GSE3189 (melanoma) (Talantov *et al.*, 2005) and GDS2545 dataset (prostate cancer) (Chandran *et al.*, 2007). These datasets were generated using Affymetrix HG U133 Plus 2, U133A, and U95A arrays, respectively. The MDA-9 protein expression levels were also assessed in the Human Protein Atlas Database (<http://www.proteinatlas.org/>). Immuno-histochemical images and other data were downloaded directly from the website. CpG methylation data (Illumina 450K) for cell lines

included in the ENCODE project, generated from Hudson Alpha Inst. (R. M. Myers lab; Huntsville, AL) were downloaded through the UCSC Genome Browser. The UCSC Genome Browser was also used to access information from the ENCODE project, such as the ChIP Seq-derived quantification of DNA-bound modified histones (B.E. Bernstein lab, Broad Inst., Cambridge, MA).

3.2. Analytical tools

A number of genomic and statistical tools were employed for this study. Initial manipulations of the downloaded datasets were performed using Gene-E (Broad Institute, Cambridge, MA). Statistical analyses were performed using JMP Pro 10 software (SAS, Cary, NC), Gene-E and Gene Set Enrichment Analysis (Subramanian *et al.*, 2005) (GSEA; Broad Inst., Cambridge, MA). Also used during the later stages of analyses were Cytoscape (Shannon *et al.*, 2003) and the Cytoscape Enrichment Map Plug-in (Merico *et al.*, 2010). Crucial to the analyses was information gathered from the UCSC Genome Browser, UCSC Cancer Genomics Browser, Human Protein Atlas, MsigDB (www.broadinstitute.org/msigdb) (Liberzon *et al.*, 2011), Reactome (www.reactome.org) (Joshi-Tope *et al.*, 2005), KEGG pathway database (<http://www.genome.jp/kegg/>) and Biocarta (www.biocarta.com). Accessed through the UCSC Genome Browser were results from ENCODE Chromatin State Segmentation using Hidden Markov Modeling (Ernst and Kellis, 2010; Ernst *et al.*, 2011). Necessary gene and probeset annotations were downloaded from Human Genome Organisation Gene Nomenclature Committee (HGNC) (<http://www.genenames.org/>) and NCBI-GEO websites.

4. The patterns of *mda-9* expression during cancer progression and their clinical implications in different cancer types

4.1. Glioma

In our recent report, we demonstrated how *mda-9* is up-regulated as tumor grade progresses. The analysis of expression datasets might provide a more quantitative assessment about *mda-9*'s expression pattern in glioma. The first dataset we examined was GSE4290 (Sun *et al.*, 2006), generated using the Affymetrix platform HG U133 Plus 2. As shown in Figure 2A, *mda-9* expression levels increase as sample progresses from lower towards higher tumor grades, as compared to normal tissues (ANOVA, $p < 0.001$). The same trend can be observed upon analysis of the combined TCGA GBM and LGG datasets, which were generated using an entirely different platform (i.e., Illumina HiSeq 2000) (ANOVA, $p < 0.001$). However, unlike the GSE4290, the TCGA glioma dataset did not include normal tissue samples. As we have previously pointed out, the expression level of *mda-9* can also be a marker of prognosis in glioma (Kegelman *et al.*, 2014). In the Kaplan-Meier graph shown in Figure 2B, the TCGA glioma samples were divided into 2 subgroups according to *mda-9* expression levels: below median (L), and above median (H). Results show that the H subgroup had much worse overall survival rate compared to the L subgroup. Unfortunately, the GSE4290 was not annotated with follow up records, thus could not be subjected to survival analysis.

4.2. Melanoma, prostate cancer, liver hepatocellular carcinoma and kidney renal papillary carcinoma

The upregulation (relative to normals) of *mda-9* is also evident in other tumor types, as shown in Figure 3. The distribution of *mda-9* expression levels in tumor and normal tissue groups for the four cancer types are shown in order of increasing z score (relative expression), calculated as $(I_T - \text{Average } I_{\text{norm}}) / \text{standard dev}_{\text{norm}}$, where *n* refers to every sample (including tumors), while *norm* refers to normal samples only. Student t-test indicates that *mda-9* expression levels in tumors are significantly higher compared to normals. The *mda-9* levels for Prostate Cancer, Liver Hepatocellular Carcinoma and Kidney Renal Papillary Carcinoma were all extracted from the TCGA RNA Seq datasets. The TCGA dataset for melanoma (SKCM) was not used for this particular part of the analysis because the dataset did not include normal samples. Most of the TCGA SKCM samples were actually classified as metastasis (which will be discussed as a component of the TCGA Pan Cancer dataset). Instead, the dataset GSE3189 (Talantov *et al.*, 2005) was analyzed to differentiate the expression levels of malignant melanoma (n = 45) to a limited number of normal skin (n = 7) samples. The original GSE3189 dataset includes nevi samples, which were not included in this analysis.

4.3. *mda-9* expression in metastasis samples

The data described above were those of *mda-9* transcription levels in primary tumors. Given that recent experimental results point to *mda-9*'s role in invasion, it would be of interest to examine its expression pattern in a cohort, which includes metastasis samples. Shown in Figure 4 is data taken from the GDS2545 dataset (Chandran *et al.*, 2007) generated using Affymetrix U95A array. As the graph indicates, there *mda-9* level is significantly higher in metastasis samples compared to primary tumors and normal samples.

5. *mda-9* expression correlates with both gene copy number and the methylation status of cg17197774

5.1. *mda-9* RNA expression generated Illumina HiSeq data

The total RNA for each TCGA tissue sample was quantitated using the Illumina HiSeq 2000 RNA Sequencing platform (performed at a TCGA Genome Characterization Center). The matrix form dataset (which is the merging of individual level 3 processed dataset) was downloaded from the UCSC Cancer Genomics Browser and the samples annotated with the accompanying clinical data. Each dataset included the expression levels for 20,501 genes. From these datasets, the transcript levels of *mda-9* were derived and further analyzed.

5.2. *mda-9* copy number as determined by Affymetrix SNP array

The genome-wide copy number data for TCGA samples were also generated in TCGA Genome Characterization Centers, using Affymetrix Genome-wide Human SNP 6.0 Array, consisting of probes for more than 906,600 SNPs, in addition to more than 946,000 probes for the detection of copy number variations. The experimental protocol described in the user manual can be downloaded from the company website (www.affymetrix.com). The TCGA FIREHOSE pipeline (Broad Institute, Cambridge, MA) employed the GISTIC2 algorithm

(Mermel *et al.*, 2011) to generate copy number estimates (CN) for the genes mapped in the genome. The GISTIC2 estimate for *mda-9* was then extracted from the dataset, and converted to copy number estimate, calculated using the formula: $CN = 2^{(GISTIC2 + 1)}$

5.3. *mda-9* CpG sites interrogated in Illumina 450K array

The genome-wide CpG methylation data for TCGA samples were generated using the Illumina Infinium Human Methylation 450K array, which is designed to interrogate more than 480,000 CpG sites in the entire human genome (Sandoval *et al.*, 2011). Bead Studio software was then employed to generate beta values, which range from 0 (fully unmethylated) to 1 (fully methylated). For this report, we used the UCSC offset value in which -0.5 was subtracted from the original scale, resulting in beta values ranging from -0.5 (fully unmethylated) to 0 (50% methylated) to 0.5 (fully methylated). A total of 13 *mda-9* locus CpG sites covered in the Illumina 450K array registered beta values in the TCGA datasets (see Table 2). Ten of the probes are part of the CpG island group located in the promoter region, with the first five located within the transcription start sites (TSS1500, TSS200), and the next five as part of the 5' UTR exon (See Figure 5). The next 5 CpG sites are located in the intervening introns and the last one within the 3' UTR exon. The distributions of the beta values, corresponding to each of the 13 CpG sites for each of the 6 TCGA cancer datasets, are shown in Figure 6. Except for two (and cg10129404), the CpG sites were mostly unmethylated across all the cancer datasets. This suggests that only the variation in the beta values of the two CpG sites translate to variation in *mda-9* expression levels. The CpG site is exactly 1105 bases from the 3' edge of the CpG island (Gardiner-Garden and Frommer, 1987), while cg10129404 is part of the 3' UTR, making it less likely for the latter to be a factor in transcription of *mda-9* (this will be clarified in the next subsection). For glioma, methylation at the CpG sites decreases as the tumor grade progresses. For liver cancer, colon adenocarcinoma and kidney papillary carcinoma, methylation at these sites decreases during the transformation from solid normals to primary tumor. There are only a few normal samples in the SKCM dataset, but it is very clear that primary and metastatic SKCM samples had the lowest levels of methylation (compared to the other 5 datasets). There was only a minimal change in methylation in primary prostate cancer relative to solid normals.

5.4. Which amongst the two CpG sites influence *mda-9* expression?

As mentioned above, it is unlikely that the 3' UTR CpG site cg10129404 influences *mda-9* transcript levels. A simple analysis was conducted by plotting *mda-9* expression vs the beta value for or cg10129404 for two select TCGA datasets (glioma and KIRP) (Figure 7). In glioma, it is clear that the methylation at may influence *mda-9* expression ($R^2 = 0.38$; linear regression). In contrast, cg10129404 appears to have a negligible effect on *mda-9* expression ($R^2 = 0.051$). Similar analysis was conducted for the TCGA KIRP dataset. Linear regression analyses indicate that the R^2 value for the plot of *mda-9* expression vs. methylation is 0.25, while that of *mda-9* expression vs. cg10129404 is almost zero. Overall, these analyses indicate that cg10129404's influence on *mda-9* expression is unlikely.

5.5. Glioma

Having established that *mda-9* expression correlates with methylation at , the next step was to analyze the dual contribution of both methylation and copy number in *mda-9* expression for each of the 6 datasets included in this study. The relationships between *mda-9* expression, copy number and the methylation status of are illustrated in Figure 8. For glioma (whose *mda-9* copy number is mostly neutral), we can see the effect of both copy number and methylation status. Copy number is a likely factor ($R = 0.26$; expression vs. copy number). However, it is also clear that among the samples with only two copies of *mda-9*, those with low degree of methylation at the cg17197774 tend to have a higher *mda-9* expression level ($R = -0.61$; expression vs. cg17197774 beta value). By itself, it appears that cg17197774 methylation status may be a reliable marker of survival in glioma (Figure 9).

5.6. Melanoma, prostate cancer, liver hepatocellular carcinoma and kidney renal papillary carcinoma

On average, SKCM samples have the lowest beta values for cg17197774, at -0.295 (Table 3). This may also explain why among the six TCGA cancer datasets, SKCM tumors have the highest expression levels for *mda-9* (at more than 20,000 units; inv log 2 scale). Nonetheless, the influence of both factors is evident in Figure 8C (correlation coefficients of -0.35 and 0.58 for expression vs. methylation and expression vs. copy number, respectively). The tumor sample with the highest overall *mda-9* expression level (at more than 190,000 units) has an *mda-9* copy number estimate of 25 and cg17197774 beta value of -0.45 . For the TCGA COAD dataset, both copy number and cg17197774 methylation appear to factor in *mda-9* expression, with the former ($R = 0.47$) having greater influence than the latter (-0.22). A great majority of KIRP samples have a neutral copy number (CN = 2) at the *mda-9* locus. Not surprisingly, the elevated *mda-9* expression is primarily due to hypomethylation at cg17197774 (*mda-9* expression vs. cg17197774 methylation correlation coefficient = -0.5). In contrast to KIRP, the mode of *mda-9* dysregulation in prostate cancer samples is primarily through copy number gain ($R = 0.67$), with cg17197774 methylation apparently lacking any effect on the gene's expression level. Among LIHC primary tumors, it is clear that both copy number ($R = 0.55$) and cg17197774 methylation ($R = -0.45$) are factors influencing *mda-9* RNA levels.

5.7. Combination of the six cancer datasets

For a unified view on the effects of both copy number and cg17197774 methylation on *mda-9* expression, we used the PANCAN-normalized expression value for *mda-9*. As explained earlier, the PANCAN dataset is the merging of all the 22 TCGA RNASeq datasets (Chang *et al.*, 2013). According to the information provided by the UCSC Cancer Genomics Browser (from which the dataset was downloaded), the original level 3 RNASeq V2 datasets were downloaded from TCGA, log (base 2) transformed, then mean-normalized across all the cohorts. The normalization across all the cohorts provides more reliable relative values for gene expression. Figures 10 A–B show the same Cartesian plot (i.e., PANCAN-normalized *mda-9* expression value vs. methylation at cg17197774) with varying information for each data point. As these figures indicate, there is a clear inverse exponential relationship between the two variables. As discussed previously, the highest *mda-9*

expression levels were those of SKCM samples, owing to the low beta values for cg17197774. The effect of copy number on *mda-9* expression is illustrated in Figure 10B, wherein solid red circles (those samples whose *mda-9* copy number is 6 or higher) mostly appear in the upper edges of the graph. This indicates that copy number can elevate *mda-9* expression levels irrespective of the methylation status of cg17197774. Overall, the correlation coefficients (Multivariate REML statistics) for expression vs. methylation and expression vs. copy number are -0.61 and 0.30 , respectively. This shows that the methylation status of cg17197774 has greater influence (compared to *mda-9* copy number) towards the gene's expression level. Figure 10C shows a superimposed exponential regression model (JMP Pro 10) relating *mda-9* expression and cg17197774. Lastly, Figure 10D illustrates the relationship between *mda-9* expression and copy number (essentially a merging of Figure 8). At this point, we already know that tumor samples with neutral copy number for *mda-9* can have an elevated expression of the gene if it is hypomethylated at cg17197774.

6. *mda-9* is highly expressed in melanoma

6.1. *mda-9* RNA expression across all cancer types (PANCAN dataset)

The results illustrated above point to *mda-9* being most highly expressed in melanoma (among the 6 cancer types analyzed). We then proceeded to examine *mda-9* expression in all of the TCGA cohorts through analysis of the TCGA PANCAN dataset. Covering 22 cancer types, the dataset includes 6040 samples (4982 primary tumors, 271 metastatic, 27 recurrent tumors, 173 peripheral blood, 587 solid tissue normals). First, the 6040 samples were grouped according to TCGA-defined sample types. On average, the highest *mda-9* levels were those of metastasis samples, which were largely the melanoma (SKCM) samples (Figure 11A). When *mda-9* expression was divided according to cohort (without the normals), it is clear that *mda-9* is most elevated in melanoma samples (Figure 11B).

6.2. MDA-9 protein expression across all cancer types (The Human Protein Atlas)

TCGA also has datasets for protein expression (using Reverse Phase Protein Array). However, it only covers a few hundred select proteins (the most widely studied cancer-related genes) and does not include MDA-9. However, a very comprehensive source of genome-wide protein expression is the Human Protein Atlas (<http://www.proteinatlas.org/>), where proteins were immunohistochemically analyzed in different cancer types (note: Human Protein Atlas samples are independent of the TCGA cohorts). For MDA-9, two different antibodies were used: HPA023840 from Sigma-Aldrich and CAB012245 from Abcam. Consistent with results from the PANCAN analysis, MDA-9 according to Human Protein Atlas, is most highly expressed in melanoma (See Figure 12). As shown in the figure (bottom right), the HPA023840 signal was much stronger in melanoma compared to glioma. The staining results were evaluated by a trained pathologist, who assigned intensity (negative, weak, moderate or strong), fraction of stained cells (rare, <25%, 25–75% or >75%), as well as subcellular localization (nuclear and/or cytoplasmic/membranous).

7. Hypomethylation at cg17197774 is associated with histone modifications indicative of higher transcriptional activity

7.1. Information from the ENCODE project

The ENCODE (Encyclopedia Of DNA Elements) project was launched several years ago (through the National Human Genome Research Institute), with the primary aim of identifying all functional elements in the human genome sequence (2011). It is a consortium involving a number of laboratories mostly in the US. The investigators examined regulatory elements using various tools such as DNA methylation assays, DNA hypersensitivity assays, RNA Sequencing and chromatin immunoprecipitation (ChIP) of proteins interacting with DNA (e.g., histones, transcription factors), followed by sequencing (ChIP Seq). In a typical ChIP Seq experiment, proteins are cross-linked to cancer cell line DNA, followed by shearing of the DNA strands. An antibody to the protein of interest (e.g., TF, histones), attached to a bead will then be used to immunoprecipitate the target protein, which is linked to the DNA sequences it recognized. The protein is then unlinked from the DNA, which is then sequenced (using Next Gen sequencing) and mapped to the genome. The binding of a protein to a DNA (or RNA) sequence is directly proportional to the number of readouts for that particular sequence. In essence, a ChIP Seq experiment is supposed to: a) identify the sequence the protein binds to, and b) measure the extent of this binding. Results from numerous ENCODE projects are now publicly available through the UCSC Genome Browser, for either on site analysis, or data download for further analysis. There are various types of analyses, which can be done on these data.

7.2. How CpG methylation is associated with differential binding of modified histones

Since we have already established (from TCGA analysis) that the methylation at the CpG site cg17197774 correlates with decreased *mda-9* expression, it would be of interest to investigate how this CpG site may also affect the constitution of chromatin covering the *mda-9* locus. Take for example, the cell lines H1-hESC (human embryonic stem cell) and HUVEC (human umbilical vein endothelial cell). Both cell lines have available genome-wide CpG methylation data (Illumina 450K methylation array), as well as ChIP Seq data for various modified histones (ENCODE data available for download and analysis through UCSC Genome Browser). As indicated in Figure 13A, the MDA-9 promoter region of HUVEC (hypomethylated at cg17197774; beta value = -0.353) exhibits greater affinity (darker bands) for Histone H3 mono-, di-, or tri-methylated at K4 (H3K4me1, H3K4me2, H3K4me3), and acetylated at K9 (H3K9ac), compared to that of H1-hESC (highly methylated at cg17197774; beta value = 0.230). Those histone modifications, just like hypomethylation at cg17197774, are associated with more active transcription (Campbell and Turner, 2013; Ernst *et al.*, 2011; Ghirlando *et al.*, 2012). Four other markers (H4K20me1 which is associated with transcriptional activity; H3K36me3 which is associated with RNAPII elongation; H3K27me3 which is associated with promoter silencing; and the transcriptional repressor CTCF) were not significantly different between the two cell lines. Ernst, Kellis and colleagues devised an analytical scheme, which summarizes the collective transcriptional influences of the aforementioned histone modifications (and CTCF) in a given segment of the chromosome (Ernst and Kellis, 2010; Ernst *et al.*, 2011). A Hidden

Markov Model was used to predict the state of a chromosomal segment with respect to the activity of the promoter, strength of enhancer sequences, and openness for transcription, based on the ChIP Seq data specific for these markers. Of the nine cell lines examined using this approach (thus far), only six have corresponding Illumina 450K methylation data. In addition to H1-hESC and HUVEC, the other cell lines were: the hepatocellular carcinoma cell line HepG2, the microvascular endothelial cell line HMEC, the lymphoblastoid cell line GM12878, and the chronic myelogenous leukemia cell line K562. As shown in Figure 13B, H1-hESC, HepG2 and HMEC, which are all highly methylated at cg17197774, are predicted to exhibit less active MDA-9 promoter and transcriptional activity (interpretations for segmental colors are listed in Figure 13C). In contrast, MDA-9 expression is expected to be higher for cell lines GM12878, K562 and HUVEC, which are all weakly methylated at cg17197774.

8. Predicting genes and pathways associated with *mda-9* dysregulation in glioma

8.1. Comparing *mda-9* High and *mda-9* Low subtypes of glioma

Genome-wide expression datasets may also be useful in predicting genes or pathways associated with *mda-9* dysregulation. One approach (See Figure 14) is to start with the isolation of two subsets (out of the complete cohort of tumor samples) at the opposite tail ends of distribution according to *mda-9* expression: a) *mda-9-high*, i.e., those whose *mda-9* expression levels are at least 1 standard deviation higher than the average, and b) *mda-9-low*, i.e., those cases whose *mda-9* expression levels are at least 1 standard deviation lower than the average. The comparative genome-wide expression of these two subsets will then be examined to identify the genes, functionalities and molecular pathways, which are most likely associated with *mda-9* upregulation or *mda-9* downregulation. This comparison between the *mda-9-high* and *mda-9-low* subtypes is akin to a gene being ectopically over-expressed or repressed (with RNAi) in a cancer cell line [an experimental approach employed in one of our recent publications (Sokhi *et al.*, 2013)] with the objective of examining the molecular changes, which occur after the process. For that reason, this approach is being referred to as “**Virtual Gene Over-Expression or Repression**” or **VIGOR**. Genes will then be ranked according to a comparative statistic (either signal-to-noise ratio or fold-change) between the two groups. The **VIGOR** approach was applied to the two glioma datasets (TCGA and GSE4290) and the results were highly concordant. As shown in Figure 15, the resulting signal-to-noise ratios (for all genes) in TCGA glioma dataset directly correlate with those of the GSE4290 dataset. Another statistical tool employed is the Gene Set Enrichment Analysis (GSEA) available through the Broad Institute (www.broadinstitute.org/gsea/) (Subramanian *et al.*, 2005). GSEA analysis starts with the recognition that genes are associated with particular groups (or gene sets), such as pathways defined in Reactome, Biocarta, or KEGG databases, as well as molecular functions, biological processes and subcellular locations according to Gene Ontology. GSEA then examines the collective trends of how genes belonging to a particular gene set behave through the calculation of an Enrichment Score (ES). A high concentration of highly ranked genes (rank = 1 for the gene with the highest fold-change; *mda-9* high/*mda-9* low) belonging to a particular gene set translates to an ES value close to 1, which may be interpreted as that

gene set being likely associated with *mda-9* upregulation. An ES value close to -1 , may be interpreted as that gene set being likely associated with *mda-9* downregulation.

8.2. Genes that are most highly dysregulated in *mda-9*-high glioma

Table 4 lists the genes most likely up-regulated when *mda-9* is over-expressed and those at the bottom of the list are the genes at the opposite end of the spectrum (i.e., down-regulated when *mda-9* is elevated). When the top 300 up-regulated genes were overlapped against the MSigDB database (<http://www.broadinstitute.org/gsea/msigdb>), it was revealed that the list includes 29 cytokines and growth factors, 34 transcription factors, 29 transcription factors which are also homeodomain proteins, 23 cell differentiation markers, 9 oncogenes and 1 protein kinase. On the other hand, when the top 300 down-regulated genes were overlapped against the same database, we found that only 5 were classified as cytokines and growth factors, 16 as transcription factors, 3 as homeodomain transcription factors, 2 cell differentiation markers, 1 oncogene (*RANBP17*) and 8 as protein kinases. These results provided us with a rough clue as to the type of genes directly associated with *mda-9* expression: that they are more likely involved in cell proliferation, or more precisely to the advancement of glioma carcinogenesis. Among the top 100 upregulated genes are *LTF* (lactotransferrin), *CHI3L1/L2* (cartilage glycoproteins), *PLA2G2A* (phospholipase A2, group IIA), *POSTN* (periostin, osteoblast specific factor), *ABCC3* (ATP-binding cassette, sub-family C (CFTR/MRP), *SAA1* (serum amyloid A1), various *ILs* (interleukins) and *ILRs* (interleukin receptors), *IBSP* (integrin-binding sialoprotein), *MMPs* (matrix metalloproteinases), *TIMPs* (TIMP metalloproteinase inhibitors), *COLs* (collagens), *HOXs* (homeobox genes) and the scavenger receptor *CD163*. None of the genes in the top list are in the 8q arm, which means that the **VIGOR** approach minimized the copy number effect in gene selection. Among the most down-regulated genes are *GRIN1* (glutamate receptor, ionotropic, N-methyl D-aspartate 1), *CUX2* (cut-like homeobox 2), *INA* (internexin neuronal intermediate filament protein, alpha), *SVOP* (SV2 related protein homolog) and *CHGA* (chromogranin A (parathyroid secretory protein 1)). The *MMPs*, *TIMPs*, *COLs* and *IBSP* are all components of extracellular matrix processes crucial to cell invasion and metastasis, thus, these genes' connections to *mda-9* are quite clear (to be discussed further in subsequent sections). *ILs* and *ILRs* are components of various immuno-related pathways, and are known to play important roles in glioma progression (Yeung *et al.*, 2013). *SAA1* codes for an acute phase protein and is a potential serum-based cancer biomarker. Recently, it was found to be a transcriptional target of the metastasis-inducing S1004 (also highly up-regulated in *mda-9* high glioma) (Hansen *et al.*, 2014). HOX proteins are oftentimes transcription factors and many of them have been linked to cancer progression (Shah and Sukumar, 2010). *PLA2G2A*, which may induce astrocytoma proliferation (Hernandez *et al.*, 2010), is an enzyme which hydrolyzes phosphatidylinositol during lipid metabolism and membrane assembly (Singer *et al.*, 2002). MDA-9 may be closely associated with this particular enzyme, given that MDA-9 has been shown to bind to phosphatidylinositol 4,5-bisphosphate (and may assist in transporting proteins to membranes) (Wawrzyniak *et al.*, 2013). *ABCC3* belongs to the ABC subfamily of multi-drug resistance proteins, known to be a major detriment to glioma therapy (Decleves *et al.*, 2006). The glycoprotein *CHI3L1* has previously been linked to glioma progression through its upregulation of VEGF (and angiogenesis) (Francescone *et al.*, 2011). The stromal protein *POSTN* has also been linked

to glioma invasion and its poor clinical outcome (Mikheev *et al.*, 2014; Tian *et al.*, 2014; Wang *et al.*, 2013). Its possible association with *mda-9* is yet to be investigated. How the iron-binding and multifunctional protein LTF (Gibbons *et al.*, 2011; Ward *et al.*, 2005) may be positively associated with glioma progression or *mda-9* functionality is not yet clear. Many of the very highly down-regulated genes in *mda-9*-high glioma appear to have neuronal functions. *GRIN1*, which codes for a critical part of the NMDA (N-methyl-D-aspartate) receptor (Gibb, 2004), is among the most highly down-regulated genes in *mda-9*-high glioma. At this point it is not exactly clear how the down-regulation of this gene may be connected to glioma pathogenesis or *mda-9* expression. *CUX2* is a neuronal transcription factor crucial to regulation of dendritic branching, as well as synapses of the upper cortical neurons (Cubelos *et al.*, 2010). It is also not clear how this gene may be involved in glioma progression. *CHGA* is the precursor to vasostatin, pancreastatin, and parastatin, which are negative modulators of the neuroendocrine system. Its role in pathogenesis of glioma has not been established, but was found to be inhibitory to angiogenesis (Loh *et al.*, 2012). *INA* or *alpha internexin* is a Type IV intermediate filament involved in neuronal morphogenesis (Lepinoux-Chambaud and Eyer, 2013). Its down-regulation among *mda-9*-high glioma is consistent with recent observations that relatively low expression of the gene correlates with the absence of 1p/19q co-deletion (which are established markers of good prognosis) (Ducray *et al.*, 2009).

8.3. Pathways and gene groups identified through GSEA

Instead of inspecting the likely *mda-9*-associated genes one at a time (like what was accomplished in the previous section), GSEA analysis was able to identify the groups of genes representing molecular pathways and related functionalities. The first 6 rows of Table 5 are the top sets of genes (i.e., the highest ES values) directly associated with *mda-9*-high glioma. The last row includes the few gene sets found to be likely associated with *mda-9* down-regulation. Some of the notable gene sets are discussed below:

8.3.1 Association with extracellular matrix (ECM), cell adhesion, and migration

8.3.1.1. The metalloproteases and associated proteins: Among the gene sets exhibiting the highest ES values are those related to the extracellular matrix. These include gene groups labeled as Reactome's *degradation of extracellular matrix* (ES= 0.79), GO Molecular Function's *metalloproteinase activity* (ES= 0.84) and GO Cellular Component's *extracellular matrix* (ES = 0.77) (Figure 16). All of these gene sets essentially refer to the same group of genes, which include the matrix *metalloproteinases* (*MMPs*) and its activator proteins (*TIMPs*). *MMPs* are zinc and calcium dependent proteases, which upon secretion can degrade the extracellular matrix (VanMeter *et al.*, 2001), an essential process for the cells to accomplish invasion and metastasis. A closer look at the GSEA plot for the Reactome gene set *degradation of extracellular matrix* indicates 12 genes (out of 28 members) considered as part of core enrichment (Figure 16A). These include *MMP9* and *TIMP1*, which were up-regulated (*mda-9*-high vs. *mda-9*-low) 44-fold and 27-fold, respectively, as well as *MMP13*, *MMP1*, *TIMP1*, *MMP11*, *MMP9* and *TPSAB1* (*tryptase alpha/beta 1*). Shown in Figure 16B are specific pathways involving the aforementioned genes (taken from www.reactome.org). These are mostly associated with degradation of the glycoproteins fibrillin (left), fibronectin (center) and decorin (right). Experimentally, our group has already demonstrated that over-

expression of *mda-9* in melanoma and glioma cell lines leads to activation of c-SRC and the transcription factor NF- κ B, followed by transcriptional elevation of MMP2 (Boukerche *et al.*, 2010; Boukerche *et al.*, 2008; Das *et al.*, 2012; Kegelman *et al.*, 2014).

8.3.1.2. Collagens: Various collagen genes also registered over-expression among *mda-9* high gliomas and this is reflected in the ES values for various gene sets such as Reactome's *collagen formation* (ES= 0.78). The following genes are highly-expressed in *mda-9* high tumors: Type I (*COL1A2*, *COL1A1*), Type III (*COL3A1*), Type IV (*COL4A1*, *COL4A2*), Type V (*COL5A1*, *COL5A2*), Type VI (*COL6A3*, *COL6A2*, *COL6A1*), Type VIII (*COL8A1*, *COL8A2*), Type XII (*COL12A1*), Type XIII (*COL13A1*), Type XIV (*COL14A1*) and Type XV (*COL15A1*). Other related genes up-regulated in *mda-9* high tumors are *PCOLCE* (*procollagen C-endopeptidase enhancer*) and the chaperone protein *SERPINH1* [*serpin peptidase inhibitor, clade H (heat shock protein 47), member 1, (collagen binding protein 1)*]. Collagen fibers provide the network of roads which cells travel on during migration (Egeblad *et al.*, 2010). There has never been any report implicating *mda-9* as a regulator of collagen expression (or the reverse case).

8.3.1.3. Integrins and focal adhesion genes: Other ECM-related gene sets highly enriched in *mda-9* High glioma belong to the following categories: Reactome's *integrin cell surface interactions* and KEGG's *focal adhesion*. Among the components of these sets are genes coding for various collagen molecules (*COL1A2*, *COL1A1*, *COL4A1*, *COL4A2*), integrins (*ITGA4*, *ITGA5*, *ITGB3*, *ITGB2*), *IBSP* (integrin-binding sialoprotein), the adhesive glycoprotein *THBS1* (thrombospondin 1), the cell surface glycoprotein *ICAM1* (*intercellular adhesion molecule 1*), laminins (*LAMB1*, *LAMC1*) and fibronectin (*FNI*). There were recent reports indicating that *mda-9* is indeed involved in the activation of focal adhesion kinase (FAK). MDA-9's role in the activation of FAK has been demonstrated in breast cancer cells (through the crosstalk between Protein Kinase C α and MDA-9) (Hwangbo *et al.*, 2010), dendritic cells (Cho *et al.*, 2013), glioma (through FAK-JNK and FAK-AKT signaling) (Zhong *et al.*, 2012), and small cell lung cancer cells (Kim *et al.*, 2014). MDA-9 was also shown to positively regulate the scaffold function of ILK (integrin-linked kinase) (Hwangbo *et al.*, 2011), part of KEGG's *focal adhesion* gene set, which is also up-regulated in *mda-9* High glioma.

8.3.1.4. Neurite outgrowth: The Reactome gene set centered around *NCAM* (*neural cell adhesion molecule*), *important for neurite outgrowth* (ES = 0.76), is also enriched among *mda-9* High glioma samples. NCAM, a member of the immunoglobulin (Ig) superfamily, is a mediator of the neurite outgrowth process (Hansen *et al.*, 2008). In glioma cells, it was shown that neurite outgrowth can be promoted through a cadherin-dependent adhesion (Cifarelli *et al.*, 2011). The computational prediction of MDA-9's association with the neurite outgrowth process was primarily driven by the over-expression of various collagen genes (e.g. *COL6A3*, *COL3A1*, *COL1A2*, *COL1A1*, *COL6A2*, *COL5A1*) among *mda-9* High glioma. Chen and colleagues have demonstrated that 3-D collagen scaffolds can enhance neurite outgrowth in neuron cancer stem cells (Chen *et al.*, 2012). Another possibly important gene in the neurite outgrowth process is *ST8SIA4* (*ST8 alpha-N-acetyl-neuraminide alpha-2, 8-sialyltransferase 4*), an enzyme necessary for synthesis of polysialic

acid, which modulates NCAM1's adhesive properties (Takahashi *et al.*, 2012). The gene is also highly up-regulated in *mda-9*High tumors. The connection between *mda-9* and neurite outgrowth stemmed from earlier reports of MDA-9 physically interacting with two molecules implicated in axon (a neurite) formation: NFASC (*neurofascin*) (Koroll *et al.*, 2001) and ULK1 (*unc-51 like autophagy activating kinase 1*) (Tomoda *et al.*, 2004).

8.3.2. Association with immuno signaling pathways

8.3.2.1. Interleukins: Among the most highly up-regulated genes in *mda-9*High gliomas are, *IL8* (*interleukin 8*) (19-fold) and *IL2RA* (*interleukin 2 receptor, alpha*) (21-fold). Indeed, GSEA analysis identifies a number of IL-related gene sets directly correlating with *mda-9* up-regulation. These include the following: Biocarta gene sets belonging to the *IL17 Pathway* (ES= 0.85), *Inflammation Pathway* (ES= 0.78), *IL2 pathway* (ES = 0.74), *IL7 pathway* (ES = 0.73) and *IL12 pathways* (ES = 0.70); GO Molecular Function gene sets involving *interleukin receptor activity* (0.83) (Figure 17A) and *interleukin binding* (0.79). The upregulation of many IL-related genes and pathways is likely related to the glioma cells trying to maintain their immunosuppressive state as much as possible (Parney, 2012). The involvement of *mda-9* in these processes has been verified in a number of reports. Geijsen and colleagues showed that MDA-9 binds to the cytoplasmic tail of IL5RA, which is necessary for IL5-mediated SOX4 activation (Geijsen *et al.*, 2001). Another interleukin (*IL16*), which contains 4 PDZ domains, in theory may interact with MDA-9, as predicted in Struc2Net (<http://groups.csail.mit.edu/cb/struct2net/webserver/>) (Singh *et al.*, 2010). A recent report from our laboratory indicated that MDA-9 positively regulates the transcription of IL8 (Kegelman *et al.*, 2014). An earlier report by Stier and colleagues suggested that tumor necrosis factor-alpha (TNF α) is capable of up-regulating the expression of both IL8 and *mda-9* in endothelial cells (Stier *et al.*, 2000).

8.3.2.2. Interferon: The direct correlation between IFNs and *mda-9* can be traced back to the observation that IFN- γ was found to induce *mda-9* expression in HO-1 human melanoma cells (Lin *et al.*, 1998). Many genes involved in the IFN- γ pathway were found to be up-regulated in *mda-9*High gliomas as shown in the GSEA plot for Reactome's *IFN- γ pathway* gene set (ES = 0.73). These include the following: *major histocompatibility complex class II genes* (*HLA-DQA1*, *HLA-DQA2*, *HLA-DRB1*, *HLA-DRB5*), *guanylate binding proteins* (*GBP5*, *GBP1*, *GBP2*), CD44, the nuclear antigen *SP100*, suppressor of cytokine signaling (*SOCS3*, *SOCS1*) and *interferon gamma* (*IFNG*). The top gene on the list (most highly up-regulated in *mda-9*High glioma) is *GBP5*, proven to be up-regulated by IFN- γ induction (Kitaya *et al.*, 2007).

8.3.2.3. CTLA4 pathway: Also noticeable is the apparent activation of the CTLA4 pathway among *mda-9*High glioma. CTLA4 (fold change ~3) is a receptor on the surface of Helper T Cells, which can down-regulate the immune system. Other *mda-9*High up-regulated genes belonging to Biocarta's *CTLA4 Pathway* gene set (ES = 0.86) are genes coding for proteins that form a complex with T cell receptors (*CD3D*, *CD3E*), lymphocyte-specific protein tyrosine kinase (*LCK*), proteins forming the MHC Class II complex (*HLA-D3A*, *HLA-DRB1*), IL2-inducible T-cell kinase (*ITK*), inducible T-cell co-stimulator (*ICOS*) and membrane-bound proteins necessary for T cell activation (*CD80*, *CD86*). The identification

of these genes is consistent with the knowledge of the presence of infiltrating cells in the glioma environment (Parney, 2012). Whether or not *mda-9* participates in the CTLA4 pathway requires experimental confirmation.

8.3.2.4. Complement cascade: The complement cascade is another pathway found to be heavily dysregulated among glioma samples with elevated MDA-9. As shown in GSEA plots for complement cascade-related gene sets in both Reactome (ES = 0.84) and Biocarta (ES = 0.88) (Figure 17B), the complement component genes *CFB*, *CFD*, *C6*, *C7*, *C1S*, *C1R*, *C2*, *C1QA*, *CR1*, *C4BPA* are indeed among the most highly up-regulated genes in the MDA-9-high glioma subset. The possible contribution of the complement proteins to cancer proliferation is a relatively new concept. Nonetheless, recent evidence points to the roles of these proteins in the promotion of mitogenic signaling pathways, angiogenesis, migration, resistance to apoptosis and immunosurveillance (Rutkowski *et al.*, 2010). For now, the assumption is that glioma cells themselves are producing these proteins. It would be interesting to investigate how MDA-9 possibly interacts with these proteins.

8.3.3. Association with transcriptional activation and other signaling pathways

8.3.3.1. IGF signaling: As our GSEA analysis indicates, there is a significant dysregulation of the Reactome gene set “*regulation of insulin like growth factor IGF activity by insulin like growth factor binding proteins IGFBPs*”, among the MDA-9-high glioma group (ES = 0.82) (Figure 17C). The most highly ranked genes in this gene set include *IGF binding proteins (IGFBPs (1–5))*, *matrix metalloproteinases (MMP1, MMP2)*, *cathepsin (CTSG)*, *pappalysin 2 (PAPPA2)* and *vascular endothelial growth factor A (VEGFA)*. IGFBPs serve as modulators of IGF1/2, whose downstream transcriptional targets are pro-invasion genes such as *MMP2* and *VEGFA*. On the other hand, PAPPA2 and CTSG are proteases, which regulate IGFBPs. IGFBP2 in particular proved to promote glioma progression (Dunlap *et al.*, 2007). A recent report published by our group demonstrated the connection between MDA-9 and IGFBP2 (Das *et al.*, 2013). As described in that report, over-expression of MDA-9 in melanoma can lead to Src and FAK activation, phosphorylation of Akt, induction of HIF-1 α (hypoxia inducible factor 1- α), eventually leading to elevated IGFBP2 transcription, VEGFA production and angiogenesis.

8.3.3.2. VEGF pathway: As mentioned above, VEGFA is highly up-regulated in the glioma subset with elevated MDA-9 expression. The *VEGF signaling pathway* (KEGG) gene set is indeed enriched among MDA-9-high gliomas, according to results from the GSEA analysis (ES = 0.65) (Figure 17D). Aside from VEGFA, other highly ranked genes belonging to the gene set are several of its downstream targets including: the phospholipase genes *PLA2G2A*, *PLG2G5* and *PLG2G4A*; the adaptor protein *SH2D2A* (*SH2 domain containing 2A*, or *VRAF*); the RAS-related gene *RAC2*, *prostaglandin-endoperoxide synthase 2 (PTGS2* or *COX-2*) and the phosphatidylinositol-4,5-bisphosphate 3-kinase genes *PIK3CG* (catalytic) and *PIK3R5* (regulatory). As mentioned above, we recently demonstrated that VEGFA is a downstream target of MDA-9 (Das *et al.*, 2013).

8.3.3.3. NF- κ B activation: Our group has shown that over-expression of MDA-9 eventually leads to NF- κ B activation and the transcriptional activation of MMP2 and IL8 (Boukerche *et*

et al., 2010; Boukerche *et al.*, 2007; Kegelman *et al.*, 2014). Results from this bioinformatic analysis are consistent with those experimental observations. Three NF- κ B-specific gene sets appear to be enriched in the MDA-9 High subset. These are Reactome's "*TAK1 activates NF- κ B by phosphorylation and activation of IKKS complex*" (ES = 0.76), "*NF- κ B activation via DAI*" (ES = 0.78) and "*TRAF6 mediated NF- κ B activation*" (ES = 0.78). Common to those three pathways are two proteins (the serum amyloid protein SAA1 and the calcium binding protein S100A12), both very highly up-regulated in MDA-9 High gliomas. SAA1 is a previously identified transcriptional target of NF- κ B (Li and Liao, 1991). On the other hand, S100A12 is a known mediator of NF- κ B signaling (Hofmann *et al.*, 1999; Srikrishna *et al.*, 2010).

8.3.4. Association with phospholipid metabolism—Zimmerman and colleagues were responsible for elucidating how MDA-9's PDZ domains bind to phosphatidylinositol 4,5-bisphosphate (PIP₂), a process believed to be crucial to MDA-9's transport to the cell membrane, signal transduction, as well as nuclear assembly (Wawrzyniak *et al.*, 2013; Zimmermann, 2006; Zimmermann *et al.*, 2005). *mda-9*'s role in phospholipid metabolism is also predicted in this bioinformatic analysis. GSEA results indicate a number of enriched gene sets which refer to the five Reactome gene sets which refer to the acyl chain remodeling of phosphatidylinositol, phosphatidylglycerol, phosphatidylserine, phosphatidylcholine and phosphatidylethanolamine. A closer examination of these gene sets points to the phospholipases *PLA2G2A* and *PLA2G5* as the two highest ranked genes common to the gene sets. Another phospholipase (*PLBD1*) is specific to phosphatidylinositol. *PLA2G2A* is among the most highly up-regulated genes in *mda-9* High gliomas (at close to 100-fold). The enzyme is involved in the hydrolysis of phosphatidylinositol (PI) so it can be reacylated by acyltransferases (Singer *et al.*, 2002); a process that diversifies molecules distributed in the membrane. Although not exactly clear at this point, it would be interesting to investigate the possibility of MDA-9 interacting with *PLA2G2A*.

8.3.5. Pathways directly associated with *mda-9* down-regulation—For the *mda-9* Low subgroup, only two gene sets exhibited ES and FDR q values within the acceptable range. One is the Reactome gene set *GABA-A receptor activation*, which has ES and FDR q values of -0.88 and 0.005, respectively. The other one is the Reactome gene set *GABA synthesis release uptake and degradation* (ES = -0.64, FDR q = 0.084). For the first gene set, the core enrichment genes (i.e. those which are highly down-regulated in *mda-9* High glioma) include various GABA-A receptor genes (*GABRG3*, *GABRB1*, *GABRA6*, *GABRA2*, *GABRA3*, *GABRA4*, *GABRB3*, *GABRA5*, *GABRG2*, *GABRA1* and *GABRB2*). For the other gene set, the genes most likely to influence the pathway are *SNAP25* (*synaptosomal-associated protein, 25kDa*), *SYT1* (*synaptotagmin I*), *SLC32A1* [*solute carrier family 32 (GABA vesicular transporter), member 1*], *glutamate decarboxylase* genes (*GAD2*, *GAD1*), *RIMS* (*regulating synaptic membrane exocytosis 1*) and *STX1A* [*syntaxin 1A (brain)*]. These results may be related to several studies describing down-regulation of GABA-A receptor during glioma progression (D'Urso *et al.*, 2012; Desrues *et al.*, 2012; Smits *et al.*, 2012).

8.4. A network of GSEA-identified gene sets

The GSEA-identified gene sets (partially shown in Table 5) can be quite difficult to interpret given the volume of information. However, many of the Reactome, Biocarta, KEGG and GO gene sets are overlapping in terms of component genes. One way to simplify the results and identify gene sets (e.g., pathways, functionalities), which form clusters, is through network analysis. Useful for this purpose is Cytoscape (Shannon *et al.*, 2003) and its Enrichment Map plug-in (Merico *et al.*, 2010). Data generated from GSEA were analyzed directly using these applications. In order to find clusters, a filter was set so that only gene sets registering $p < 0.005$ and FDR $q < 0.015$. This resulted in the narrowing down of the very complex interactions into several prominent clusters including the *Extracellular Matrix Cluster* (Figure 18A) where we can see the overlap and interactions of genes defined by peptidase activity, roles in degradation of the ECM matrix, neurite outgrowth, as well as collagen formation. Another interesting cluster identified through this approach is the *Autoimmunity Disease Cluster* (Figure 18B), even though the dataset used in the analyses came from glioma. This suggests that the same sets of *mda-9*-associated genes may also be important in certain autoimmune diseases including Asthma (something that the authors are already aware of) and Type I Diabetes. This approach also identified a cytokine-related cluster (figure not shown) connecting GO's *leukocyte migration*, *leukocyte chemotaxis*, *locomotory behavior*, *chemokine activity*, *GPCR binding*, *chemokine receptor binding* and Reactome's *chemokine receptor binds chemokines*. As indicated in Table 5, these gene sets are all enriched in the *mda-9* High glioma subset.

9. Summary, Conclusion and Future Directions

Through the integrated bioinformatic analyses of publicly available cancer genomic datasets, we were able to comprehensively analyze the various epigenetic and molecular factors associated with the dysregulation of *mda-9* in various types of cancer. The following is a list of the most important conclusions obtained from this exercise:

- a. The elevation of *mda-9* expression during cancer progression correlates with both a copy number increase and reduced methylation at a key CpG site (cg17197774) located at the intron, more than 1,000 bases 3' from the CpG island. These observations derive from analyses of genome-wide expression, CpG methylation and copy number for TCGA Glioma (Glioblastoma Multiforme, Lower Grade Glioma), SKCM (Skin Cutaneous Melanoma), PRAD (Prostate Adenocarcinoma), COAD (colon adenocarcinoma), LIHC (Liver Hepatocellular Carcinoma) and KIRP (Kidney Renal Papillary Cell Carcinoma) datasets. The degree of influence of each molecular factor (copy number, methylation) towards *mda-9* expression varies among cancer types.
- b. In glioma, both *mda-9* expression level and methylation at cg17197774 can provide prognostic markers, with elevated expression and hypomethylation, respectively, correlating with poor survival.
- c. Among the cancer types included in the TCGA Pan Cancer (PANCAN) dataset, melanoma exhibits the highest level of *mda-9* (which is consistent

with information downloaded from the Human Protein Atlas), which may be explained by its generally low methylation level at the cg17197774 CpG site (which has the lowest among the 6 TCGA datasets analyzed for CpG methylation).

- d. Hypomethylation at cg17197774 is associated with higher intensity of modified histones such as H3K4 di- and tri-methylation and H3K9 acetylation in the promoter area. H3K4 methylations are characteristics of active (or poised to be activated) promoters, while H3K9 acetylation describes a likely active transcription. Data from the ENCODE project (ChIP Seq plus CpG Methylation) were used for this particular set of observations.
- e. An analytical scheme (the **Virtual Gene Over-Expression or Repression; VIGOR**) was designed to predict the groups of genes, functionalities and molecular pathways associated with *mda-9* dysregulation. The scheme started with the formation of two subsets of glioma patients, according to *mda-9* expression: 1) “*mda-9*-High” group consists of samples whose *mda-9* expression is at least 1 sd higher than the average ($z \geq 1$; upper 15.8%) and 2) “*mda-9*-Low” group consists of samples whose *mda-9* expression is at most 1 sd lower than the average ($z \leq -1$; lower 15.8%). The genome-wide expressions of the two subgroups were compared statistically. Gene Set Enrichment Analysis (GSEA) (Subramanian *et al.*, 2005) was then employed to identify gene sets likely associated with *mda-9* dysregulation. Among the top gene groups and molecular pathways identified using the scheme described above are those previously linked with both *mda-9* dysregulation and glioma progression (e.g., matrix metalloproteinases, interleukins, IGFBP2 signaling, NF- κ B activation). Other interesting immune-related pathways such as that of CTLA-4 have been associated with glioma progression but their connections to *mda-9* functionality are yet to be defined.

The driving forces behind a gene's dysregulated expression in cancer (either upregulation or downregulation) involve a multitude of factors including gene copy number variations, epigenetic regulation and transcription factors. As for *mda-9*, the gene is located in chromosome 8, which in many cancer types exhibits a gain of the entire or a greater part of the q arm (Knuutila *et al.*, 1998). A deeper reason may be a resident oncogene in the 8q arm (*MYC*), whose expression often gets elevated in various cancer types (Prochownik, 2008). The 8p loss/8q gain aberration commonly occurs in colorectal cancer (Diep *et al.*, 2006), liver cancer (van Malenstein *et al.*, 2011), prostate cancer (Brothman, 2002) and melanoma (Carlson *et al.*, 2005). In glioma and renal cancer, this specific type of chromosomal aberration is far less frequent (Gardina *et al.*, 2008; Kim *et al.*, 2009). This explains why the R values (REML analysis) for *mda-9* expression vs. *mda-9* copy number are lower in these two cancer types (compared to the other four) (Figure 8). In contrast, the copy number influence is most pronounced in PRAD samples, which are mostly highly methylated at cg17197774. The influence of cg17197774 methylation towards *mda-9* levels is readily observed in the 5 other tumor types.

Our identification of cg17197774 as the most differentially methylated (and most highly associated with *mda-9* transcript levels) is consistent with recent observations by Irizarry and colleagues (Irizarry *et al.*, 2009). In their genome-wide methylation analysis of colon cancer samples, they made the general observation that differentially methylated CpG sites tend to be located outside the CpG island group, as described by Gardiner-Garden and Frommer (Gardiner-Garden and Frommer, 1987). Instead, many of the differentially methylated CpG sites are a few kilobases 5' (referred to as N Shore) or 3' (referred to as S Shore) from the CpG island cluster. The cg17197774 is actually 1,105 bases from the 3' end of the CpG island site.

Another epigenetic factor, which can influence *mda-9* expression are the type of modifications which nucleosomes associated with a gene locus might undergo. For example, an H3 histone di- or tri-methylated at K4 is indicative of a poised or active promoter. A mono-methylated H3 histone is often associated with enhancer downstream of a start site. Transcriptional activity is characterized by the presence of H3 histones whose modifications may be acetylation at K9 or K27, or trimethylation at K36. Active transcription may also be characterized by an H4 histone methylated at K20. In contrast to the aforementioned nucleosomal markers, the locus binding of H3 histone trimethylated at K27, as well as the transcription factor CTCF are both indicative of transcriptional silencing (Campbell and Turner, 2013; Ernst *et al.*, 2011; Ghirlando *et al.*, 2012). Those were the possible scenarios considered in the chromosomal segmentation analysis developed by Ernst, Kellis and co-workers, meant to measure activity of transcription at every gene locus in the entire genome (Ernst and Kellis, 2010; Ernst *et al.*, 2011). We then scrutinized the ChIP Seq-generated ENCODE data accessible through the UCSC Genome Browser. What we observed is that cell lines weakly methylated at cg17197774 (such as GM12878, K562 and HUVEC) tend to have more intense binding by H3 methylated at K4 or acetylated at K9, and H4 methylated at K20. Results from chromosomal segmentation analysis indicate that those 3 cell lines have more active transcription of *mda-9*. For now, we can just assume that the methylation at cg17197774 may be detrimental to the binding affinity of these modified histones. However, this is a readily testable hypothesis.

The GSEA analyses of two glioma subsets at the opposite ends of distribution (*mda-9*High vs. *mda-9*low), apparently was capable of correctly predicting pathways and functionalities that have proven associations with *mda-9*. The top gene sets listed in Table 5 were predicted out of 217 Biocarta, 186 KEGG, 674 Reactome, 825 GO Biological Process, 233 GO Cellular Component and 396 Molecular Function gene sets. This suggests that other predicted pathways without any known link to *mda-9* may be worthy of further investigations.

Many of the gene sets identified by GSEA as likely associated with *mda-9* upregulation refer to extracellular matrix processes and functionalities. These results are consistent with the metastasis-promoting property of *mda-9*. Nonetheless, a closer look at those gene sets leads to questions pertaining to the exact role *mda-9* might play in these processes. For example, it would be interesting to define the details connecting *mda-9* with the various collagen molecules found to be highly up-regulated in *mda-9*High glioma. GSEA analysis also identified gene sets related to signaling pathways involving IGFBP2, NF- κ B and VEGF.

Previous reports have linked *mda-9* to these proteins. Another very interesting observation is the elevated expression of the phospholipase PLA2G2A, an enzyme involved in the acyl chain remodeling of phosphatidylinositol. The possible connection between *mda-9* and PLA2G2A and how it may relate to glioma pathogenesis is an interesting topic to pursue, since MDA-9 is known to bind to phosphoinositides. PLA2G2A is involved in the hydrolysis of phospholipids, a process crucial to the asymmetric distribution of molecules in cell membranes.

The upregulation of various immune related genes and pathways (such as interleukins, interferons and complement cascade) pertaining to the *mda-9*High glioma datasets is not surprising since glioma is known to express many immunosuppressive and immunomodulatory factors (Parney, 2012), such as IL-6, IL-10 and VEGF-A, all of which are over-expressed in *mda-9*High glioma. As we have already proven, over-expression of *mda-9* in glioma cells can indeed stimulate the expression of IL-8 (Kegelman *et al.*, 2014). However, it is important to point out that infiltrating cells such as microglia and monocytes may also contribute to the expression of these cytokines (Parney, 2012). Of additional interest is the predicted association of the CTLA-4 pathway with *mda-9*high glioma. CTLA4 (expressed in Helper T cells and a negative regulator of T lymphocytes) (Daga *et al.*, 2011) increased ~ 3-fold in *mda-9*High relative to *mda-9*Low glioma. At this point, there is still very limited knowledge relative to *mda-9*'s role in glioma immunosuppression.

There are additional interesting observations, which came out of this database analysis. These include the observation that lactotransferrin is the most over-expressed gene in *mda-9*High glioma. Its role in glioma pathogenesis and certainly its possible physical interaction with MDA-9 needs to be investigated. It is also of interest to investigate how *mda-9* contributes to the regulatory protein S100A4's already proven induction of SAA1 (serum amyloid A1) and subsequently IL-8, MMPs and metastasis (Bettum *et al.*, 2014; Hansen *et al.*, 2014; Zhang *et al.*, 2014). All of the aforementioned genes are very highly up-regulated in *mda-9*High gliomas.

What we just demonstrated in this report is the potential of bioinformatic analyses to provide potential insights into our understanding of the biology of the metastasis-promoting gene *mda-9* (its regulation and associated pathways), which could be readily applied to other genes of interest in regulating defined cellular phenotypes. It is now possible to conduct these types of analyses using integrated genomic datasets because of the availability of genomic datasets for various types of molecular profiles for a wide range of cancer types, which continues to expand. Many of the datasets (TCGA in particular) are integrated, which means that a single tumor case has been profiled for genome-wide expression, copy number, mutation and methylation. The approaches employed including **VIGOR** may also be applicable to other genes. We are not implying that the **VIGOR** approach is universally applicable (which will require additional analyses to support the universality of this approach). However, it has proven to be a very useful tool (along with GSEA) in predicting *mda-9*-associated genes and pathways. Overall, this bioinformatic exercise yielded a plethora of information regarding *mda-9*'s functionality (cancer-related or not), which is not surprising given the presence of its promiscuously interacting PDZ domains.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The present work was supported in part by National Institutes of Health Grants R01 CA134721, the National Foundation for Cancer Research (NFCR), the Samuel Waxman Cancer Research Foundation (SWCRF) and cancer research development funds from the VCU Massey Cancer Center. DS is a Harrison and Blick Scholar in the VCU Massey Cancer Center and VCU School of Medicine. PBF holds the Thelma Newmeyer Corman Chair in Cancer Research in the VCU Massey Cancer Center. We sincerely thank Mary Goldman (UCSC) and Joshua Gould (Broad Inst.) for their very helpful advice regarding the use of UCSC Cancer Genomics Browser and Gene-E, respectively.

References

- The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science*. 2004; 306(5696):636–640. [PubMed: 15499007]
- A user's guide to the encyclopedia of DNA elements (ENCODE). *PLoS Biol*. 2011; 9(4):e1001046. [PubMed: 21526222]
- Beekman JM, Coffey PJ. The ins and outs of syntenin, a multifunctional intracellular adaptor protein. *J Cell Sci*. 2008; 121(Pt 9):1349–1355. [PubMed: 18434645]
- Bettum IJ, Vasiliaskaite K, Nygaard V, Clancy T, Pettersen SJ, Tenstad E, et al. Metastasis-associated protein S100A4 induces a network of inflammatory cytokines that activate stromal cells to acquire pro-tumorigenic properties. *Cancer Lett*. 2014; 344(1):28–39. [PubMed: 24215866]
- Boukerche H, Aissaoui H, Prevost C, Hirbec H, Das SK, Su ZZ, et al. Src kinase activation is mandatory for MDA-9/syntenin-mediated activation of nuclear factor-kappaB. *Oncogene*. 2010; 29(21):3054–3066. [PubMed: 20228839]
- Boukerche H, Su ZZ, Emdad L, Baril P, Balme B, Thomas L, et al. mda-9/Syntenin: a positive regulator of melanoma metastasis. *Cancer Res*. 2005; 65(23):10901–10911. [PubMed: 16322237]
- Boukerche H, Su ZZ, Emdad L, Sarkar D, Fisher PB. mda-9/Syntenin regulates the metastatic phenotype in human melanoma cells by activating nuclear factor-kappaB. *Cancer Res*. 2007; 67(4):1812–1822. [PubMed: 17308124]
- Boukerche H, Su ZZ, Prevot C, Sarkar D, Fisher PB. mda-9/Syntenin promotes metastasis in human melanoma cells by activating c-Src. *Proc Natl Acad Sci U S A*. 2008; 105(41):15914–15919. [PubMed: 18832467]
- Brothman AR. Cytogenetics and molecular genetics of cancer of the prostate. *Am J Med Genet*. 2002; 115(3):150–156. [PubMed: 12407695]
- Campbell MJ, Turner BM. Altered histone modifications in cancer. *Adv Exp Med Biol*. 2013; 754:81–107. [PubMed: 22956497]
- Carlson JA, Ross JS, Slominski A, Linette G, Mysliborski J, Hill J, et al. Molecular diagnostics in melanoma. *J Am Acad Dermatol*. 2005; 52(5):743–775. quiz 775-748. [PubMed: 15858465]
- Chandran UR, Ma C, Dhir R, Bisceglia M, Lyons-Weiler M, Liang W, et al. Gene expression profiles of prostate cancer reveal involvement of multiple molecular pathways in the metastatic process. *BMC Cancer*. 2007; 7:64. [PubMed: 17430594]
- Chang K, Creighton CJ, Davis C, Donehower L, Drummond J, Wheeler D, et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet*. 2013; 45(10):1113–1120. [PubMed: 24071849]
- Chen CH, Kuo SM, Liu GS, Chen WN, Chuang CW, Liu LF. Enhancement of neurite outgrowth in neuron cancer stem cells by growth on 3-D collagen scaffolds. *Biochem Biophys Res Commun*. 2012; 428(1):68–73. [PubMed: 23047009]
- Cho W, Kim H, Lee JH, Hong SH, Choe J. Syntenin is expressed in human follicular dendritic cells and involved in the activation of focal adhesion kinase. *Immune Netw*. 2013; 13(5):199–204. [PubMed: 24198745]

- Cifarelli CP, Titus B, Yeoh HK. Cadherin-dependent adhesion of human U373MG glioblastoma cells promotes neurite outgrowth and increases migratory capacity. Laboratory investigation. *J Neurosurg.* 2011; 114(3):663–669. [PubMed: 20415527]
- Cubelos B, Sebastian-Serrano A, Beccari L, Calcagnotto ME, Cisneros E, Kim S, et al. Cux1 and Cux2 regulate dendritic branching, spine morphology, and synapses of the upper layer neurons of the cortex. *Neuron.* 2010; 66(4):523–535. [PubMed: 20510857]
- D'Urso PI, D'Urso OF, Storelli C, Mallardo M, Gianfreda CD, Montinaro A, et al. miR-155 is up-regulated in primary and secondary glioblastoma and promotes tumour growth by inhibiting GABA receptors. *Int J Oncol.* 2012; 41(1):228–234. [PubMed: 22470130]
- Daga A, Bottino C, Castriconi R, Gangemi R, Ferrini S. New perspectives in glioma immunotherapy. *Curr Pharm Des.* 2011; 17(23):2439–2467. [PubMed: 21827420]
- Das SK, Bhutia SK, Azab B, Kegelman TP, Peachy L, Santhekadur PK, et al. MDA-9/syntenin and IGFBP-2 promote angiogenesis in human melanoma. *Cancer Res.* 2013; 73(2):844–854. [PubMed: 23233738]
- Das SK, Bhutia SK, Kegelman TP, Peachy L, Oyesanya RA, Dasgupta S, et al. MDA-9/syntenin: a positive gatekeeper of melanoma metastasis. *Frontiers in bioscience : a journal and virtual library.* 2012; 17:1–15.
- Dasgupta S, Menezes ME, Das SK, Emdad L, Janjic A, Bhatia S, et al. Novel role of MDA-9/syntenin in regulating urothelial cell proliferation by modulating EGFR signaling. *Clin Cancer Res.* 2013; 19(17):4621–4633. [PubMed: 23873690]
- Decleves X, Amiel A, Delattre JY, Scherrmann JM. Role of ABC transporters in the chemoresistance of human gliomas. *Curr Cancer Drug Targets.* 2006; 6(5):433–445. [PubMed: 16918310]
- Desrues L, Lefebvre T, Lecointre C, Schouft MT, Leprince J, Compere V, et al. Down-regulation of GABA(A) receptor via promiscuity with the vasoactive peptide urotensin II receptor. Potential involvement in astrocyte plasticity. *PLoS One.* 2012; 7(5):e36319. [PubMed: 22563490]
- Diep CB, Kleivi K, Ribeiro FR, Teixeira MR, Lindgjaerde OC, Lothe RA. The order of genetic events associated with colorectal cancer progression inferred from meta-analysis of copy number changes. *Genes Chromosomes Cancer.* 2006; 45(1):31–41. [PubMed: 16145679]
- Ducray F, Criniere E, Idhahbi A, Mokhtari K, Marie Y, Paris S, et al. alpha-Internexin expression identifies 1p19q codeleted gliomas. *Neurology.* 2009; 72(2):156–161. [PubMed: 19139367]
- Dunlap SM, Celestino J, Wang H, Jiang R, Holland EC, Fuller GN, et al. Insulin-like growth factor binding protein 2 promotes glioma development and progression. *Proc Natl Acad Sci U S A.* 2007; 104(28):11736–11741. [PubMed: 17606927]
- Egeblad M, Rasch MG, Weaver VM. Dynamic interplay between the collagen scaffold and tumor evolution. *Curr Opin Cell Biol.* 2010; 22(5):697–706. [PubMed: 20822891]
- Ernst J, Kellis M. Discovery and characterization of chromatin states for systematic annotation of the human genome. *Nat Biotechnol.* 2010; 28(8):817–825. [PubMed: 20657582]
- Ernst J, Kheradpour P, Mikkelsen TS, Shores N, Ward LD, Epstein CB, et al. Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature.* 2011; 473(7345):43–49. [PubMed: 21441907]
- Fernandez-Larrea J, Merlos-Suarez A, Urena JM, Baselga J, Arribas J. A role for a PDZ protein in the early secretory pathway for the targeting of proTGF- α to the cell surface. *Mol Cell.* 1999; 3(4):423–433. [PubMed: 10230395]
- Fisher PB, Prignoli DR, Herms H Jr, Weinstein IB, Pestka S. Effects of combined treatment with interferon and mezerein on melanogenesis and growth in human melanoma cells. *J Interferon Res.* 1985; 5(1):11–22. [PubMed: 3989332]
- Francescone RA, Scully S, Faibish M, Taylor SL, Oh D, Moral L, et al. Role of YKL-40 in the angiogenesis, radioresistance, and progression of glioblastoma. *J Biol Chem.* 2011; 286(17):15332–15343. [PubMed: 21385870]
- Gardina PJ, Lo KC, Lee W, Cowell JK, Turpaz Y. Ploidy status and copy number aberrations in primary glioblastomas defined by integrated analysis of allelic ratios, signal ratios and loss of heterozygosity using 500K SNP Mapping Arrays. *BMC Genomics.* 2008; 9:489. [PubMed: 18928532]

- Gardiner-Garden M, Frommer M. CpG islands in vertebrate genomes. *J Mol Biol.* 1987; 196(2):261–282. [PubMed: 3656447]
- Geijsen N, Uings IJ, Pals C, Armstrong J, McKinnon M, Raaijmakers JA, et al. Cytokine-specific transcriptional regulation through an IL-5Ralpha interacting protein. *Science.* 2001; 293(5532): 1136–1138. [PubMed: 11498591]
- Ghirlando R, Giles K, Gowher H, Xiao T, Xu Z, Yao H, et al. Chromatin domains, insulators, and the regulation of gene expression. *Biochim Biophys Acta.* 2012; 1819(7):644–651. [PubMed: 22326678]
- Gibb AJ. NMDA receptor subunit gating--uncovered. *Trends Neurosci.* 2004; 27(1):7–10. discussion 10. [PubMed: 14698601]
- Gibbons JA, Kanwar RK, Kanwar JR. Lactoferrin and cancer in different cancer models. *Front Biosci (Schol Ed).* 2011; 3:1080–1088. [PubMed: 21622257]
- Goldman M, Craft B, Swatloski T, Ellrott K, Cline M, Diekhans M, et al. The UCSC Cancer Genomics Browser: update 2013. *Nucleic Acids Res.* 2013; 41(Database issue):D949–D954. [PubMed: 23109555]
- Gordon-Alonso M, Rocha-Perugini V, Alvarez S, Moreno-Gonzalo O, Ursa A, Lopez-Martin S, et al. The PDZ-adaptor protein syntenin-1 regulates HIV-1 entry. *Mol Biol Cell.* 2012; 23(12):2253–2263. [PubMed: 22535526]
- Hansen MT, Forst B, Cremers N, Quagliata L, Ambartsumian N, Grum-Schwensen B, et al. A link between inflammation and metastasis: serum amyloid A1 and A3 induce metastasis, and are targets of metastasis-inducing S100A4. *Oncogene.* 2014
- Hansen SM, Berezin V, Bock E. Signaling mechanisms of neurite outgrowth induced by the cell adhesion molecules NCAM and N-cadherin. *Cell Mol Life Sci.* 2008; 65(23):3809–3821. [PubMed: 18791849]
- Hao JM, Chen JZ, Sui HM, Si-Ma XQ, Li GQ, Liu C, et al. A five-gene signature as a potential predictor of metastasis and survival in colorectal cancer. *J Pathol.* 2010; 220(4):475–489. [PubMed: 20077526]
- Hernandez M, Martin R, Garcia-Cubillas MD, Maeso-Hernandez P, Nieto ML. Secreted PLA2 induces proliferation in astrocytoma through the EGF receptor: another inflammation-cancer link. *Neuro-oncology.* 2010; 12(10):1014–1023. [PubMed: 20639215]
- Hirbec M, Martin S, Henley JM. Syntenin is involved in the developmental regulation of neuronal membrane architecture. *Mol Cell Neurosci.* 2005; 28(4):737–746. [PubMed: 15797720]
- Hofmann MA, Drury S, Fu C, Qu W, Taguchi A, Lu Y, et al. RAGE mediates a novel proinflammatory axis: a central cell surface receptor for S100/calgranulin polypeptides. *Cell.* 1999; 97(7):889–901. [PubMed: 10399917]
- Hwangbo C, Kim J, Lee JJ, Lee JH. Activation of the integrin effector kinase focal adhesion kinase in cancer cells is regulated by crosstalk between protein kinase Calpha and the PDZ adapter protein mda-9/Syntenin. *Cancer Res.* 2010; 70(4):1645–1655. [PubMed: 20145126]
- Hwangbo C, Park J, Lee JH. mda-9/Syntenin protein positively regulates the activation of Akt protein by facilitating integrin-linked kinase adaptor function during adhesion to type I collagen. *J Biol Chem.* 2011; 286(38):33601–33612. [PubMed: 21828040]
- Irizarry RA, Ladd-Acosta C, Wen B, Wu Z, Montano C, Onyango P, et al. The human colon cancer methylome shows similar hypo- and hypermethylation at conserved tissue-specific CpG island shores. *Nat Genet.* 2009; 41(2):178–186. [PubMed: 19151715]
- Ivarsson Y. Plasticity of PDZ domains in ligand recognition and signaling. *FEBS Lett.* 2012; 586(17): 2638–2647. [PubMed: 22576124]
- Jiang H, Lin JJ, Su ZZ, Goldstein NI, Fisher PB. Subtraction hybridization identifies a novel melanoma differentiation associated gene, mda-7, modulated during human melanoma differentiation, growth and progression. *Oncogene.* 1995; 11(12):2477–2486. [PubMed: 8545104]
- Joshi-Tope G, Gillespie M, Vastrik I, D'Eustachio P, Schmidt E, de Bono B, et al. Reactome: a knowledgebase of biological pathways. *Nucleic Acids Res.* 2005; 33(Database issue):D428–D432. [PubMed: 15608231]
- Kaiser J. National Institutes of Health. NCI gears up for cancer genome project. *Science.* 2005; 307(5713):1182. [PubMed: 15731412]

- Kegelman TP, Das SK, Emdad L, Hu B, Menezes ME, Bhoopathi P, et al. Targeting tumor invasion: the roles of MDA-9/Syntenin. Expert opinion on therapeutic targets. 2015; 19(1):97–112. [PubMed: 25219541]
- Kegelman TP, Das SK, Hu B, Bacolod MD, Fuller CE, Menezes ME, et al. MDA-9/syntenin is a key regulator of glioma pathogenesis. Neuro-oncology. 2014; 16(1):50–61. [PubMed: 24305713]
- Kim HJ, Shen SS, Ayala AG, Ro JY, Truong LD, Alvarez K, et al. Virtual-karyotyping with SNP microarrays in morphologically challenging renal cell neoplasms: a practical and useful diagnostic modality. Am J Surg Pathol. 2009; 33(9):1276–1286. [PubMed: 19461508]
- Kim WY, Jang JY, Jeon YK, Chung DH, Kim YG, Kim CW. Syntenin increases the invasiveness of small cell lung cancer cells by activating p38, AKT, focal adhesion kinase and SP1. Exp Mol Med. 2014; 46:e90. [PubMed: 24722482]
- Kitaya K, Yasuo T, Yamaguchi T, Fushiki S, Honjo H. Genes regulated by interferon-gamma in human uterine microvascular endothelial cells. Int J Mol Med. 2007; 20(5):689–697. [PubMed: 17912462]
- Knuutila S, Bjorkqvist AM, Autio K, Tarkkanen M, Wolf M, Monni O, et al. DNA copy number amplifications in human neoplasms: review of comparative genomic hybridization studies. Am J Pathol. 1998; 152(5):1107–1123. [PubMed: 9588877]
- Koo TH, Lee JJ, Kim EM, Kim KW, Kim HD, Lee JH. Syntenin is overexpressed and promotes cell migration in metastatic human breast and gastric cancer cell lines. Oncogene. 2002; 21(26):4080–4088. [PubMed: 12037664]
- Koroll M, Rathjen FG, Volkmer H. The neural cell recognition molecule neurofascin interacts with syntenin-1 but not with syntenin-2, both of which reveal self-associating activity. J Biol Chem. 2001; 276(14):10646–10654. [PubMed: 11152476]
- Lee H, Kim Y, Choi Y, Choi S, Hong E, Oh ES. Syndecan-2 cytoplasmic domain regulates colon cancer cell migration via interaction with syntenin-1. Biochem Biophys Res Commun. 2011; 409(1):148–153. [PubMed: 21569759]
- Lepinoux-Chambaud C, Eyer J. Review on intermediate filaments of the nervous system and their pathological alterations. Histochem Cell Biol. 2013; 140(1):13–22. [PubMed: 23749407]
- Li XX, Liao WS. Expression of rat serum amyloid A1 gene involves both C/EBP-like and NF kappa B-like transcription factors. J Biol Chem. 1991; 266(23):15192–15201. [PubMed: 1869549]
- Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP. Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011; 27(12):1739–1740. [PubMed: 21546393]
- Lin JJ, Jiang H, Fisher PB. Melanoma differentiation associated gene-9, mda-9, is a human gamma interferon responsive gene. Gene. 1998; 207(2):105–110. [PubMed: 9511750]
- Liu X, Zhang X, Lv Y, Xiang J, Shi J. Overexpression of syntenin enhances hepatoma cell proliferation and invasion: Potential roles in human hepatoma. Oncol Rep. 2014
- Loh YP, Cheng Y, Mahata SK, Corti A, Tota B. Chromogranin A and derived peptides in health and disease. J Mol Neurosci. 2012; 48(2):347–356. [PubMed: 22388654]
- Merico D, Isserlin R, Stueker O, Emili A, Bader GD. Enrichment map: a network-based method for gene-set enrichment visualization and interpretation. PLoS One. 2010; 5(11):e13984. [PubMed: 21085593]
- Mermel CH, Schumacher SE, Hill B, Meyerson ML, Beroukhim R, Getz G. GISTIC2.0 facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers. Genome Biol. 2011; 12(4):R41. [PubMed: 21527027]
- Mikheev AM, Mikheeva SA, Trister AD, Tokita MJ, Emerson SN, Parada CA, et al. Periostin is a novel therapeutic target that predicts and regulates glioma malignancy. Neuro-oncology. 2014
- Parney IF. Basic concepts in glioma immunology. Adv Exp Med Biol. 2012; 746:42–52. [PubMed: 22639158]
- Prochownik EV. c-Myc: linking transformation and genomic instability. Curr Mol Med. 2008; 8(6): 446–458. [PubMed: 18781952]
- Qian XL, Li YQ, Yu B, Gu F, Liu FF, Li WD, et al. Syndecan binding protein (SDCBP) is overexpressed in estrogen receptor negative breast cancers, and is a potential promoter for tumor proliferation. PLoS One. 2013; 8(3):e60046. [PubMed: 23533663]

- Rutkowski MJ, Sughrue ME, Kane AJ, Mills SA, Parsa AT. Cancer and the complement cascade. *Mol Cancer Res.* 2010; 8(11):1453–1465. [PubMed: 20870736]
- Sala-Valdes M, Gordon-Alonso M, Tejera E, Ibanez A, Cabrero JR, Ursa A, et al. Association of syntenin-1 with M-RIP polarizes Rac-1 activation during chemotaxis and immune interactions. *J Cell Sci.* 2012; 125(Pt 5):1235–1246. [PubMed: 22349701]
- Sandoval J, Heyn H, Moran S, Serra-Musach J, Pujana MA, Bibikova M, et al. Validation of a DNA methylation microarray for 450,000 CpG sites in the human genome. *Epigenetics.* 2011; 6(6):692–702. [PubMed: 21593595]
- Sarkar D, Boukerche H, Su ZZ, Fisher PB. mda-9/Syntenin: more than just a simple adapter protein when it comes to cancer metastasis. *Cancer Res.* 2008; 68(9):3087–3093. [PubMed: 18451132]
- Shah N, Sukumar S. The Hox genes and their roles in oncogenesis. *Nat Rev Cancer.* 2010; 10(5):361–371. [PubMed: 20357775]
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* 2003; 13(11):2498–2504. [PubMed: 14597658]
- Singer AG, Ghomashchi F, Le Calvez C, Bollinger J, Bezzine S, Rouault M, et al. Interfacial kinetic and binding properties of the complete set of human and mouse groups I, II, V, X, and XII secreted phospholipases A2. *J Biol Chem.* 2002; 277(50):48535–48549. [PubMed: 12359733]
- Singh R, Park D, Xu J, Hosur R, Berger B. Struct2Net: a web service to predict protein-protein interactions using a structure-based approach. *Nucleic Acids Res.* 2010; 38(Web Server issue):W508–W515. [PubMed: 20513650]
- Smits A, Jin Z, Elsir T, Pedder H, Nister M, Alafuzoff I, et al. GABA-A channel subunit expression in human glioma correlates with tumor histology and clinical outcome. *PLoS One.* 2012; 7(5):e37041. [PubMed: 22615883]
- Sokhi UK, Bacolod MD, Dasgupta S, Emdad L, Das SK, Dumur CI, et al. Identification of genes potentially regulated by human polynucleotide phosphorylase (hPNPase old-35) using melanoma as a model. *PLoS One.* 2013; 8(10):e76284. [PubMed: 24143183]
- Srikrishna G, Nayak J, Weigle B, Temme A, Foell D, Hazelwood L, et al. Carboxylated N-glycans on RAGE promote S100A12 binding and signaling. *J Cell Biochem.* 2010; 110(3):645–659. [PubMed: 20512925]
- Stier S, Totzke G, Grunewald E, Neuhaus T, Fronhoffs S, Sachinidis A, et al. Identification of syntenin and other TNF-inducible genes in human umbilical arterial endothelial cells by suppression subtractive hybridization. *FEBS Lett.* 2000; 467(2–3):299–304. [PubMed: 10675558]
- Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A.* 2005; 102(43):15545–15550. [PubMed: 16199517]
- Sun L, Hui AM, Su Q, Vortmeyer A, Kotliarov Y, Pastorino S, et al. Neuronal and glioma-derived stem cell factor induces angiogenesis within the brain. *Cancer Cell.* 2006; 9(4):287–300. [PubMed: 16616334]
- Takahashi K, Mitoma J, Hosono M, Shiozaki K, Sato C, Yamaguchi K, et al. Sialidase NEU4 hydrolyzes polysialic acids of neural cell adhesion molecules and negatively regulates neurite formation by hippocampal neurons. *J Biol Chem.* 2012; 287(18):14816–14826. [PubMed: 22393058]
- Talantov D, Mazumder A, Yu JX, Briggs T, Jiang Y, Backus J, et al. Novel genes associated with malignant melanoma but not benign melanocytic lesions. *Clin Cancer Res.* 2005; 11(20):7234–7242. [PubMed: 16243793]
- Tian B, Zhang Y, Zhang J. Periostin is a new potential prognostic biomarker for glioma. *Tumour Biol.* 2014; 35(6):5877–5883. [PubMed: 24719188]
- Tomoda T, Kim JH, Zhan C, Hatten ME. Role of Unc51.1 and its binding partners in CNS axon outgrowth. *Genes Dev.* 2004; 18(5):541–558. [PubMed: 15014045]
- Uhlen M, Oksvold P, Fagerberg L, Lundberg E, Jonasson K, Forsberg M, et al. Towards a knowledge-based Human Protein Atlas. *Nat Biotechnol.* 2010; 28(12):1248–1250. [PubMed: 21139605]
- van Malenstein H, van Pelt J, Verslype C. Molecular classification of hepatocellular carcinoma anno 2011. *Eur J Cancer.* 2011; 47(12):1789–1797. [PubMed: 21612914]

- VanMeter TE, Rooprai HK, Kibble MM, Fillmore HL, Broaddus WC, Pilkington GJ. The role of matrix metalloproteinase genes in glioma invasion: co-dependent and interactive proteolysis. *J Neurooncol.* 2001; 53(2):213–235. [PubMed: 11716072]
- Wang H, Wang Y, Jiang C. Stromal protein periostin identified as a progression associated and prognostic biomarker in glioma via inducing an invasive and proliferative phenotype. *Int J Oncol.* 2013; 42(5):1716–1724. [PubMed: 23467707]
- Ward PP, Paz E, Conneely OM. Multifunctional roles of lactoferrin: a critical overview. *Cell Mol Life Sci.* 2005; 62(22):2540–2548. [PubMed: 16261256]
- Wawrzyniak AM, Kashyap R, Zimmermann P. Phosphoinositides and PDZ domain scaffolds. *Adv Exp Med Biol.* 2013; 991:41–57. [PubMed: 23775690]
- Yeung YT, McDonald KL, Grewal T, Munoz L. Interleukins in glioblastoma pathophysiology: implications for therapy. *Br J Pharmacol.* 2013; 168(3):591–606. [PubMed: 23062197]
- Zhang W, Liu Y, Wang CW. S100A4 promotes squamous cell laryngeal cancer Hep-2 cell invasion via NF-kB/MMP-9 signal. *Eur Rev Med Pharmacol Sci.* 2014; 18(9):1361–1367. [PubMed: 24867514]
- Zhong D, Ran JH, Tang WY, Zhang XD, Tan Y, Chen GJ, et al. Mda-9/syntenin promotes human brain glioma migration through focal adhesion kinase (FAK)-JNK and FAK-AKT signaling. *Asian Pac J Cancer Prev.* 2012; 13(6):2897–2901. [PubMed: 22938480]
- Zhu J, Sanborn JZ, Benz S, Szeto C, Hsu F, Kuhn RM, et al. The UCSC Cancer Genomics Browser. *Nat Methods.* 2009; 6(4):239–240. [PubMed: 19333237]
- Zimmermann P. The prevalence and significance of PDZ domain-phosphoinositide interactions. *Biochim Biophys Acta.* 2006; 1761(8):947–956. [PubMed: 16884951]
- Zimmermann P, Tomatis D, Rosas M, Grootjans J, Leenaerts I, Degeest G, et al. Characterization of syntenin, a syndecan-binding PDZ protein, as a component of cell adhesion sites and microfilaments. *Mol Biol Cell.* 2001; 12(2):339–350. [PubMed: 11179419]
- Zimmermann P, Zhang Z, Degeest G, Mortier E, Leenaerts I, Coomans C, et al. Syndecan recycling [corrected] is controlled by syntenin-PIP2 interaction and Arf6. *Dev Cell.* 2005; 9(3):377–388. [PubMed: 16139226]

Figure 1.

Flowchart depicting the objectives and schemes of the present study.

Figure 2.

A. The progressive increase in the expression level of *mda-9/syntenin* in glioma as observed in the TCGA Glioma dataset (left) and GEO dataset GSE4290 (right). **B.** The overall survival of glioma patients, grouped according to *mda-9* expression level.

Figure 3.

The expression of *mda-9* (relative to normal samples) in melanoma (**A**), prostate cancer (**B**), liver cancer (**C**), and renal cancer (**D**). The expression values were derived from public genome-wide expression datasets. The relative expression (z) is calculated as $z = (I_{i,n} - \text{Ave. } I_{norm}) / sd_{norm}$, where n refers to every sample (including tumors), while $norm$ refers to normal samples only. Inset are the resulting p values for t-tests comparing the normal and tumor sample groups.

Figure 4. Comparative *mda-9* expression profiles of normal prostate, primary prostate tumors and metastatic prostate cancer.

Figure 5.

A. The locus map of *mda-9* with the 9 identified exons in RefSeq. UCSC Genes prediction identified a 10th exon, after the 5' UTR exon. **B.** Higher resolution of the promoter region, indicating the locations of the interrogated CpG sites. Coordinates were derived from GRCh37/hg19 assembly.

Figure 6.

The degree of methylation, ranging from totally unmethylated (−0.5) to totally methylated (0.5), at the 13 CpG sites interrogated in Illumina 450K array, for tissue samples belonging to 6 TCGA cancer groups. Of the 13 sites, the most differentially methylated is cg179774 (dark blue), followed by cg10129404 (dark red).

Figure 7.

In both TCGA glioma (A) and KIRP (B), *mda-9* expression is clearly influenced by the methylation status of cg17197774 (the CpG site close to 5' UTR) and not by cg10129404 (the CpG site at 3' UTR).

Figure 8.

The expression of *mda-9* as a function of its copy number and methylation at cg17197774, among six TCGA datasets.

Figure 9.

Methylation at cg17197774 is a marker of clinical outcome in glioma. Patients were grouped into two, according to % methylation (**A**), or methylation value (Beta value) relative to median (**B**).

Figure 10. *mda-9* expression (PANCAN-normalized) vs. cg17197774 Beta values with data points labeled according to TCGA cohort (**A**), and copy number (**B**). **C**. The same dataset with the exponential 3P regression plot. **D** *mda-9* expression vs. Copy Number, with data points marked according to cg17197774 Beta value.

Figure 11.

The Expression of *mda-9* (PANCAN-normalized) with samples grouped according to sample type (A) and TCGA cohort (solid normals not included) (B). The cohorts are the following: A. Acute Myeloid Leukemia [LAML], B. Bladder Urothelial Carcinoma [BLCA], C. Breast invasive carcinoma [BRCA], D. Cervical squamous cell carcinoma and endocervical adenocarcinoma (Francescone *et al.*), E. Colon adenocarcinoma [COAD], F. Uterine Corpus Endometrial Carcinoma [UCEC], G. Glioblastoma multiforme [GBM], H. Head and Neck squamous cell carcinoma [HNSC], I. Kidney Chromophobe [KICH], J. Kidney renal clear cell carcinoma [KIRC], K. Kidney renal papillary cell carcinoma [KIRP], L. Liver hepatocellular carcinoma [LIHC], M. Brain Lower Grade Glioma [LGG], N. Lung adenocarcinoma [LUAD], O. Lung squamous cell carcinoma [LUSC], P. Skin Cutaneous Melanoma [SKCM], Q. Ovarian serous cystadenocarcinoma [OV], R. Pancreatic adenocarcinoma [PAAD], S. Prostate adenocarcinoma [PRAD], T. Rectum adenocarcinoma [READ], U. Sarcoma [SARC], V. Thyroid carcinoma [THCA].

Figure 12.

MDA-9 protein levels, as assessed by Immunohistochemistry (semi-quantitative) using two different antibodies. The bars indicate that melanoma has the highest MDA-9 expression among different cancer types. Insets are the immunohistochemical images of melanoma and glioma sections, indicative of the former's stronger MDA-9 signal. All of the images above were taken from the Protein Atlas (<http://www.proteinatlas.org>).

Figure 13.
A. The “affinity” of modified histones and CTCF towards the *mda-9* promoter region of the HUVEC and H1-hESC cell lines. **B.** Chromatin state segmentation at the *mda-9* locus, for 6 cell lines (using the method of *Ernst and Kellis, 2010*). Cell lines with low Beta values at cg1719774 tend to have a more active promoter (dark red) and transcription (dark green). **C.** Color codes for the different chromosomal states (copied from UCSC Genome Browser; <http://genome.ucsc.edu>).

Figure 14. Virtual Gene Over-expression or Repression (VIGOR)

An analytical scheme to identify genes and pathways associated with *mda-9* dysregulation in glioma.

Figure 15.

There is a great degree of similarity between TCGA Glioma and GSE4290 datasets, in terms of signal to noise ratio (*mda-9*High vs. *mda-9*Low) for every gene.

Figure 16.
A. The enrichment of genes involved in extracellular matrix degradation in *mda-9* High glioma (relative to *mda-9* Low glioma). **B.** Snapshots of the pathways as shown in Reactome (www.reactome.org). Red markers indicate the involvement of core enrichment genes in the degradation of fibrillin (left), fibronectin (center) and decorin (right).

Figure 17.

Four representative pathways and processes, which exhibited gene enrichment in GSEA analysis. Indicated are the core enrichment genes in order of decreasing rank.

Figure 18.

Two of the prominent clusters of GSEA-identified gene sets.

Table 1

List of publicly available genomic datasets analyzed for the study

Dataset ID	Cancer Type	Platform	No. Cancer	No. Normal
A. Genome-wide Expression				
TCGA_GBM ^a	Glioma (Glioblastoma Multiforme)	Illumina Hi Seq 2000	168	0
TCGA_LGG	Glioma (Lower Grade Glioma)	Illumina Hi Seq 2000	205	0
TCGA_LIHC	Liver Hepatocellular Carcinoma	Illumina Hi Seq 2000	69	36
TCGA_PRAD	Prostate Adenocarcinoma	Illumina Hi Seq 2000	142	37
TCGA_KIRP	Kidney Renal Papillary Cell Carcinoma	Illumina Hi Seq 2000	76	25
TCGA_PANCAN ^b	Pan Cancer (Combination of 22 Cancer Types)	Illumina Hi Seq 2000	5453	587
GSE4290b ^c	Glioma (all grades)	Affymetrix HG U133 Plus 2 Array	153	23
GSE3189 ^c	Melanoma	Affymetrix HG U133A Array	45	7
GDS2545 ^{c,d}	Prostate Cancer	Affymetrix HG U95A Array	90	81
B. Genome-wide copy number				
TCGA_GBM	Glioma (Glioblastoma Multiforme)	Affymetrix SNP 6 Array	544	N/A
TCGA_LGG	Glioma (Lower Grade Glioma)	Affymetrix SNP 6 Array	206	N/A
TCGA_LIHC	Liver Hepatocellular Carcinoma	Affymetrix SNP 6 Array	97	N/A
TCGA_COAD	Colon Adenocarcinoma	Affymetrix SNP 6 Array	413	N/A
TCGA_PRAD	Prostate Adenocarcinoma	Affymetrix SNP 6 Array	187	N/A
TCGA_KIRP	Kidney Renal Papillary Cell Carcinoma	Affymetrix SNP 6 Array	127	N/A
TCGA_SKCM	Skin Cutaneous Melanoma	Affymetrix SNP 6 Array	260	N/A
C. Genome-wide CpG methylation				
TCGA_GBM	Glioma (Glioblastoma Multiforme)	Illumina Methylation 450K Beadchip Array	121	0
TCGA_LGG	Glioma (Lower Grade Glioma)	Illumina Methylation 450K Beadchip Array	204	0
TCGA_LIHC	Liver Hepatocellular Carcinoma	Illumina Methylation 450K Beadchip Array	98	50
TCGA_COAD	Colon Adenocarcinoma	Illumina Methylation 450K Beadchip Array	258	38
TCGA_PRAD	Prostate Adenocarcinoma	Illumina Methylation 450K Beadchip Array	192	49
TCGA_KIRP	Kidney Renal Papillary Cell Carcinoma	Illumina Methylation 450K Beadchip Array	111	45

Dataset ID	Cancer Type	Platform	No. Cancer	No. Normal
TCGA_SKCM	Skin Cutaneous Melanoma	Illumina Methylation 450K Beadchip Array	338 ^d	1

Notes:

^a All of TCGA datasets and accompanying clinico-pathological information were downloaded as pre-processed data, through the UCSC Cancer Genomics Browser (<https://genome-cancer.ucsc.edu>).

^b The TCGA Pan Cancer (PANCAN) dataset is mean-normalized, consisting of 22 TCGA datasets. Normals were not analyzed for this study.

^c These datasets were downloaded from the NCBI Gene Expression Omnibus (GEO) site (<http://www.ncbi.nlm.nih.gov/geo/>).

^d Tumor samples include 267 metastatic and 71 primary tumors.

Table 2

List of the *MDA-9* CpG sites interrogated in the Illumina Infinium 450K Beadchip array

Probe ID	Strand	Chr	Coordinate (build 37)	Relation to UC SC CpG Island	UCSC RefGene Group	Forward Sequence
cg20052227	R	8	59465520	Island	TSS1500	GTGGGTGGCA[CG]GGGCCCCGCGG
cg02896624	R	8	59465528	Island	TSS1500	GCACGGGGCC[CG]CGGGCACGAA
cg17984783	R	8	59465536	Island	TSS200	CCCCGGGCA[CG]AACAGCCGAA
cg26656684	F	8	59465596	Island	TSS200	CAGCGGACAG[CG]GGCGGCATGA
cg11550426	F	8	59465608	Island	TSS200	CGGCATGAAC[CG]CCCCACTTTG
cg27280034	F	8	59465624	Island	TSS200	CCCACTTTGC[CG]GATACCTGGA
cg06294637	F	8	59465768	Island	1stExon,5'UTR	GCCTCGGGGG[CG]GTCTTCGGGC
cg07798892	F	8	59465776	Island	1stExon,5'UTR	GGTCTTCGGG[CG]CGCACCGCTC
cg20848390	F	8	59465780	Island	1stExon,5'UTR	TCTTCGGGG[CG]CACCCGCTCTC
cg02103294	R	8	59465972	Island	5'UTR	GCATCCTGGT[CG]CAGCCGTTTT
cg12046629	R	8	59466300	S_Shore	5'UTR	TCCCAGTGT[CG]GCGTTTCTAG
cg17197774	F	8	59467208	S_Shore	5'UTR	TAATGGTTGC[CG]GTTAAATGTA
cg10129404	R	8	59494304		3'UTR	TTAAAAATTCA[CG]GCACCATGGA

Table 3

Comparison of *MDA-9* copy number, CpG methylation (at cg17197774), RNA level, and protein level among six TCGA cohorts. Included in the analyses are tumor samples having all the three molecular data (copy number, expression, and methylation).

Tumor Cohort (TCGA)	No. of tumor samples	Copy number estimate (lowest, highest)	CpG Methylation at cg17197774 (–0.5 to 0.5; low to high)	Expression level
SKCM	241	2.80 (1.15, 24.2)	–0.295	23062 ± 20569
PRAD	175	2.18 (1.20, 8.82)	0.195	3039 ± 943
COAD	188	2.82 (1.11, 17.93)	–0.019	4172 ± 1514
GBM/LGG	257	2.09 (1.09, 5.56)	0.209	4308 ± 1500
LIHC	68	2.92 (1.09, 14.55)	–0.088	4623 ± 2505
KIRP	60	2.16 (0.84)	–0.014	5553 ± 3268

Table 4

The Top 200 upregulated and Top 50 down-regulated genes in MDA9-high glioma

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
A. Genes highly upregulated in MDA9-high glioma							
<i>LTF</i>	lactotransferrin	3p21.31	175.41	1.29	2.0E-03	202018_s_at	68.19
<i>CHI3L1</i>	chitinase 3-like 1 (cartilage glycoprotein-39)	1q22.1	110.21	1.26	2.0E-03	209396_s_at	16.94
<i>PLA2G2A</i>	phospholipase A2, group IIA (platelets, synovial fluid)	1p35	98.04	1.26	2.0E-03	203649_s_at	7.45
<i>POSTN</i>	periostin, osteoblast specific factor	13q13.3	86.88	0.98	2.0E-03	210809_s_at	12.53
<i>ABCC3</i>	ATP-binding cassette, sub-family C (CFTR/MRP), member 3	17q21	59.53	1.33	2.0E-03	208161_s_at	23.81
<i>SAA1</i>	serum amyloid A1 (SAA1), transcript variant 3	11p15.1	54.64	1.05	2.0E-03	208607_s_at	10.29
<i>IBSP</i>	integrin-binding sialoprotein	4q21.1	51.14	1.08	2.0E-03	236028_at	11.05
<i>MMP9</i>	matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)	20q12-q13	44.21	1.01	2.0E-03	203936_s_at	2.65
<i>NNMT</i>	nicotinamide N-methyltransferase	11q23.1	37.14	1.24	2.0E-03	202238_s_at	18.72
<i>CLEC5A</i>	C-type lectin domain family 5, member A	7q34	33.97	1.38	2.0E-03	219890_at	5.46
<i>COL3A1</i>	collagen, type III, alpha 1	2q32.2	30.99	0.99	2.0E-03	211161_s_at	16.7
<i>COL1A1</i>	collagen, type I, alpha 1	17q21.33	30.08	0.99	2.0E-03	202310_s_at	17.27
<i>TREML1</i>	triggering receptor expressed on myeloid cells 1	6p21.1	30.06	1.14	2.0E-03	219434_at	5.12
<i>HMXC10</i>	homeobox C10	12q13.13	28.58	0.9	2.0E-03	218959_at	2.38
<i>CHI3L2</i>	chitinase 3-like 2	1p13.3	27.18	1.07	2.0E-03	213060_s_at	7.83
<i>TIMP1</i>	TIMP metalloproteinase inhibitor 1	Xp11.3-p11.23	27.17	1.29	2.0E-03	201666_at	10.89
<i>CD163</i>	CD163 molecule	12p13	26.87	1.3	2.0E-03	215049_x_at	10.62
<i>SHOX2</i>	short stature homeobox 2	3q25.32	26.33	0.96	2.0E-03	210135_s_at	7.65
<i>FCGR2B</i>	Fc fragment of IgG, low affinity IIb, receptor (CD32)	1q23	25.18	1.25	2.0E-03	210889_s_at	3.49
<i>PI3</i>	peptidase inhibitor 3, skin-derived	20q13.12	24.76	0.84	2.0E-03	203691_at	3.8
<i>SRPX2</i>	sushi-repeat-containing protein, X-linked 2	Xq21.33-q23	23.25	1.29	2.0E-03	205499_at	11.53
<i>GDF15</i>	growth differentiation factor 15	19p13.11	23.08	1.23	2.0E-03	221577_x_at	6.27
<i>MEOX2</i>	mesenchyme homeobox 2	7p22.1-p21.3	22.45	0.84	2.0E-03	206201_s_at	9.29
<i>SERPINA5</i>	serpin peptidase inhibitor, clade A (alpha-1 antitrypsin, antitrypsin),	14q32.1	22.32	1.33	2.0E-03	209443_at	3.67

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
	member 5						
<i>HOXA10</i>	homeobox A10	7p15.2	21.96	0.88	2.0E-03	213150_at	6.32
<i>HOXD13</i>	homeobox D13	2q31.1	21.57	0.86	2.0E-03	207397_s_at	3.73
<i>TNFSF12-TNFSF13</i>	TNFSF12-TNFSF13 readthrough	17p13.1	21.56	1.15	2.0E-03	209499_x_at	1.60
<i>HOXB3</i>	homeobox B3	17q21.32	21.52	0.9	2.0E-03	228904_at	4.18
<i>PDPN</i>	podoplanin	1p36.21	21.39	1.2	2.0E-03	204879_at	27.01
<i>LIF</i>	leukemia inhibitory factor (cholinergic differentiation factor)	22q12.2	21.38	1.05	2.0E-03	205266_at	3.82
<i>SERPINE1</i>	serpin peptidase inhibitor, clade E (nexin, plasminogen activator inhibitor type 1), member 1	7q22.1	21.1	1.17	2.0E-03	202627_s_at	8.69
<i>IL2RA</i>	interleukin 2 receptor, alpha	10p15-p14	20.86	1.12	2.0E-03	206341_at	1.49
<i>HOXD11</i>	homeobox D11	2q31.1	20.3	0.93	2.0E-03	214604_at	3.85
<i>ESM1</i>	endothelial cell-specific molecule 1	5q11	19.95	1.02	2.0E-03	208394_x_at	3.89
<i>ANXA1</i>	annexin A1	9q21.13	19.6	1.55	2.0E-03	201012_at	10.05
<i>HOXA3</i>	homeobox A3	7p15.2	19.2	0.93	2.0E-03	235521_at	4.21
<i>LOX</i>	lysyl oxidase	5q23.3-q31.2	19.07	1.25	2.0E-03	215446_s_at	14.4
<i>EMP3</i>	epithelial membrane protein 3	19q13.3	18.95	1.26	2.0E-03	203729_at	9.7
<i>IL8</i>	interleukin 8	4q13-q21	18.93	0.79	2.0E-03	211506_s_at	7.86
<i>HOXD10</i>	homeobox D10	2q31.1	18.87	0.89	2.0E-03	229400_at	4.74
<i>SPOCD1</i>	SPOC domain containing 1	1p35.1	18.77	0.97	2.0E-03	235417_at	14.27
<i>GPX8</i>	glutathione peroxidase 8 (putative)	5q11.2	18.76	1.23	2.0E-03	228141_at	7.38
<i>HOXA4</i>	homeobox A4	7p15.2	18.33	0.82	2.0E-03	206289_at	2.21
<i>CXCL10</i>	chemokine (C-X-C motif) ligand 10	4q21	18.28	1.01	2.0E-03	204533_at	2.73
<i>HOXA7</i>	homeobox A7	7p15.2	18.2	0.7	2.0E-03	235753_at	2.63
<i>HOXA5</i>	homeobox A5	7p15.2	17.67	0.95	2.0E-03	213844_at	3.83
<i>GALNT5</i>	UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylglucosaminyltransferase 5 (GalNAc-T5)	2q24.1	17.34	1.14	2.0E-03	236129_at	1.76
<i>CA3</i>	carbonic anhydrase III, muscle specific	8q21.2	17.26	0.92	2.0E-03	204865_at	6.67
<i>STC1</i>	stanniocalcin 1	8p22-p12	17.13	1.08	2.0E-03	204597_x_at	10.02
<i>PTX3</i>	pentraxin 3, long	3q25	16.81	0.98	2.0E-03	206157_at	9.7

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
<i>CSTA</i>	cystatin A (stefin A)	3q21	16.67	1.37	2.0E-03	204971_at	3.71
<i>METTL7B</i>	methyltransferase like 7B	12q13.2	16.52	1.14	2.0E-03	227055_at	4.37
<i>S100A4</i>	S100 calcium binding protein A4	1q12-q22	16.37	1.19	2.0E-03	203186_s_at	6.94
<i>TRPM8</i>	transient receptor potential cation channel, subfamily M, member 8	2q37	16.14	1.11	2.0E-03	243483_at	3.94
<i>IGFBP2</i>	insulin-like growth factor binding protein 2, 36kDa	2q33-q34	16.03	0.98	2.0E-03	202718_at	19.33
<i>FMOD</i>	fibromodulin	1q32	15.94	1.06	2.0E-03	202709_at	3.63
<i>H19</i>	H19, imprinted maternally expressed transcript (non-protein coding)	11p15.5	15.94	0.62	2.0E-03	224646_x_at	4.4
<i>IL13RA2</i>	interleukin 13 receptor, alpha 2	Xq13.1-q28	15.87	0.71	2.0E-03	206172_at	6.65
<i>HPD</i>	4-hydroxyphenylpyruvate dioxygenase	12q24.31	15.45	1.08	2.0E-03	206024_at	1.53
<i>SERPINA3</i>	serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 3	14q32.1	15.08	1.04	2.0E-03	202376_at	6.22
<i>COL6A3</i>	collagen, type VI, alpha 3	2q37	14.99	0.89	2.0E-03	201438_at	8.45
<i>SLC47A2</i>	solute carrier family 47, member 2	17p11.2	14.93	0.95	2.0E-03	231068_at	3.24
<i>WISP1</i>	WNT1 inducible signaling pathway protein 1	8q24.22	14.67	1.04	2.0E-03	206796_at	3.64
<i>ANXA2P2</i>	annexin A2 pseudogene 2	9p13.3	14.52	1.3	2.0E-03	208816_x_at	4.43
<i>CHRNA9</i>	cholinergic receptor, nicotinic, alpha 9	4p14	14.42	0.89	2.0E-03	221107_at	2.97
<i>F13A1</i>	coagulation factor XIII, A1 polypeptide	6p24.2-p23	14.38	0.93	2.0E-03	203305_at	3.38
<i>TNFRSF12A</i>	tumor necrosis factor receptor superfamily, member 12A	16p13.3	14.35	1.17	2.0E-03	218368_s_at	7.47
<i>ADM</i>	adrenomedullin	11p15.4	14.32	1.01	2.0E-03	202912_at	7.57
<i>COL8A1</i>	collagen, type VIII, alpha 1	3q11.1-q13.2	14.29	1	2.0E-03	226237_at	9.88
<i>MOXD1</i>	monooxygenase, DBH-like 1	6q23.2	14.24	0.83	2.0E-03	1554474_a_at	3.6
<i>COL6A2</i>	collagen, type VI, alpha 2	21q22.3	13.96	1	2.0E-03	209156_s_at	9.88
<i>SOC3</i>	suppressor of cytokine signaling 3	17q25.3	13.77	1.08	2.0E-03	227697_at	10.04
<i>EMRI</i>	egf-like module containing, mucin-like, hormone receptor-like 1	19p13.3	13.7	1.08	2.0E-03	207111_at	1.2
<i>TCTEX1D1</i>	Tetex1 domain containing 1	1p31.2	13.69	0.91	2.0E-03	1553635_s_at	2.27
<i>NKX2-5</i>	NK2 transcription factor related, locus 5 (Drosophila)	5q34	13.66	0.77	2.0E-03	206578_at	3
<i>COL5A1</i>	collagen, type V, alpha 1	9q34.2-q34.3	13.22	0.87	2.0E-03	203325_s_at	5.38
<i>PLEK2</i>	pleckstrin 2	14q23.3	13.11	1.17	2.0E-03	218644_at	1.24
<i>HOXB4</i>	homeobox B4	17q21.32	13.06	0.87	2.0E-03	231767_at	1.69

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
<i>CLCF1</i>	cardiotrophin-like cytokine factor 1	11q13.3	12.92	1.12	2.0E-03	219500_at	2.52
<i>CA9</i>	carbonic anhydrase IX	9p13.3	12.85	0.67	2.0E-03	205199_at	1.4
<i>LYZ</i>	lysozyme	12q15	12.75	1.01	2.0E-03	213975_s_at	8.16
<i>HOTAIR</i>	hox transcript antisense RNA (non-protein coding)	12q13.13	12.71	0.88	2.0E-03	239153_at	4.45
<i>MSR1</i>	macrophage scavenger receptor 1	8p22	12.66	1.35	2.0E-03	214770_at	4.12
<i>GPRC5A</i>	G protein-coupled receptor, family C, group 5, member A	12p13-p12.3	12.59	1.07	2.0E-03	203108_at	3.36
<i>PLAU</i>	plasminogen activator, urokinase	10q24	12.59	1.08	2.0E-03	205479_s_at	5.57
<i>EN1</i>	engrailed homeobox 1	2q14.2	12.44	0.72	2.0E-03	220559_at	3.92
<i>FCGBP</i>	Fc fragment of IgG binding protein	19q13.2	12.41	1.04	2.0E-03	203240_at	6.56
<i>ADAM12</i>	ADAM metalloproteinase domain 12	10q26	12.39	1.06	2.0E-03	202952_s_at	5.39
<i>IFI30</i>	interferon, gamma-inducible protein 30	19p13.1	12.33	1.21	2.0E-03	201422_at	4.83
<i>PAX3</i>	paired box 3	2q36.1	12.23	0.78	2.0E-03	231666_at	2.35
<i>DDIT4L</i>	DNA-damage-inducible transcript 4-like	4q23	12.23	0.91	2.0E-03	228057_at	2.58
<i>COL4A1</i>	collagen, type IV, alpha 1	13q34	12.02	1.12	2.0E-03	211981_at	15.36
<i>LUM</i>	lumican	12q21.3-q22	11.92	0.9	2.0E-03	201744_s_at	7.11
<i>FCGR2C</i>	Fc fragment of IgG, low affinity IIc, receptor for (CD32) (gene/pseudogene)	1q23	11.84	1.26	2.0E-03	210992_x_at	11.95
<i>S100A9</i>	S100 calcium binding protein A9	1q21	11.82	0.83	2.0E-03	203535_at	6.04
<i>CCL20</i>	chemokine (C-C motif) ligand 20	2q36.3	11.8	1	2.0E-03	205476_at	3.21
<i>HOXB2</i>	homeobox B2	17q21.32	11.73	0.8	2.0E-03	205453_at	3.17
<i>NCRNA00152</i>	non-protein coding RNA 152		11.73	1.02	2.0E-03	1552258_at	1.53
<i>HOXA2</i>	homeobox A2	7p15.2	11.68	0.78	2.0E-03	214457_at	3.04
<i>ADAMDECI</i>	ADAM-like, decysin 1	8p12	11.66	0.76	2.0E-03	206134_at	2.41
<i>CCL2</i>	chemokine (C-C motif) ligand 2	17q11.2-q21.1	11.51	0.96	2.0E-03	216598_s_at	8.73
<i>G0S2</i>	G0/G1switch 2	1q32.2-q41	11.49	0.92	2.0E-03	213524_s_at	1.9
<i>HOXA1</i>	homeobox A1	7p15.2	11.49	1.16	2.0E-03	214639_s_at	5.7
<i>HOXC6</i>	homeobox C6	12q13.13	11.39	0.88	2.0E-03	206858_s_at	3.08
<i>SH2D4A</i>	SH2 domain containing 4A	8p21	11.38	1.35	2.0E-03	219749_at	1.35
<i>COL1A2</i>	collagen, type I, alpha 2	7q21.3	11.3	0.94	2.0E-03	202404_s_at	11.33

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
<i>DPEP1</i>	dipeptidase 1 (renal)	16q24	11.2	0.75	2.0E-03	205983_at	1.08
<i>MMP7</i>	matrix metalloproteinase 7 (matrilysin, uterine)	11q21-q22	11.2	0.66	2.0E-03	204259_at	1.8
<i>S100A8</i>	S100 calcium binding protein A8	1q12-q22	11.08	0.81	2.0E-03	202917_s_at	5.21
<i>PTPN22</i>	protein tyrosine phosphatase, non-receptor type 22 (lymphoid)	1p13.2	11.04	1.29	2.0E-03	236539_at	2.91
<i>MARCO</i>	macrophage receptor with collagenous structure	2q14.2	10.95	0.77	2.0E-03	205819_at	3.02
<i>HLA-DQA1</i>	major histocompatibility complex, class II, DQ alpha 1	6p21.3	10.77	0.87	2.0E-03	203290_at	3.12
<i>STEAP3</i>	STEAP family member 3	2q14.2	10.74	1.28	2.0E-03	1554830_a_at	9.76
<i>IGFBP3</i>	insulin-like growth factor 2 mRNA binding protein 3	7p15.3	10.71	0.88	2.0E-03	203819_s_at	6.63
<i>CTHRC1</i>	collagen triple helix repeat containing 1	8q22.3	10.66	1.02	2.0E-03	225681_at	4.52
<i>C7orf57</i>	chromosome 7 open reading frame 57	7p12.3	10.64	0.81	2.0E-03	1557636_a_at	6.67
<i>TGFBI</i>	transforming growth factor, beta-induced, 68kDa	5q31	10.63	0.98	2.0E-03	201506_at	6.66
<i>MYBPH</i>	myosin binding protein H	1q32.1	10.61	0.86	2.0E-03	206304_at	1.42
<i>HOXB7</i>	homeobox B7	17q21.32	10.59	0.87	2.0E-03	204778_x_at	2.95
<i>HOXC13</i>	homeobox C13	12q13.13	10.58	0.71	2.0E-03	219832_s_at	1.53
<i>TNFRSF11B</i>	tumor necrosis factor receptor superfamily, member 11b	8q24	10.56	1.03	2.0E-03	204933_s_at	12.3
<i>FPR3</i>	formyl peptide receptor 3	19q13.3-q13.4	10.35	0.92	2.0E-03	230422_at	3.26
<i>MGP</i>	matrix Gla protein	12p12.3	10.31	0.95	2.0E-03	202291_s_at	5.36
<i>IGFBP2</i>	insulin-like growth factor 2 mRNA binding protein 2	3q27.2	10.3	0.86	2.0E-03	218847_at	1.53
<i>CCDC109B</i>	coiled-coil domain containing 109B	4q25	10.29	1.38	2.0E-03	218802_at	3.55
<i>TUBA1C</i>	tubulin, alpha 1c	12q13.12	10.27	1.12	2.0E-03	209251_x_at	1.02
<i>MS4A6A</i>	membrane-spanning 4-domains, subfamily A, member 6A	11q12.1	10.27	1.06	2.0E-03	224356_x_at	4.92
<i>GPR65</i>	G protein-coupled receptor 65	14q31-q32.1	10.26	1.24	2.0E-03	214467_at	3.96
<i>FAM20A</i>	family with sequence similarity 20, member A	17q24.3	10.06	1.03	2.0E-03	226804_at	3.31
<i>GPR82</i>	G protein-coupled receptor 82	Xp11.4	10.03	1.1	2.0E-03	244434_at	3.85
<i>ANXA2</i>	annexin A2	15q22.2	10.02	1.28	2.0E-03	213503_x_at	6.09
<i>C15orf48</i>	chromosome 15 open reading frame 48	15q21.1	10.02	0.94	2.0E-03	223484_at	6.09
<i>HAMP</i>	hepcidin antimicrobial peptide	19q13.1	9.96	0.85	2.0E-03	220491_at	2.03
<i>MYO1G</i>	myosin IG	7p13-p11.2	9.93	1.09	2.0E-03	227799_at	1.22

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
<i>SERPINA1</i>	serpin peptidase inhibitor, clade A (alpha-1 antitrypsin, antitrypsin), member 1	14q32.1	9.84	1.19	2.0E-03	202833_s_at	4.61
<i>AEBP1</i>	AE binding protein 1	7p	9.81	1.04	2.0E-03	201792_at	5.23
<i>ADAM6</i>	ADAM metalloproteinase domain 6 (pseudogene)	14q32.33	9.81	0.68	2.0E-03	237909_at	-2.17
<i>PLAUR</i>	plasminogen activator, urokinase receptor	19q13	9.79	1.23	2.0E-03	211924_s_at	2.82
<i>TNFAIP6</i>	tumor necrosis factor, alpha-induced protein 6	2q23.3	9.77	0.91	2.0E-03	206025_s_at	3.38
<i>MAP1LC3C</i>	microtubule-associated protein 1 light chain 3 gamma	1q43	9.73	1.17	2.0E-03	221697_at	2.64
<i>LOC541471</i>	Hypothetical LOC541471		9.65	0.96	2.0E-03	1562876_s_at	3.06
<i>HMG2</i>	high mobility group AT-hook 2	12q15	9.63	0.74	2.0E-03	1558682_at	-1.67
<i>RARRES2</i>	retinoic acid receptor responder (azartene induced) 2	7q36.1	9.6	0.82	2.0E-03	209496_at	4.01
<i>GPR1</i>	G protein-coupled receptor 1	2q33.3	9.53	0.9	2.0E-03	214605_x_at	1.6
<i>DPYD</i>	dihydropyrimidine dehydrogenase	1p22	9.44	1.42	2.0E-03	1554536_at	4.89
<i>RNASE2</i>	ribonuclease, RNase A family, 2 (liver, eosinophil-derived neurotoxin)	14q24-q31	9.36	1.07	2.0E-03	206111_at	2.52
<i>CD48</i>	CD48 molecule	1q21.3-q22	9.26	0.99	2.0E-03	204118_at	2
<i>IGFBP3</i>	insulin-like growth factor binding protein 3	7p13-p12	9.25	0.87	2.0E-03	210095_s_at	6.64
<i>PDLIM1</i>	PDZ and LIM domain 1	10q23.1	9.24	1.14	2.0E-03	208690_s_at	4.53
<i>DES</i>	desmin	2q35	9.23	0.71	2.0E-03	202222_s_at	1.72
<i>GNPMB</i>	glycoprotein (transmembrane) nmb	7p	9.2	0.97	2.0E-03	1554018_at	5.2
<i>LRRN4CL</i>	LRRN4 C-terminal like	11q12.3	9.19	1.07	2.0E-03	1556427_s_at	2.61
<i>THBS1</i>	thrombospondin 1	15q15	9.09	0.88	2.0E-03	201109_s_at	8.69
<i>IDO1</i>	indoleamine 2,3-dioxygenase 1	8p12-p11	9.04	0.88	2.0E-03	210029_at	1.48
<i>CLEC12A</i>	C-type lectin domain family 12, member A	12p13.31	9.03	0.97	2.0E-03	1552398_a_at	1.82
<i>OSR2</i>	odd-skipped related 2 (Drosophila)	8q22.2	9.02	0.87	2.0E-03	213568_at	2.12
<i>DMRTA2</i>	DMRT-like family A2	1p33	9.01	0.65	2.0E-03	1558856_at	2.21
<i>IL1RN</i>	interleukin 1 receptor antagonist	2q14.2	9.01	0.89	2.0E-03	212657_s_at	2.12
<i>CCR2</i>	chemokine (C-C motif) receptor 2	3p21	8.97	1.07	2.0E-03	206978_at	1.43
<i>HOXD9</i>	homeobox D9	2q31.1	8.97	0.64	2.0E-03	205604_at	1.95
<i>KDELR3</i>	KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 3	22q13	8.97	1.15	2.0E-03	204017_at	2.33

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
<i>IL1R2</i>	interleukin 1 receptor, type II	2q12	8.96	0.85	2.0E-03	205403_at	2.62
<i>STAC</i>	SH3 and cysteine rich domain	3p22.3	8.95	0.72	2.0E-03	205743_at	1.55
<i>AREG</i>	amphiregulin	4q13.3	8.87	0.84	2.0E-03	205239_at	1.9
<i>ACTG2</i>	actin, gamma 2, smooth muscle, enteric	2p13.1	8.84	0.75	2.0E-03	202274_at	1.91
<i>ANXA2P1</i>	annexin A2 pseudogene 1	4q31.3	8.81	1.13	2.0E-03	210876_at	2.65
<i>SPAG4</i>	sperm associated antigen 4	20q11.2	8.8	0.98	2.0E-03	219888_at	3.73
<i>PITXI</i>	paired-like homeodomain 1	5q31.1	8.75	0.71	2.0E-03	208502_s_at	3.39
<i>CLIC1</i>	chloride intracellular channel 1	6p21.3	8.72	1.34	2.0E-03	208659_at	4.19
<i>C21orf7</i>	chromosome 21 open reading frame 7		8.69	1.2	2.0E-03	221211_s_at	2.8
<i>LSP1</i>	lymphocyte-specific protein 1	11p15.5	8.66	1.09	2.0E-03	203523_at	5.02
<i>HOXC11</i>	homeobox C11	12q13.13	8.65	0.73	2.0E-03	206745_at	1.44
<i>PLBD1</i>	phospholipase B domain containing 1	12p13.1	8.63	1.33	2.0E-03	218454_at	2.21
<i>VDR</i>	vitamin D (1,25- dihydroxyvitamin D3) receptor	12q12-q14	8.61	1.1	2.0E-03	204254_s_at	1.88
<i>HOXC9</i>	homeobox C9	12q13.13	8.61	0.77	2.0E-03	231936_at	1.5
<i>CP</i>	ceruloplasmin (ferroxidase)	3q23-q25	8.6	0.85	2.0E-03	1558034_s_at	6.76
<i>BCL2A1</i>	BCL2-related protein A1	15q24.3	8.58	0.9	2.0E-03	205681_at	3.23
<i>TYMP</i>	thymidine phosphorylase	22q13	8.58	1.06	2.0E-03	204858_s_at	2.47
<i>SFRP4</i>	secreted frizzled-related protein 4	7p14.1	8.57	0.98	2.0E-03	204051_s_at	3.64
<i>HLA-DRA</i>	major histocompatibility complex, class II, DR alpha	6p21.3	8.56	1.12	2.0E-03	210982_s_at	4.69
<i>SLC11A1</i>	solute carrier family 11 (proton-coupled divalent metal ion transporters), member 1	2q35	8.54	1.1	2.0E-03	217507_at	3.23
<i>PTPN7</i>	protein tyrosine phosphatase, non-receptor type 7	1q32.1	8.51	1.15	2.0E-03	204852_s_at	1.23
<i>C1R</i>	complement component 1, r subcomponent	12p13.31	8.45	1.22	2.0E-03	212067_s_at	4.36
<i>IL6</i>	interleukin 6 (interferon, beta 2)	7p21-p15	8.44	0.73	2.0E-03	205207_at	2.15
<i>PDLIM4</i>	PDZ and LIM domain 4	5q31.1	8.43	0.83	2.0E-03	211564_s_at	6.01
<i>CASP4</i>	caspase 4, apoptosis-related cysteine peptidase	11q22.2-q22.3	8.42	1.38	2.0E-03	209310_s_at	3.59
<i>GZMA</i>	granzyme A (granzyme 1, cytotoxic T-lymphocyte-associated serine esterase 3)	5q11-q12	8.38	1.01	2.0E-03	205488_at	1.25
<i>FPR2</i>	formyl peptide receptor 2	19q13.3-q13.4	8.36	0.93	2.0E-03	210772_at	2.61

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
<i>GBP5</i>	guanylate binding protein 5	1p22.2	8.36	1.03	2.0E-03	229625_at	4.48
<i>F2RL2</i>	coagulation factor II (thrombin) receptor-like 2	5q13	8.32	0.9	2.0E-03	230147_at	7.55
<i>PCOLCE</i>	procollagen C-endopeptidase enhancer	7q22	8.29	0.86	2.0E-03	202465_at	5.24
<i>TMEM71</i>	transmembrane protein 71	8q24.22	8.29	1.08	2.0E-03	238429_at	10.76
<i>SPP1</i>	Secreted phosphoprotein 1	4q22.1	8.29	0.89	2.0E-03	1568574_x_at	2.21
<i>KYNU</i>	kynureninase (L-kynurenine hydrolase)	2q22.2	8.29	1.17	2.0E-03	210663_s_at	3.45
<i>PDCD1LG2</i>	programmed cell death 1 ligand 2	9p24.2	8.28	1.18	2.0E-03	220049_s_at	1.36
<i>FAP</i>	fibroblast activation protein, alpha	2q23	8.24	1.01	2.0E-03	209955_s_at	2.45
<i>OR51E1</i>	olfactory receptor, family 51, subfamily E, member 1	11p15.4	8.22	0.86	2.0E-03	229768_at	1.08
<i>RBPI</i>	retinol binding protein 1, cellular	3q21-q23	8.21	0.89	2.0E-03	203423_at	2.71
<i>DSG2</i>	desmoglein 2	18q12.1	8.21	0.64	2.0E-03	217901_at	2.18
<i>OTP</i>	orthopedia homeobox	5q14.1	8.19	0.66	2.0E-03	223835_x_at	1.85
<i>NOX4</i>	NADPH oxidase 4	11q14.2-q21	8.18	0.96	2.0E-03	219773_at	2.9
<i>KRT75</i>	keratin 75	12q13.13	8.17	0.74	2.0E-03	207065_at	1.59
<i>B. Genes highly downregulated in MDA9-high glioma</i>							
<i>GRIN1</i>	glutamate receptor, ionotropic, N-methyl D-aspartate 1	9q34.3	-47.19	-1.03	2.0E-03	210781_x_at	-5.19
<i>CUX2</i>	cut-like homeobox 2	12q24.12	-30.25	-1.23	2.0E-03	213920_at	-6.58
<i>INA</i>	intemexin neuronal intermediate filament protein, alpha	10q24	-28.94	-1.11	2.0E-03	204465_s_at	-6.42
<i>SVOP</i>	SV2 related protein homolog (rat)	12q24.11	-27.91	-1.02	2.0E-03	229818_at	-7.29
<i>TNR</i>	tenascin R (restrictin, janusin)	1q24	-27.59	-0.96	2.0E-03	1564897_at	-1.87
<i>CHGA</i>	chromogranin A (parathyroid secretory protein 1)	14q32	-26.84	-0.99	2.0E-03	204697_s_at	-7.24
<i>LRTM2</i>	leucine-rich repeats and transmembrane domains 2	12p13.33	-26.48	-1.15	2.0E-03	1558530_at	-2.86
<i>SLC1A6</i>	solute carrier family 1 (high affinity aspartate/glutamate transporter), member 6	19p13.12	-26.13	-1.18	2.0E-03	1554593_s_at	-7.03
<i>JPH3</i>	junctophilin 3	16q24.3	-25.52	-1.27	2.0E-03	229294_at	-10.33
<i>C2orf85</i>	chromosome 2 open reading frame 85		-24.44	-1.29	2.0E-03	1560035_at	-1.69
<i>CACNG2</i>	calcium channel, voltage-dependent, gamma subunit 2	22q13.1	-24.06	-1.19	2.0E-03	214495_at	-7.69
<i>FAM123C</i>	family with sequence similarity 123C		-23.91	-1.15	2.0E-03	244709_at	-5.02
<i>PRLHR</i>	prolactin releasing hormone receptor	10q25.3-q26	-23.89	-1.01	2.0E-03	231805_at	-2.34

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
<i>C5MD3</i>	CUB and Sushi multiple domains 3	8q23.3	-23.02	-0.96	2.0E-03	240228_at	-5.32
<i>RIMS2</i>	regulating synaptic membrane exocytosis 2	8q22.3	-22.68	-1.09	2.0E-03	229823_at	-7.59
<i>CPLX2</i>	complexin 2	5q35.2	-22.32	-0.95	2.0E-03	225815_at	-8.52
<i>CACNA1B</i>	calcium channel, voltage-dependent, N type, alpha 1B subunit	9q34	-22.31	-1.03	2.0E-03	235781_at	-4.37
<i>MYT1L</i>	myelin transcription factor 1-like	2p25.3	-22.01	-0.91	2.0E-03	210016_at	-9.13
<i>SSTR1</i>	somatostatin receptor 1	14q13	-21.55	-0.99	2.0E-03	235591_at	-7.38
<i>PTPRT</i>	protein tyrosine phosphatase, receptor type, T	20q12-q13	-21.4	-1.19	2.0E-03	205948_at	-8.59
<i>PCDH15</i>	protocadherin-related 15	10q21.1	-21.24	-0.91	2.0E-03	1553344_at	-7.92
<i>SCRT1</i>	scratch homolog 1, zinc finger protein (Drosophila)	8q24.3	-21.23	-1.16	2.0E-03	228761_at	-5.7
<i>CBLN1</i>	cerebellin 1 precursor	16q12.1	-20.32	-1.25	2.0E-03	205747_at	-2.33
<i>VSTM2A</i>	V-set and transmembrane domain containing 2A	7p11.2	-20.15	-0.94	2.0E-03	1554530_at	-8.52
<i>HRH3</i>	histamine receptor H3	20q13.33	-20.07	-1.13	2.0E-03	220447_at	-6.95
<i>ST8SIA3</i>	ST8 alpha-N-acetyl-neuraminide alpha-2,8-sialyltransferase 3	18q21.31	-20.05	-0.99	2.0E-03	230262_at	-4.69
<i>GABRG2</i>	gamma-aminobutyric acid (GABA) A receptor, gamma 2	5q34	-19.84	-0.8	2.0E-03	1568612_at	-9.37
<i>SNAP91</i>	synaptosomal-associated protein, 91kDa homolog (mouse)	6q15	-19.6	-1.05	2.0E-03	204953_at	-10.7
<i>SPHKAP</i>	SPHK1 interactor, AKAP domain containing	2q36.3	-19.43	-0.94	2.0E-03	228509_at	-3.95
<i>UNC13C</i>	unc-13 homolog C (C. elegans)	15q21.3	-19.11	-0.91	2.0E-03	1556095_at	-11.56
<i>PCDH11X</i>	protocadherin 11 X-linked	Xq21.3	-18.92	-1.05	2.0E-03	208366_at	-1.03
<i>CHRNA4</i>	cholinergic receptor, nicotinic, alpha 4	20q13.33	-18.91	-1.12	2.0E-03	206736_x_at	-1.63
<i>HMP19</i>	HMP19 protein		-18.87	-1.04	2.0E-03	218623_at	-7.22
<i>KSR2</i>	kinase suppressor of ras 2	12q24.22-q24.23	-18.45	-0.95	2.0E-03	230551_at	-4.61
<i>ELFN2</i>	extracellular leucine-rich repeat and fibronectin type III domain containing 2	22q13.1	-18.03	-1.09	2.0E-03	1559072_a_at	-5.83
<i>GABRA1</i>	gamma-aminobutyric acid (GABA) A receptor, alpha 1	5q34	-17.92	-0.71	2.0E-03	244118_at	-17.68
<i>PAK7</i>	p21 protein (Cdc42/Rac)-activated kinase 7	20p12	-17.89	-1.11	2.0E-03	210721_s_at	-8.97
<i>WSCD2</i>	WSC domain containing 2	12q23.3	-17.87	-1.01	2.0E-03	229032_at	-4.8
<i>SLC8A2</i>	solute carrier family 8 (sodium/calcium exchanger), member 2	19q13.32	-17.81	-1.06	2.0E-03	215267_s_at	-7.58
<i>ACTL6B</i>	actin-like 6B	7q22	-17.65	-1	2.0E-03	206013_s_at	-11.81

Id	Gene Title	Chromosome	Fold change (TCGA)	S-to-N ratio (TCGA)	p-value (TCGA)	Affymetrix ID	Fold change (GSE4290)
<i>PCK2</i>	proprotein convertase subtilisin/kexin type 2	20p11.2	-17.5	-0.85	2.0E-03	204870_s_at	-10.56
<i>GLRA3</i>	glycine receptor, alpha 3	4q34.1	-17.2	-1.06	2.0E-03	207928_s_at	-2.7
<i>USH1C</i>	Usher syndrome 1C (autosomal recessive, severe)	11p14.3	-17.12	-0.87	2.0E-03	205137_x_at	-3.81
<i>HPSE2</i>	heparanase 2	10q23-q24	-17.07	-0.93	2.0E-03	220927_s_at	-2.73
<i>CHRM1</i>	cholinergic receptor, muscarinic 1	11q12-q13	-17.04	-0.95	2.0E-03	231783_at	-9.4
<i>KCNT1</i>	potassium channel, subfamily T, member 1	9q34.3	-16.93	-0.95	2.0E-03	1563608_a_at	-2.52
<i>OPALIN</i>	oligodendrocytic myelin paranodal and inner loop protein	10q23-q24	-16.92	-0.66	2.0E-03	239575_at	-3.82
<i>CHD5</i>	chromodomain helicase DNA binding protein 5	1p36.3	-16.68	-0.93	2.0E-03	213965_s_at	-4.45
<i>CACNG3</i>	calcium channel, voltage-dependent, gamma subunit 3	16p12.1	-16.05	-0.78	2.0E-03	206384_at	-9.57
<i>SHANK2</i>	SH3 and multiple ankyrin repeat domains 2	11q13.2	-15.85	-1.15	2.0E-03	243681_at	-6.99
<i>ZDHHC22</i>	zinc finger, DHHC-type containing 22	14q24.3	-15.52	-1.23	2.0E-03	229875_at	-15.29

Table 5

The top gene sets found to be enriched in *MDA-9* High (A–F) and *MDA9* Low (G) glioma according to GSEA analysis. The gene sets represent molecular pathways, biological process, gene functionalities, subcellular locations, as well as other diseases. Indicates next to a gene set is an Enrichment Score (ES).

<i>Gene Set Source</i>	Gene Sets Enriched in MDA-9 High Glioma
A. Reactome	translocation of ZAP70 to immunological synapse (0.89), phosphorylation of CD3 and TCR zeta chains (0.88), acyl chain remodelling of PI (0.88), PD1 signaling (0.88), initial triggering of complement (0.87), acyl chain remodelling of PG (0.87), regulation of complement cascade (0.86), acyl chain remodelling of PS (0.85), NCAM1 interactions (0.85), acyl chain remodelling of PC (0.84), acyl chain remodelling of PE (0.84), advanced glycosylation endproduct receptor signaling (0.84), complement cascade (0.83), degradation of the extracellular matrix (0.82), regulation of insulin like growth factor IGF activity by insulin like growth factor binding proteins IGFbps (0.82), generation of second messenger molecules (0.82), latent infection of homo sapiens with mycobacterium tuberculosis (0.8), extracellular matrix organization (0.79), synthesis of PA (0.79), RIP mediated NFKB activation via DAI (0.78), extrinsic pathway for apoptosis (0.78), TRAF6 mediated NFKB activation (0.78), collagen formation (0.78), chemokine receptors bind chemokines (0.77), trafficking and processing of endosomal TLR (0.77), TAK1 activates NFKB by phosphorylation and activation of IKKS complex (0.76), immunoregulatory interactions between a lymphoid and a non lymphoid cell (0.76), NCAM signaling for neurite out growth (0.76), peptide ligand binding receptors (0.73), interferon gamma signaling (0.73), IL receptor SHC signaling (0.72), GPVI mediated activation cascade (0.72), common pathway (0.71), integrin cell surface interactions (0.71), amyloids (0.7), costimulation by the CD28 family (0.69), highly calcium permeable postsynaptic nicotinic acetylcholine receptors (0.69), reversible hydration of carbon dioxide (0.69), cell surface interactions at the vascular wall (0.69), keratan sulfate biosynthesis (0.68)
B. Biocarta	CTL pathway (0.89), classic pathway (0.89), T Helper pathway (0.88), T cytotoxic pathway (0.88), complement pathway (0.87), antigen dependent B cell activation pathway (0.87), granulocytes pathway (0.86), CTLA4 pathway (0.85), LAIR pathway (0.85), IL17 pathway (0.84), TH1TH2 pathway (0.83), NO2IL12 pathway (0.81), fibrinolysis pathway (0.81), stem pathway (0.79), TOB1 pathway (0.78), inflammation pathway (0.78), platelet APP pathway (0.74), NKT pathway (0.74), CSK pathway (0.73), IL2 pathway (0.73), D4GDI pathway (0.73), IL7 pathway (0.72), cytokine pathway (0.71), IL12 pathway (0.71), DC pathway (0.7), IL10 pathway (0.69), STATHMIN pathway (0.69), IL2RB pathway (0.69)
C. KEGG	alpha linolenic acid metabolism (0.86), allograft rejection (0.86), asthma (0.86), graft versus host disease (0.84), type I diabetes mellitus (0.82), linoleic acid metabolism (0.8), autoimmune thyroid disease (0.78), ether lipid metabolism (0.78), complement and coagulation cascades (0.77), intestinal immune network for IGA production (0.77), ECM receptor interaction (0.76), hematopoietic cell lineage (0.76), Leishmania infection (0.74), renin angiotensin system (0.72), phenylalanine metabolism (0.71), cytokine cytokine receptor interaction (0.7), viral myocarditis (0.7), cell adhesion molecules cams (0.68), NOD like receptor signaling pathway (0.68), arachidonic acid metabolism (0.67), long term depression (0.67), pantothenate and CoA biosynthesis (0.67), tryptophan metabolism (0.67), Fc epsilon RI signaling pathway (0.67), primary immunodeficiency (0.67), chemokine signaling pathway (0.65), VEGF signaling pathway (0.64), focal adhesion (0.63), other glycan degradation (0.63), antigen processing and presentation (0.61), systemic lupus erythematosus (0.61), glycosaminoglycan degradation (0.61), nicotinate and nicotinamide metabolism (0.6), bladder cancer (0.6), toll like receptor signaling pathway (0.6), O -glycan biosynthesis (0.6), prion diseases (0.6)
D. GO Biological Processes	positive regulation of cell adhesion (0.87), leukocyte chemotaxis (0.87), leukocyte migration (0.85), keratinocyte differentiation (0.81), locomotory behavior (0.8), humoral immune response (0.79), regulation of cell adhesion (0.78), cell maturation (0.78), developmental maturation (0.76), immune response (0.75), cytokine and chemokine mediated signaling pathway (0.74), negative regulation of cytokine biosynthetic process (0.74), inflammatory response (0.74), negative regulation of phosphorylation (0.74), immune system process (0.73), cellular defense response (0.73), behavior (0.73), vitamin transport (0.73), cellular cation homeostasis (0.72), regulation of defense response (0.72), response to wounding (0.72), cation

<i>Gene Set Source</i>	Gene Sets Enriched in MDA-9 High Glioma
	homeostasis (0.72), regulation of viral reproduction (0.72), calcium mediated signaling (0.71), response to external stimulus (0.71), negative regulation of phosphate metabolic process (0.71), positive regulation of peptidyl tyrosine phosphorylation (0.71), T cell differentiation (0.71), defense response (0.7), cell recognition (0.7), regulation of JAK STAT cascade (0.7), regulation of response to external stimulus (0.7), defense response to virus (0.7), innate immune response (0.7), positive regulation of response to stimulus (0.69), regulation of hormone secretion (0.69), positive regulation of cell cycle (0.69), interferon gamma biosynthetic process (0.69), adaptive immune response (0.69), positive regulation of immune response (0.69)
E. GO Molecular Function	phospholipase A2 activity (0.9), G protein coupled receptor binding (0.85), chemokine receptor binding (0.84), chemokine activity (0.84), metalloendopeptidase activity (0.84), interleukin receptor activity (0.83), cysteine type endopeptidase activity (0.81), interleukin binding (0.79), endopeptidase activity (0.79), serine type endopeptidase activity (0.78), hematopoietin interferon class (D200 domain) cytokine receptor activity (0.78), serine type peptidase activity (0.77), extracellular matrix structural constituent (0.77), serine hydrolase activity (0.77), carboxylesterase activity (0.77), cysteine type peptidase activity (0.76), glycosaminoglycan binding (0.76), polysaccharide binding (0.76), collagen binding (0.76), metallopeptidase activity (0.75), growth factor binding (0.75), inositol or phosphatidylinositol phosphodiesterase activity (0.74), heparin binding (0.73), peptidase activity (0.73), cytokine binding (0.73), cytokine activity (0.73), phospholipase activity (0.73), phospholipase c activity (0.72), pattern binding (0.71), lipase activity (0.71), carbohydrate binding (0.7), serine type endopeptidase inhibitor activity (0.68), sugar binding (0.68), protease inhibitor activity (0.67), sodium channel activity (0.65), caspase regulator activity (0.65), arylsulfatase activity (0.64), carbonate dehydratase activity (0.64), oxidoreductase activity acting on the aldehyde or oxo group of donors (0.64), acetylgalactosaminyltransferase activity (0.64)
F. GO Cellular Component	collagen (0.87), proteinaceous extracellular matrix (0.77), extracellular matrix (0.77), extracellular matrix part (0.76), integrin complex (0.74), cornified envelope (0.71), extracellular region part (0.71), extracellular region (0.71), extracellular space (0.69), external side of plasma membrane (0.69)
	Gene Sets Enriched in MDA-9 Low Glioma
Reactome	GABA A receptor activation (−0.88), GABA synthesis release reuptake and degradation (−0.64)