Published in final edited form as:

J Trauma Stress. 2016 August ; 29(4): 379–383. doi:10.1002/jts.22122.

The Role of Posttraumatic Growth in a Randomized Controlled Trial of Cognitive-Behavioral Conjoint Therapy for PTSD

Anne C. Wagner¹, Lindsey Torbit¹, Tiffany Jenzer², Meredith S. H. Landy¹, Nicole D. Pukay-Martin³, Alexandra Macdonald⁴, Steffany J. Fredman⁵, and Candice M. Monson¹

¹Department of Psychology, Ryerson University, Toronto, Ontario, Canada

²Department of Psychology, The University at Buffalo, State University of New York, Buffalo, New York, United States of America

³Trauma Recovery Center, Cincinnati VA Medical Center, Cincinnati, Ohio, United States of America

⁴The Citadel, Charleston, South Carolina, United States of America

⁵Department of Human Development and Family Studies The Pennsylvania State University, University Park, Pennsylvania, United States of America

Abstract

Posttraumatic growth (PTG) is defined as a positive psychological change that can emerge following a highly challenging life event. Although documented in noninterventional studies of traumatized individuals, there is scant data on the potential for therapy to induce or improve PTG. Thus, the primary goal of this study was to examine changes in PTG in a waitlist controlled trial of cognitive-behavioral conjoint therapy for posttraumatic stress disorder (CBCT for PTSD; Monson & Fredman, 2012). We also examined whether pre-treatment relationship satisfaction and PTSD symptomatology moderated PTG change. There were 40 couples (75% with a female partner with PTSD) who were randomized to either immediate CBCT for PTSD or a 3-month waitlist (WL). Compared to WL, individuals who received CBCT for PTSD demonstrated a significant increase in PTG, with a moderate effect size between-group difference (Hedge's $g = 0.45$). There was a nonsignificant relationship with a moderate effect size (Hedge's $g = 0.65$) for the positive effect of pretreatment relationship satisfaction on the trajectory of PTG, but no effect of pretreatment PTSD symptoms. Results suggested that CBCT for PTSD facilitates PTG, even with a limited focus on PTG in this conjoint intervention. Future research should target PTG as a treatment goal and further examine the role of close others in facilitating its development.

Posttraumatic growth (PTG) refers to positive psychological change that can emerge following exposure to trauma, which may be the result of having survived the trauma and seeing new possibilities in life (e.g., thinking about how best to live; Calhoun & Tedeschi, 2014). Although PTG is thought to reflect a cognitive adaptation process among those who experience posttraumatic stress disorder (PTSD) in response to a trauma (Dekel, Mandl, &

Solomon, 2011), little is known about the potential for PTSD treatments to enhance PTG. Thus, the purpose of this study was to examine changes in PTG in a waitlist controlled trial of Cognitive-Behavioral Conjoint Therapy (CBCT) for PTSD (Monson & Fredman, 2012) and to examine whether pre-treatment relationship satisfaction and PTSD symptomatology moderated PTG change.

PTG is an important and desirable occurrence, as it is positively associated with hope, interpersonal relationship functioning, meaning-making of the traumatic event, and the ability to cope with future stressors and negatively associated with avoidance (Prati & Pietrantonio, 2009; Tedeschi & Calhoun, 2004). Research directly evaluating PTG as an outcome of PTSD treatment is scant; findings from one treatment trial of individual CBT for PTSD found increases in some (e.g., new possibilities and personal strength) but not other areas of PTG (relations to others, spiritual change, and appreciation of life; Zoellner, Rabe, Karl, & Maercker, 2011).

Interpersonal factors have a strong association with PTG and may be key elements in the facilitation of its development (e.g., Prati & Pietrantonio, 2009). Responses of close others to trauma disclosure, positive role modeling of PTG, and shared perceptions/interpretations/beliefs with close others about the trauma are positively associated with PTG (e.g., Calhoun & Tedeschi, 2014). Given the importance of social factors in PTG, dyadic PTSD interventions may be particularly well-suited to enhancing the development of PTG in individuals with PTSD. CBCT for PTSD (Monson & Fredman, 2012) is a trauma-focused conjoint therapy that was developed to both treat PTSD symptoms and enhance relationship functioning. CBCT for PTSD has a session specifically devoted to PTG in the protocol. Thus, the goals of the current study were to examine the effects of CBCT for PTSD on PTG and to examine the association of PTG with pre-treatment relationship satisfaction (to determine whether satisfaction influences growth) with data from a waitlist (WL) controlled trial. An additional exploratory goal was to examine PTG's association with pre-treatment PTSD symptomatology. We hypothesized that (a) PTG would increase more with treatment than without (WL); (b) pre-treatment relationship satisfaction would moderate the trajectory of PTG over the course of treatment, such that higher pre-treatment relationship satisfaction would be positively associated with PTG outcomes; (c) PTG outcomes would not be moderated by pre-treatment PTSD symptomatology because the literature suggests that PTSD symptoms and PTG are not necessarily associated (Calhoun & Tedeschi, 2014).

Method

Participants and Procedure

Participants were 40 intimate couples who were randomized to immediate CBCT for PTSD or a 3-month WL (see Monson et al., 2012 for further details (107 potential couples were self or clinician-referred, with the most common reason for not participating being electing to not give consent ($n = 31$ couples)), and a full description of demographic characteristics). Participants with PTSD had a mean age of 37.10 years ($SD = 11.26$), 30 (75.0%) were female, 29 (72.5%) identified as Non-Hispanic White, 3 (7.5%) as Black, 2 (5.0%) as Hispanic, and 6 (15.0%) as other ethnicity or race. Thirteen (32.5%) were married, and the average length of relationship was 6.85 years ($SD = 7.50$). The majority were employed ($n =$

24, 60%), mean years of education was 15.80 ($SD = 2.70$, Median = 16.00, Range = 7 to 21), and median household income was between \$35,000 and \$49,999 per year. Participants were recruited from a U.S. Veteran's Affairs outpatient clinic in Boston, MA, and a university-based research center in Toronto, ON. No site differences existed on the study variables. To be eligible for participation, both partners had to be between 18 and 75 years of age and if on psychotropic medication, to have been on a stable regimen for at least two months. One partner in each couple had to meet diagnostic criteria for current PTSD according to the Clinician-Administered PTSD Scale (CAPS; Blake et al., 1995). Trauma experiences were heterogeneous, including interpersonal (e.g., sexual assault), combat, and other experiences. Exclusion criteria included imminent suicidal, homicidal, or self-injurious behavior within the previous 6 months, meeting diagnostic criteria for current substance dependence, having an uncontrolled psychotic or bipolar disorder as assessed by the Structured Clinical Interview DSM-IV-Patient Version (SCID-P; First, Spitzer, Gibbon, & Williams, 2002), both partners meeting criteria for current PTSD, or endorsement of severe intimate partner sexual or physical aggression in the past 12 months.

Ethics approval was received from Ryerson University, the VA Boston Healthcare System, and the Boston University School of Medicine. Participants were recruited via posters and postings online and were screened by telephone. Interested participants provided written informed consent in person, and their eligibility was subsequently assessed. Master's and doctoral-level clinicians who were blind to study condition conducted the assessments, which took on average 2-3 hours to complete. Assessments were conducted at baseline (T1), mid-treatment (after session 7 of treatment) (T2), and post-treatment (within one month of ending treatment) (T3) for the treatment immediately group. Assessments were conducted at baseline (T1), 4 weeks waiting (T2), and 12 weeks waiting (T3) (which then served as their pre-treatment assessment) for WL participants. WL participants then went through treatment, and had an additional assessment at the end of 15 sessions of treatment (WL post-treatment assessment). Participants received \$25 compensation following each assessment.

Cognitive-Behavioral Conjoint Therapy for PTSD (CBCT) is a three phase, 15-session, manualized, trauma-focused conjoint therapy designed to simultaneously reduce PTSD symptoms and enhance intimate relationship functioning (Monson & Fredman, 2012). In the first phase, the treatment protocol and rationale are presented, and psychoeducation about PTSD symptoms and relationships is provided. The second phase is focused on improving communication, decreasing emotional numbing and avoidance associated with PTSD, and increasing relationship satisfaction. The third phase is focused on making meaning of the traumatic event by re-evaluating the trauma to arrive at an adaptive interpretation of the event. In addition, the treatment involves the completion of assignments between sessions to facilitate skill acquisition. Session 14 of the protocol specifically addresses PTG and barriers to PTG, including thoughts such as, "If I experience positive outcomes after the trauma, then what I experienced was not actually traumatic." Growth and change following the trauma, both individually and as a couple, are discussed, and both partners' beliefs are examined using a dyadic cognitive process. Sessions are conducted twice per week in Phases 1 and 2 and weekly thereafter. Participants who were unable to attend sessions twice per week in Phases 1 and 2 were seen weekly throughout treatment.

Measures

Current PTSD diagnostic status and symptom severity were determined using the CAPS (Blake et al., 1995). Ten percent of the CAPS conducted at each site in this study were randomly sampled and evaluated by an independent reliability monitor, yielding an intraclass correlation of .99 for total symptom severity and a κ of 1.00 for diagnostic status. The SCID-P (First et al., 2002) was used to assess DSM-IV Axis 1 disorders other than PTSD and to determine exclusion criteria. The reliability of the SCID-P diagnoses was excellent ($\kappa = .71-1.00$) across all conditions except mood disorders, where reliability was moderate ($\kappa = .60$).

The Post-Traumatic Growth Inventory (PTGI; Tedeschi & Calhoun, 1996) was completed by the patient with PTSD. The PTGI is a 21-item self-report measure of posttraumatic growth in the following five domains: relations to others, perceptions of new possibilities, personal strength, spiritual change, and appreciation of life. Each item is scored on a 6-point scale, assessing currently, ranging from I did not experience this change as a result of my crisis to I experienced this change to a very great degree as a result of my crisis. The PTGI demonstrated good internal consistency in this sample across pre-, mid- and post-treatment assessment points ($\alpha = .94, .93$ and $.94$, respectively).

The Dyadic Adjustment Scale (DAS; Spanier, 1976), a 32-item self-report measure of relationship quality in romantic dyads, was used to assess relationship satisfaction. Scores on the DAS can range from 0 to 151; a score of 98 or higher is considered satisfied. The internal consistency in this study was $\alpha = .96$.

Data Analysis

Intention-to-treat principles were used in conducting the analyses. Hierarchical linear modeling (HLM; Raudenbush & Bryk, 2002) was used, with PTG as the outcome variable and time, condition, and the time by condition interaction as independent variables to test the first hypothesis. The first analysis used baseline (T1), mid-treatment (treatment immediately)/4 weeks of waiting (WL) (T2), and post-treatment (treatment immediately)/12 weeks of waiting (WL) assessment time points (T3). The groups did not differ at baseline on any of the key variables (CAPS, DAS or PTGI). To test the potential for pre-treatment relationship satisfaction and PTSD symptoms to moderate PTG outcomes, all participants who were offered therapy (immediately or after waiting) were collapsed into a single sample ($n = 37$ due to missing data for one participant on baseline DAS and CAPS and two participants on PTGI). For participants in the immediate treatment condition, assessments during treatment were analyzed for pre-, mid- and post-treatment and for the waitlist condition assessments were analyzed for pre- and post-treatment, as a mid-treatment assessment was not conducted for PTG. HLM models were then conducted with pre-treatment DAS and CAPS scores predicting PTG trajectory, respectively. Hedge's g (Hedges & Olkin, 1985), a measure of effect size, was calculated and interpreted according to Cohen's (1988) descriptions of 0.20 small, 0.50 medium, and over 0.80 large.

Results

Mean and standard deviations were calculated for the PTGI, CAPS, and DAS at each time point (see Table 1). Scores demonstrated that, on average, patients were relatively satisfied in their relationships across the study period and were experiencing moderately severe PTSD symptoms at baseline. The interaction between time and treatment condition was significant in the first model. Compared with WL, individuals receiving CBCT for PTSD demonstrated a significant increase in PTG over time, with a between-group moderate effect size difference ($g = 0.45$; see Table 2 and Figure 1). Additionally, the HLM model predicting PTG over time based on pre-treatment DAS revealed a positive, albeit not significant, association between baseline DAS and increasing PTG and was of a moderate size ($g = 0.65$; see Table 1). The HLM model predicting PTG over time as a function of pre-treatment CAPS was not significant.

Discussion

The results of this study reveal that CBCT for PTSD facilitates PTG despite the relatively brief focus on PTG in this intervention. Participation in the intervention led to a moderate effect size increase in PTG scores. Consistent with prior research and theory, the improvements found in PTG may be associated with the interpersonal nature of this conjoint intervention, along with reducing avoidance and the process of meaning-making associated with trauma-focused work. Previous research on individual CBT for PTSD found a small effect size for change in PTG (Zoellner et al., 2011), compared to the moderate effect size in the current study. These findings suggest that the presence of supportive intimate relationships in trauma-focused interventions, as well as the explicit focus on PTG in treatment, may help foster growth. Previous research has found that presence of a relationship is not directly related to PTG; rather, it is the quality of relationships that seems to be of greater value (Barsakova & Oesterreich, 2009). It is possible that discussing the trauma together, as well as conjointly addressing barriers to PTG and ways that the participants may have grown individually and as a couple as a result of the trauma and its treatment, in the context of the therapeutic relationship promoted PTG. Findings from this study were in line with previous research suggesting that treatment that focuses on making meaning of the trauma and modifying maladaptive beliefs about events, behaviors, or symptoms are critical for the development of PTG (Tedeschi, 1999). The finding that pre-treatment PTSD symptoms were not associated with PTG is consistent with previous research suggesting that symptoms and growth seem to be independent constructs (e.g., Calhoun & Tedeschi, 2014).

There are several limitations of this study. First, the sample size used in this study was small and were relatively relationally satisfied at baseline; these findings should be replicated in larger and more distressed samples, including those with greater racial/ethnic and gender diversity. Nevertheless, PTG significantly increased, with a moderate effect size. Additionally, only one member of the dyad had PTSD and completed the measure of PTG. Future work should include additional measures of this construct to be completed by both members of the dyad and in dual PTSD cases, should examine whether low versus high pre-treatment relationship satisfaction moderates the relationship between time and treatment,

and should include longitudinal assessment of PTG at long-term follow-up to determine if gains are maintained, as previous research indicates that posttraumatic growth has been maintained over time (e.g., Linley & Joseph, 2004). These limitations notwithstanding, these initial findings suggest that trauma-focused couple therapy can help promote PTG. Further targeting PTG in the context of dyadic PTSD treatment should be considered in the future in order to capitalize on the fertile ground of intimate relationships to promote growth.

Acknowledgments

This research was supported in part by a grant from the National Institute of Mental Health to Candice M. Monson (R34 MH076813). We gratefully acknowledge Philippe Shnaider for his assistance with data analysis.

References

- Barskova T, Oesterreich R. Post-traumatic growth in people living with a serious medical condition and its relations to physical and mental health: A systematic review. *Disability & Rehabilitation*. 2009; 31:1709–1733. DOI: 10.1080/09638280902738441 [PubMed: 19350430]
- Blake DD, Weathers FW, Nagy LM, Kaloupek DG, Gusman FD, Charney DS, Keane TM. The development of a clinician-administered PTSD scale. *Journal of Traumatic Stress*. 1995; 8:75–90. DOI: 10.1002/jts.2490080106 [PubMed: 7712061]
- Calhoun, LG.; Tedeschi, RG. The foundations of posttraumatic growth: An expanded framework. In: Calhoun, LG.; Tedeschi, RG., editors. *Handbook of posttraumatic growth: Research and practice*. New York, NY: Lawrence Erlbaum Associates; 2014. p. 1-23.
- Cohen, J. *Statistical power analysis for the behavioral sciences*. 2nd. Hillsdale, NJ: Lawrence Erlbaum Associates; 1988.
- Dekel S, Mandl C, Solomon Z. Shared and unique predictors of posttraumatic growth and distress. *Journal of Clinical Psychology*. 2011; 67:241–252. DOI: 10.1002/jclp.20747 [PubMed: 21254052]
- First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. *Structured Clinical Interview for DSM-IV Axis I Disorders, Research Version, Patient Edition (SCID-I/P)*. New York: Biometrics Research Dept, New York State Psychiatric Institute; 2002.
- Hedges, LV.; Olkin, I. *Statistical methods for meta-analysis*. London, England: Academic Press; 1985.
- Linley PA, Joseph S. Positive change following trauma and adversity: A review. *Journal of Traumatic Stress*. 2004; 17:11–21. DOI: 10.1023/B:JOTS.0000014671.27856.7 [PubMed: 15027788]
- Monson CM, Fredman SJ, Macdonald A, Pukay-Martin ND, Resick PA, Schnurr PP. Effects of a cognitive-behavioral couple therapy for PTSD. *Journal of the American Medical Association*. 2012; 308:700–709. DOI: 10.1001/jama.2012.9307 [PubMed: 22893167]
- Monson, CM.; Fredman, SJ. *Cognitive-behavioral conjoint therapy for posttraumatic stress disorder: Therapist's manual*. New York, NY: Guilford Press; 2012.
- Prati G, Pietrantonio L. Optimism, social support, and coping strategies as factors contributing to posttraumatic growth: A meta-analysis. *Journal of Loss and Trauma*. 2009; 14:364–388. DOI: 10.1080/15325020902724271
- Raudenbush, SW.; Bryk, AS. *Hierarchical Linear Models*. Second. Thousand Oaks: Sage; 2002.
- Spanier GB. Measuring dyadic adjustment: New scales for assessing the quality of marriage and similar dyads. *Journal of Marriage and the Family*. 1976; 38:15–28. DOI: 10.2307/350547
- Tedeschi RG. Violence transformed: Posttraumatic growth in survivors and their societies. *Aggression and Violent Behavior*. 1999; 4:319–341. DOI: 10.1016/S1359-1789(98)00005-6
- Tedeschi RG, Calhoun LG. The Posttraumatic Growth Inventory: Measuring the positive legacy of trauma. *Journal of Traumatic Stress*. 1996; 9:455–471. DOI: 10.1002/jts.2490090305 [PubMed: 8827649]
- Tedeschi RG, Calhoun LG. Posttraumatic growth: Conceptual foundations and empirical evidence. *Psychological Inquiry*. 2004; 15:1–18. DOI: 10.1207/s15327965pli1501_01

Zoellner T, Rabe S, Karl A, Maercker A. Post-traumatic growth as outcome of a cognitive-behavioural therapy trial for motor vehicle accident survivors with PTSD. *Psychology and Psychotherapy: Theory, Research and Practice*. 2011; 84:201–213. DOI: 10.1348/147608310X520157

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.

Change in posttraumatic growth as a function of treatment group, cognitive-behavioral conjoint therapy for posttraumatic stress disorder (CBCT for PTSD) versus waitlist (WL) assessed at three time points, baseline (T1), mid-treatment/4-week wait (T2), and post-treatment/12-week wait (T3) ($n = 40$).

Table 1
Means and Standard Deviations by Time Separately for Treatment Condition

Variable	Baseline			Mid-Tx/4wkwait			Post-Tx/12wkwait			Post-Tx WL		
	M	SD	Range	M	SD	Range	M	SD	Range	M	SD	Range
				Immediate CBCT								
PTGI	46.88	25.14	2-104	51.33	24.25	6-100	60.80	27.15	7-105	-	-	-
CAPS	69.45	12.74	45-92	50.72	21.44	4-88	32.31	24.81	0-80	-	-	-
DAS	101.29	22.25	25-124	102.71	17.74	64-130	117.15	10.21	98-131	-	-	-
				Waitlist								
PTGI	43.47	24.13	0-82	44.37	24.24	0-83	46.37	27.55	0-99	53.00	25.52	10-91
CAPS	74.80	14.96	54-114	63.72	23.58	19-111	63.11	26.02	27-116	35.00	23.46	3-68
DAS	98.12	20.78	50-127	98.34	21.66	43-126	101.77	20.41	43-132	106.59	24.80	52-138

Note. Mid-Tx/4wkwait = Mid-Treatment (immediate treatment group) or 4 weeks of waiting (WL group); Post-Tx/12wkwait = Post-Treatment (treatment immediately group) or 12 weeks of waiting (WL group); Post-Tx WL = Post-Treatment (WL group); PTGI = Posttraumatic Growth Inventory; CAPS = Clinician Administered PTSD Scale; DAS = Dyadic Adjustment Scale. Sample sizes vary due to participants not completing all measures at each time point. For Immediate CBCT: PTGI (Baseline $n = 17$, Mid-Tx $n = 18$, Post-Tx $n = 15$); CAPS (Baseline $n = 20$, Mid-Tx $n = 18$, Post-Tx $n = 16$); DAS (Baseline $n = 20$, Mid-Tx $n = 17$, Post-Tx $n = 13$). For Waitlist: PTGI (Baseline $n = 19$, 4 week wait $n = 17$, 12 week wait $n = 19$, Post-Tx $n = 14$); CAPS (Baseline $n = 20$, 4 week wait $n = 18$, 12 week wait $n = 19$, Post-Tx $n = 14$); DAS (Baseline $n = 20$, 4 week wait $n = 18$, 12 week wait $n = 19$, Post-Tx $n = 14$).

Table 2
Two-level HLM Models (Time Nested Within Individual) Predicting PTGI

Variable	Coefficient	SE	<i>t</i>	<i>df</i>
Waitlist versus treatment immediately				
Intercept	45.19	5.27	8.57**	42.1
Time	0.31	1.74	0.18	36.3
Condition	0.06	7.58	0.01	42.6
Time × Condition	6.24	2.60	2.40*	38.8
Pre-treatment DAS				
Intercept	65.32	14.08	4.64**	36
DAS	-0.19	0.14	-1.41	36
Time	-8.90	6.94	-1.28	36
Time × DAS	0.14	0.07	2.01	36

Note. For the time x condition interaction, Hedge's *g* between group effect size = 0.45. For the time × DAS interaction, Hedge's *g* effect size = 0.65. PTGI = Posttraumatic Growth Inventory; DAS = Dyadic Adjustment Scale.

*
 $p < .05$.

**
 $p < .01$.