Successful use of dexamethasone when prednisolone is not tolerated

We have read with interest the article by Al-Ozairi *et al.*¹ in the March issue of the journal emphasizing the value and safety of prednisolone in the management of hyperemesis gravidarum. We recently cared for a woman in whom prednisolone failed to control vomiting because of drug intolerance but parenteral dexamethasone succeeded, even as an outpatient. Furthermore, her ketosis-prone type 1 diabetes mellitus was not aggravated by her dexamethasone therapy.

The 23-year-old primigravid patient with type 1 diabetes and controlled Grave's disease for nine years, managed with insulin aspart thrice daily, insulin glargine nocte and neomercazole, developed progressively severe vomiting from five weeks gestation. Despite a range of treatments including B vitamins, ondansetron, promethazine, prochlorperazine and intermittent trials of prednisolone, she lost 14% of her prepregnant body weight, reaching a nadir of 51 kg by 13 weeks. Four hospital admissions for rehydration, intravenous hydrocortisone and insulin infusion for diabetic ketoacidosis occurred over this period, due to a combination of hyperemesis and patient compliance issues with self-discharges. At 13 weeks as an ambulatory patient by her choice, a further attempt of oral prednisolone 25 mg twice daily was unsuccessful, leading to a trial of intramuscular dexamethasone 4 mg daily for four days followed by oral administration for a further week. This dramatically arrested her vomiting and allowed a tapering low dose of prednisolone (initially 10 mg daily) to be re-introduced, ceasing at 16 weeks with only residual occasional nausea. Her glycaemic control stabilized with HbA1c being 7.7% at 10 weeks and 7.4% at 20 weeks. Body weight progressively increased, albeit suboptimally, almost reaching her prepregnant weight by 29 weeks. She remained biochemically euthyroid throughout the pregnancy. A healthy female infant (3210 g) was delivered vaginally at 36 weeks after premature rupture of membranes.

Although there was a hesitancy to use dexamethasone, given its placental transfer with incomplete inactivation² and unknown teratogenic risk, it was demonstrated here to be very useful and apparently safe but it would be prudent to restrict its use to after eight weeks gestation when lip fusion has been completed. We speculate that its superior effect here may be due to its initial parenteral use, greater potency, prolonged action and possibly better blood-brain barrier penetration, given its likely medullary site of action demonstrated experimentally in cancer chemotherapy-induced emesis.

V J Heazlewood*, M Ratnapala† and L Cochrane†

*Caboolture Hospital – Medicine;

†Caboolture Hospital – Obstetrics and Gynaecology, Caboolture, Queensland, Australia

Correspondence to: Dr V J Heazlewood

Email: vernon_heazlewood@health.qld.gov.au

DOI: 10.1258/om.2009.090026

REFERENCES

- 1 Al-Ozairi E, Waugh JJS, Taylor R. Termination is not the treatment of choice for severe hyperemesis gravidarum: successful management using prednisolone. *Obstet Med* 2009;2:34–7
- 2 Diederich S, Eigendorff E, Burkhardt P *et al.* 11-beta-hydroxysteroid dehydrogenase types 1 and 2: an important pharmacokinetic determinant for the activity of synthetic mineralo- and glucocorticoids. *J Clin Endocrinol Metab* 2002;87:5695–701

Intrauterine growth rates in diabetic pregnancies and clinical outcomes

In Lim *et al.*'s recent article, 'Intrauterine growth rate in pregnancies complicated by type 1, type 2 and gestational diabetes',¹ it was interesting to note the similar growth rates and rates of macrosomia in the three types of diabetes. From a clinical point of view in diabetes we are interested in predictors of adverse outcomes such as shoulder dystocia, birth trauma and stillbirth. Although the data set may not be large enough to look at growth curves for some of these outcomes it would be interesting to examine whether the growth curves for these outcomes were different from those for pregnancies with good outcomes. Such information may then help us identify those diabetic pregnancies that are at even higher risk and adjust care accordingly.

Michael Peek and Ralph Nanan

University of Sydney, Nepean Clinical School, Level 5, South Block Nepean Hospital, Penrith, New South Wales, 2750 Australia

Email: mpeek@med.usyd.edu.au

DOI: 10.1258/om.2009.090011

REFERENCE

- 1 Lim EL, Burden T, Marshall SM, *et al.* Intrauterine growth rate in pregnancies complicated by type 1, type 2 and gestational diabetes. *Obstet Med* 2009;2:21–5