Published in final edited form as:

Clin Imaging. 2016 ; 40(2): 228–231. doi:10.1016/j.clinimag.2015.11.001.

Pancreatic Neuroendocrine Tumor with Aneurysms of the Gastroduodenal Artery: a Case Report

Karim Rebeiz¹, Ali Shamseddine¹, Ralph Hachem¹, Karen Brown^{2,3}, Deborah Mukherji¹, Walid Faraj¹, Ghassan K. Abou-Alfa^{2,3}, Layla A. Nasr¹, and Ali A Haydar¹

¹American University of Beirut Medical Center, Beirut, Lebanon

²Memorial SloanKettering Cancer Center, New York, NY, USA

³Weill Cornell Medical College, New York, NY, USA

Abstract

Gastroduodenal artery (GDA) aneurysm is a very rare condition. It is divided into false and true aneurysms where pseudoaneurysms are associated with pancreatitis and true aneurysms are secondary to celiac trunk stenosis. We report a 24 year old female patient who was diagnosed with pancreatic head neuroendocrine tumor and was incidentally found to have multiple GDA aneurysms in the absence of celiac artery stenosis. A decision to embolize the aneurysms was taken in view of the presumed high risk of bleeding. The procedure was successful with a follow up CT scan showing no recurrence.

Keywords

Aneurysm; Embolization; Vascular intervention

Introduction

The gastroduodenal artery (GDA) typically arises from the common hepatic artery of the celiac trunk and supplies blood to the proximal part of the duodenum, pylorus and pancreatic head. It first branches into the supraduodenal artery, followed by the posterior superior pancreaticoduodenal artery and terminates as the right gastroepiploic artery and the anterior superior pancreaticoduodenal artery. These branches form anastomoses with the superior mesenteric artery (SMA) through the anterior and posterior pancreaticoduodenal arcades providing an important source of collateral circulation.

Corresponding Author: Ali Haydar, MD, American University of Beirut Medical Centre, Beirut, Lebanon, Tel: 00 961 1 350000 ext 5027, ah24@aub.edu.lb.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Ethical Approval: For this type of study formal consent is not required.

Informed Consent: Informed consent was obtained from the patient participating in this study.

Conflict of Interest Statement: The authors of this study declare no conflict of interest.

Funding: No funding was required for this study.

GDA aneurysm is uncommon (1) comprising only 1.5% of all reported visceral artery aneurysms (VAA) (2). The majority of those are pseudoaneurysms occurring in the setting of pancreatitis (3). True aneurysms are most commonly associated with middle aged men (4) and atheromatous celiac trunk stenosis (5). We report a 24 year old woman, presenting with a pancreatic head neuroendocrine tumor and multiple GDA aneurysms in the absence of celiac artery stenosis.

Case report

A 24 year old female patient presented to our institution with severe epigastric pain. She was diagnosed with a pancreatic head neuroendocrine tumor.

Laboratory investigation revealed hypercalcemia with a calcium level of 20.7 mg/dL and aspartate aminotransferase (AST) and alanine aminotransferase (ALT) around 130–140 IU/L (normal ranges of ALT (7–55 IU/L), AST (8–48 IU/L)).

Computed tomography (CT) angiography of the abdomen showed a heterogeneously enhancing pancreatic head mass with areas of necrosis and calcification and multiple arterial aneurysmal dilatations in the pancreaticoduodenal arcade, the largest of which measured 1.5, cm arising from the anterior superior pancreaticoduodenal artery with peripheral calcification. Multiple metastases to liver and bone were also identified. There were no CT findings suggestive of pancreatitis, or celiac artery stenosis (Figure 1)

Digital subtraction arteriogram (DSA) after embolization of the right hepatic artery for treatment of hepatic metastases showed multiple gastroduodenal arcade aneurysms (Figure 2)

A decision to coil the aneurysms was taken in order to decrease the risk of rupture and bleeding. The pancreaticoduodenal artery was catheterised and angiography confirmed multiple aneurysms in the pancreaticoduodenal arcade. These aneurysms were excluded from the circulation using multiple microcoils and microparticles.

Final angiogram post-embolization did not show any further filling of the aneurysms. (Figure 3) Follow up CT angiography 5 days post-procedure confirmed absence of flow in the aneurysms.

Discussion

GDA aneurysms are divided into true and false aneurysm, the latter being more common and frequently seen in the setting of acute and chronic pancreatitis. Periarterial inflammation caused by pancreatic proteolytic enzymes secondary to pancreatitis is an important cause of GDA pseudoaneurysms (6). Atherosclerosis, congenital absence of celiac axis and autoimmune diseases like Systemic lupus erythematosus, Wegener's granulomatosis and polyarteritis nodosa are possible etiologies of true GDA aneurysms (7).

The pancreaticoduodenal arcade is the main collateral pathway between the celiac axis and the superior mesenteric artery. Increased blood flow in the this arcade, as compensation for celiac artery stenosis, is associated with pancreaticoduodenal artery aneurysm (8).

The same theory suggests that occlusion or stenosis of the superior mesenteric artery or celiac axis could predispose to the formation of a gastroduodenal artery aneurysm (9). After researching online databases, including PUBMED and Medline, we did not find any reports of pancreatic head neuroendocrine tumor associated with GDA aneurysms. It is possible that in this case, the presence of the hypervascular neuroendocrine tumor induced an increased flow through the GDA arcade, leading to aneurysms formation.

Another possible etiology is the tumor induced neovascularization. Pancreatic neuroendocrine tumors exhibit a high degree of vascularization (10) and are also known to secrete many proangiogenic molecules, in particular vascular endothelial growth factor (VEGF) (11). This could lead to the formation of numerous friable vessels prone to aneurysmal dilatation secondary to increased blood flow. A third possible etiology is that the tumor may have an aneurysmal component secondary to its histological origin. This is similar to an angiomyolipoma which derives from perivascular epithelioid cells. It contains fat, smooth muscles and vessels, the latter of which develop aneurysms. Even though there is a possibility of a potential common pathogenesis of neuroendocrine tumors and aneurysms, to our knowledge, we found no evidence linking the two pathologies in the literature.

The gold standard diagnostic test is visceral angiography, serving both diagnostic and therapeutic purposes by delineating the arterial anatomy and allowing therapeutic intervention (12). It has the highest sensitivity (100%) followed by computed tomography (CT) (67%) (12).

Ruptured GDA aneurysm may lead to death in up to 40% of cases (13). Open surgical and endovascular techniques are available to treat GDA aneurysms. Endovascular techniques are more often used to treat visceral artery aneurysms with very high success rate and low recurrence, morbidity and mortality (1). Endovascular treatment aims at excluding aneurysms from the parent circulation. Treatment options include: coiling (such as in our patient), stent graft placement or balloon occlusion, and embolization with liquid agents such as glue, onyx, or thrombin (14). We found no evidence that favors one method over the other. However, the use of stents and balloon occlusion in a tortuous and tapered artery is challenging (15). Also, when injecting liquid agents, there is possibility of reflux into the patent parent arteries causing thrombosis (15). For these reasons, and given the complexity of this case and the size of the tumor, decision favored endovascular coiling. The procedure was successful with complete occlusion of the aneurysms.

Conclusion

GDA aneurysms are uncommon and there are no known associations between GDA aneurysms and pancreatic neuroendocrine tumors. We describe a patient with PNET with multiple aneurysms successfully treated with endovascular therapy and present possible flow and tumor related etiologies for this unusual presentation.

References

1. Fankhauser GT, Stone WM, Naidu SG, Oderich GS, Ricotta JJ, Bjarnason H, et al. The minimally invasive management of visceral artery aneurysms and pseudoaneurysms. *Journal of vascular surgery*. 2011; 53(4):966–970. [PubMed: 21216559]
2. Carr SC, Mahvi DM, Hoch JR, Archer CW, Turnipseed WD. Visceral artery aneurysm rupture. *Journal of vascular surgery*. 2001; 33(4):806–811. [PubMed: 11296336]
3. White AF, Baum S, Buranasiri S. Aneurysms secondary to pancreatitis. *AJR American journal of roentgenology*. 1976; 127(3):393–396. [PubMed: 183522]
4. Morita Y, Kawamura N, Saito H, Shinohara M, Irie G, Okushiba S, et al. [Diagnosis and embolotherapy of aneurysm of the gastroduodenal artery]. *Rinsho hoshasen Clinical radiography*. 1988; 33(5):555–561. [PubMed: 3418920]
5. Tori M, Nakahara M, Akamatsu H, Ueshima S, Shimizu M, Nakao K. Significance of intraoperative monitoring of arterial blood flow velocity and hepatic venous oxygen saturation for performing minimally invasive surgery in a patient with multiple calcified pancreaticoduodenal aneurysms with celiac artery occlusion. *Journal of hepato-biliary-pancreatic surgery*. 2006; 13(5):472–476. [PubMed: 17013726]
6. Bergert H, Hinterseher I, Kersting S, Leonhardt J, Bloomenthal A, Saeger HD. Management and outcome of hemorrhage due to arterial pseudoaneurysms in pancreatitis. *Surgery*. 2005; 137(3):323–328. [PubMed: 15746787]
7. Pasha SF, Gloviczki P, Stanson AW, Kamath PS. Splanchnic artery aneurysms. *Mayo Clinic proceedings*. 2007; 82(4):472–479. [PubMed: 17418076]
8. Quandalle P, Chambon JP, Marache P, Saudemont A, Maes B. Pancreaticoduodenal artery aneurysms associated with celiac axis stenosis: report of two cases and review of the literature. *Annals of vascular surgery*. 1990; 4(6):540–545. [PubMed: 2261321]
9. Schweizer W, Gries NC, Maddern G, Triller J. Splenic infarction complicating ligation of a gastroduodenal artery aneurysm. *Digestive surgery*. 1999; 16(3):236–237. [PubMed: 10436373]
10. Terris B, Scoazec JY, Rubbia L, Bregeaud L, Pepper MS, Ruzniewski P, et al. Expression of vascular endothelial growth factor in digestive neuroendocrine tumours. *Histopathology*. 1998; 32(2):133–138. [PubMed: 9543669]
11. Missiaglia E, Dalai I, Barbi S, Beghelli S, Falconi M, della Peruta M, et al. Pancreatic endocrine tumors: expression profiling evidences a role for AKT-mTOR pathway. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2010; 28(2):245–255. [PubMed: 19917848]
12. Yeh TS, Jan YY, Jeng LB, Hwang TL, Wang CS, Chen MF. Massive extra-enteric gastrointestinal hemorrhage secondary to splanchnic artery aneurysms. *Hepato-gastroenterology*. 1997; 44(16): 1152–1156. [PubMed: 9261616]
13. Lee PC, Rhee RY, Gordon RY, Fung JJ, Webster MW. Management of splenic artery aneurysms: the significance of portal and essential hypertension. *Journal of the American College of Surgeons*. 1999; 189(5):483–490. [PubMed: 10549737]
14. Chadha M, Ahuja C. Visceral artery aneurysms: diagnosis and percutaneous management. *Seminars in Interventional Radiology*. 2009; 26(3):196–206. ISSN 0739-9529. [PubMed: 21326564]
15. Moses V, Keshava SN, Chiramel GK, Mammen S, Narayanam SRS. Endovascular management of splanchnic aneurysms. *Vascular Disease Management*. 2013; 10(1):E5–E10.

a.

b.

Figure 1.
1A CT angiography showing two aneurysms (straight arrows) within the pancreatic head, one of which is partially calcified **1B.** Axial CT in the arterial phase showing a heterogeneously enhancing mass (arrow) in the head of the pancreas compatible with neuroendocrine tumor

A

B and C

D

Figure 2.

2A Digital subtraction arteriogram of the celiac artery after right hepatic artery embolization showing an enlarged GDA origin (arrow). **2B** and **C** Early and late GDA selective angiography showing multiple aneurysms (arrows). **2D** Selective angiography distal to the aneurysms, showing a normal caliber gastroepiploic artery.

A

B

Figure 3.
3A Mid treatment angiogram shows coils in the distal gastroduodenal and gastroepiploic arteries (Arrows). 3B Post treatment angiogram showing no flow within the aneurysms.