Published in final edited form as:

Biomol NMR Assign. 2009 December ; 3(2): 191–194. doi:10.1007/s12104-009-9172-6.

¹H, ¹³C, and ¹⁵N NMR assignments for the *Bacillus subtilis* yndB START domain

Kelly A. Mercier^{1,2}, Geoffrey A. Mueller³, Thomas B. Acton⁴, Rong Xiao⁴, Gaetano T. Montelione⁴, and Robert Powers^{1,*}

¹Department of Chemistry, University of Nebraska-Lincoln, Lincoln, NE 68588-0304

³Laboratory of Structural Biology, National Institute of Environmental Health Sciences, 111 T. W. Alexander Drive, Durham, NC, 27713

⁴Center for Advanced Biotechnology and Medicine, Department of Molecular Biology and Biochemistry and Northeast Structural Genomics Consortium, Rutgers University, Piscataway, NJ 08854, USA

Abstract

The steroidogenic acute regulatory-related lipid transfer (START) domain is found in both eukaryotes and prokaryotes, with putative functions including signal transduction, transcriptional regulation, GTPase activation and thioester hydrolysis. Here we report the near complete ¹H, ¹⁵N and ¹³C backbone and side chain NMR resonance assignments for the *Bacillus subtilis* START domain protein yndB.

Keywords

Bacillus subtilis; protein yndB; START domain; lipid binding; NMR; resonance assignments

Biological context

Bacillus subtilis protein yndB is a 16.1 kDa monomer annotated as a ‘conserved hypothetical protein’ in major protein databases: SWISS-PROT (YND_BACSU, O31806), PIR (C69889) and GENBANK (NP_389656). Yndb is a structural target (NESG_ID SR211) from the Northeast Structural Genomic Consortium (NESG; www.nesg.org).

The solution structures for two related functionally uncharacterized proteins, *Bacillus cereus* protein BC4709 (PDB ID: 1xn6) and *Bacillus halodurans* protein BH1534 (PDB ID: 1xn5), were previously solved by NMR spectroscopy (Liu, et al. 2005). The amino-acid sequences for BC4709 and BH1534 are 64% and 57% homologous to yndB, respectively. The BC4709 and BH1534 NMR structures were shown to belong to the steroidogenic acute regulatory-related lipid transfer (START) domain superfamily (CATH: 3.30.530.20). The START

*To whom correspondence should be addressed: Department of Chemistry, 722 Hamilton Hall, University of Nebraska-Lincoln, Lincoln, NE 68588, Tel: (402) 472-3073; Fax (402) 472-9402, rpowers3@unl.edu.

²Current address: Laboratory of Structural Biology, National Institute of Environmental Health Sciences, 111 T. W. Alexander Drive, Durham, NC, 27713

domain is found in both eukaryotes and prokaryotes, with putative functions that include signal transduction, transcriptional regulation, GTPase activation and thioester hydrolysis. Currently, there are 16 START domain proteins in the PDB, several of which contain bound lipid ranging in mass from 397.6 to 790.2 Da.

START domains exhibit a range of functions, but share a common fold and ligand-binding mode (Iyer, et al. 2001). The START domain superfamily binds a diverse array of ligands, so that the ligands that interact with a specific START domain are a major determinant of the protein's biological function. Sequence modifications contribute to variations in the size, depth and enzymatic activity of the ligand-binding pocket, that has led to an evolutionary divergence in ligand specificity and protein function among the START domain superfamily members. Currently, there is very little information in the literature describing ligand specificity and biological activity for the START domain superfamily. Thus, the experimental structure of *B. subtilis* protein yndB will be an important asset in our efforts to decipher the relationships between structure, function and ligand specificity within the START domain superfamily. By comparing the three-dimensional structures of available START domain proteins along with the determination of ligand binding profiles, functional classifications and ligand specificity may be inferred for other members of the START domain superfamily. In order to provide a basis for studies of the structure, ligand binding specificity, and function of *B. subtilis* protein yndB, we report the near complete ^1H , ^{15}N and ^{13}C backbone and side chain NMR resonance assignments for yndB.

Methods and Experiments

Expression and purification of yndB

Uniformly ^{13}C , ^{15}N -enriched yndB (152 amino acids) was cloned, expressed and purified following standard protocols used in the NESG consortium (Acton, et al. 2005). Briefly, the full length gene yndB or BSU17730 from *Bacillus subtilis* was cloned into a pET21d (Novagen) derivative, yielding the plasmid pSR211-21.2. The resulting yndB open reading frame contains eight nonnative residues at the C-terminus (LEHHHHHH) of the protein. *Escherichia coli* BL21 (DE3) pMGK cells, a rare codon enhanced strain, were transformed with pSR211-21.2, and cultured in MJ9 minimal medium containing $(^{15}\text{NH}_4)_2\text{SO}_4$ and $[\text{U-}^{13}\text{C}]\text{-D-glucose}$ as sole nitrogen and carbon sources (Jansson, et al. 1996). Initial growth was carried out at 37°C until the OD_{600} of the culture reached ~ 0.8 units. The incubation temperature was then decreased to 17°C and protein expression was induced by the addition of IPTG (isopropyl- β -D-thiogalactopyranoside) at a final concentration of 1 mM. Following overnight incubation at 17°C , the cells were harvested by centrifugation and lysed by sonication. $[\text{U-}^{13}\text{C}, ^{15}\text{N}]$ yndB was purified in a two-step protocol consisting of Ni-NTA affinity column (Qiagen) and gel filtration column (HiLoad 26/60 Superdex 75 pg, Amersham Biosciences) chromatography. These samples were determined to be largely ($> 98\%$) monomeric by analytical gel filtration (20 mM MES, 100 mM NaCl, 5 mM CaCl_2 , 10 mM DTT, pH6.5) followed by static light scattering measurements (as described in Acton et al, 2005). The final yield of pure $[\text{U-}^{13}\text{C}, ^{15}\text{N}]$ yndB (estimated $> 97\%$ by SDS-PAGE; 17993.4 Da by MALDI-TOF mass spectrometry) was approximately 32.4 mg from a 1 L culture.

NMR spectroscopy

Samples of [U- ^{13}C , ^{15}N] yndB for NMR spectroscopy were prepared at a protein concentration of 1.0 mM in 95% H_2O /5% D_2O solution containing 20 mM MES, 100 mM NaCl, 10 mM DTT, 5 mM CaCl_2 , 0.02% NaN_3 at pH 6.5 (uncorrected) in a sealed Shigemi tube (Shigemi Inc, Allison Park, PA).

All NMR experiments used for the protein backbone assignments of yndB were collected at 25°C on a 5 channel 600 MHz Bruker Avance spectrometer equipped with a 5 mm TXI probe. NMR experiments used for the protein side chain resonances were collected at the Pacific Northwest National Laboratory on a 4 channel 750 MHz Varian INOVA spectrometer equipped with a 5 mm HCN probe and at the National Institute of Environmental Health Sciences on a 600 MHz Varian INOVA spectrometer also equipped with a 5 mm HCN probe. Assignments of the backbone and side chain resonances were obtained from the following spectra: ^{15}N -HSQC, ^{13}C -HSQC, HNCO, HNCA, CBCACONH, CBCANH, HNHA, HBHACONH, C(CO)NH-TOCSY, H(CCO)NH-TOCSY, HCCH-TOCSY, and H(CCH)-COSY (Ferentz and Wagner 2000). The NMR data was processed using NMRpipe (Delaglio, et al. 1995) and analyzed using PIPP (Garrett, et al. 1991) and NMRViewJ (Johnson 2004).

Assignments and Data Deposition

Extensive assignments for the backbone and side-chain resonances of yndB were obtained (123/143 HN, 123/144 Ca, 140/168 Ha, 121/132 C β , 168/229 H β , 88/134 C γ , 33/75 C δ , 110/144 C', excluding the LEHHHHHH purification tag). NMR assignments are based on 3D heteronuclear experiments performed on the $^{13}\text{C}/^{15}\text{N}$ -labeled yndB sample using NMRView and NVAssign (Kirby, et al. 2004). Figure 1 shows an annotated 2D ^1H - ^{15}N HSQC spectrum of yndB. The terminal residues were not observed in the ^1H - ^{15}N HSQC spectrum. Many of the unassigned peaks in the ^1H - ^{15}N HSQC spectrum are near random-coil chemical shifts and likely residues in the loop regions. Several loops (based on sequence similarity prediction) were difficult to assign because of either missing resonances, degenerate shifts, or a lack on connectivity in the triple-resonance experiments.

The secondary structure determination of yndB was based on characteristic NOE data involving the HN, Ha and H β protons from ^{15}N -edited NOESY-HMQC and ^{13}C -edited NOESY-HMQC spectra, $^3\text{JHN}\alpha$ coupling constants from HNHA, slowly exchanging HN protons and ^{13}Ca and $^{13}\text{C}\beta$ secondary chemical shifts (Marion, et al. 1989, Wagner and Wuthrich 1986). The yndB NMR structure is composed of 6 β strands forming an anti-parallel β -sheet and 3 α -helices: 1 long and 2 short. Residues K37 to M40, H53 to S61 and K106 to E117 yielded limited assignments presumably because of local flexibility. The residues that are unassigned in these loops were: K37, W38, F39, M40, H53, L54, Q55, S56, P57, P60, S61, K106, E107, P108, N109, G113, K114, N116, and E117. Figure 2 shows the predicted secondary structure of yndB and the consensus secondary structure of START domain proteins. The unassigned residues correspond to the predicted loop regions of the START domain fold. The identification and addition of a ligand to the yndB NMR sample may improve the quality of the NMR data for these loop regions by reducing the inherent flexibility and inducing a defined conformation. The sequence specific ^1H , ^{13}C , and ^{15}N

resonance assignments of the yndB were deposited in the BioMagResBank under accession number 16043.

Acknowledgments

We thank Prof. Diana Murray for stimulating discussions regarding the NESG START domain project and Dr. John Cort at PNNL for providing the 750 MHz NMR data. This work was supported by a grant from the Protein Structure Initiative of the National Institutes of Health (U54 GM074958 to G.T.M.), Nebraska Tobacco Settlement Biomedical Research Development Fund and by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences. Research was performed in facilities renovated with support from the NIH under Grant RR015468-01. NMR spectra were acquired at the U.S. Department of Energy Office of Biological and Environmental Research) located at Pacific Northwest National Laboratory and operated for DOE by Battelle (contract KP130103).

References

- Acton TB, Gunsalus KC, Xiao R, Ma LC, Aramini J, Baran MC, Chiang YW, Climent T, Cooper B, Denissova NG, Douglas SM, Everett JK, Ho CK, Macapagal D, Rajan PK, Shastry R, Shih LY, Swapna GV, Wilson M, Wu M, Gerstein M, Inouye M, Hunt JF, Montelione GT. Robotic cloning and Protein Production Platform of the Northeast Structural Genomics Consortium. *Methods Enzymol.* 2005; 394:210–243. [PubMed: 15808222]
- Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A. NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR.* 1995; 6:277–293. [PubMed: 8520220]
- Ferentz AE, Wagner G. NMR spectroscopy: a multifaceted approach to macromolecular structure. *Q Rev Biophys.* 2000; 33:29–65. [PubMed: 11075388]
- Garrett DS, Powers R, Gronenborn AM, Clore GM. A Common Sense Approach to Peak Picking in Two-, Three-, and Four-Dimensional Spectra Using Automatic Computer Analysis of Contour Diagrams. *Journal of Magnetic Resonance.* 1991; 95:214–220.
- Iyer LM, Koonin EV, Aravind L. Adaptations of the helix-grip fold for ligand binding and catalysis in the START domain superfamily. *Proteins.* 2001; 43:134–144. [PubMed: 11276083]
- Jansson M, Li YC, Jendeborg L, Anderson S, Montelione BT, Nilsson B. High-level production of uniformly ^{15}N - and ^{13}C -enriched fusion proteins in *Escherichia coli*. *J Biomol NMR.* 1996; 7:131–141. [PubMed: 8616269]
- Johnson BA. Using NMRView to visualize and analyze the NMR spectra of macromolecules. *Methods Mol Biol.* 2004; 278:313–352. [PubMed: 15318002]
- Kirby NI, DeRose EF, London RE, Mueller GA. NvAssign: protein NMR spectral assignment with NMRView. *Bioinformatics.* 2004; 20:1201–1203. [PubMed: 14871875]
- Liu G, Shen Y, Atreya HS, Parish D, Shao Y, Sukumaran DK, Xiao R, Yee A, Lemak A, Bhattacharya A, Acton TA, Arrowsmith CH, Montelione GT, Szyperski T. NMR data collection and analysis protocol for high-throughput protein structure determination. *Proc Natl Acad Sci U S A.* 2005; 102:10487–10492. [PubMed: 16027363]
- Marion D, Driscoll PC, Kay LE, Wingfield PT, Bax A, Gronenborn AM, Clore GM. Overcoming the overlap problem in the assignment of ^1H NMR spectra of larger proteins by use of three-dimensional heteronuclear ^1H - ^{15}N Hartmann-Hahn-multiple quantum coherence and nuclear Overhauser-multiple quantum coherence spectroscopy: application to interleukin 1 beta. *Biochemistry.* 1989; 28:6150–6156. [PubMed: 2675964]
- McGuffin LJ, Bryson K, Jones DT. The PSIPRED protein structure prediction server. *Bioinformatics.* 2000; 16:404–405. [PubMed: 10869041]
- Wagner G, Wuthrich K. Observation of internal motility of proteins by nuclear magnetic resonance in solution. *Methods Enzymol.* 1986; 131:307–326. [PubMed: 3773764]
- Wishart DS, Sykes BD, Richards FM. Relationship between nuclear magnetic resonance chemical shift and protein secondary structure. *J Mol Biol.* 1991; 222:311–333. [PubMed: 1960729]

Fig. 1.
 2D ^1H - ^{15}N HSQC spectrum (A) and zoomed region (B) of uniformly ^{13}C , ^{15}N -labeled yndB in 1.0 mM in 95% H_2O /5% D_2O at pH 6.5 (uncorrected). The spectrum was collected on a Bruker Avance 600 MHz spectrometer equipped with a 5 mm TXI probe. The assignments are annotated by the sequence number.

Fig. 2.

The ^{15}N , $^{13}\text{C}\alpha$, and $^{13}\text{C}\beta$ chemical shift index (Wishart, et al. 1991) of yndB, where blue circles imply a β -strand region and red circles suggest an α -helix. The position of each circle along the bar indicates the relative consistency of the chemical shifts with either a β -sheet or α -helix. The lack of a circle indicates a missing NMR assignment. A consensus secondary structure (McGuffin, et al. 2000) for the START domain proteins is shown below the chemical shift index, where an arrow indicates a β -strand and a ribbon indicates an α -helix.