Published in final edited form as:

Am J Prev Med. 2016 September ; 51(3): 327–335. doi:10.1016/j.amepre.2016.04.021.

Prospective Changes in Healthy Lifestyle Among Midlife Women: When Psychological Symptoms Get in the Way

Claudia Trudel-Fitzgerald, PhD¹, Shelley S. Tworoger, PhD^{2,3}, Elizabeth M. Poole, PhD²,
David R. Williams, PhD, MPH¹, and Laura D. Kubzansky, PhD, MPH¹

¹Department of Social and Behavioral Sciences, Harvard T.H. Chan School of Public Health, Boston Massachusetts

²Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston Massachusetts

³Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts

Abstract

Introduction—Anxiety and depression are linked to increased risk of cardiometabolic disease and mortality, and unhealthy behaviors may be a key mechanism underlying these associations. Although higher levels of psychological symptoms are associated with individual unhealthy behaviors (e.g., physical inactivity, smoking), their role on overall lifestyle remains understudied.

Methods—Midlife women (N=55,395) from the cohort Nurses' Health Study reported anxiety and depression symptoms in 1988 and 1992, respectively. Health behaviors (i.e., physical activity, BMI, diet, alcohol and tobacco consumption) were measured in self-administered questionnaires in 1988 or 1992, and every 4 years until the last assessment available (2010; follow-up, 18–22 years). Data were analyzed in 2014–2015. Women were categorized according to initial level of psychological symptoms (e.g., lower versus higher anxiety symptoms).

Results—Despite slight improvements in healthy lifestyle over time among women with higher versus lower anxiety ($\beta_{\text{interaction}}=0.002$, 95% CI=0.001, 0.003), those experiencing more severe symptoms had a consistently less healthy lifestyle over time ($p<0.0001$). Each SD increase in anxiety symptoms was related to a decrease in healthy lifestyle score throughout follow-up ($\beta_{\text{pooled}}=-0.09$, 95% CI= -0.09, -0.08). Women with higher versus lower anxiety symptoms also had decreased odds of having a healthy lifestyle in 2010 (AOR=0.78, 95% CI=0.75, 0.81),

Address correspondence to: Claudia Trudel-Fitzgerald, PhD, Department of Social and Behavioral Sciences, Harvard T.H. Chan School of Public Health, Landmark Center, 401 Park Drive, 4th floor, Boston MA 02215. ctrudel@hsph.harvard.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Author's contributions: study concept and design, CTF, SST, EMP, and LDK; interpretation of data, all authors; drafting of the manuscript, CTF and LDK; critical revision of the manuscript for important intellectual content, all authors; administrative support, SST; statistical analysis, CTF, EMP, and LDK.

Preliminary data have been presented in posters at the Integrative Medicine Research Forum, Boston, Massachusetts (November 2014) and the Annual Scientific Meeting of the Society of Behavioral Medicine, San Antonio, Texas (April 2015).

No financial disclosures were reported by the authors of this paper.

particularly among women with an initially unhealthy lifestyle ($p_{\text{interaction}} = 0.0001$). Comparable patterns were observed with depression symptoms.

Conclusions—Among midlife women, anxiety and depression symptoms were associated with unhealthier lifestyle throughout follow-up and reduced odds of having a healthy lifestyle 20 years later. Treating psychological symptoms may promote healthier lifestyles.

Introduction

Unfavorable habits like physical inactivity, unhealthy diet, and smoking are critical risk factors for cardiovascular diseases, diabetes, and cancer.^{1–4} Anxiety and depression have also been linked with increased risk of cardiometabolic disease and mortality,^{5, 6} and various unhealthy behaviors have been proposed as potential mechanisms. Accordingly, biobehavioral theories have suggested psychological symptoms of anxiety and depression may be critical determinants of whether individuals engage in an integrated healthy lifestyle.^{6, 7} However, studies examining the influence of these symptoms on future adherence to a favorable lifestyle have been conducted primarily in patient populations (e.g., cardiac patients^{8, 9}). Thus, it is unclear if psychological symptoms influence adherence to an overall healthy lifestyle over time among adults without chronic illness.

Prospective studies have examined whether psychological symptoms increase future risk of single unfavorable behaviors among apparently healthy individuals.^{10–18} For instance, in adults with initially low physical activity levels, higher anxiety and depression symptoms were related to 79% increased odds of being inactive 3 and 6 years later.¹¹ Other studies have demonstrated associations of anxiety and depression symptoms with subsequently developing excess adiposity and becoming obese^{10, 13, 18} and with poor food choice, factors that may be used to cope with negative feelings.^{19, 20} A recent meta-analysis showed clinical anxiety disorders were related to a 40% increased risk of becoming a regular smoker,¹⁴ while women with moderate versus low levels of psychological symptoms had five fold increased odds of developing alcohol dependence problem over 18 months.¹⁷

Thus, anxiety and depression symptoms appear to be associated with increased risk of engaging in specific unhealthy behaviors later on but less is known about their impact on lifestyle globally. Because behaviors tend to cluster^{4, 21} and have a multiplicative impact on mortality compared with each individual behavior,²² the contribution of psychological symptoms to shaping lifestyle over time needs to be examined. This study investigated whether anxiety or depression symptoms lead to worsening lifestyle or reduce the odds of subsequently having a healthy lifestyle in midlife women, who generally report more psychological symptoms than men of similar age.^{16, 18, 23} The authors hypothesized that higher levels of anxiety/depression symptoms would be associated with unhealthier lifestyle over time and a lower odds of adopting a healthy lifestyle 20 years later.

Methods

Participants

The Nurses' Health Study is an ongoing cohort comprised of 121,700 U.S. female married nurses, aged 30–55 years at study inception in 1976.²⁴ They have completed biennial

questionnaires on lifestyle, medical history, and newly diagnosed medical conditions, with a high response rate of 90% since 1976.²⁵ The present study includes women who responded to questionnaires including an assessment of anxiety (1988) and depression (1992) symptoms (differences between included and excluded women in Appendix Table 1).

Figure 1 describes the two samples used for the analyses with anxiety and depression, respectively. Briefly, participants reporting a major medical condition (i.e., cancer, diabetes, heart disease) or those missing data on anxiety or depression symptoms, health behaviors, and confounders at baseline were not eligible. In analyses estimating changes in lifestyle score over time, women with data for only one of the four (for depression symptom analyses) or five (for anxiety symptoms analyses) follow-up lifestyle assessments were excluded (analytic samples: $n_{\text{anxiety}}=55,395$, $n_{\text{depression}}=51,979$). Hence, women with lifestyle data on at least two follow-up assessments (74.17% to 96.37% across time points) were included in the models. In analyses evaluating the odds of a healthy lifestyle at follow-up, women missing data on lifestyle in 2010 were excluded (analytic samples: $n_{\text{anxiety}}=41,125$, $n_{\text{depression}}=41,343$). Differences between excluded and included women are detailed in Appendix Table 1. The study protocol was approved by the IRB of Brigham and Women's Hospital and the Harvard T.H. Chan School of Public Health.

Measures

Anxiety symptoms were assessed in 1988 using the validated Crown–Crisp index.²⁶ The eight items are scored as 0 for *never*, 1 for *sometimes*, or 2 for *always*, with a derived sum ranging from 0 (no anxiety) to 16 (high anxiety). Internal consistency reliability in this sample was acceptable ($\alpha=0.61$). Anxiety symptoms were dichotomized into higher (≥ 4) versus lower (<4) levels.²⁷

Depression symptoms were measured in 1992 using the five-item Mental Health Index from the Medical Outcomes Study Short-Form 36 Health Status Survey.²⁸ All items, with responses ranging from 1 for *all the time* to 6 for *none of the time*, were summed and the total was scaled from 0 (high symptoms) to 100 (low symptoms). Internal consistency reliability was high ($\alpha=0.82$). Following previous work,²⁹ the score was dichotomized whereby ≤ 60 indicated high and >60 lower depression symptoms. Though not assessed at the same time, measures of both anxiety and depression symptoms were available in the two analytic samples (i.e., anxiety sample using 1988 baseline and depression sample using 1992 baseline). In each sample, the two measures were modestly correlated ($r_{\text{anxiety}}=0.30$ and $r_{\text{depression sample}}=0.29$).

Consistent with a lifestyle composite index used in previous studies and with available prevention cancer and cardiovascular guidelines,^{4, 30–34} the lifestyle score included five behavior-related factors: physical activity, diet, BMI, and alcohol and tobacco consumption. Individual behaviors, obtained via self-report in 1998 or 1992, and every 4 years (last time point available, 2010), were first dichotomized according to whether individuals met recommended guidelines or not (1/0). Component scores were then summed to create a healthy lifestyle score, ranging from 0 for “less healthy” to 5 for “most healthy.” Healthy lifestyle was also dichotomized at endorsement of 4 or 5 healthy behaviors, as this score has been associated with approximately 50% decreased risk of stroke in this cohort.³⁴

Physical activity was assessed with a validated questionnaire.³⁵ A score of 1 was assigned when women reported 150 minutes per week of moderate to vigorous activity (e.g., brisk walking, running, bicycling). BMI was derived using women's self-reported initial height and updated weight. Previous work with the cohort has shown self-reported weight is highly correlated with weight measured by study staff ($r=0.96$).³⁶ Optimal weight was defined as BMI ≤ 25 kg/m² (score of 1).

Dietary information was obtained from the 131 item Food Frequency Questionnaire, which has high reproducibility and validity when compared with 1-week diet records and biochemical markers.^{37, 38} This study's summary score encompasses the following components of the Alternative Healthy Eating Index, adapted from the U.S. Department of Agriculture Healthy Eating Index³⁹: Higher intakes of vegetables, fruit, nuts, soy, and cereal fiber; high ratio of chicken plus fish to red meat and polyunsaturated to saturated fat; low intake of trans fat; and multivitamin use of 5 years. The score for each dietary component ranges from 0 to 10 (optimal dietary behavior). Within this cohort, researchers have used the highest 40% of the diet score distribution as a cut off,^{34, 40} a range that was strongly related to a lower risk of several diseases, including stroke, diabetes, and cancer.⁴¹ Accordingly, a healthy diet (score of 1) was defined as a score in the top 40% of the current cohort distribution, updated at each time point. Healthy alcohol consumption (score of 1) was defined as drinking one drink or fewer per day.^{42, 43} Finally, women received a score of 1 if they reported being a non-current smoker. Scores were updated at each 4-year assessment.

Selected potential confounders included age (continuous), education level (registered nurses, bachelor, master, doctorate), and physical exam in the last two years (yes/no), as prior work has found receiving advice from a clinician is associated with weight loss, physical activity, and dietary changes.⁴⁴ Age and physical exam were self-reported at study baseline (anxiety analyses, 1988; depression analyses, 1992). Education was reported in 1992.

Statistical Analysis

Initial sets of analyses used linear mixed models for repeated measures to evaluate changes over time in lifestyle. These models use maximum likelihood estimate techniques that are robust to incomplete data on repeated measures.^{45, 46} For analyses considering anxiety, in the first model, anxiety symptoms were included as a continuous measure (standardized) along with a time variable (continuous measure in years). This model estimated change in lifestyle associated with: (1) each 1-SD change in anxiety symptoms; and (2) each additional year across the follow-up period. The second model used a dichotomous anxiety measure and estimated whether the lifestyle score significantly differed comparing women with lower versus higher anxiety symptoms. Both models included an interaction term to test whether the rate of change in lifestyle score depended on level of anxiety (e.g., decreased more rapidly among women with higher versus lower symptoms). Further analyses evaluated the main effect of anxiety symptoms on healthy lifestyle by pooling lifestyle scores across time points after taking into account any interaction effect. Stratified analyses tested if associations between anxiety symptoms and subsequent lifestyle scores differed depending on initial lifestyle status (healthy/unhealthy). Analogous models were conducted using depression symptoms.

The association of baseline levels of psychological symptoms (measured continuously [standardized] and categorically [lower versus higher anxiety/depression symptoms]) with the odds of reporting a healthy lifestyle at the end of follow-up was estimated with logistic regression models. A healthy lifestyle was defined as endorsing four or more healthy behaviors at the end of follow-up, otherwise (three or fewer healthy behaviors) categorized as unhealthy. Sensitivity analyses stratified logistic regression models by baseline lifestyle score (healthy/unhealthy).

Because unhealthy participants might be more likely to drop out of the study, inverse probability weights were developed and included in the models.⁴⁷ Specifically, the probability of participating at each time point was modeled based on the exposure and confounders of interest among participants who were included at baseline, and then a weight that corresponded to the inverse probability of participating was created. As results from age-adjusted models were similar to models including age, education level, and physical exam, only fully adjusted results are presented. Analyses were conducted in 2014–2015 using SAS, version 9.3 with a 5% level of significance.

Results

At the 1988 baseline (for the anxiety analysis; 22 years of follow-up), women were aged 54.07 (SD=7.07; range, 41–69) years on average; most were registered nurses (69.1%) and reported a recent physical exam (82.4%). The majority reported low alcohol consumption (88.1%), were non-smokers (83.2%), and had a healthy BMI (56.7%). One third engaged in moderate to vigorous physical activity (34.5%) and reported high anxiety levels at baseline (35.0%). Distribution of characteristics were similar at the 1992 baseline (for the depression analysis; mean age, 57.61; RN degree, 68.8%; physical exam, 87.9%; low alcohol consumption, 87.5%; non-smoker, 87.4%; healthy BMI, 51.4%; moderate to vigorous level of physical activity, 41.2%; 18 years of follow-up), although high depression symptom levels occurred in 15.3% of women only. Table 1 presents the distribution of confounders and health behaviors across baseline anxiety and depression symptom levels.

Both anxiety models suggested a small but significant interaction effect, although not in the expected direction (Table 2). For instance, Model 2 revealed that the rate of change in mean lifestyle score significantly differed over time, with more lifestyle improvement for the women with higher versus lower anxiety symptoms ($\beta_{\text{interaction}}=0.002$, 95% CI=0.001, 0.003). This small improvement associated with more anxiety symptoms appeared mainly among women who had unhealthy lifestyle at baseline (Appendix Table 2). For these women, stratified analyses revealed a main effect of time such that lifestyle improved slightly over follow-up ($\beta_{\text{time}}=0.005$, 95% CI=0.004, 0.006); however, it is important to note they never achieved a healthy lifestyle, adopting fewer than three healthy behaviors throughout follow-up. By contrast, in women with healthy lifestyle at baseline, lifestyle score worsened over follow-up ($\beta_{\text{time}}=-0.02$, 95% CI= -0.02, -0.02). In these stratified analyses, higher versus lower anxiety symptoms were associated with similar decreases in lifestyle scores in women with healthy or unhealthy lifestyle at the outset ($\beta_{\text{higher anxiety symptoms/healthy}}=-0.09$, 95% CI= -0.11, -0.07; $\beta_{\text{higher anxiety symptoms/unhealthy}}=-0.09$, 95% CI= -0.10, -0.07). When considering anxiety

effects in the full sample and after controlling for the interaction effect, mean healthy lifestyle scores of women with higher anxiety symptoms remained lower than those of women with lower anxiety symptoms throughout follow-up (pooled lifestyle score, lower symptoms=3.22 versus higher symptoms=3.07; $p<0.0001$); similarly, each 1-SD increase in anxiety symptoms was associated with a significant decrease in lifestyle score ($\beta_{\text{pooled}}=-0.09$, 95% CI= -0.09, -0.08).

Although the time effects were null, the interaction of depression symptoms with time was modestly but significantly associated with lifestyle score in both models. For example, in Model 4, the lifestyle score reported by women with higher versus lower symptoms slightly improved over time ($\beta_{\text{interaction}}=0.003$, 95% CI=0.001, 0.005), but this was primarily evident among women with initially unhealthy lifestyle scores (Appendix Table 2). As observed in anxiety models, when stratified by initial lifestyle score, effects of time on lifestyle score were in opposite direction ($\beta_{\text{time among healthy}}=-0.02$, 95% CI= -0.02, -0.02; $\beta_{\text{time among unhealthy}}=0.01$, 95% CI=0.01, 0.01). Nonetheless, the effect of depression symptoms on lifestyle was similar across strata ($\beta_{\text{higher depression symptoms/healthy}}=-0.07$, 95% CI= -0.10, -0.04; $\beta_{\text{higher depression symptoms/unhealthy}}=-0.05$, 95% CI= -0.07, -0.03). When considering depression effects in the full sample, after accounting for any interaction effect, unhealthy lifestyle was associated with higher versus lower depression symptoms levels (pooled lifestyle score, lower symptoms=3.18 versus higher symptoms=3.05; $p<0.0001$); similarly, lifestyle worsened over time with each 1-SD increase in depression symptoms ($\beta_{\text{pooled}}=-0.07$, 95% CI= -0.07, -0.06).

Every SD increase in anxiety symptoms was associated with 13% reduced odds (OR=0.87, 95% CI=0.85, 0.88) of reporting a healthy lifestyle in 2010 (Table 3). Likewise, women with higher versus lower symptoms had 22% reduced odds (OR=0.78, 95% CI=0.75, 0.81) of endorsing a healthy lifestyle in this period. In analyses stratified according to initial lifestyle score (healthy/unhealthy), higher levels of anxiety symptoms had a negative impact on future healthy lifestyle for women with initially healthy lifestyle (OR=0.87, 95% CI=0.82, 0.93), and even more strongly for women with initially unhealthy lifestyle (OR=0.82, 95% CI=0.78, 0.86, $p_{\text{interaction}}=0.0001$).

Similarly, there was 12% decreased odds of reporting a healthy lifestyle at the end of follow-up for every SD increase in depression symptoms (OR=0.88, 95% CI=0.87, 0.90). Likewise, women with higher versus lower symptoms had 21% reduced odds (OR=0.79, 95% CI=0.75, 0.83) of having a healthy lifestyle in 2010. Moreover, higher versus lower levels of depression symptoms negatively impacted future healthy lifestyle, especially among women with initially unhealthy lifestyle scores (OR_{healthy}=0.91, 95% CI=0.83, 0.99; OR_{unhealthy}=0.86, 95% CI=0.81, 0.92; $p_{\text{interaction}}=0.0001$).

Discussion

This research investigated whether anxiety or depression symptoms influence lifestyle over time and the odds of having a healthy lifestyle during adulthood. Consistent with prior work on individual health behaviors,^{11, 13, 48, 49} results adjusting for relevant confounders indicated that every SD increase in anxiety or depression symptoms was associated with a

significant decrease in the healthy lifestyle score over 20 years and with 12%–13% reduced odds of endorsing a healthy lifestyle at the end of follow-up, suggesting that even less severe symptoms may alter behaviors. Furthermore, women with higher versus lower levels of symptoms had a sustained unhealthier lifestyle score across time points and 21%–22% lower odds of reporting a healthy lifestyle 20 years later. It is particularly noteworthy that this effect was independent of levels of healthy lifestyle at study inception: Regardless of lifestyle score at the outset, women with higher versus lower psychological symptom levels had lower lifestyle scores throughout follow-up and lower odds of having a healthy lifestyle 20 years later.

The decreases in the healthy lifestyle score evident in relation to psychological symptoms are meaningful for health. For example, an unadjusted 0.13–0.18-point decrease in the mean healthy lifestyle score was observed among women reporting higher levels of anxiety/depression symptoms at any given point in time (Appendix Discussion). These decreases translate into a 2.1%–2.9% increased risk of stroke among women with higher versus lower levels of psychological symptoms, according to a study conducted among the same cohort where an average of 16.2% increased risk of stroke was obtained for each 1-point decrease on the healthy lifestyle score (Appendix Discussion).³⁴ This adds to initial risk already incurred, and may be a conservative estimate given that this study assessed additional risk starting when women were already at midlife.

Somewhat surprisingly, there was a non-significant effect of time on lifestyle when considering depressive symptoms. This might be related to the shorter follow-up period for this sample. Stratified analyses also provided some insight. Effects of time on lifestyle score among those with initially healthy versus unhealthy lifestyle were in the opposite direction and of similar magnitude, which could result in a null finding in the overall model. Unexpectedly, mixed models showed a slight tendency to adopt a healthier lifestyle over time among women with higher versus lower levels of psychological symptoms. Specifically, the number of cigarettes and drinks per day decreased over time, as other authors have also noted.⁵⁰ This could reflect some efficiency from public health policies in recent years or stem from methodologic issues, like “regression toward the mean” for the more anxious/depressed women or “ceiling effect” for the less anxious/depressed women. However, it was the women with both more severe anxiety/depression symptoms and unhealthy lifestyles at baseline who mainly drove this unexpected result; despite some improvements, these women generally maintained fewer than three healthy behaviors throughout follow-up. Finally, although this study found relatively small changes or differences between groups, these associations are similar in magnitude and direction to those found in other studies examining psychological states in relation to health behavior changes over time.^{51–53} Moreover, as noted above, prior work suggests that even small changes in behaviors are of clinical relevance and can impact health.

Limitations

Findings are from a sample of midlife women free of chronic illnesses at baseline, and cannot be generalized to other populations. Subjective reports of health behaviors may be vulnerable to social desirability bias, which can lead to overestimation of activity level and

underestimation of consumed calories.^{54, 55} However, even if absolute levels are not reported completely accurately, behaviors would likely still be categorized appropriately as either healthy/unhealthy. Likewise, anxiety or depression symptoms might affect health behavior reports; however, women were not aware that these factors would be considered together. Strengths include the use of a prospective design over 20 years. The composite healthy lifestyle score also facilitates a comprehensive perspective on how anxiety and depression symptoms influence multiple behaviors that jointly matter for health.

Conclusions

Together, these results suggest that psychological symptoms influence subsequent lifestyle changes. Anxiety and depression levels indeed affected maintenance (or not) of health behaviors over time and the odds of having a future favorable lifestyle, even in a healthy population and regardless of initial behaviors. Though it may not be surprising that feeling more anxious or depressed impedes attaining or maintaining favorable habits, it is noteworthy that psychological symptoms can also lead to worse lifestyle over time in women with initially healthy lifestyle. Hence, screening for psychological symptoms may be an effective primary prevention strategy, even when subclinical levels of symptoms or healthy habits are reported, enabling identification of middle-aged women who are at risk not only for common mental disorders but also for subsequently developing adverse physical health outcomes. Interestingly, some components of evidence-based treatments for anxiety/depression are directly related to lifestyle (e.g., increasing physical activity in cognitive behavioral therapy).⁵⁶ Thus, if confirmed by future clinical studies, implementing such treatments for psychological symptoms might be of particular interest.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The Nurses' Health Study is supported by grant UM1 CA186107. CTF received a postdoctoral fellowship from the Canadian Institutes of Health Research. This work was conducted with support from Dr. Henry Feldman as well as the Harvard Catalyst | The Harvard Clinical and Translational Science Center (National Center for Research Resources and the National Center for Advancing Translational Sciences, NIH Award UL1 TR001102) and financial contributions from Harvard University and its affiliated academic healthcare centers. The content is solely the responsibility of the authors and does not necessarily represent the official views of Harvard Catalyst, Harvard University and its affiliated academic healthcare centers, or NIH. CTF, EMP, and SST had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Wenger NK. What do the 2011 American Heart Association guidelines tell us about prevention of cardiovascular disease in women? *Clin Cardiol.* 2011; 34(9):520–523. <http://dx.doi.org/10.1002/clc.20940>. [PubMed: 21905040]
2. Klein WM, Bloch M, Hesse BW, et al. Behavioral research in cancer prevention and control: a look to the future. *Am J Prev Med.* 2014; 46(3):303–311. <http://dx.doi.org/10.1016/j.amepre.2013.10.004>. [PubMed: 24512871]
3. Narayan KM, Kanaya AM, Gregg EW. Lifestyle intervention for the prevention of type 2 diabetes mellitus: putting theory to practice. *Treat Endocrinol.* 2003; 2(5):315–320. <http://dx.doi.org/10.2165/00024677-200302050-00003>. [PubMed: 15981948]

4. WHO. Global status report on noncommunicable diseases. WHO; 2014.
5. Kubzansky, LD.; Winning, A. Mental disorders and the emergence of physical disorders. In: Koenen, KC.; Rudenstine, S.; Susser, E.; Galea, S., editors. A lifecourse approach to mental disorders. Oxford, U.K.: Oxford University Press; 2013. p. 291-304. <http://dx.doi.org/10.1093/acprof:oso/9780199657018.003.0026>
6. Kubzansky, LD.; Winning, A.; Kawachi, I. Affective states and health. In: Berkman, LF.; Kawachi, I.; Glymour, MM., editors. Social Epidemiology. 2. New York: Oxford University Press; 2014. p. 320-364. <http://dx.doi.org/10.1093/med/9780195377903.003.0009>
7. Baum A, Posluszny DM. Health psychology: mapping biobehavioral contributions to health and illness. *Annu Rev Psychol.* 1999; 50:137–163. <http://dx.doi.org/10.1146/annurev.psych.50.1.137>. [PubMed: 10074676]
8. Benyamini Y, Roziner I, Goldbourt U, Drory Y, Gerber Y. Israel Study Group on First Acute Myocardial I. Depression and anxiety following myocardial infarction and their inverse associations with future health behaviors and quality of life. *Ann Behav Med.* 2013; 46(3):310–321. <http://dx.doi.org/10.1007/s12160-013-9509-3>. [PubMed: 23645421]
9. Murray J, Craigs CL, Hill KM, Honey S, House A. A systematic review of patient reported factors associated with uptake and completion of cardiovascular lifestyle behaviour change. *BMC Cardiovasc Disord.* 2012; 12:120. <http://dx.doi.org/10.1186/1471-2261-12-120>. [PubMed: 23216627]
10. Brumpton B, Langhammer A, Romundstad P, Chen Y, Mai XM. The associations of anxiety and depression symptoms with weight change and incident obesity: The HUNT Study. *Int J Obes (Lond).* 2013; 37(9):1268–1274. <http://dx.doi.org/10.1038/ijo.2012.204>. [PubMed: 23229732]
11. Azevedo Da Silva M, Singh-Manoux A, Brunner EJ, et al. Bidirectional association between physical activity and symptoms of anxiety and depression: the Whitehall II study. *Eur J Epidemiol.* 2012; 27(7):537–546. <http://dx.doi.org/10.1007/s10654-012-9692-8>. [PubMed: 22623145]
12. Bulloch A, Lavorato D, Williams J, Patten S. Alcohol consumption and major depression in the general population: the critical importance of dependence. *Depress Anxiety.* 2012; 29(12):1058–1064. <http://dx.doi.org/10.1002/da.22001>. [PubMed: 22987574]
13. Hawkins MA, Stewart JC. Do negative emotional factors have independent associations with excess adiposity? *J Psychosom Res.* 2012; 73(4):243–250. <http://dx.doi.org/10.1016/j.jpsychores.2012.07.009>. [PubMed: 22980527]
14. Jiang F, Li S, Pan L, Zhang N, Jia C. Association of anxiety disorders with the risk of smoking behaviors: a meta-analysis of prospective observational studies. *Drug Alcohol Depend.* 2014; 145:69–76. <http://dx.doi.org/10.1016/j.drugalcdep.2014.10.022>. [PubMed: 25456325]
15. Pinto Pereira SM, Geoffroy MC, Power C. Depressive symptoms and physical activity during 3 decades in adult life: bidirectional associations in a prospective cohort study. *JAMA Psychiatry.* 2014; 71(12):1373–1380. <http://dx.doi.org/10.1001/jamapsychiatry.2014.1240>. [PubMed: 25321867]
16. Weinberger AH, Pilver CE, Desai RA, Mazure CM, McKee SA. The relationship of dysthymia, minor depression, and gender to changes in smoking for current and former smokers: longitudinal evaluation in the U.S. population. *Drug Alcohol Depend.* 2013; 127(1–3):170–176. <http://dx.doi.org/10.1016/j.drugalcdep.2012.06.028>. [PubMed: 22809897]
17. Haynes JC, Farrell M, Singleton N, et al. Alcohol consumption as a risk factor for anxiety and depression: results from the longitudinal follow-up of the National Psychiatric Morbidity Survey. *Br J Psychiatry.* 2005; 187:544–551. <http://dx.doi.org/10.1192/bjp.187.6.544>. [PubMed: 16319407]
18. Chiriboga DE, Ma Y, Li W, et al. Gender differences in predictors of body weight and body weight change in healthy adults. *Obesity (Silver Spring).* 2008; 16(1):137–145. <http://dx.doi.org/10.1038/oby.2007.38>. [PubMed: 18223626]
19. Singh M. Mood, food, and obesity. *Front Psychol.* 2014; 5:925. <http://dx.doi.org/10.3389/fpsyg.2014.00925>. [PubMed: 25225489]
20. Schellekens H, Finger BC, Dinan TG, Cryan JF. Ghrelin signalling and obesity: at the interface of stress, mood and food reward. *Pharmacol Ther.* 2012; 135(3):316–326. <http://dx.doi.org/10.1016/j.pharmthera.2012.06.004>. [PubMed: 22749794]

21. Spring B, King AC, Pagoto SL, Van Horn L, Fisher JD. Fostering multiple healthy lifestyle behaviors for primary prevention of cancer. *Am Psychol*. 2015; 70(2):75–90. <http://dx.doi.org/10.1037/a0038806>. [PubMed: 25730716]
22. Shaw BA, Agahi N. A prospective cohort study of health behavior profiles after age 50 and mortality risk. *BMC Public Health*. 2012; 12:803. <http://dx.doi.org/10.1186/1471-2458-12-803>. [PubMed: 22989155]
23. Pickering RP, Goldstein RB, Hasin DS, et al. Temporal relationships between overweight and obesity and DSM-IV substance use, mood, and anxiety disorders: results from a prospective study, the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry*. 2011; 72(11):1494–1502. <http://dx.doi.org/10.4088/JCP.10m06077gry>. [PubMed: 21457678]
24. Willett WC, Green A, Stampfer MJ, et al. Relative and absolute excess risks of coronary heart disease among women who smoke cigarettes. *N Engl J Med*. 1987; 317(21):1303–1309. <http://dx.doi.org/10.1056/NEJM198711193172102>. [PubMed: 3683458]
25. Colditz GA, Manson JE, Hankinson SE. The Nurses' Health Study: 20-year contribution to the understanding of health among women. *J Womens Health*. 1997; 6(1):49–62. <http://dx.doi.org/10.1089/jwh.1997.6.49>. [PubMed: 9065374]
26. Crown S, Crisp AH. A short clinical diagnostic self-rating scale for psychoneurotic patients. The Middlesex Hospital Questionnaire (M.H.Q.). *Br J Psychiatry*. 1966; 112(490):917–923. <http://dx.doi.org/10.1192/bjp.112.490.917>. [PubMed: 5970912]
27. Albert CM, Chae CU, Rexrode KM, Manson JE, Kawachi I. Phobic anxiety and risk of coronary heart disease and sudden cardiac death among women. *Circulation*. 2005; 111(4):480–487. <http://dx.doi.org/10.1161/01.CIR.0000153813.64165.5D>. [PubMed: 15687137]
28. Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care*. 1992; 30(6):473–483. <http://dx.doi.org/10.1097/00005650-199206000-00002>. [PubMed: 1593914]
29. Rumpf HJ, Meyer C, Hapke U, John U. Screening for mental health: validity of the MHI-5 using DSM-IV Axis I psychiatric disorders as gold standard. *Psychiatry Res*. 2001; 105(3):243–253. [http://dx.doi.org/10.1016/S0165-1781\(01\)00329-8](http://dx.doi.org/10.1016/S0165-1781(01)00329-8). [PubMed: 11814543]
30. Gingras D, Beliveau R. Colorectal cancer prevention through dietary and lifestyle modifications. *Cancer Microenviron*. 2011; 4(2):133–139. <http://dx.doi.org/10.1007/s12307-010-0060-5>. [PubMed: 21909875]
31. Khan MU. Lifestyle Modification in the Prevention of Type II Diabetes Mellitus. *Oman Med J*. 2012; 27(2):170–171. <http://dx.doi.org/10.5001/omj.2012.36>. [PubMed: 22496947]
32. Chiuve SE, Cook NR, Shay CM, et al. Lifestyle-based prediction model for the prevention of CVD: the Healthy Heart Score. *J Am Heart Assoc*. 2014; 3(6):e000954. <http://dx.doi.org/10.1161/JAHA.114.000954>. [PubMed: 25398889]
33. Sun Q, Shi L, Prescott J, et al. Healthy lifestyle and leukocyte telomere length in U.S. women. *PLoS One*. 2012; 7(5):e38374. <http://dx.doi.org/10.1371/journal.pone.0038374>. [PubMed: 22675460]
34. Chiuve SE, Rexrode KM, Spiegelman D, Logroscino G, Manson JE, Rimm EB. Primary prevention of stroke by healthy lifestyle. *Circulation*. 2008; 118(9):947–954. <http://dx.doi.org/10.1161/CIRCULATIONAHA.108.781062>. [PubMed: 18697819]
35. Chasan-Taber S, Rimm EB, Stampfer MJ, et al. Reproducibility and validity of a self-administered physical activity questionnaire for male health professionals. *Epidemiology*. 1996; 7(1):81–86. <http://dx.doi.org/10.1097/00001648-199601000-00014>. [PubMed: 8664406]
36. Rimm EB, Stampfer MJ, Colditz GA, Chute CG, Litin LB, Willett WC. Validity of self-reported waist and hip circumferences in men and women. *Epidemiology*. 1990; 1(6):466–473. <http://dx.doi.org/10.1097/00001648-199011000-00009>. [PubMed: 2090285]
37. Rimm EB, Giovannucci EL, Stampfer MJ, Colditz GA, Litin LB, Willett WC. Reproducibility and validity of an expanded self-administered semiquantitative food frequency questionnaire among male health professionals. *Am J Epidemiol*. 1992; 135(10):1114–1126. discussion 1127–1136. [PubMed: 1632423]
38. Giovannucci E, Colditz G, Stampfer MJ, et al. The assessment of alcohol consumption by a simple self-administered questionnaire. *Am J Epidemiol*. 1991; 133(8):810–817. [PubMed: 2021148]

39. McCullough ML, Feskanich D, Stampfer MJ, et al. Diet quality and major chronic disease risk in men and women: moving toward improved dietary guidance. *Am J Clin Nutr.* 2002; 76(6):1261–1271. [PubMed: 12450892]
40. Stampfer MJ, Hu FB, Manson JE, Rimm EB, Willett WC. Primary prevention of coronary heart disease in women through diet and lifestyle. *N Engl J Med.* 2000; 343(1):16–22. <http://dx.doi.org/10.1056/NEJM200007063430103>. [PubMed: 10882764]
41. Chiuve SE, Fung TT, Rimm EB, et al. Alternative dietary indices both strongly predict risk of chronic disease. *J Nutr.* 2012; 142(6):1009–1018. <http://dx.doi.org/10.3945/jn.111.157222>. [PubMed: 22513989]
42. Kushi LH, Doyle C, McCullough M, et al. American Cancer Society Guidelines on nutrition and physical activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J Clin.* 2012; 62(1):30–67. <http://dx.doi.org/10.3322/caac.20140>. [PubMed: 22237782]
43. Mosca L, Benjamin EJ, Berra K, et al. Effectiveness-based guidelines for the prevention of cardiovascular disease in women-2011 update: a guideline from the American Heart Association. *J Am Coll Cardiol.* 2011; 57(12):1404–1423. <http://dx.doi.org/10.1016/j.jacc.2011.02.005>. [PubMed: 21388771]
44. Rose SA, Poynter PS, Anderson JW, Noar SM, Conigliaro J. Physician weight loss advice and patient weight loss behavior change: a literature review and meta-analysis of survey data. *Int J Obes (Lond).* 2013; 37(1):118–128. <http://dx.doi.org/10.1038/ijo.2012.24>. [PubMed: 22450855]
45. Allison, P. Handling missing data by maximum likelihood. *SAS Global Forum*; 2012.
46. Fitzmaurice, GM.; Laird, NM.; Ware, JH. *Applied longitudinal analysis*. 2nd. New Jersey: John Wiley & Sons; 2011.
47. Hernan MA, Hernandez-Diaz S, Robins JM. A structural approach to selection bias. *Epidemiology.* 2004; 15(5):615–625. <http://dx.doi.org/10.1097/01.ede.0000135174.63482.43>. [PubMed: 15308962]
48. McKenzie SH, Harris MF. Understanding the relationship between stress, distress and healthy lifestyle behaviour: a qualitative study of patients and general practitioners. *BMC Fam Pract.* 2013; 14:166. <http://dx.doi.org/10.1186/1471-2296-14-166>. [PubMed: 24175998]
49. Haller CS, Etter JF, Courvoisier DS. Trajectories in cigarette dependence as a function of anxiety: a multilevel analysis. *Drug Alcohol Depend.* 2014; 139:115–120. <http://dx.doi.org/10.1016/j.drugalcdep.2014.03.016>. [PubMed: 24703608]
50. Epstein EE, Fischer-Elber K, Al-Otaiba Z. Women, aging, and alcohol use disorders. *J Women Aging.* 2007; 19(1–2):31–48. http://dx.doi.org/10.1300/J074v19n01_03. [PubMed: 17588878]
51. Holahan CK, Holahan CJ, Powers DA, Hayes RB, Marti CN, Ockene JK. Depressive symptoms and smoking in middle-aged and older women. *Nicotine Tob Res.* 2011; 13(8):722–731. <http://dx.doi.org/10.1093/ntr/ntr066>. [PubMed: 21504881]
52. Hingle MD, Wertheim BC, Tindle HA, et al. Optimism and diet quality in the Women's Health Initiative. *J Acad Nutr Diet.* 2014; 114(7):1036–1045. <http://dx.doi.org/10.1016/j.jand.2013.12.018>. [PubMed: 24556429]
53. Olthuis JV, Watt MC, Mackinnon SP, Stewart SH. CBT for high anxiety sensitivity: Alcohol outcomes. *Addict Behav.* 2015; 46:19–24. <http://dx.doi.org/10.1016/j.addbeh.2015.02.018>. [PubMed: 25753932]
54. Adams SA, Matthews CE, Ebbeling CB, et al. The effect of social desirability and social approval on self-reports of physical activity. *Am J Epidemiol.* 2005; 161(4):389–398. <http://dx.doi.org/10.1093/aje/kwi054>. [PubMed: 15692083]
55. Hebert JR, Ebbeling CB, Matthews CE, et al. Systematic errors in middle-aged women's estimates of energy intake: comparing three self-report measures to total energy expenditure from doubly labeled water. *Ann Epidemiol.* 2002; 12(8):577–586. [http://dx.doi.org/10.1016/S1047-2797\(01\)00297-6](http://dx.doi.org/10.1016/S1047-2797(01)00297-6). [PubMed: 12495831]
56. Barlow, DH. *Clinical Handbook of Psychological Disorders: A Step-by-step Treatment Manual*. 5th. Guilford Press; 2014.

Figure 1.
Flowchart of the different analytic samples.

Distribution of Confounders and Health Behaviors According to Baseline Anxiety (1988) and Depression (1992) Symptoms

Table 1

	Anxiety symptoms				Depression symptoms			
	Lower sympto ms (n=13,579)	Higher sympto ms (n=5,901)	Lower sympto ms (n=22,458)	Higher sympto ms (n=13,457)	Lower sympto ms (n=16,500)	Higher sympto ms (n=2,298)	Lower sympto ms (n=27,512)	Higher sympto ms (n=5,669)
	Healthy	Unhealthy	Healthy	Unhealthy	Healthy	Unhealthy	Healthy	Unhealthy
Mean age (SD)	54.62 (7.07)	54.90 (7.08)	53.69 (7.04)	53.80 (7.04)	58.35 (7.03)	57.02 (7.06)	57.54 (6.93)	56.02 (6.82)
Education level (%)								
Registered nurse	8,421 (62.01)	4,186 (70.94)	15,295 (68.10)	10,360 (76.99)	10,613 (64.32)	1,475 (64.19)	19,589 (71.20)	4,102 (72.36)
Bachelors	3,279 (24.15)	1,160 (19.66)	4,680 (20.84)	2,246 (16.69)	3,788 (22.96)	566 (24.63)	5,376 (19.54)	1,109 (19.56)
Masters	1,657 (12.20)	510 (8.64)	2,232 (9.94)	787 (5.85)	1,866 (11.31)	229 (9.97)	2,343 (8.52)	420 (7.41)
Doctorate	222 (1.63)	45 (0.76)	251 (1.12)	64 (0.48)	233 (1.41)	28 (1.22)	204 (0.74)	38 (0.67)
Physical exam in the last two years (%)								
Yes	11,446 (84.29)	5,016 (85.00)	18,177 (80.94)	10,984 (81.62)	14,669 (88.90)	2,076 (90.34)	24,002 (87.24)	4,959 (87.48)
Health behaviors (%)								
Healthy diet (top 40%)	10,194 (75.07)	4,407 (74.68)	4,870 (21.68)	2,723 (20.23)	12,221 (74.01)	1,656 (72.06)	5,836 (21.21)	1,108 (19.54)
Healthy alcohol consumption (1 drink/day)	12,912 (95.09)	5,648 (95.71)	18,806 (83.74)	11,416 (85.17)	15,571 (94.37)	2,182 (94.95)	22,899 (83.23)	4,852 (85.59)
Healthy smoking status (non-smoker currently)	13,052 (96.32)	5,711 (96.90)	17,213 (76.76)	10,055 (74.81)	16,053 (97.44)	2,222 (96.90)	22,654 (82.45)	4,428 (78.22)

	Anxiety symptoms				Depression symptoms			
	Lower sympto ms (n=13,579)	Higher sympto ms (n=5,901)	Lower sympto ms (n=22,458)	Higher sympto ms (n=13,457)	Lower sympto ms (n=16,500)	Higher sympto ms (n=2,298)	Lower sympto ms (n=27,512)	Higher sympto ms (n=5,669)
	Healthy		Unhealthy		Healthy		Unhealthy	
Healthy level of physical activity (150min/ week)	9,136 (73.93)	3,681 (71.05)	3,363 (15.59)	1,768 (13.72)	12,400 (79.02)	1,595 (74.88)	5,799 (21.49)	932 (16.75)
Healthy BMI (< 25 kg/m ²)	10,581 (84.37)	4,502 (83.08)	9,326 (43.56)	5,208 (40.61)	12,418 (81.14)	1,743 (82.84)	9,189 (34.71)	1,989 (36.68)

Note. Percentages refer to the row percentage of individuals within each psychological symptoms/lifestyle category with that characteristic.

Table 2

Linear Mixed Models With Baseline Anxiety (1988) and Depression (1992) Symptoms, Predicting Evolution of the Mean Continuous Lifestyle Score Between 1994 or 1998 and 2010

Exposures	Estimates (β)	95% CI	
Anxiety (N=55,395)			
Model 1			
Anxiety symptoms (per 1 SD change)	-0.11 ****	-0.12	-0.10
Time (years)	-0.003 ****	-0.004	-0.003
Interaction (anxiety symptoms × time)	0.001 ****	0.001	0.002
Model 2			
Higher vs. lower anxiety symptoms	-0.18 ****	-0.20	-0.16
Time (years)	-0.004 ****	-0.005	-0.003
Interaction (anxiety symptoms level × time)	0.002 ****	0.001	0.003
Depression (N=51,979)			
Model 3			
Depression symptoms (per 1 SD change)	-0.08 ****	-0.07	-0.07
Time (years)	0.0001	-0.001	0.001
Interaction (depression symptoms × time)	0.002 ****	0.001	0.002
Model 4			
Higher vs. lower depression symptoms	-0.17 ****	-0.20	-0.14
Time (years)	-0.001	-0.001	0.0003
Interaction (depression symptoms level × time)	0.003 **	0.001	0.005

Note. Boldface indicates statistical significance

*
p 0.05;

**
p 0.01;

p 0.001;

p 0.0001.

To ease interpretation, the depression score was reversed (0=no depressive symptoms, 100=highest depressive symptoms) and then standardized. Because results from age-adjusted models were similar to models including age, education level and physical exam, only fully-adjusted results are presented.

Table 3

Logistic Regressions Models Evaluating the Association of Baseline Anxiety (1988) and Depression (1992) Symptoms With the Odds of Having a Healthy Lifestyle in 2010

Main models				
	Anxiety		Depression	
	Continuous (1 SD)	Dichotomized (Higher vs. lower symptoms)	Continuous (1 SD)	Dichotomized (Higher vs. lower symptoms)
Sample size (number of women with healthy lifestyle at follow-up)	41,125 (13,433)	41,125 (13,433)	41,434 (13,692)	41,434 (13,692)
OR (95% CI)	0.87 **** (0.85–0.88)	0.78 **** (0.75–0.81)	0.88 **** (0.87–0.90)	0.79 **** (0.75–0.83)
Stratified Analyses				
	Anxiety		Depression	
	Healthy (Higher vs. lower symptoms)	Unhealthy (Higher vs. lower symptoms)	Healthy (Higher vs. lower symptoms)	Unhealthy (Higher vs. lower symptoms)
Sample size (number of women with healthy lifestyle at follow-up)	14,865 (7,641)	26,260 (5,792)	15,315 (8,181)	26,119 (5,511)
OR (95% CI)	0.87 **** (0.82–0.93)	0.82 **** (0.78–0.86)	0.91 * (0.83–0.99)	0.86 **** (0.81–0.92)

Notes. Boldface indicates statistical significance

* p 0.05;

** p 0.01;

*** p 0.001;

**** p 0.0001.

To ease interpretation, the depression score was reversed (0=no depressive symptoms, 100=highest depressive symptoms) and then standardized. Because results from age-adjusted models were similar to models including age, education level and physical exam, only fully-adjusted results are presented. An odd ratio <1 reflects *lower* odds of adopting a healthy lifestyle at the 2010 follow-up time assessment.