Published in final edited form as:

Am J Prev Med. 2015 September ; 49(3): 414–418. doi:10.1016/j.amepre.2015.01.020.

Randomized Trial of a Fitbit-Based Physical Activity Intervention for Women

Lisa A. Cadmus-Bertram, Ph.D.¹, Bess H. Marcus, Ph.D.², Ruth E. Patterson, Ph.D.^{2,3}, Barbara A. Parker, M.D.³, and Brittany L. Morey, M.P.H.⁴

¹Department of Kinesiology, University of Wisconsin – Madison

²Department of Family Medicine and Public Health, University of California, San Diego

³Department of Medicine, University of California, San Diego

⁴Moore's Cancer Center, University of California, San Diego

Abstract

Background—Direct-to-consumer *mHealth* devices are a potential asset to behavioral research but are rarely tested as intervention tools. This trial examined the accelerometer-based Fitbit tracker and website as a basis for a low-touch physical activity intervention.

Purpose—To evaluate, within a randomized controlled trial, the feasibility and preliminary efficacy of integrating the Fitbit tracker and website into a physical activity intervention for postmenopausal women.

Methods—Fifty-one inactive, postmenopausal women with BMI ≥ 25.0 kg/m² were randomized to a 16-week web-based self-monitoring intervention (N=25) or a comparison group (N=26). Those in the Web-Based Tracking Group received a Fitbit, an instructional session, and a follow-up call at 4 weeks. The comparison group received a standard pedometer. All were asked to perform 150 min/week of moderate-to-vigorous physical activity (MVPA). Physical activity outcomes were measured by the ActiGraph GT3X+ accelerometer.

Results—Data were collected and analyzed in 2013–2014. Participants were 60 ± 7 years old with BMI = 29.2 ± 3.5 kg/m². Relative to baseline, the Web-Based Tracking Group increased MVPA by 62 ± 108 min/week ($p < .01$), MVPA in 10-min bouts by 38 ± 83 min/week ($p = .008$), and steps by $789 \pm 1,979$ ($p = .01$), compared to non-significant increases in the Pedometer Group (between-group *p*-values: .11, .28 and .30, respectively). The Web-Based Tracking Group wore the tracker on 95% of intervention days; 96% reported liking the website and 100% liked the tracker.

Address correspondence to: Lisa Cadmus-Bertram, Ph.D., University of Wisconsin - Madison, 2000 Observatory Drive, Madison, WI 53706, Phone: (858) 265-5946, Fax: (608) 262-1656, cadmusbertra@wisc.edu.

The authors have no conflicts of interest to report.

Lisa Cadmus-Bertram has no financial disclosures.

Bess H. Marcus has no financial disclosures.

Ruth E. Patterson has no financial disclosures.

Barbara A. Parker has no financial disclosures.

Brittany L. Morey has no financial disclosures.

Registration

ClinicalTrials.gov identifier: NCT01837147.

Conclusions—The Fitbit was well-accepted in this sample of women and was associated with increased physical activity at 16 weeks. By leveraging direct-to-consumer *mHealth* technologies that align with behavior change theories, researchers can strengthen physical activity interventions.

Given the established relationship between inactivity and chronic disease, physical activity (PA) promotion is a priority for postmenopausal women.^{1–4} Accelerometer-based tracking devices and apps are of interest for use in scalable PA interventions because they can encourage the use of theory-driven self-regulation skills known to be associated with behavior change success.^{5, 6} One example is the Fitbit tracker, which collects activity data, uploads it to the web, and produces simple graphs and charts. Published studies examining the Fitbit as an intervention tool are currently limited to two single-arm studies^{7, 8} and one randomized trial.⁹ The aims of this trial were to examine (a) the acceptability of the Fitbit among postmenopausal, overweight/obese women and (b) the effect of a Fitbit-based intervention on PA.

Methods

Participants

Participants were 51 overweight, postmenopausal women performing <60 minutes/week of moderate-to-vigorous intensity physical activity (MVPA)¹⁰ (to ensure they were well below the recommended 150 min/week¹¹) who were regular internet users, owned a computer/tablet, and could exercise safely.¹²

Study Visits

Participants attended three visits at the University of California, San Diego (UCSD). Procedures were approved by the UCSD Human Research Protections Program. At the initial visit, informed written consent was obtained and the participant received baseline assessments. At the second visit, a web app assigned her with equal probability (blocked on age and BMI) to the Web-Based Tracking Group or the Pedometer Group. The participant was then oriented her group. Participants returned for a Final Visit at 16 weeks and received \$20 for study completion.

Measures

Physical Activity—The ActiGraph GT3X+ (ActiGraph, Pensacola, FL) was worn on the hip for 7 days at baseline and 16 weeks. Standard thresholds were used to aggregate data into sedentary, light, moderate, and vigorous activity.¹³

Anthropometrics—Height and weight were measured using standard procedures.

Questionnaire—The Baseline Questionnaire included demographics and technology use (pewinternet.org). The Final Questionnaire evaluated the assigned study arm. For participants in the Web-Based Tracking Group, this included frequency of Fitbit wear, use of the website, perceived helpfulness of specific features, and perceived benefits/barriers.

Intervention Groups

This study compared (a) a low-touch, Fitbit-based PA intervention focused on self-monitoring/self-regulation skills^{5, 6} (Web-Based Tracking Group) and (b) provision of a basic step-counting pedometer (Pedometer Group). Both were asked to perform 150 minutes/week of MVPA and walk 10,000 steps/day.

Web-Based Tracking Group—Each participant received a Fitbit One, an accelerometer-based device that clips to the waistband or bra, or is placed in a pocket. Summary data are shown on the tracker's display and PA intensities and temporal patterns are available on the website. Prior to randomization, the study coordinator initialized a tracker for each participant (in case she was randomized to the Fitbit group) and modified the online “dashboard” to display only PA data. At the Randomization Visit, the study coordinator demonstrated how to download/install the software and use the website. This training was reinforced with a study-specific handbook. The intervention was based on the CALO-RE framework, which identifies self-monitoring, combined with other self-regulatory skills (e.g., goal-setting, frequent behavioral feedback) as the most important theory-driven component of successful behavior change^{5, 6}. Individualized goals were set for the first 4 weeks of the study (using data observed on the baseline ActiGraph) and the participant committed to a specific plan to achieve these. A follow-up call at 4 Weeks was used to evaluate progress and refine goals.

Pedometer Group—These participants received a basic pedometer and printed materials with tips for increasing steps. They also completed a brief goal-setting process, based on steps observed on the ActiGraph.

Data management and statistical analysis

Data were collected/managed in 2013–2014¹⁴ and analyzed in 2014 using SAS 9.4, according to the intent-to-treat principle. Baseline characteristics were compared using chi-square and t-tests. ActiGraph data were adjusted for number of valid days (95% had 7 valid days; 5% had 5–6). Baseline-to-16-Week PA changes were assessed using repeated-measures ANCOVA, adjusted for age and ActiGraph daily wear time to address potential residual confounding. Missing data (n=2 from the Pedometer Group) were imputed by carrying forward baseline values.

Results

Participants (N=51) were 60.0±7.1 years of age with a body mass index of 29.2±3.5 kg/m² and groups were comparable on key characteristics (Table 1). At baseline, women were performing 33±56 min/week of MVPA in bouts 10 minutes in length (as specified by the PA guidelines¹⁵) and accumulating 5,866±2,195 steps/day.

Baseline to 16-Week Changes in Physical Activity

Relative to baseline, the Web-Based Tracking Group increased MVPA in bouts by 38±83 min/week ($p<.01$), increased total MVPA by 62±108 min/week ($p=.008$), and increased

steps/day by $789 \pm 1,979$ ($p=.01$) (Table 2). The Pedometer Group experienced non-significant increases in PA. No adverse events occurred.

Usability and feasibility of Fitbit tracker and website

96% of participants reported wearing the tracker 4 days/week. Wear time was corroborated by Fitbit data, which showed the median days of any wear (2,000 steps accrued) was 106 of 112 prescribed days. 88% used the website, with 52% logging in 2–3 days/week. Participants were most engaged with the device itself (72% viewing tracker data 1 time/day), using a passive approach to the website (viewing feedback but not changing goals or manually logging behaviors).

Barriers were low: 80% had no computer issues, 80% reported no technical difficulty with the tracker, and 84% had no issues with a lost/broken tracker. 96% of women liked the Fitbit website and 100% liked the tracker. 76% said that they would recommend the Fitbit to a friend. When asked about future preferences, 56% of participants preferred a clip-on tracker, 20% preferred a wrist-worn tracker, and 24% had no preference. 96% rated the Fitbit as “somewhat or very” helpful for increasing PA, compared to only 32% of the Pedometer Group who found the basic pedometer “somewhat or very” helpful.

Discussion

In this study, a Fitbit-based intervention was associated with increased PA, with no change observed in the Pedometer Group. Although between-group tests did not reach significance, this may be due to sample size. The increase of 62 min/week of MVPA observed in the Fitbit arm is substantial, particularly if maintained over time. This maintenance would likely require progressive goal increases established via periodic contact with investigators via web or telephone. This study’s findings differ from those of Thompson, et al., who found that a Fitbit with feedback did not increase PA among older adults.⁹ This may be partly because of the ~20-year difference in participant age.

Barriers to Fitbit use were low and technical issues were resolved quickly. Participants adhered well to wearing and viewing feedback on the tracker. Participant feedback indicated that women liked the website but either found the tracker sufficient for their needs or were otherwise unmotivated to explore the website beyond its basic functions; several mentioned the desire for additional hands-on training. Such training may help increase effect size by enhancing participants’ engagement with the website.

Strengths of this study include the use of the ActiGraph, detailed participant feedback, and use of Fitbit data to corroborate adherence. Limitations include a small sample, short duration, and lack of generalizability. Further research is needed to better understand the potential of direct-to-consumer devices for PA promotion, particularly over the long term.

Acknowledgments

This study was funded by NIH (1R03CA168450) and recruitment supported by the Athena Breast Health Network.

This research was supported by the National Cancer Institute (1R03CA168450). This research was collected at the UC San Diego while Dr. Cadmus-Bertram was on the faculty there.

References

1. Eliassen AH, Hankinson SE, Rosner B, Holmes MD, Willett WC. Physical activity and risk of breast cancer among postmenopausal women. *Arch Intern Med*. 2010; 170(19):1758–64. [PubMed: 20975025]
2. Ratnasinghe LD, Modali RV, Seddon MB, Lehman TA. Physical activity and reduced breast cancer risk: a multinational study. *Nutr Cancer*. 2010; 62(4):425–35. [PubMed: 20432163]
3. Oguma Y, Shinoda-Tagawa T. Physical activity decreases cardiovascular disease risk in women: review and meta-analysis. *Am J Prev Med*. 2004; 26(5):407–18. [PubMed: 15165657]
4. Hsia J, Wu L, Allen C, Oberman A, Lawson WE, Torrens J, et al. Physical activity and diabetes risk in postmenopausal women. *Am J Prev Med*. 2005; 28(1):19–25. [PubMed: 15626551]
5. Michie S, Abraham C, Whittington C, McAteer J, Gupta S. Effective techniques in healthy eating and physical activity interventions: a meta-regression. *Health Psychol*. 2009; 28(6):690–701. [PubMed: 19916637]
6. Michie S, Ashford S, Sniehotta FF, Dombrowski SU, Bishop A, French DP. A refined taxonomy of behaviour change techniques to help people change their physical activity and healthy eating behaviours: the CALO-RE taxonomy. *Psychol Health*. 2011; 26(11):1479–98. [PubMed: 21678185]
7. Kurti AN, Dallery J. Internet-based contingency management increases walking in sedentary adults. *J Appl Beh Anal*. 2013; 46(3):568–81.
8. Washington WD, Banna KM, Gibson AL. Preliminary efficacy of prize-based contingency management to increase activity levels in healthy adults. *J Appl Behav Anal*. 2014; 47(2):231–45. [PubMed: 24740477]
9. Thompson WG, Kuhle CL, Koepp GA, McCrady-Spitzer SK, Levine JA. “Go4Life” exercise counseling, accelerometer feedback, and activity levels in older people. *Arch Gerontol Geriatr*. 2014; 58(3):314–9. [PubMed: 24485546]
10. Sallis JF, Haskell WL, Wood PD, Fortmann SP, Rogers T, Blair SN, et al. Physical activity assessment methodology in the Five-City Project. *Am J Epidemiol*. 1985; 121(1):91–106. [PubMed: 3964995]
11. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc*. 2007; 39(8):1423–34. [PubMed: 17762377]
12. Shephard RJ. PAR-Q, Canadian Home Fitness Test and exercise screening alternatives. *Sports Med*. 1988; 5(3):185–95. [PubMed: 3368685]
13. Freedson PS, Melanson E, Sirard J. Calibration of the Computer Science and Applications, Inc. accelerometer. *Med Sci Sports Exerc*. 1998; 30(5):777–81. [PubMed: 9588623]
14. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009; 42(2):377–81. [PubMed: 18929686]
15. Winett RA, Tate DF, Anderson ES, Wojcik JR, Winett SG. Long-term weight gain prevention: a theoretically based Internet approach. *Prev Med*. 2005; 41(2):629–41. [PubMed: 15917062]

Table 1

Baseline characteristics of study participants.

	Web-based Tracking Group (N=25)	Pedometer Group (N=26)	<i>p</i> for diff.
	Mean (SD) or %	Mean (SD) or %	
Age	58.6 (6.5)	61.3 (7.5)	.17
Weight (kg)	82.4 (14.7)	79.3 (12.2)	.42
BMI ^a (kg/m ²)	29.2 (3.8)	29.1 (3.2)	.94
Non-Hispanic White	23 (92%)	23 (88%)	.61
College degree or higher	14 (56%)	18 (69%)	.33
Moderate-vigorous physical activity^b			
Performed in Freedson bouts (min/wk)	24 (39)	42 (68)	.26
Meeting physical activity guidelines	0%	8%	.15
Total moderate-vigorous activity (min/wk)	172 (83)	176 (117)	.89
Steps^b			
Average steps per day	5,906 (1,964)	5,823 (2,431)	.90
% Walking 10,000 steps/day	4%	8%	.99
Technology use			
Daily internet user	84%	88%	.64
Comfortable using computers and the Internet ^c			.99
Neutral	4%	0%	
Somewhat comfortable	12%	15%	
Very comfortable	84%	85%	
Enjoys using computers and the Internet ^d			.38
Neutral	17%	4%	
Somewhat enjoy	21%	23%	
Very much enjoy	62%	73%	
Type of primary computer			.38
Desktop	32%	50%	
Laptop	64%	42%	
Tablet	4%	8%	
Operating system of primary computer			.99
Windows	64%	65%	
Mac	36%	31%	
Other (e.g., Linux)	0%	4%	

^aBody mass index^bAs measured by ActiGraph GT3X+ accelerometer^cWomen who responded "Very uncomfortable" or "Somewhat uncomfortable" were ineligible for this study.^dWomen who responded "Very much do not enjoy" or "Somewhat do not enjoy" were ineligible for this study.

Table 2

Baseline to 16-week changes in objectively-measured physical activity and body weight.

	Web-Based Tracking Group N=25				Pedometer Group N=26				Effect size (Cohen's <i>d</i>)
	Baseline	16 weeks	Change	<i>p</i>	Baseline	16 weeks	Change	<i>p</i>	
Min/week of physical activity									
Mod-vig. intensity (total)	172 (83)	234 (119)	62 (108)	.008	176 (117)	189 (99)	13 (98)	.51	.48
Mod-vig. intensity (in bouts)	24 (39)	62 (82)	38 (83)	.01	42 (68)	57 (69)	16 (76)	.26	.28
Light intensity	1276 (311)	1262 (320)	-14 (204)	.49	1284 (383)	1252 (317)	-33 (225)	.82	.09
Average steps per day	5,906 (1,968)	6,695 (2,708)	789 (1,979)	.01	5,827 (2,431)	6,188 (2,423)	362 (1,605)	.17	.24
Body weight (kg)	82.4 (14.7)	82.2 (16.0)	-0.3 (2.4)	.49	79.3 (12.2)	79.2 (13.2)	0.01 (2.3)	.98	.06