Published in final edited form as:

Sci Transl Med. 2016 April 27; 8(336): 336ra61. doi:10.1126/scitranslmed.aad9370.

MDM2 Inhibition rescues neurogenic and cognitive deficits in fragile X mice

Yue Li¹, Michael E. Stockton¹, Ismat Bhuiyan¹, Brian E. Eisinger¹, Yu Gao¹, Jessica L. Miller¹, Anita Bhattacharyya¹, and Xinyu Zhao^{1,2}

¹Waisman Center, University of Wisconsin-Madison, Madison, WI 53705, USA

²Department of Neuroscience, University of Wisconsin-Madison, Madison, WI 53705, USA

Abstract

Fragile X syndrome, the most common form of inherited intellectual disability, is caused most often by a lack of fragile X mental retardation protein (FMRP). However, the mechanism remains unclear and effective treatment is lacking. Here we show that a loss of FMRP leads to activation of adult neural stem cells (NSCs) and a subsequent reduction in neuronal production. We identified ubiquitin ligase MDM2 as a target of FMRP. FMRP regulates *Mdm2* mRNA stability, and loss of FMRP results in elevated mRNA and MDM2 protein levels. We further found that increased MDM2 levels lead to reduced P53 in NSCs, which alters NSC proliferation and differentiation. Treatment with Nutlin-3, a small molecule undergoing clinical trials for cancer, specifically

Corresponding author: Xinyu Zhao (Waisman Center and Department of Neuroscience, University of Wisconsin-Madison School of Medicine and Public Health, Madison, WI 53705, USA; Phone: (608) 263-9906; Xinyu.zhao@wisc.edu.

SUPPLEMENTARY MATERIALS

fig. S1. FMRP deficiency does not affect DG volume but leads to increased number of activated cells in the adult DG and increased numbers of GFP+ cell, NSCs, IPCs in both dorsal and ventral DG.

fig. S2. Conditional deletion of FMRP leads to increased NSC numbers in the adult DG without affecting overall DG volume.

fig. S3. Adult NPCs derived from *Fmr1* KO mice exhibit increased proliferation, reduced neuronal differentiation, but increased astroglial differentiation.

fig. S4. Assessment of optimal protein loading amount for protein quantification using Western blotting.

fig. S5. Quantitative tables for p-MDM2 level assessment using western blot analysis.

fig. S6. Mdm2 mRNA stability analysis in WT and *Fmr1* KO aNPCs

fig. S7. *Fmr1* KO aNPCs do not exhibit increased cell death.

fig. S8. Both acute knockdown of MDM2 and treatment with MK-2206 rescue astroglial differentiation of *Fmr1* KO aNPCs.

fig. S9. Inverse correlation of FMRP and p-MDM2 expression levels in *Fmr1*^{+/−}::Nestin-GFP female brains.

fig. S10. Selective deletion of FMRP from Nestin-positive aNSCs and their progenies resulted in cognitive deficits, without affecting locomotor activities.

fig. S11. Nutlin-3 treatment does not affect general health and activities of mice

fig. S12. Increased MDM2 mRNA and p-MDM2 protein levels in the hippocampal and cortical tissues from *Fmr1* KO mice and postmortem brains of FXS patients.

fig. S13. P53 protein levels did not show significant changes in the hippocampal and cortical tissues from *Fmr1* KO mice

Table S1. FMRP Candidate Targets (provide as a separate Excel file).

Table S2. Neurogenic Regulators that Regulate Adult Neural Stem Cell Quiescence and Activation

Table S3. All known physical interactions with proteins known to be important in NSC activation/proliferation (provide as a separate Excel file).

Table S4. Number of physical interactions between FMRP Candidate Targets identified by PAR-CLIP-seq (Table S1) and Neurogenic Regulators with role in NSC activation obtained from MANGO database and literature search (Table S2).

Author Contributions: Y.L. and X.Z. conceived the concept, designed the experiments, analyzed data, and wrote the manuscript. Y.L. performed all in vitro and in vivo experiments; M.E.S. performed behavioral and cell fate counting. I.B. and J.L.M. performed cell fate counting. B.E. performed bioinformatics analysis. Y.G. created the lentiviral vectors. A.B. participated in human tissue analysis.

Competing interests: The authors declare no conflict of interest.

Data and materials availability: All materials generated are available for distribution.

inhibits MDM2 and P53 interaction, and rescues the neurogenic and cognitive deficits in FMRP-deficient mice. Our data unveil a regulatory role for FMRP and a potential new treatment for fragile X syndrome.

INTRODUCTION

Fragile X syndrome (FXS) affects 1 in 4000 males and 1 in 6000 females and is the most common form of inherited intellectual disability(1, 2). FXS arises largely due to mutations in the fragile X mental retardation (*FMR1*) gene, resulting in a functional deficiency of fragile X mental retardation protein (FMRP), a brain-enriched RNA-binding protein. Genome-wide analyses predict that FMRP regulates many mRNAs(3, 4). FMRP is highly expressed in neurons, and studies show that a loss of FMRP leads to altered expression of many neuronal genes. Therefore a number neurotransmitters and neuronal signaling pathways have been intensely investigated as potential therapeutic targets, including pathways associated with group 1 metabotropic glutamate receptor type 5 (mGluR5), N-methyl-D-aspartate receptor subunits, GABA_A receptor, mammalian target of rapamycin, tuberous sclerosis complex 2, and glycogen synthase kinase-3 β , etc. (reviewed by (1, 5)). These studies led to a number of possible FXS treatments (6). Unfortunately, disappointing results from recent clinical trials of inhibitors for mGluR5, the best known FMRP drug target, underscore a pressing need for innovation in terms of both target selection and consideration of CNS cell types beyond neurons alone(7). Despite the intense interest in FXS, especially because of its potential as a gateway to understanding autism, the precise mechanisms behind this unique syndrome remain unclear, and there is still no FDA-approved treatment. Exploring novel mechanisms to explain the etiology of FXS is therefore essential if we are to develop a treatment.

Adult neurogenesis has been explored because of its potential to influence information processing associated with cognitive outcomes(8). Impaired adult neurogenesis is implicated in both neuropsychiatric disorders, such as depression and schizophrenia, and neurodegenerative diseases, such as Alzheimer's and Huntington's disease(9). Interventions aimed at regulating adult neurogenesis are thus being evaluated as potential therapeutic strategies. In many adult organs, both quiescent and activated (actively dividing) stem cells coexist, and the regulation of stem cell activation plays an essential role in tissue maintenance, regeneration, function, plasticity, aging, and disease(10). In the adult hippocampus, quiescent neural stem cells (NSCs) dynamically integrate both extrinsic and intrinsic signals to either maintain their dormant state, or to become activated and give rise to intermediate progenitor cells (IPCs), which subsequently differentiate into postmitotic dentate gyrus (DG) granule neurons or astrocytes(9). Many studies over the last 20 years have uncovered intracellular factors that regulate the activation of DG NSCs. Although it is clear that maintaining a balance of quiescence and activation is important for the preservation of adult NSCs, whether this balance directly impacts neuronal production has yet to be explored.

We have shown that FMRP deficiency in adult NSCs leads to increased proliferation but reduced neuronal differentiation and impaired hippocampus-dependent learning abilities,

which can be rescued by either restoring FMRP expression in adult NSCs, or by treating mice with a small molecule that promotes neuronal differentiation(11–13). Several key questions nonetheless remained. (1) Does FMRP play a role in regulating the activation of adult NSCs? (2) Can manipulating NSC activation directly impact neuronal production? (3) Will rebalancing adult NSC quiescence and activation rescue aberrant hippocampal neurogenesis and the associated FXS cognitive deficits?

Here we tested the hypothesis that FMRP deficiency leads to activation of adult NSCs, and that rebalancing NSC quiescence and activation rescues cognitive deficits in a FXS mouse model. We show that a loss of FMRP leads to greater activation of adult NSCs. We identified mouse double minute 2 homolog (MDM2), an E3 ubiquitin ligase, as a mediator of FMRP action in adult NSCs. FMRP regulates the expression levels of MDM2 by reducing its mRNA stability. In addition, the absence of FMRP also leads to elevated AKT signaling in NSCs, resulting in enhanced phosphorylation of MDM2. Thus, a lack of FMRP results in increased levels of phosphorylated (activated) MDM2 via two distinct mechanisms. Phosphorylated MDM2 targets and degrades tumor suppressor protein P53. We found that specific inhibition of MDM2-P53 interaction with a small molecule undergoing cancer clinical trial rescued both the neurogenic and cognitive deficits in FXS mice. Thus our data not only unveil a regulatory mechanism for FMRP during neuronal development, but also point the way to a promising new therapy for treating FXS.

RESULTS

FMRP deficiency leads to increased NSC activation

To investigate the role of FMRP in regulating NSC activation, we crossed *Fmr1* mutant (KO) mice with mice expressing GFP driven by the promoter of a neural progenitor cell marker, Nestin (*Nestin-GFP*), to create both *Fmr1* KO (*Fmr1*^{-y::Nestin-GFP}) and WT (*Fmr1*^{+y::Nestin-GFP}) mice, in which adult NSCs and IPCs are labeled by GFP (Fig. 1A, and fig S1A). We found that the *Fmr1* KO mice had significantly increased number (Fig. 1B) and density (Fig. 1C) of GFP⁺ cells in the DG compared to their WT littermates without significant changes in the overall volume of the DG (Fig. S1, C and D). We next analyzed different populations of GFP⁺ cells in the subgranular zone by using radial glia marker GFAP for type 1 radial glia-like NSCs (GFP⁺GFAP⁺) and TBR2 for IPCs (TBR2⁺ or GFP⁺GFAP⁻). *Fmr1* KO mice had significantly more NSCs (Fig. 1, D and E) and IPCs (Fig. 1F–H) compared to their WT littermates. We then used MCM2, a marker for cell cycle initiation and stem cell activation to distinguish activated NSCs (GFP⁺GFAP⁺MCM2⁺) from quiescent NSCs (GFP⁺GFAP⁺MCM2⁻) (Fig. 1D and fig. S1E). *Fmr1* KO mice had a significantly higher percentage of activated NSCs (Fig. 1I) and a higher number of MCM2⁺ cells (fig. S1F) in the adult DG compared to their WT littermates. In addition, the total numbers of GFP⁺ cells, NSCs, and IPCs, as well as activated (MCM2⁺) cells were elevated in both dorsal and ventral DG (fig. S1, G to O). As expected, both NSCs and progenitors exhibited greater proliferation as assessed by cell cycle marker Ki67 in *Fmr1* KO versus WT (Fig. 1, J to L). Thus, a loss of FMRP leads to increased activation and proliferation of NSCs in the adult DG.

To determine whether FMRP regulation of NSC activation is intrinsic to NSCs, we created a triple transgenic line (cKO::Cre::Ai14) by crossing *Fmr1*-floxed (cKO) mice with inducible *Nestin* promoter (Nes)-driven Cre transgenic mice (*Nes-CreER^{T2}*) and *Rosa26-STOP-tdTomato* (Ai14) reporter mice as described previously (12) (Fig. S2A). We injected adult triple transgenic mice (cKO::Cre::Ai14) and control littermates (Cre::Ai14) with tamoxifen to achieve targeted deletion of FMRP specifically in adult NSCs and their subsequent progenies, and at 4 weeks later analyzed the steady-state level of NSC activation by assessing the number of tdTomato-positive (tdT⁺) cells that both expressed GFAP and met the morphological criteria of NSCs. We found that the cKO::Cre::Ai14 mice had significantly more NSCs (tdT⁺GFAP⁺) (Fig. 1, M and N and fig. S2B) and elevated NSC activation compared to Cre::Ai14 littermates (Fig. 1O). Again, the overall volume of the DG was unchanged (fig. S2C). Our data suggest that FMRP regulates NSC activation via an intrinsic mechanism within NSCs.

FMRP regulates MDM2 expression in adult neural stem/progenitor cells

FMRP is known to bind a subset of specific mRNAs in the brain and regulate their translation (4, 14). Since we aim to discover FMRP targets that regulate NSC activation, we used published FMRP PAR-CLIP data on proliferating cells (HEK 293 cells) (3) to select candidate FMRP targets. Given that the number of FMRP-associated reads precipitated for each mRNA reflects the likelihood of interaction, we chose an arbitrary threshold of 1000 reads, yielding 417 prospective FMRP-regulated mRNAs (Fig. 2A, “FMRP Candidate Targets”, Table S1). Meanwhile, we searched the Mammalian Adult Neurogenesis Gene Ontology (MANGO) database (15) and literature to curate a set of 16 “Neurogenic Regulators” with well-established roles in adult NSC quiescence and activation (Fig. 2A; Table S2). Among them, only MDM2 (murine double minute-2 protein) also appeared as one of the 417 FMRP Candidate Targets. Since FMRP may influence neurogenesis by acting either directly on these 16 Neurogenic Regulators or on related proteins participating in their functional pathways, we used the BioGRID 3.4 protein interaction database and GeneMania to identify 322 proteins that have literature-supported physical interactions with the 16 Neurogenic Regulators, including some of the core Neurogenic Regulators themselves (Table S3). Comparing these 335 “Neurogenic Interactors” to the 417 candidate FMRP targets revealed an intersection of 29 putative FMRP targets known to participate in neurogenic pathways. Candidates were ranked by the number of pathways in which they participate, and MDM2 stood alone with four interactions supported by twenty-five publications (Table S4). Importantly, the known interaction between MDM2 and its target, P53, is a proven mechanism for regulating NSC activation (16). Because MDM2 was among our core Neurogenic Regulators and exhibited the highest degree of interaction with strongly supported neurogenic pathways, we chose to focus on MDM2 as the mediator of FMRP-regulated aNSC activation.

Next we used primary neural stem/progenitor cells (aNPCs) isolated from the adult DG as a model system to interrogate the mechanism behind FMRP’s regulation of aNSC activation. We confirmed that *Fmr1* KO aNPCs indeed exhibit increased proliferation and astroglial production but reduced neuronal differentiation (Fig. S3), as described previously (11–13). To determine whether FMRP regulates MDM2, we performed RNA binding protein

immunoprecipitation and identification of co-precipitated RNA (RNA-IP) with an FMRP antibody coupled with real-time PCR analysis of *Mdm2* mRNA. We confirmed that FMRP associated with *Mdm2* mRNA in aNPCs (Fig. 2B). We also detected the association of FMRP with *Gsk3 β* and *Map1b* mRNAs, two known targets of FMRP (11, 17), but not to *Gapdh* mRNA (18) (Fig. 2, C to E), validating the specificity of our assay.

We next found that both *Mdm2* mRNA and MDM2 protein levels were higher in *Fmr1* KO aNPCs compared to WT aNPCs (Fig. 2, F to H; fig. S4 and fig. S5A). The higher *Mdm2* mRNA levels could be due to either increased gene transcription or increased mRNA stability. Because FMRP is known to associate with target mRNAs and modulate mRNA stability (19), we evaluated *Mdm2* mRNA decay kinetics over an 8-hour period in *Fmr1* KO and WT aNPCs in the presence of an inhibitor of transcription actinomycin D. We found that *Mdm2* mRNA exhibited fast decay kinetics, with most (> 90%) RNA disappearing within 4 hours in both WT and KO NSCs (fig. S6A). We then repeated this experiment by analyzing mRNA over a 120-min period. We found that that *Mdm2* mRNA had a longer half-life in *Fmr1* KO ($T_{1/2}$ = 58.7 – 61.3 minutes) versus WT ($T_{1/2}$ = 44.4 – 45.0 minutes) aNPCs (Fig. 2I, fig. S6, B and C), whereas the decay of control *Gapdh* and *Ywhaz* mRNA exhibited no difference between genotypes (fig. S6, D and E). Therefore, FMRP controls MDM2 levels in aNPCs at least in part by reducing the half-life of *Mdm2* mRNA.

FMRP regulates MDM2 phosphorylation through AKT pathway

The elevated MDM2 protein levels in *Fmr1* KO aNPCs prompted us to assess the levels of phosphorylated MDM2 (p-MDM2), the active form of MDM2 (Fig. 3A). MDM2 has multiple phosphorylation sites. Phosphorylation on serine 395 and 407 is known to regulate cell death (20), whereas phosphorylation of MDM2 on serine 166 and 186 is related to cell growth (21). Since we found no altered cell death or survival in *Fmr1* KO aNPCs compared to WT aNPCs (fig. S7), we assessed the phosphorylation of serine 166, which is known to be simultaneously phosphorylated with serine 186 by p-AKT in response to cell growth signals (22). We found that *Fmr1* KO aNPCs had more p-MDM2 (Fig. 3, B and C) and higher p-MDM2 to total MDM2 ratio (Fig. 3D) compared to WT cells, suggesting that increased p-MDM2 in *Fmr1* KO aNPCs levels may result from both increased total MDM2 protein levels and elevated phosphorylation. Activated AKT (AKT phosphorylated at serine 473 or threonine 308) is known to phosphorylate MDM2 at serine 166 and 168 (22), and FMRP deficiency results in increased AKT phosphorylation at Serine 473 in neurons(23). Elevated PI3K/AKT signaling in adult NSCs leads to increased NSC activation (24). We found that *Fmr1* KO aNPCs had a higher ratio of p-AKT to total AKT than WT aNPCs (Fig. 3, E and F). Thus *Fmr1* KO aNPCs have increased p-MDM2 levels due to both increased total MDM2 levels and elevated phosphorylation, possibly by p-AKT.

MDM2 phosphorylation on Ser166 is necessary in response to growth factor and mitogenic signaling, leading to increased MDM2-mediated ubiquitylation and degradation of P53 (25, 26). P53 is known to repress NSC activation (27), and we found that *Fmr1* KO aNPCs had reduced levels of P53 (Fig. 3, G and H). Treatment with MK-2206, a specific inhibitor of AKT phosphorylation (28) (Fig. 3A) led to reduced p-AKT (Fig. 3, I and J) and reduced p-MDM2 (Fig. 3, I and K), without affecting the total MDM2 level (Fig. 3, I and L) in both

KO and WT aNPCs. In addition, MK-2206 treatment enhanced the levels of P53 in *Fmr1* KO aNPCs (Fig. 3, I and M). Thus, in addition to controlling MDM2 expression levels through mRNA stability, FMRP also restricts p-MDM2 levels through the AKT pathway. The absence of FMRP leads to increased p-MDM2 and subsequently reduced P53 levels in aNPCs.

The levels of MDM2 and p-MDM2 directly impact the proliferation and differentiation of aNPCs

To investigate the effect of elevated p-MDM2 levels on the proliferation and differentiation of aNPCs, we acutely knocked down MDM2 expression in aNPCs using lentivirus expressing shRNA against *Mdm2* (*shMdm2*, Fig. 4A and fig. S8A). Lenti-*shMdm2*-treated aNPCs had decreased total MDM2 and p-MDM2 levels (Fig. 4, B to E) and increased P53 levels (Fig. 4, F and G) compared to lenti-*shNC* (non-silencing control shRNA)-treated aNPCs. We next assessed the impact of MDM2 knockdown on aNPC proliferation as assessed by BrdU pulse-labeling, and differentiation as assessed by cell lineage-specific antibodies, β -tubulin III (Tuj1) for neurons and GFAP for astrocytes. Acute knockdown of MDM2 (*shMdm2*) led to reduced cell proliferation in both WT and *Fmr1* KO aNPCs (Fig. 4, H and I), increased neuronal differentiation (Fig. 4, J and K) and reduced astroglial differentiation (Fig. S8, B and C) in both WT and *Fmr1* KO aNPCs. More importantly, the *shMdm2* treatment restored the proliferation and differentiation of *Fmr1* KO aNPCs to WT levels (Lenti-*shNC*-treated) aNPCs (Fig. 4, I and K). Thus, reducing MDM2 levels in aNPCs rescues proliferation and differentiation deficits of FMRP-deficient aNPCs.

We next investigated whether reducing MDM2 phosphorylation alone could also rescue the proliferation and differentiation of *Fmr1* KO aNPCs. We found that MK-2206 treatment resulted in decreased cell proliferation (Fig. 4, L and M) and astroglial differentiation (Fig. S8, D and E), but increased neuronal differentiation (Fig. 4, N and O) in both WT and *Fmr1* KO aNPCs. More importantly, MK-2206-treated *Fmr1* KO aNPCs exhibit similar proliferation and differentiation levels as vehicle-treated WT aNPCs. Therefore, both reducing the expression levels of MDM2 and inhibition of MDM2 phosphorylation can rescue the proliferation and differentiation deficits of *Fmr1* KO aNPCs.

Nutlin-3 treatment rescues the proliferation and differentiation of aNPCs

Although MK-2206 could correct the deficits of *Fmr1* KO aNPCs, its effects on WT aNPCs raised concerns over potentially broad effects resulting from AKT pathway inhibition, with obvious potential therapeutic complications. We therefore searched for a specific inhibitor for MDM2 and found Nutlin-3, a potent MDM2 inhibitor that specifically interrupts MDM2 and P53 interaction (29, 30). Nutlin-3 has been evaluated extensively for its therapeutic potential and mechanism of action in human cancer and is currently in a phase 1 clinical trial for the treatment of retinoblastoma(31). We found that Nutlin-3 treatment did not affect P-MDM2 levels in either *Fmr1* KO or WT cells (Fig. 5, A to C), but led to increased P53 levels in KO cells with no significant effect on WT cells (Fig. 5, D and E). Next we found that Nutlin-3 treatment reduced the proliferation and differentiation of *Fmr1* KO aNPCs without significant effect on WT aNPCs (Fig. 5, F to K). Nutlin-3 also did not exhibit significant effect on cell death and survival of either KO or WT aNPCs (fig. S7, D to F). Thus Nutlin-3,

a specific inhibitor of MDM2-P53 interaction, can rescue the proliferation and differentiation of *Fmr1* KO aNPCs without affecting WT aNPCs.

Nutlin-3 treatment rescues neurogenic deficits in *Fmr1* KO mice

We next investigated whether Nutlin-3 could rescue the neurogenic deficits in *Fmr1* KO mice. We found that in *Fmr1* KO mice, the percentage of p-MDM2⁺ cells among GFP⁺ cells was significantly higher than that in *Fmr1* WT mice (Fig. 6, A to C). In addition, we found that in the brain of a female *Fmr1*^{+/-}::*Nestin-GFP* mouse in which half of its cells expressed FMRP while the other half of the cells did not due to X chromosome inactivation, FMRP and p-MDM2 expression levels exhibited an inverse relationship (fig. S9). Nutlin-3 has been used in mice at high dosages (50 to 200 mg/kg) for 2 weeks or longer to examine its inhibition of cancer cell growth *in vivo* (32–35). Since our goal was to correct rather than totally abolish NSC activation in the adult brain, we chose a 10 mg/kg dosage, which is the lowest dosage with known effects on *in vivo* cell proliferation (36). We also chose a shortened injection period based on literature (37) and our experience (13). We treated *Fmr1* KO and littermate WT mice with either vehicle or Nutlin-3 (10mg/kg) every other day for 5 injections over 9 days. We found that Nutlin-3 treatment specifically reduced NSC activation in *Fmr1* KO mice and brought it to WT levels, with no significant effect on WT mice (Fig. 6, E and F). Since FMRP deficiency impairs adult neurogenesis (11, 12), we next determined whether rebalancing NSC activation via Nutlin-3 treatment could rescue neuronal production (Fig. 6G). Mice received both Nutlin-3 treatment and BrdU injections to label dividing neural progenitors and were analyzed 4 weeks after the last Nutlin-3 treatment. Nutlin-3 treatment rescued both neuronal and astroglial differentiation in *Fmr1* KO mice and brought them to levels similar to those of WT (vehicle-treatment) mice, without significant effect on WT mice (Fig. 6, H to K). These results suggest that Nutlin-3 treatment can correct NSC activation and rescue adult neurogenesis deficits in *Fmr1* KO mice.

Nutlin-3 treatment rescues cognitive deficits in *Fmr1* KO mice

We next investigated whether Nutlin-3 could reverse the behavioral deficits of *Fmr1* KO mice. We showed previously that *Fmr1* KO mice exhibited deficits in two hippocampus-dependent learning tasks: the delayed non-match to place radial arm maze (RAM) and trace conditioning (TC) (12, 13). However, these tests require stressful food restriction or electric shock, so we decided to employ two cognitive tests with minimal stress on the mice: the novel object recognition test (NOR) and novel location test (NLT). First, using *Fmr1* *cKO*::*Cre*::*Ai14* triple transgenic mice (Fig. S2), we confirmed that mice with selective deletion of FMRP from adult NSCs and their progenies exhibited deficits in NLT tests (fig. S10). This result corroborated our previous findings using RAM test (12, 13) and validated the use of NLT for assessing spatial memory of FMRP deficient mice. We then treated littermate WT or *Fmr1* KO mice with either vehicle or Nutlin-3 and analyzed their cognition 4 weeks later (Fig. 7A). Nutlin-3 had no significant effect on overall health and body weight (fig. S11, A and B) or locomotor activity and anxiety (fig. S11, C and D). Consistent with previous studies (18, 38, 39), *Fmr1* KO mice exhibited both impaired NLT spatial learning (Fig. 7, B and C) and defective NOR learning (Fig. 7, D and E) compared to WT mice. Administration of Nutlin-3 rescued both NTL (Fig. 7C) and NOR (Fig. 7E) in *Fmr1* KO mice, with no significant effect on WT mice. Therefore, Nutlin-3 treatment could rescue

cognitive deficits of *Fmr1* KO mice. We found that, similar to *Fmr1* KO mice, postmortem hippocampal and cortical tissues from FXS patients also exhibited higher *MDM2* mRNA and p-MDM2 protein levels compared to those from age and gender-matched controls (fig. S12). Therefore inhibition of p-MDM2 inhibition might be a potential therapy for treating FXS (Fig. 7F).

DISCUSSION

In this study we show that FMRP plays an important role in adult NSC activation via MDM2 level control. We further demonstrate that specific inhibition of MDM2-P53 interaction with a small molecule undergoing clinical trials for cancer rescues both the neurogenic and cognitive deficits of FXS mice. Our findings show that manipulating NSC activation can impact the production of mature neurons contributing to cognition. More importantly, since most adult brain cells are postmitotic, targeting a cell proliferation pathway in NSCs provides much-needed specificity for therapeutic application.

Despite massive efforts, there is no FDA-approved treatment for FXS. Since FMRP is highly expressed in neurons, extensive research has naturally focused on the neuron and identified a number of neurotransmitter receptors and signaling pathways that are either targets or regulators of FMRP(2, 5). These discoveries led to the exciting development of pharmaceuticals for FXS. For example, lithium attenuates GSK3 β activity, which reduces group 1 mGluR-dependent activation of protein translation (40); GSK3 β inhibitors improve cognition in a mouse model of FXS(13, 41); GABA_A receptor agonists reduce anxiety and audiogenic seizures in *Fmr1* KO mice(42); and several mGluR5 antagonists, including Fenobam, RG7090 (Roche), AFQ056 (Novartis), CTEP, MPEP, and STX107 (Seaside Therapeutics), reverse multiple phenotypes in FXS mice (6). However human clinical trials have yielded disappointing results. Although earlier open labeled studies (e.g. Fenobam) showed promising effects, recent double-blind clinical trials for inhibitors of mGluR5 (RG7090, AFQ056, STX107), the best known fragile X drug target, yielded inconclusive results (summarized by(7)), highlighting the importance of searching for other cellular mechanisms and pathways. The contributions to FXS of other cell types, such as glia and immune cells, are being actively pursued (1, 43). We have focused on the involvement of adult neurogenesis in FXS. Although aNSCs constitute a small number of cells in the adult brain, FMRP deficiency in adult-born new neurons directly impacts cognition and restoration of FMRP in adult new neurons rescues these deficits (12) (fig. S8B). We further showed that inhibition of GSK3 β , an FMRP target, rescues adult neurogenesis and learning deficits(44). These data suggest that targeting adult neurogenesis might be an effective treatment for certain cognition deficits in adult FXS. However, GSK3 β is involved in diverse biological processes in many types of brain cells, making drug specificity a major concern. A critical question has been whether treatment aimed at targeting adult NSCs can be an effective therapy for FXS. In this study, we discovered that administration of a small molecule designed for cancer treatment, but used at much lower dosage, could rebalance NSC activation and rescue the cognitive deficits of FXS mice with no effect on WT mice. Adult NSC activation is highly regulated; studies over the past two decades have identified a number of intrinsic regulators and niche factors (8, 9). Most studies have focused on the role of NSC activation in NSC maintenance; the impact of NSC activation on mature neuron

production has gone unexplored. We found that rebalancing NSC activation during the initial neurogenic phase could correct terminal differentiation into neurons and astrocytes (analyzed 30 days later). Therefore, “correcting” NSC activation during the early stage of neurogenesis has long-lasting effects on cell fate. Since most adult brain cells are postmitotic, targeting a cell proliferation pathway may restrict drug action largely to NSCs without significantly affecting postmitotic cells, at last yielding the desired specificity for therapy. The results of this study should be interpreted with consideration of several potential limitations. Complex social and language deficits of FXS are ambiguously modeled in mice, and the effect of Nutlin-3 on these behaviors has not been evaluated. Although *Fmr1* KO mice used in our study recapitulate several key deficits in human FXS patients, the neurocognitive impairment of FXS mouse models is much milder than those seen in human patients, making it difficult to predict the therapeutic efficacy of Nutlin-3 treatment in preclinical trials.

MDM2 is an E3 ubiquitin ligase with many biological functions, from wound healing and carcinogenesis to tissue regeneration. MDM2 is activated via phosphorylation at multiple sites and the sites of phosphorylation are linked to distinct functions (45). Phosphorylation of MDM2 at Tyr394, Ser395, and Ser407 by ATM/ATR is stimulated by DNA damage, leading to inhibition of MDM2, which results in apoptosis. We found no significant difference in cell death between *Fmr1* WT and KO aNPCs; therefore, we decided that MDM2 phosphorylation by ATM/ATR may have no significant role in *Fmr1* KO NPCs. On the other hand, growth factor and mitogenic signaling trigger AKT phosphorylation of MDM2 at Ser166 and Ser186, leading to MDM2 activation and subsequent P53 ubiquitylation and degradation (25, 45). AKT activity is elevated in FMRP-deficient mouse brains and *Drosophila* neuroblasts (46–48), as well as FXS patient-derived lymphoblastoid cells (47). Here, we found a dysregulated AKT-MDM2-P53 pathway in *Fmr1* KO aNPCs, and that inhibiting either AKT or MDM2-P53’s interaction restored the P53 levels and rescued NPC proliferation and differentiation deficits in *Fmr1* KO NPCs. AKT signaling is frequently dysregulated in human cancers, and MK-2206, a potent allosteric inhibitor of all AKT isoforms, has been used to treat patients with tumors in several clinical trials (examples(49, 50)). Although our study might extend the reach of MK-2206 to FXS, one concern is the molecule’s broad effects on many cellular processes in many types of cells. This concern is substantiated by our own observation that, although MK-2206 is highly effective in rescuing *Fmr1* KO NPCs, it also has potent effects on WT aNPCs.

The MDM2-P53 interaction is not only necessary for early embryonic development (51, 52), but also crucial for stem cell maintenance in adults (20). Particularly relevant to the present study, MDM2 is known to inhibit P53 activity and regulate adult NSC activation (16). Besides its well-described proapoptotic effect, P53 is implicated in nervous system development(53), and P53 null mice showed deficits of learning and memory and behavioral alterations (54). We selected Nutlin-3 because it is a potent inhibitor specifically designed to inhibit MDM2-P53 interaction (29), has been evaluated extensively for its therapeutic potential and mechanism of action in human cancer (55), and is currently in phase 1 clinical trials for retinoblastoma(31). To limit the effect of Nutlin-3 on normally proliferating cells in mice, we chose a significantly lower dosage (10× lower) and shorter treatment scheme compared to those used for cancer treatment, and our mice showed no obvious changes in

health and behavior. Another exciting observation is that, unlike with MK-2206, Nutlin-3 treatments exhibit no effect on WT mice or NPCs. Interestingly, although both *Fmr1* KO mice and human FXS individuals exhibit elevated P-MDM2 levels in hippocampal and cortical brain tissues, similar to what we have seen in aNSCs, P53 levels exhibited no difference in mouse cortex and hippocampal tissues (fig. S13) and were too low to be reliably assessed in human tissues. Therefore, it is likely that P53 is not a major target for MDM2 in mature neurons, which comprise a large proportion of cells in heterogeneous brain tissue. It is not surprising that we found very low levels of P53 in cortex and hippocampus, especially in humans, because it has been shown that P53 expression is low in mature neurons and that elevated P53 levels in response to stress signal can lead to neuronal death(56). Therefore, our discovery of the widespread upregulation of P-MDM2 in FXS human and mouse tissues with no apparent impact on apoptotic P53 in neurons suggest that targeting p-MDM2 might be a promising new therapeutic method for FXS. MDM2 has other substrates(57), such as regulating PSD95 in neurons(58). Comprehensive identification of P-MDM2 targets in neurons and specific inhibitors for p-MDM2 interaction with these targets will be a fruitful area of FXS translational research. We do not rule out that in Nutlin-3 treated mice, Nutlin-3 may inhibit p-MDM2 interaction with targets other than P53 and exert effects on other types of cells beside aNSCs, including mature neurons. However, Nutlin-3 was designed specifically to inhibit MDM2-P53 interaction, and much higher dosage is needed for its effect on other MDM2 targets(57). Since we used a low dose of Nutlin-3 in our study, the effects we observed are most likely to be through P53 inhibition in aNSCs compared to other targets or in mature cell types. Cell type-specific targeting of Nutlin-3 action will be required to validate the direct link between restoration of aNSC activity and cognitive rescue, however such method is not yet available. The high specificity of Nutlin-3 towards KO mice, along with the fact that most brain cells are postmitotic and therefore likely to be less sensitive to Nutlin-3, make it an attractive candidate for FXS treatment.

MATERIALS AND METHODS

Study design

The purpose of this study was to investigate the role of FMRP in adult NSC activation and whether rebalancing NSC quiescence and activation could rescue both neurogenesis and cognitive deficits in a FXS mouse model. Based on our publications and power analysis, at least 3 to 6 biological replicates were used for each *in vitro* or *in vivo* biochemical and histological analysis, whereas a sample size of 9 to 21 per group was used for behavioral testing. The aNPCs used for *in vitro* analyses were isolated independently from different *Fmr1* KO and WT littermates and are therefore biological replicates. Data collection was carried out for a predetermined period of time, as dictated by literature- or core facility-based standards, and no exclusion criteria were applied. For drug treatment, animals were randomly assigned to treatment arms with approximately equivalent numbers in each group. All cell counting and behavioral analyses were performed by experimenters who were blind to the identity of the samples. All analyses were performed by scientists blinded to genotype and /or treatment arm.

Mice and treatment

All animal procedures were performed according to protocols approved by the University of Wisconsin-Madison Care and Use Committee. All mice were on C57B/L6 genetic background. The *Fmr1* KO/Nestin-GFP mice (*Fmr1*^{-/-}::*Nestin-GFP*) were created by crossing female *Fmr1* heterozygous KO mice (*Fmr1*^{+/-}) (59) with homozygous Nestin-GFP transgenic males (60). Generation of FMRP inducible conditional mutant mice (*Fmr1*^{loxP/y}::*Nestin-CreERT2*::*Rosa26-tdt*) and Tamoxifen injections to induce recombination were as performed described (12). To induce recombination, mice (6-weeks old) received Tamoxifen (Sigma-Aldrich) daily for 5 d (160 mg/kg) as described (12). Nutlin-3 (10mg/kg) were given to 7–8 weeks old mice through IP injections every other day for 5 injections and sacrificed at either 24 hours or 4 weeks after the last injection. For aNSC differentiation analysis, mice also received four BrdU injections (100mg/kg) between day 2 and 3 of Nutlin-3 injections.

Tissue preparation and immunohistochemistry

Brain tissue processing and histological analysis of mouse brains were performed as described in our publications (11, 12, 61). Briefly mice were euthanized by IP injection of sodium pentobarbital and then transcardially perfused with saline followed by 4% PFA. Brains were dissected out, post-fixed overnight in 4% PFA, and equilibrated in 30% sucrose. 40 µm brain sections were generated using a sliding microtome and stored as floating sections in cryoprotectant solution (glycerol, ethylene glycol, and 0.1M phosphate buffer, pH 7.4, 1:1:2 by volume). We performed immunohistological analysis on 1-in-6 serial floating brain sections (240 µm apart). After staining with primary and fluorescent secondary antibodies (Supplementary Methods), sections were counterstained with DAPI (4', 6-diamidino-2'-phenylindole dihydrochloride, 1:2000, Roche Applied Science, Indianapolis, IN) and then mounted, coverslipped, and maintained at 4°C in the dark until analysis.

In vivo cell quantification

The total numbers of GFP⁺ or tdT⁺ cells in the DG of each animal were counted on a Zeiss Apotome microscope using unbiased stereology (StereoInvestigator software, MBF Biosciences, Inc). A 5-µm guard zone and exhaustive counting method were used (62). DG volume was measured using StereoInvestigator as described (11, 12). Dorsal (bregma -0.94 to -2.30 mm) and ventral (bregma -2.30 to -3.80 mm) hippocampus regions were defined based on literature (63). Phenotypic analysis was performed as described (12). Briefly, at least 50 GFP⁺ or BrdU⁺ cells in the DG were randomly selected, and their colocalization with cell-lineage markers was determined using a Nikon A1 confocal microscope and quantified using ImageJ software (NIH).

Bioinformatics analysis

To select candidate FMRP targets, we utilized FMRP CLIP data from Ascano et al.(3). mRNA transcripts were ranked by the number of reads precipitated in the experiment, and an arbitrary threshold of 1000 reads was chosen to determine the most likely targets of FMRP. This cutoff yielded 417 top genes, or approximately 2% of all probes in the

experiment. 16 genes with known roles in adult NSC activation or proliferation were selected from the Mammalian Adult Neurogenesis Gene Ontology (MANGO(<http://mango.adult-neurogenesis.de>)) database and literature, and were named “Neurogenic Regulators”. For each of these “Neurogenic Regulators”, all known physical interactions were mapped by BioGrid 3.4 (<http://thebiogrid.org>) and GeneMania (<http://www.genemania.org>), producing 16 networks of 335 unique proteins that interact with neurogenic regulators (“Neurogenic Interactors”). Additionally, the number of articles supporting each interaction as listed by BioGrid 3.4 was used as an indicator of the strength of a given interaction. Overlap was then assessed between candidate FMRP targets and both “Neurogenic Regulators” and “Neurogenic Interactors” to find proteins likely to be regulated by FMRP that influence adult NSC activation.

Adult NPC analyses

NPCs were isolated from the DG of 8 to 10-week-old male *Fmr1* KO mice and wild-type (WT) littermate controls based on our published methods(12, 64). Independently isolated cells serve as biological replicates. Proliferation and differentiation of aNPCs was analyzed as described (11, 12, 61). We used only early passage cells (between passage 4 and 10) and only the same passage numbers of wild type (WT) and *Fmr1* KO cells. For each experiment, triplicate wells of cells were analyzed, and results were averaged as one data point (n = 1). At least 3 independent biological replicates were used (n = 3) for statistical analyses.

Chemical treatment

For proliferation assay *in vitro*, 1 μ M MK2206 (Selleckchem, S1078) (AKT phosphorylation inhibitor), or 2 μ M Nutlin-3 (Nt3, Selleckchem, S1061) (Mdm2 inhibitor) was added to proliferating aNPCs at 6 h post-plating. Cells were then pulse labeled 18 hours later with 5 μ M 5-bromo-2'-deoxyuridine (BrdU, Sigma-Aldrich, B5002) and incubated for 6 hours in the presence of inhibitor, followed by fixation, staining, and quantification. For differentiation assay *in vitro*, 1 μ M MK2206 and 2 μ M Nt3 were added to aNPCs upon initiation of differentiation with RA/Fsk for 4 days, followed by fixation, staining and stereological quantification.

RNA immunoprecipitation

RNA-IP was performed as described (11, 61). Briefly, WT and *Fmr1* KO NPCs were harvested and homogenized in 1ml of ice-cold lysis buffer (10 mM Hepes [pH 7.4], 200 mM NaCl, 30 mM EDTA, and 0.5% Triton X-100) with 2 $\times$ complete protease inhibitors (Boehringer-Mannheim). Nuclei and debris were pelleted at 3,000 $\times$ g for 10 min; the supernatant was collected and raised to 300 mM NaCl, and clarified at 14,000 $\times$ g for 30 min. The resulting supernatant was pre-cleared for 1 h with 100 μ L recombinant protein G agarose (Invitrogen) (pre-washed with lysis buffer). An aliquot of pre-cleared input was saved for RNA extraction (200 μ L) and protein analysis (100 μ L). A monoclonal antibody against FMRP (7G1-1, DSHB) or a monoclonal antibody against IgG (5415S, Cell Signaling) was incubated with recombinant protein G dynabeads at 4°C for 2 h and washed 3 times with lysis buffer. RNase Inhibitors (Roche) were added to the remaining lysates. The pre-cleared lysates were immunoprecipitated with antibody-coated recombinant protein G agarose at 4°C for 2 hours. After three washes with lysis buffer, 10% of immunoprecipitate

was saved for protein analysis. The remainder was washed one more time and the immunoprecipitate was re-suspended into Trizol (Invitrogen) for RNA isolation.

Real-time PCR assay

Real-time PCR was performed using standard methods as described (61). The first-strand cDNA was generated by reverse transcription with Oligo (dT) primer (Roche). To quantify the mRNA levels using real-time PCR, aliquots of first-stranded cDNA were amplified with gene-specific primers and Power SYBR Green PCR Master Mix (Bio-Rad) using a Step-1 Real-Time PCR System (Applied Biosystems). The PCR reactions contained 1 µg of cDNA (except the cDNA for the IP, for which 5% of the cDNA was used for each gene examined), Universal Master Mix (Applied Biosystems), and 10 µM of forward and reverse primers in a final reaction volume of 20 µL. The mRNA level of different samples was calculated by the data analysis software built in with the 7300 Real-Time PCR System. For RIP-real time PCR, cDNA from IP and input was used and IP samples were normalized to Input samples. The Sequences of primers are provided in supplementary methods.

Mdm2 mRNA Stability Assay

DG-NPCs from WT and *Fmr1* KO mice were grown in proliferating condition (see above). 10 µg/ml of Actinomycin D (Sigma-Aldrich) was added based on published protocol (61) and total RNA was isolated at various time intervals. The *Mdm2* mRNA level was normalized to *Gapdh* or *Ywhaz* mRNA as measured by real time PCR. RNA decay kinetics and half-life were analyzed using published methods (62). Briefly, we used the exponential function $M = M_0 e^{-\lambda t}$ (M : amount of mRNA at t time, M_0 : amount of mRNA at t time). $\lambda = (\ln)/T_{1/2}$ ($T_{1/2}$ is the half-life of the mRNA).

Western blotting analyses

Protein samples were separated on SDS-PAGE gels (Bio-Rad), transferred to PVDF membranes (Millipore), and incubated with primary antibodies (see Supplementary Methods). After incubation with fluorescence-labeled secondary antibodies (Li-CoR), the membranes were imaged using Li-CoR and quantification was performed using ImageJ software. At least $n=3$ independent blots were used for statistical analysis.

Human tissues

Human tissue was obtained from University of Maryland Brain and Tissue Bank, a Brain and Tissue Repository of the NIH NeuroBioBank. Frozen hippocampal and cortical tissues from two age, gender, and postmortem time-matched FXS (80 and 85 years old) and two control (82 and 84 years old) male Caucasian individuals were used for *MDM2* mRNA and p-MDM2 (S166) analysis.

Novel Location Test (NLT)

This test measures spatial memory through an evaluation of the ability of mice to recognize the new location of a familiar object with respect to spatial cues. The experimental procedure was performed as previously described (65). The mice were exposed to 2 identical objects for 6 minutes during the acquisition phase 3 times and tested at 3 minutes later.

Object preference was evaluated during this session. During the trial session, one of the objects was moved to a novel location, and the mice were allowed to explore the objects for 6 minutes, and the total time spent exploring each object was measured. Exploration was considered as any investigative behavior (i.e., head orientation, sniffing occurring within < 1.0 cm) or deliberate contact occurring with each object. The discrimination index was calculated as the percentage of time spent investigating the object in the new location minus the percentage of time spent investigating the object in the old location: discrimination index = (Novel Location exploration time/total exploration time × 100) – (old Location exploration time/total exploration time × 100). All experiments were videotaped and scored by scientists who were blinded to experimental conditions.

Novel Object Recognition (NOR) Test

This test is based on the natural propensity of rodents to preferentially explore novel objects over familiar ones. The experimental procedure was performed as previously described (38). On the first day, mice were habituated for 10 min to the maze in which the task was performed. On the second day, mice were put back in the maze for 10 min, and two identical objects were presented. 24 hours later, one of the familiar objects was replaced with a novel object, and the mice were again placed in the maze and were allowed to explore for 10 min, and the total time spent exploring each of the two objects (novel and familiar) was measured. The discrimination index was calculated as the difference between the percentages of time spent investigating the novel object and the time spent investigating the familiar objects: discrimination index = (novel object exploration time/total exploration time × 100) – (familiar object exploration time/total exploration time × 100). All experiments were videotaped and scored by scientists who were blinded to experimental conditions.

Statistical analysis

All experiments were randomized and blinded to scientists who performed quantification. Statistical analysis was performed using ANOVA and Student's t-test, unless specified, with the GraphPad software. Two tailed and unpaired t-test was used to compare two conditions. One-Way ANOVA was used for comparison among multiple experimental conditions. Bonferroni post hoc test was used when comparing among each condition. For *Mdm2* mRNA stability analysis, Two-Way ANOVA was used for comparison of the different decay rates. All data were shown as mean with standard error of mean (mean ± SEM). Probabilities of $P < 0.05$ were considered as significant.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Cheryl T. Strauss for editing, Yina Xing, and Erich Berndt for technical assistance, the Zhao lab members for discussion. We thank John Cottrell, Robert Johnson and the NICDH Brain and Tissue Bank at the University of Maryland for providing human postmortem tissues.

Funding: Supported by grants from the NIH (R01MH080434, R01MH078972, and R21NS095632 to X.Z.) and a Center grant from the NIH to the Waisman Center (P30HD03352).

REFERENCES AND NOTES

1. Hagerman RJ, Polussa J. Treatment of the psychiatric problems associated with fragile X syndrome. *Curr Opin Psychiatry*. 2015; 28:107–112. [PubMed: 25602250]
2. Wang T, Bray SM, Warren ST. New perspectives on the biology of fragile X syndrome. *Curr Opin Genet Dev*. 2012; 22:256–263. [PubMed: 22382129]
3. Ascano M, Mukherjee N, Bandaru P, Miller JB, Nusbaum JD, Corcoran DL, Langlois C, Munschauer M, Dewell S, Hafner M, Williams Z, Ohler U, Tuschl T. FMRP targets distinct mRNA sequence elements to regulate protein expression. *Nature*. 2012; 492:382–+. [PubMed: 23235829]
4. Darnell JC, Van Driesche SJ, Zhang CL, Hung KYS, Mele A, Fraser CE, Stone EF, Chen C, Fak JJ, Chi SW, Licatalosi DD, Richter JD, Darnell RB. FMRP Stalls Ribosomal Translocation on mRNAs Linked to Synaptic Function and Autism. *Cell*. 2011; 146:247–261. [PubMed: 21784246]
5. Contractor A, Klyachko VA, Portera-Cailliau C. Altered Neuronal and Circuit Excitability in Fragile X Syndrome. *Neuron*. 2015; 87:699–715. [PubMed: 26291156]
6. Berry-Kravis E. Mechanism-Based Treatments in Neurodevelopmental Disorders: Fragile X Syndrome. *Pediatr Neurol*. 2014; 50:297–302. [PubMed: 24518745]
7. Mullard A. Fragile X disappointments upset autism ambitions. *Nat Rev Drug Discov*. 2015; 14:151–153. [PubMed: 25722228]
8. Jessberger S, Gage FH. Adult neurogenesis: bridging the gap between mice and humans. *Trends Cell Biol*. 2014; 24:558–563. [PubMed: 25124338]
9. Christian KM, Song HJ, Ming GL. Functions and Dysfunctions of Adult Hippocampal Neurogenesis. *Annu Rev Neurosci*. 2014; 37:243–262. [PubMed: 24905596]
10. Li LH, Clevers H. Coexistence of Quiescent and Active Adult Stem Cells in Mammals. *Science*. 2010; 327:542–545. [PubMed: 20110496]
11. Luo YP, Shan G, Guo WX, Smrt RD, Johnson EB, Li XK, Pfeiffer RL, Szulwach KE, Duan RH, Barkho BZ, Li WD, Liu CM, Jin P, Zhao XY. Fragile X Mental Retardation Protein Regulates Proliferation and Differentiation of Adult Neural Stem/Progenitor Cells. *PLoS Genet*. 2010; 6
12. Guo WX, Allan AM, Zong RT, Zhang L, Johnson EB, Schaller EG, Murthy AC, Goggin SL, Eisch AJ, Oostra BA, Nelson DL, Jin P, Zhao XY. Ablation of Fmrp in adult neural stem cells disrupts hippocampus-dependent learning. *Nat Med*. 2011; 17:559–U575. [PubMed: 21516088]
13. Guo WX, Murthy AC, Zhang L, Johnson EB, Schaller EG, Allan AM, Zhao XY. Inhibition of GSK3 beta improves hippocampus-dependent learning and rescues neurogenesis in a mouse model of fragile X syndrome. *Hum Mol Genet*. 2012; 21:681–691. [PubMed: 22048960]
14. Brown V, Jin P, Ceman S, Darnell JC, O'Donnell WT, Tenenbaum SA, Jin XK, Feng Y, Wilkinson KD, Keene JD, Darnell RB, Warren ST. Microarray identification of FMRP-associated brain mRNAs and altered mRNA translational profiles in fragile X syndrome. *Cell*. 2001; 107:477–487. [PubMed: 11719188]
15. Overall RW, Paszkowski-Rogacz M, Kempermann G. The mammalian adult neurogenesis gene ontology (MANGO) provides a structural framework for published information on genes regulating adult hippocampal neurogenesis. *PLoS One*. 2012; 7:e48527. [PubMed: 23139788]
16. Francoz S, Froment P, Bogaerts S, De Clercq S, Maetens M, Doumont G, Bellefroid E, Marine JC. Mdm4 and Mdm2 cooperate to inhibit p53 activity in proliferating and quiescent cells in vivo. *Proc Natl Acad Sci U S A*. 2006; 103:3232–3237. [PubMed: 16492744]
17. Lu R, Wang HP, Liang Z, Ku L, O'Donnell WT, Li W, Warren ST, Feng Y. The fragile X protein controls microtubule-associated protein 1B translation and microtubule stability in brain neuron development. *Proc Natl Acad Sci U S A*. 2004; 101:15201–15206. [PubMed: 15475576]
18. Udagawa T, Farny NG, Jakovcevski M, Kaphzan H, Alarcon JM, Anilkumar S, Ivshina M, Hurt JA, Nagaoka K, Nalavadi VC, Lorenz LJ, Bassell GJ, Akbarian S, Chattarji S, Klann E, Richter JD. Genetic and acute CPEB1 depletion ameliorate fragile X pathophysiology. *Nat Med*. 2013; 19:1473–+. [PubMed: 24141422]
19. De Rubeis S, Bagni C. Fragile X mental retardation protein control of neuronal mRNA metabolism: Insights into mRNA stability. *Molecular and Cellular Neuroscience*. 2010; 43:43–50. [PubMed: 19837168]

20. Gannon HS, Woda BA, Jones SN. ATM phosphorylation of Mdm2 Ser394 regulates the amplitude and duration of the DNA damage response in mice. *Cancer Cell*. 2012; 21:668–679. [PubMed: 22624716]
21. Gottlieb TM, Leal JFM, Seger R, Taya Y, Oren M. Cross-talk between Akt, p53 and Mdm2: possible implications for the regulation of apoptosis. *Oncogene*. 2002; 21:1299–1303. [PubMed: 11850850]
22. Mayo LD, Donner DB. A phosphatidylinositol 3-kinase/Akt pathway promotes translocation of Mdm2 from the cytoplasm to the nucleus. *Proc Natl Acad Sci U S A*. 2001; 98:11598–11603. [PubMed: 11504915]
23. Gross C, Nakamoto M, Yao XD, Chan CB, Yim SY, Ye KQ, Warren ST, Bassell GJ. Excess Phosphoinositide 3-Kinase Subunit Synthesis and Activity as a Novel Therapeutic Target in Fragile X Syndrome. *Journal of Neuroscience*. 2010; 30:10624–10638. [PubMed: 20702695]
24. Bonaguidi MA, Wheeler MA, Shapiro JS, Stadel RP, Sun GJ, Ming GL, Song H. In vivo clonal analysis reveals self-renewing and multipotent adult neural stem cell characteristics. *Cell*. 2011; 145:1142–1155. [PubMed: 21664664]
25. Ogawara Y, Kishishita S, Obata T, Isazawa Y, Suzuki T, Tanaka K, Masuyama N, Gotoh Y. Akt enhances Mdm2-mediated ubiquitination and degradation of p53. *Journal of Biological Chemistry*. 2002; 277:21843–21850. [PubMed: 11923280]
26. Zhou BP, Liaq Y, Xia WY, Zou YY, Spohn B, Hung MC. HER-2/neu induces p53 ubiquitination via Akt-mediated MDM2 phosphorylation (vol 3, pg 973, 2001). *Nat Cell Biol*. 2002; 4:736–736.
27. Wang Y, Yang J, Zheng HR, Tomasek GJ, Zhang P, McKeever PE, Lee EYHP, Zhu Y. Expression of Mutant p53 Proteins Implicates a Lineage Relationship between Neural Stem Cells and Malignant Astrocytic Glioma in a Murine Model. *Cancer Cell*. 2009; 15:514–526. [PubMed: 19477430]
28. Rafalski VA, Ho PP, Brett JO, Ucar D, Dugas JC, Pollina EA, Chow LML, Ibrahim A, Baker SJ, Barres BA, Steinman L, Brunet A. Expansion of oligodendrocyte progenitor cells following SIRT1 inactivation in the adult brain. *Nat Cell Biol*. 2013; 15:614–+. [PubMed: 23644469]
29. Manfredi JJ. The Mdm2-p53 relationship evolves: Mdm2 swings both ways as an oncogene and a tumor suppressor. *Genes Dev*. 2010; 24:1580–1589. [PubMed: 20679392]
30. Vassilev LT, Vu BT, Graves B, Carvajal D, Podlaski F, Filipovic Z, Kong N, Kammlott U, Lukacs C, Klein C, Fotouhi N, Liu EA. In vivo activation of the p53 pathway by small-molecule antagonists of MDM2. *Science*. 2004; 303:844–848. [PubMed: 14704432]
31. Secchiero P, Bosco R, Celeghini C, Zauli G. Recent advances in the therapeutic perspectives of Nutlin-3. *Current pharmaceutical design*. 2011; 17:569–577. [PubMed: 21391907]
32. Zheng T, Yin D, Lu Z, Wang J, Li Y, Chen X, Liang Y, Song X, Qi S, Sun B, Xie C, Meng X, Pan S, Liu J, Jiang H, Liu L. Nutlin-3 overcomes arsenic trioxide resistance and tumor metastasis mediated by mutant p53 in Hepatocellular Carcinoma. *Mol Cancer*. 2014; 13:133. [PubMed: 24884809]
33. Kunkele A, De Preter K, Heukamp L, Thor T, Pajtler KW, Hartmann W, Mittelbronn M, Grotzer MA, Deubzer HE, Speleman F, Schramm A, Eggert A, Schulte JH. Pharmacological activation of the p53 pathway by nutlin-3 exerts anti-tumoral effects in medulloblastomas. *Neuro Oncol*. 2012; 14:859–869. [PubMed: 22591662]
34. Tovar C, Rosinski J, Filipovic Z, Higgins B, Kolinsky K, Hilton H, Zhao X, Vu BT, Qing W, Packman K, Myklebost O, Heimbrook DC, Vassilev LT. Small-molecule MDM2 antagonists reveal aberrant p53 signaling in cancer: implications for therapy. *Proc Natl Acad Sci U S A*. 2006; 103:1888–1893. [PubMed: 16443686]
35. Ye F, Lattif AA, Xie J, Weinberg A, Gao S. Nutlin-3 induces apoptosis, disrupts viral latency and inhibits expression of angiopoietin-2 in Kaposi sarcoma tumor cells. *Cell Cycle*. 2012; 11:1393–1399. [PubMed: 22421142]
36. Mouraret N, Marcos E, Abid S, Gary-Bobo G, Saker M, Houssaini A, Dubois-Rande JL, Boyer L, Boczkowski J, Derumeaux G, Amsellem V, Adnot S. Activation of lung p53 by Nutlin-3a prevents and reverses experimental pulmonary hypertension. *Circulation*. 2013; 127:1664–1676. [PubMed: 23513067]

37. Zhou MO, Li WD, Huang S, Song J, Kim JY, Tian XL, Kang EC, Sano Y, Liu C, Balaji J, Wu SM, Zhou Y, Zhou Y, Parivash SN, Ehninger D, He L, Song HJ, Ming GL, Silva AJ. mTOR Inhibition Ameliorates Cognitive and Affective Deficits Caused by Disc1 Knockdown in Adult-Born Dentate Granule Neurons. *Neuron*. 2013; 77:647–654. [PubMed: 23439118]
38. Busquets-Garcia A, Gomis-Gonzalez M, Guegan T, Agustin-Pavon C, Pastor A, Mato S, Perez-Samartin A, Matute C, de la Torre R, Dierssen M, Maldonado R, Ozaita A. Targeting the endocannabinoid system in the treatment of fragile X syndrome. *Nat Med*. 2013; 19:603–607. [PubMed: 23542787]
39. King MK, Jope RS. Lithium treatment alleviates impaired cognition in a mouse model of fragile X syndrome. *Genes Brain and Behavior*. 2013; 12:723–731.
40. Yuskaitis CJ, Mines MA, King MK, Sweatt JD, Miller CA, Jope RS. Lithium ameliorates altered glycogen synthase kinase-3 and behavior in a mouse model of Fragile X syndrome. *Biochem Pharmacol*. 2010; 79:632–646. [PubMed: 19799873]
41. Franklin AV, King MK, Palomo V, Martinez A, McMahon LL, Jope RS. Glycogen synthase kinase-3 inhibitors reverse deficits in long-term potentiation and cognition in fragile X mice. *Biol Psychiatry*. 2014; 75:198–206. [PubMed: 24041505]
42. Heulens I, D'Hulst C, Van Dam D, De Deyn PP, Kooy RF. Pharmacological treatment of fragile X syndrome with GABAergic drugs in a knockout mouse model. *Behav Brain Res*. 2012; 229:244–249. [PubMed: 22285772]
43. Li Y, Zhao X. Concise review: Fragile X proteins in stem cell maintenance and differentiation. *Stem Cells*. 2014; 32:1724–1733. [PubMed: 24648324]
44. Guo W, Murthy AC, Zhang L, Johnson EB, Schaller EG, Allan AM, Zhao X. Inhibition of GSK3beta improves hippocampus-dependent learning and rescues neurogenesis in a mouse model of fragile X syndrome. *Hum Mol Genet*. 2012; 21:681–691. [PubMed: 22048960]
45. Meek DW, Hupp TR. The regulation of MDM2 by multisite phosphorylation--opportunities for molecular-based intervention to target tumours? *Semin Cancer Biol*. 2010; 20:19–28. [PubMed: 19897041]
46. Gross C, Raj N, Molinaro G, Allen AG, Whyte AJ, Gibson JR, Huber KM, Gourley SL, Bassell GJ. Selective role of the catalytic PI3K subunit p110beta in impaired higher order cognition in fragile X syndrome. *Cell Rep*. 2015; 11:681–688. [PubMed: 25921527]
47. Gross C, Bassell GJ. Excess protein synthesis in FXS patient lymphoblastoid cells can be rescued with a p110beta-selective inhibitor. *Mol Med*. 2012; 18:336–345. [PubMed: 22207187]
48. Callan MA, Clements N, Ahrendt N, Zarnescu DC. Fragile X Protein is required for inhibition of insulin signaling and regulates glial-dependent neuroblast reactivation in the developing brain. *Brain Res*. 2012; 1462:151–161. [PubMed: 22513101]
49. Ahn DH, Li J, Wei L, Doyle A, Marshall JL, Schaaf LJ, Phelps MA, Villalona-Calero MA, Bekaii-Saab T. Results of an abbreviated phase-II study with the Akt Inhibitor MK-2206 in Patients with Advanced Biliary Cancer. *Sci Rep*. 2015; 5:12122. [PubMed: 26161813]
50. Lara PN Jr, Longmate J, Mack PC, Kelly K, Socinski MA, Salgia R, Gitlitz B, Li T, Koczywas M, Reckamp KL, Gandara DR. Phase II Study of the AKT inhibitor MK-2206 plus Erlotinib in Patients with Advanced Non-Small Cell Lung Cancer who Previously Progressed on Erlotinib. *Clin Cancer Res*. 2015
51. Jones SN, Roe AE, Donehower LA, Bradley A. Rescue of embryonic lethality in Mdm2-deficient mice by absence of p53. *Nature*. 1995; 378:206–208. [PubMed: 7477327]
52. Montes de Oca Luna R, Wagner DS, Lozano G. Rescue of early embryonic lethality in mdm2-deficient mice by deletion of p53. *Nature*. 1995; 378:203–206. [PubMed: 7477326]
53. Tedeschi A, Di Giovanni S. The non-apoptotic role of p53 in neuronal biology: enlightening the dark side of the moon. *EMBO Rep*. 2009; 10:576–583. [PubMed: 19424293]
54. Amson R, Lassalle JM, Halley H, Prieur S, Lethrosne F, Roperch JP, Israeli D, Gendron MC, Duyckaerts C, Checler F, Dausset J, Cohen D, Oren M, Telerman A. Behavioral alterations associated with apoptosis and down-regulation of presenilin 1 in the brains of p53-deficient mice. *Proc Natl Acad Sci U S A*. 2000; 97:5346–5350. [PubMed: 10805794]
55. Carry JC, Garcia-Echeverria C. Inhibitors of the p53/hdm2 protein-protein interaction-path to the clinic. *Bioorg Med Chem Lett*. 2013; 23:2480–2485. [PubMed: 23541651]

56. Morrison RS, Kinoshita Y. The role of p53 in neuronal cell death. *Cell Death Differ.* 2000; 7:868–879. [PubMed: 11279532]
57. Wang, S.; Zhao, Y.; Bernard, D.; Aguilar, A.; Kumar, S. Protein-Protein Interactions. Wendt, MD., editor. Berlin Heidelberg: Springer; 2012. p. 57-79.
58. Tsai NP, Wilkerson JR, Guo W, Maksimova MA, DeMartino GN, Cowan CW, Huber KM. Multiple autism-linked genes mediate synapse elimination via proteasomal degradation of a synaptic scaffold PSD-95. *Cell.* 2012; 151:1581–1594. [PubMed: 23260144]
59. T. D.-B. F. X. Consortium. Fmr1 knockout mice: a model to study fragile X mental retardation. The Dutch-Belgian Fragile X Consortium. *Cell.* 1994; 78:23–33. [PubMed: 8033209]
60. Yamaguchi M, Saito H, Suzuki M, Mori K. Visualization of neurogenesis in the central nervous system using nestin promoter-GFP transgenic mice. *Neuroreport.* 2000; 11:1991–1996. [PubMed: 10884058]
61. Guo WX, Zhang L, Christopher DM, Teng ZQ, Fausett SR, Liu CM, George OL, Klingensmith J, Jin P, Zhao XY. RNA-Binding Protein FXR2 Regulates Adult Hippocampal Neurogenesis by Reducing Noggin Expression. *Neuron.* 2011; 70:924–938. [PubMed: 21658585]
62. Guo WX, Polich ED, Su J, Gao Y, Christopher DM, Allan AM, Wang M, Wang FF, Wang GF, Zhao XY. Fragile X Proteins FMRP and FXR2P Control Synaptic GluA1 Expression and Neuronal Maturation via Distinct Mechanisms. *Cell Rep.* 2015; 11:1651–1666. [PubMed: 26051932]
63. Eadie BD, Zhang WN, Boehme F, Gil-Mohapel J, Kainer L, Simpson JM, Christie BR. Fmr1 knockout mice show reduced anxiety and alterations in neurogenesis that are specific to the ventral dentate gyrus. *Neurobiol Dis.* 2009; 36:361–373. [PubMed: 19666116]
64. Guo WX, Patzlaff NE, Jobe EM, Zhao XY. Isolation of multipotent neural stem or progenitor cells from both the dentate gyrus and subventricular zone of a single adult mouse. *Nat Protoc.* 2012; 7:2005–2012. [PubMed: 23080272]
65. Contestabile A, Greco B, Ghezzi D, Tucci V, Benfenati F, Gasparini L. Lithium rescues synaptic plasticity and memory in Down syndrome mice. *J Clin Invest.* 2013; 123:348–361. [PubMed: 23202733]

Figure 1. FMRP deficiency leads to increased NSC activation in the adult DG

(A) Schematic diagram showing stages of neurogenesis in the adult DG and cell lineage-specific markers. qNSC, quiescent NSC. aNSC, activated NSC. (B and C) Quantitative comparison of the numbers of total GFP⁺ cells (B) and the density of GFP⁺ Cells (C) in the DG of adult *Fmr1* KO and WT littermate control mice. (D) Sample confocal images used in E, F and I for identifying NSCs (GFP⁺GFAP⁺), progenitors (GFP⁺GFAP⁻), activated NSCs (GFP⁺GFAP⁺MCM2⁺) in the DG of adult *Fmr1* KO and WT mice bred onto a *Nestin-GFP* mouse line. Blue, DAPI; Green, GFP; Red, MCM2; white, GFAP. Scale bars: 20μm. (E and

F) Quantitative comparison of the numbers of NSCs (E) and IPCs (F) in the DG. (**G, H**), Sample confocal images (G) and quantitative comparison of the numbers of IPCs (GFP⁺TBR2⁺) in the adult DG. (**I**) Comparison of the percentage of activated NSCs (GFP⁺GFAP⁺MCM2⁺) among the total NSCs (GFP⁺GFAP⁺) in the DG. (**J**) Sample confocal images used in K and L for identifying proliferating (Ki67⁺) NSCs (GFP⁺GFAP⁺Ki67⁺) and progenitors (GFP⁺GFAP⁺Ki67⁺). Green, GFP; Red, Ki67; White, GFAP; Arrow heads indicate proliferating aNSCs; Arrows indicate proliferating IPCs; Scale bars: 20μm. (**K**) Comparison of the percentage of proliferating among total NSCs in the DG. (**L**) Comparison of the percentage of proliferating among total progenitors. (**M**) Sample confocal images from *Fmr1*-cKO mice used for identifying activated NSCs (tdT⁺GFAP⁺MCM2⁺). Green, MCM2; Red, tdT; White, GFAP. Scale bars: 20μm. (**N**) Quantitative comparison of the total numbers of activated NSCs in the DG of cKO::Cre::Ai14 mice and Cre;Ai14 control mice. (**O**) Comparison of the percentage of activated among total NSCs in the DG (n = 6 per genotype). Student's t-tests were used for all data in this figure. **P < 0.01, ***P < 0.001. From A to N, n = 5 per genotype. Data are presented as mean ± SEM.

Figure 2. FMRP regulates MDM2 expression in adult NPCs

(A) Workflow for selecting potential FMRP targeted genes that may regulate adult NSC activation. (B to E) FMRP RNA-IP followed by quantitative real-time PCR analyses for *Mdm2* (B), *GSK3β* (C), *Map1b* (D), and *Gapdh* (E, negative control) mRNAs in *Fmr1* WT and KO aNPCs (n = 3). The levels of each mRNA in IP samples were normalized to that in the input samples. (F) Schematic model showing that FMRP inhibits the levels of *Mdm2* mRNA, which in turn leads to reduced MDM2 protein levels in aNPCs. (G and H) Quantification analyses of *Mdm2* mRNA (G) and MDM2 protein (H) levels in WT and *Fmr1* KO aNPCs (n = 3). *Gapdh* was used as the internal control for qPCR analysis, and GAPDH was used as a loading control for western blot analysis. (I) *Mdm2* mRNA stability in *Fmr1* WT and KO aNPCs treated with Actinomycin D to inhibit transcription. The percentage of *Mdm2* mRNA remaining in aNPCs was quantified using real-time PCR. Comparisons of the different decay rates were performed by Two-Way ANOVA: (n = 3). Half-life of decay was calculated after ln2 transformation. Except for Figure 2I, Student's t-

tests were used for data analyses in this figure. Data are presented as mean $\pm$ SEM. * $P < 0.05$; ** $P < 0.01$. n.s., no significant difference.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 3. FMRP regulates MDM2 phosphorylation via the AKT pathway

(A) Schematic drawing showing that, in addition to inhibiting *Mdm2* mRNA levels, FMRP can also inhibit MDM2 phosphorylation at Ser166/168 through AKT signaling. MK2206, a selective AKT inhibitor. (B to D) Western blot analyses (B) of total MDM2 (C) and phosphorylated MDM2 at Ser166 (p-MDM2, D) in *Fmr1* WT and KO aNPCs, (n = 3). (E and F) Western blot analyses of total and phosphorylated AKT at Ser473 (p-AKT) (n = 3). (G and H) Western blot analyses of P53 showing that P53 levels were decreased in *Fmr1* KO aNPCs compared with WT cells (n = 3). GAPDH was used as a loading control. (I to

M) Representative western blot image (I) and quantitative analyses of p-AKT (J), p-MDM2 (K), total MDM2 (L), and P53 (M) levels in *Fmr1* KO and WT aNPCs treated with MK2206 or vehicle (Veh). Total AKT was used as a loading control for p-AKT, and GAPDH was used as a loading control for the others (n = 3). *P< 0.05; **P<0.01; ***P< 0.001; n.s., no significant difference; student's t-tests were used in C–H; One-Way ANOVA were used in J–M. Data are presented as mean ± SEM.

Figure 4. The levels of MDM2 and p-MDM2 directly impact the proliferation and differentiation of aNPCs

(A) Schematic model showing that both acute knockdown of MDM2 and treatment with MK-2206 should repress MDM2 levels, relieve P53 repression, and rescue proliferation and differentiation of *Fmr1* KO aNPCs. (B to E) Western blot analyses (B) of MDM2 (C) and p-MDM2 (D) in aNPCs with acute knockdown of MDM2 using *shMdm2*, $n = 3$, normalized to GAPDH or total MDM2. (F and G) Western blot analyses (F) of P53 in cells with acute knockdown of MDM2 by *shMdm2* (G), ($n = 3$). GAPDH was used as a loading control. (H)

Representative images showing that both *Fmr1* WT and KO aNPCs incorporated BrdU (Red) under proliferating conditions, with or without *shMdm2* treatment (Green). Scale bars: 20µm. **(I)** Quantitative analysis showing that acute knockdown of MDM2 by *shMdm2* reduced the proliferation rate of both *Fmr1* WT and KO aNPCs (n = 3). **(J and K)** Acute knockdown of MDM2 by *shMdm2* rescued neuronal differentiation phenotypes of *Fmr1* KO aNPCs, as assessed by neuronal marker Tuj1⁺ (J, Red for Tuj1; Green for *shMdm2* or *shNC* viral-infected cells) and quantification (K). (n = 3). **(L)** Both WT and KO aNPCs incorporated BrdU (Red) under proliferating conditions. Scale bars: 20µm. **(M)** MK-2206 treatment reduced the proliferation rate of both *Fmr1* WT and KO aNPCs (n = 3). **(N and O)** MK-2206 treatment rescued neuronal differentiation phenotypes of *Fmr1* KO aNPCs, (n = 3)., One-Way ANOVA was used for all data analyses in this figure. *P < 0.05; **P < 0.01; ***P < 0.001. Data are presented as mean ± SEM.

Figure 5. Nutlin-3 treatment rescues the proliferation and differentiation of aNPCs *in vitro*
(A) Schematic drawing showing that Nutlin-3 inhibits the interaction between p-MDM2(S166/S168) and P53 to relieve the repression of P53 and rescue the proliferation and neuronal differentiation of aNPCs. **(B to E)** Western blot analyses of protein levels of p-MDM2 (**B, C**) and P53 (**D, E**) in *Fmr1* WT and KO aNPCs with Nutlin-3 treatment showing Nutlin-3 treatment had no significant effect on p-MDM2 levels in both KO and WT cells but specifically rescued the protein levels of P53 in *Fmr1* KO aNPCs without affecting WT cells. (n = 3). GAPDH was used as a loading control in western blot analysis. **(D)** Both *Fmr1*

WT and KO aNPCs incorporated BrdU (Red) under proliferating conditions. Scale bars: 20 μ m. **(E)** Quantitative analysis showing that Nutlin-3 treatment specifically reduced the high proliferation rate of *Fmr1* KO aNPCs (n = 3) without affecting WT cells. **(F and G)** Nutlin-3 treatment rescued cell proliferation phenotype of *Fmr1* KO aNPCs, as demonstrated by immunostaining cells using cell proliferation marker BrdU (F, Red), followed by quantitative analysis of BrdU⁺ cells (G), (n=3). **(H and I)** Nutlin-3 treatment rescued neuronal differentiation phenotypes of *Fmr1* KO aNPCs, as assessed by a neuronal marker Tuj1⁺(H, Red) and quantitative analysis. (I, n = 3). **(J and K)** Nutlin-3 treatment specifically rescued astroglial differentiation phenotypes of *Fmr1* KO aNPCs, as assessed by astroglial marker GFAP⁺ (J, Green) followed by quantitative analysis (K, n = 3). **P<0.01; ***P< 0.001. One-Way ANOVA was used for all data analyses in this figure. Data are presented as mean $\pm$ SEM.

Figure 6. Nutlin-3 treatment can rescue neurogenesis deficits in *Fmr1* KO mice

(A to C) Sample confocal images of $GFP^+p\text{-MDM2}^+$ (A) and $GFP^+p\text{-MDM2}^-$ (B) cells in the adult DG and quantitative comparison of the percentage of $p\text{-MDM2}^+$ among GFP^+ cells between *Fmr1* KO and WT mice (C, $n = 3$, $**P < 0.01$, Student's t-test). Blue, DAPI; Green, GFP; Red, p-MDM2(S166); Scale bars: 20 μm . (D) Experimental scheme for assessing aNSC activation in *Fmr1* WT and KO mice treated with Nutlin-3 or vehicle. (E) Sample confocal images of activated NSCs ($GFP^+GFAP^+MCM2^+$) in the DG of adult *Fmr1* WT/KO-*Nestin-GFP* mice. Scale bar: 20 μm . (F) Comparison of the percentage of activated among total NSCs in the DG of *Fmr1* KO mice and WT mice with or without Nutlin-3

treatment (n = 3 per group). **(J)** Experimental scheme for assessing aNSC differentiation in *Fmr1* WT and KO mice treated with Nutlin-3 or vehicle. **(H and I)** Nutlin-3 treatment rescued neuronal differentiation in *Fmr1* KO mice, as assessed by immunostaining with neuronal marker NeuN (Green) and BrdU (Red) (H, Scale bar: 20μm) followed by quantification of percentage of neurons among BrdU⁺ cells (I), (n = 3–5 per group). **(J and K)** Nutlin-3 treatment rescued astroglial differentiation specifically in *Fmr1* KO mice, as assessed by an astroglial marker S100β (Green) and BrdU (Red) (J, Scale bar: 20μm) followed by quantification of the percentages of astrocytes among BrdU⁺ cells (K), (n = 3–5 per group). **P < 0.01; ****P < 0.0001; One-Way ANOVA. Data are presented as mean ± SEM. n.s.; no significant difference.

Figure 7. Nutlin-3 treatment rescues cognitive functions in *Fmr1* KO mice

(A) Experimental scheme for assessing cognitive functions in *Fmr1* WT and KO mice with Nutlin-3 treatment. (B) Schematic drawing of NLT for assessing spatial learning. (C) Nutlin-3 treatment fully rescued spatial memory deficits in *Fmr1* KO mice in the NLT test (n = 9–13 mice per group). (D) Schematic drawing of NOR test for novel object recognition. (E) Nutlin-3 treatment fully rescued NOR deficits in *Fmr1* KO mice in the NOR test (n = 8–14 mice per group), **P < 0.01; ***P < 0.001, One-Way ANOVA was used for all data analyses in this figure. Data are presented as mean ± SEM. (F) Models for FMRP regulation of the MDM2 and P53 pathway, which significantly impacts adult neurogenesis and cognition.