Published in final edited form as:

Int J Gynecol Cancer. 2015 October ; 25(8): 1461–1467. doi:10.1097/IGC.0000000000000519.

Complementary prognostic value of pelvic MRI and whole-body FDG PET/CT in the pretreatment assessment of patients with cervical cancer

Evis Sala, MD, PhD¹, Maura Micco, MD¹, Irene A. Burger, MD¹, Derya Yakar, MD¹, Marisa A. Kollmeier, MD², Debra A. Goldman, MS³, Mithat Gonen, PhD³, Kay J. Park, MD⁴, Nadeem R. Abu-Rustum, MD⁵, Hedvig Hricak, MD, PhD¹, and Hebert Alberto Vargas, MD¹

¹Department of Radiology, Memorial Sloan-Kettering Cancer Center, 1275 York Ave, New York, NY 10065

²Department of Radiation Oncology, Memorial Sloan-Kettering Cancer Center, 1275 York Ave, New York, NY 10065

³Department of Epidemiology-Biostatistics, Memorial Sloan-Kettering Cancer Center, 307 E 63rd Street, New York, NY 10065

⁴Department of Pathology, Memorial Sloan-Kettering Cancer Center, 1275 York Ave, New York, NY 10065

⁵Department of Surgery, Gynecologic Oncology Service, Memorial Sloan-Kettering Cancer Center, 1275 York Ave, New York, NY 10065

Abstract

Objective—To evaluate the incremental prognostic value of pelvic magnetic resonance imaging (MRI) and whole-body 18F-fluorodeoxyglucose (FDG) positron-emission tomography/computed tomography (PET/CT) findings compared to clinical-histopathologic factors in patients with newly-diagnosed cervical cancer.

Methods—The institutional review board approved this retrospective study of 114 patients (median age 40.6 years) with International Federation of Gynecology and Obstetrics (FIGO) stage I-IVB cervical cancer who underwent pre-treatment MRI and PET/CT. All scans were reviewed for locoregional tumor extent, pelvic or/and paraaortic lymphadenopathy, and distant metastases. Univariate Cox proportional hazard regression was performed to evaluate associations between clinical-histopathologic factors, imaging findings and progression-free survival (PFS). Multivariate models were built using independent predictors for PFS. Harrell's C was used to measure concordance (C-index).

Results—Forty patients progressed within a median time of 10.4 months (range: 0.4–40.3 months). At univariate analysis, age, FIGO stage, tumor histology, tumor grade and all MRI and PET/CT features were significantly associated with PFS ($p < 0.0001$ – $p = 0.0474$). A multivariate

model including clinical and imaging parameters (parametrial invasion on MRI and paraaortic lymphadenopathy/distant metastases on PET/CT) had significantly higher concordance for predicting PFS than a model including clinical parameters only (C-index: 0.81 [95%CI: 0.75–0.87] vs. 0.68 [95% CI: 0.59–0.78], $p<0.001$). The comparison of C-indices for the combined clinical and imaging model approached significance when compared to a FIGO stage model (C-index: 0.81 [95%CI: 0.75–0.87] vs. 0.75 [95% CI: 0.69–0.82, $p=0.058$).

Conclusion—In patients with newly-diagnosed cervical cancer, a prognostic model including combined MRI and PET/CT findings provides information that complements clinical and histopathologic factors.

Introduction

In patients with newly-diagnosed cervical cancer, the prediction of prognosis is currently based on the International Federation of Gynecologists and Obstetricians (FIGO) clinical staging system, which has its known limitations^{1,2}. The survival rate of patients with the same FIGO stage may vary³, therefore, FIGO stage alone may not be entirely satisfactory to accurately provide prognostic information in patients with cervical cancer. The prognostic relevance of other pretreatment clinical-histopathological variables such as patient's age, tumor histology and grade, tumor diameter, deep stromal and parametrial invasion, lymph vascular invasion and lymph node (LN) metastases has also been evaluated. In a study of 187 patients with stage Ib₂–IIb cervical cancer, pathologically assessed parametrial invasion and LN status were predictors of disease-free survival and overall survival⁴ independent of other clinical-histopathological factors.

Few previous reports have simultaneously analyzed the prognostic value of multiple variables (including imaging findings) in patients with cervical cancer to create a nomogram able to provide individualized prediction of outcome^{5–9}. These nomograms which include age, histologic type, tumor size, parametrial involvement, pelvic organ invasion and LN status, have been shown to compare favorably with FIGO staging system. In patients with locally advanced cervical cancer (LACC), Kang et al.⁵ developed a four-parameter risk assessment model to predict recurrence, which included pelvic/paraortic LN metastasis detected on 18F-fluorodeoxyglucose (FDG) positron-emission tomography/computed tomography (PET/CT) and clinical-histopathological factors. In contrast, Shim SH et al.⁹ introduced a three variable risk assessment model to predict 5-year survival in patients with LACC, which incorporated histology, tumor size, and paraaortic LN metastasis detected on MRI. Both nomograms showed high accurate prediction and their use as an alternative to FIGO stage system has been suggested. However, the prognostic value of combining information derived from MRI, FDG PET/CT and clinical-histopathological parameters in patients with all-stage cervical cancer has not been reported before. This information is particularly important with the recent introduction of integrated PET/MRI scanners in clinical use. PET/MRI is able to provide integrated anatomic, metabolic, and functional information. In the near future, specific PET/MRI acquisition and interpretation protocols that provide information useful to predict clinical outcomes in patients with cervical cancer need to be developed. Hence, the aim of our study was to 1) evaluate the prognostic value of pelvic MRI, whole body FDG PET/CT and clinical-histopathologic factors in the

pretreatment assessment of patients with FIGO stage I to IV cervical cancer, and 2) assess the added value of a combined MRI and whole body FDG PET/CT qualitative imaging parameters and clinical-histopathological factors, compared to clinical-histopathological factors alone.

Materials and Methods

The Institutional Review Board approved and issued a waiver of informed consent for this retrospective study, which was compliant with the Health Insurance Portability and Accountability Act.

Patient Cohort

Review of our prospectively maintained uterine cervical cancer database identified 116 patients with pathologically-proven cervical cancer who underwent pelvic MRI and FDG PET/CT for initial staging prior to primary treatment (surgery and/or chemoradiotherapy) at our institution between November 2005 and May 2012. Two patients were excluded from the analysis because no follow-up information was available; thus, our study cohort included 114 patients. 49 patients have been included in a previous published study evaluating the use of quantitative FDG uptake (PET) and diffusion (MRI) metrics in the assessment of cervical cancer⁷.

Imaging Protocol

MR imaging was performed on a 1.5-Tesla (Signa; GE Medical Systems, Milwaukee, WI) or 3-Tesla (Signa HDxt or Discovery MR750; GE Medical Systems, Milwaukee, WI) system using a body coil for excitation and a multichannel phased-array coil for signal reception. Specific MR imaging parameters varied during the study period, following changes in our standard clinical protocol, but all examinations included an axial T1-weighted spin-echo (SE) sequence and T2-weighted fast spin-echo (FSE) sequences in three imaging planes (axial/sagittal/axial oblique perpendicular to the cervix). Contrast-enhanced MR images were obtained with a three-dimensional spoiled gradient-pulse T1-weighted LAVA (liver acquisition volume acceleration) sequence before and after administration of 0.1 mmol/kg of gadolinium at a rate of 2 mL/sec, which was followed by a 20-mL saline bolus injection.

All patients were scanned on dedicated PET/CT systems (Discovery STE, LS or 690 [GE Medical Systems], Biograph 16 [Siemens Medical Systems]). A clinical imaging protocol was applied with injection of 400–455 MBq FDG after 6 h fasting and blood glucose level <200 mg/dL followed by approximately 60 minute uptake period. Subsequently, a low-dose, attenuation correction CT scan (120–140 kV, 80 mA) was acquired, followed by PET emission images from the mid thighs to skull base.

Images analysis and interpretation

MRI and FDG PET/CT exams were retrospectively reviewed by a radiologist (blinded for review) with 5 years of experience in gynecologic cancer imaging, using our institution's Picture Archiving and Communication System (PACS) (Centricity, GE Medical Systems, Milwaukee, WI, USA). The reader was blinded to clinical and histopathological

information. The reader evaluated all MRIs, followed by the PET/CTs. The reader recorded orthogonal tumor diameters, the distance between the tumor and internal cervical os, the presence of cervical stromal invasion, parametrial involvement, vaginal extension (lower 1/3, and upper 2/3), bladder, rectum and/or pelvic sidewall invasion, presence of pelvic and/or paraaortic lymphadenopathy and distant metastasis. The three orthogonal tumor diameters were measured on axial and sagittal FSE T2-weighted images (WI). On MRI, diagnostic criteria of parametrial invasion were: full-thickness disruption of the normal T2WI hypointense cervical stroma, spiculated tumor–parametrium interface (referred to as spiculated tumor), soft-tissue extension into the parametria, or encasement of the periuterine vessels. Vaginal extension was diagnosed when an interruption of the normal T2WI hypointense vaginal wall was observed. On MRI, bladder and/or rectal involvement was diagnosed when an interruption of the normal T2WI hypointense bladder or rectal wall was observed, with tumor nodules seen in the mucosal layer on T2WI. On MRI, diagnostic criteria for pelvic sidewall invasion was tumor extending within 3 mm and abutting the obturator internus or piriformis muscles, with concomitant loss of the normally present intervening fat planes.

On PET/CT, bladder or rectum involvement was diagnosed when direct contact between the FDG-avid tumor and the organ was observed, with concomitant absence of a fat plane evidenced on the CT component of the examination. Similarly, pelvic sidewall invasion was diagnosed in the presence of direct contact between the FDG-avid tumor and obturator internus and/or piriformis muscle.

On MRI, pelvic and paraaortic LN were considered metastatic if their short axis were more than 1 cm, or more than 0.8 cm if they had irregular borders and round shape. LN showing FDG-uptake above the background activity on PET images was considered metastatic. Focal areas of FDG-uptake above the background activity located in liver, lung, and thorax were considered distant metastasis. Each of the imaging features pertaining to the regional tumor spread, nodal involvement, and distant metastases were assessed on a qualitative 1–5 scale as follows: 1=definitely absent, 2=probably absent, 3=indeterminate, 4= probably present, and 5=definitely present. Any imaging feature scored as 4 or 5 was considered a positive finding.

Outcome data

One of the authors (blinded for review) reviewed patients' electronic medical records. All patients were evaluated after treatment every 3 months until death or last contact. Progression of disease was defined as the presence of tumor at biopsy or disease progression on follow-up imaging according to RECIST 1.1¹⁰.

Statistical Methods

Descriptive statistics were calculated, including frequencies and percents for categorical variables and medians and ranges for continuous variables. For the purpose of analysis, FIGO stage was grouped into I–IIa vs. IIB–IVB, grade (G) was grouped as 1/2 vs. 3 and histology was dichotomized as squamous cell carcinoma (SCC) vs. non-squamous cell carcinoma (non-SCC). Pathological FIGO stage was used in patients who underwent surgery instead of pretreatment clinical FIGO stage. Age at diagnosis, maximum tumor diameter on

MRI, and tumor volume were evaluated as continuous measurements. Organ involvement on MRI and PET/CT were categorized as positive or negative. Paraaortic LN metastasis and distant metastases detected on PET/CT were combined to indicate disease outside of the pelvis. Imaging parameters with 10 or more positive cases were included in the analysis.

Univariate Cox proportional hazard regression was used to assess the relationship between imaging parameters and progression free survival (PFS). Given the limited number of events, only a bivariate model could be built for OS, which would not allow for sufficient assessment of imaging parameters' incremental value. Therefore, only PFS was assessed at the univariate and multivariate level. The two modalities, MR and PET were analyzed separately. We used methods of Lin, Wei, and Ying¹¹ to assess the proportional assumption. PFS was defined as the interval between first imaging date and date of last follow up, date of death, or date of first progression. Patients free of pelvic and extrapelvic disease and alive at last follow up were censored. P-values less than 0.05 were considered statistically significant.

Multicollinearity was assessed using spearman correlations. Any two variables with a correlation, $\rho > 0.50$ were considered non-independent and not included in multivariate analysis. Variables significant at $p < 0.05$ on univariate analyses and clinically meaningful prognostic indicators were considered for multivariate analysis. Multivariate analyses were performed for progression free survival using Cox proportional hazards regression. We used the standard event per variable (EPV) rule developed by Peduzzi et al.¹³ to determine the maximum number of parameters allowed in the multivariate model.

Based on prior clinical prognostic indicators, age and grade comprised our baseline clinical model. Parametrial invasion on MRI and paraaortic lymphadenopathy/distant metastasis were added to create the full model. The multivariate models' discriminatory ability was calculated using the concordance index (C-index), Harrell's C, with jackknife 95% confidence intervals. We compared the C-indices of the multivariate model including clinical and imaging parameters with the model including clinical parameters only and the FIGO stage model. As this was a retrospective, exploratory study, no corrections were made for multiple hypothesis testing. All analyses were performed using SAS 9.2 (SAS Institute, Cary, NC) and Stata SE 12 (StataCorp, College Station, TX USA).

Results

Patients Characteristics

Patient characteristics are summarized in Table 1. The median age at diagnosis was 40.1 years (range: 18.2–88.9 years) and the median time between MRI and PET/CT was 3 days (range: 0–50 days). Patients were treated per standard of care at our institution with surgery alone (32.5 %), chemoradiation-therapy alone (CRT) (43.9%) or surgery and CRT (21.0%). The median follow-up time was 25.8 months (range: 0.3–84.8 months). The median time to progression was 10.4 months (range: 0.4–40.3 months). At the end of follow up, 24 patients had died of disease (21.1%) and an additional 16 patients progressed but were alive (14.0%).

Progression-Free Survival

Univariate Analyses—FIGO stage ($p<0.0001$), age at diagnosis ($p=0.0001$), tumor grade ($p=0.0455$), and histology ($p=0.0017$) were significantly associated with PFS. Patients with FIGO stage IIB–IVB, tumor grade 3, and increased age were at increased risk for progression. All MRI and PET/CT parameters assessed were associated with PFS: distance of tumor to the internal cervical os on MRI ($p=0.0474$), maximum diameter on MRI ($p<0.0001$), tumor volume ($p<0.0001$) on MRI, parametrial invasion on MRI ($p<0.0001$), spiculated tumor on MRI ($p<0.0001$), vessel encasement on MRI ($p<0.0001$), vaginal invasion on MRI ($p<0.0001$), bladder invasion on MRI ($p<0.0001$), pelvic side wall invasion on PET and MRI ($p<0.0001$ for both), pelvic lymphadenopathy on PET and MRI ($p<0.001$ for both), and paraaortic lymphadenopathy/distant metastases on PET ($p<0.0001$). Presence of features or positivity of invasion increased the risk of progression, (Table 2).

Multivariate Analyses—Our statistical design allowed a maximum of four variables to be used in multivariate analysis (Table 3). Multicollinearity was present among pelvic side wall invasion, parametrial invasion, pelvic lymphadenopathy, spiculated tumor, tumor volume, maximum diameter and vaginal involvement location, indicating a lack of independence among these parameters. Therefore, based on clinical significance and prognostic indication on univariate analyses, parametrial invasion was chosen to represent the MRI parameters. Paraortic lymphadenopathy/distant metastasis assessed on PET/CT was chosen to represent PET/CT parameters given its relative independence and clinical importance.

Based on prior clinical prognostic indicators, age and grade comprised our baseline clinical model. Age ($p=0.0001$) and grade ($p=0.0315$) were both significant in the multivariate model with a c-index of 0.68 [95% CI: 0.59–0.78]. When parametrial invasion on MRI and paraaortic lymphadenopathy/distant metastasis were added to create the full model, only grade ($p=0.0375$) parametrial invasion ($p=0.005$) and paraaortic lymphadenopathy/distant metastasis ($p=0.0395$) remained significant. The concordance index rose to 0.81 [95% CI: 0.75–0.87], which was significantly different from the baseline clinical model ($p<0.001$). This indicated that imaging parameters add significant incremental value to clinical prognostic indicators. We then compared the concordance index of the model of FIGO Stage alone to the full model, and this difference approached significance ($p=0.058$).

Discussion

In this study, we demonstrated a significant incremental prognostic value of pre-treatment MRI and FDG PET/CT findings to clinical and histopathologic prognostic indicators in patients with newly-diagnosed cervical cancer. When imaging findings were incorporated into a multivariate model together with clinical-histopathological factors, the concordance index was significantly higher than a model with clinical factors alone, indicating that the full model that includes pre-treatment imaging findings had superior alignment between the predicted and observed survival times. Similar to our findings, previous studies showed more accurate predictive ability of models that included qualitative imaging parameters and clinical-pathological factors compared to FIGO stage alone. However, these studies focused only on high-risk cohorts of patients with LACC treated with definitive chemoradiation-

therapy^{5,14} thus unable to construct a prognostic model which is, by definition, independent of treatment received. Only one previous study constructed and validated a prognostic model of clinical-pathological factors alone (with no added imaging findings) in a large cohort of cervical cancer patients including FIGO stage I to IV⁸. In another study⁴ of 187 patients with stage Ib₂-IIb cervical cancer, Gadduci et al found that *pathologically* assessed parametrial invasion and LN status were predictors of disease-free survival and overall survival independent of other clinical-histopathological factors. In contrast, in our study *MRI assessment of parametrial invasion and LN status on PET/CT* were predictors of PFS. This is important as MRI is widely used to assess parametrial invasion and PET/CT to assess LN status prior to treatment decision (surgery versus radiotherapy). For example, patients with findings of parametrial invasion on MRI don't usually undergo surgery; they have radiotherapy instead so histological confirmation is not possible in all cases. To the best of our knowledge, our study is the first to investigate the incremental prognostic value of combined MRI and FDG PET/CT qualitative parameters and clinical-pathological factors compared to clinical-pathological factors alone in the pre-treatment assessment of patients with all-stage cervical cancer.

In patients with newly diagnosed cervical cancer the prognostic prediction is currently based on FIGO staging system. FIGO stage is based upon clinical examination and the system's limitations have been described^{1,2}. Moreover, the risk of recurrence in patients with early stage of disease has been reported to range between 5% to 30%¹⁵. Polterauer and colleagues⁸ built a nomogram, incorporating FIGO stage, patient's age, tumor histology, LN status, tumor size and parametrial invasion to predict survival in patients with stage I-IV. We found that the prognostic ability of models including MRI and FDG PET/CT parameters were higher than FIGO stage alone. Though the difference in concordance indices between the FIGO stage and full model incorporating imaging parameters (0.75 vs 0.81), did not reach significance (p=0.058), this level of improvement in C-indices is considered substantial and lends credence to the possibility that the small number of events may explain the lack of significance. Previous studies have indicated that FIGO staging may be influenced by MRI findings especially in patients who undergo radiotherapy and parametrial invasion cannot be assessed at pathology, thus, it may be difficult to retrospectively assess the independent predictive ability of these variables¹.

Maximum tumor diameter was an independent prognostic factor for PFS in our study. Tumor diameter is an established prognostic parameter^{16,17} in cervical cancer, and correlates well with other pathological prognostic factors, especially with parametrial involvement¹⁸. Parametrial involvement significantly influences outcome of patients with cervical cancer¹⁹, and the assessment of parametrial involvement is crucial for the primary treatment. Therefore, it was included in our prognostic four-factor model. Parametrial spread may be detected at histopathology even in women with early FIGO stage²⁰, and it has been reported to be related with other adverse prognostic factors, such as higher histologic grade, tumor size, and metastasis to pelvic LN¹⁸. In early invasive cervical cancers extended to parametria and paraaortic LN at surgical staging, patients' survival was reported to be irrespective of other clinical-pathological factors, such as histologic subtypes¹⁹. Our study supports the previous results^{8,19}. Importantly, we showed that parametrial invasion assessed

non-invasively on MRI was the most robust prognostic factor for PFS, exceeding the prognostic ability of other clinical-histopathological factors.

In our study, the presence of pelvic LN metastasis on imaging was a predictor PFS in the univariate analysis, and paraaortic lymphadenopathy/distant metastases was a significant predictor of PFS on both univariate and multivariate analyses. Many authors have reported that the presence of pelvic/paraaortic LN metastases has an effect on patient's survival. Nonetheless, the prognostic importance of LN status assessed on imaging is still controversial in cervical cancer^{5,9}. Some studies show that pelvic/paraaortic LN metastases assessed on FDG PET/CT is an independent predictive factor for PFS in LACC^{5,14}. Kang et al. introduced a model to predict distant recurrence in patients with LACC, including histology, tumor size, and pelvic/paraaortic LN metastasis on FDG PET/CT. The concordance index (CI: 0.73) was superior to that of FIGO stage alone (CI: 0.57). Shim et al showed that paraaortic LN metastasis on MRI, but not 18F (FDG) PET/CT, were independent predictor of OS in patients with cervical cancer. The concordance index of this model (CI: 0.69) was superior to that of FIGO stage alone (CI: 0.59). It is important to note the C-indices in our study were much higher than those previously published.

Combined MRI and FDG PET/CT have been shown to improve triage to the most appropriate primary therapy in patients with early stage cervical cancer²¹. The prediction of progression or recurrence in individual patients before any therapy has started would add clinical value as a 'risk assessment model' including MRI and FDG PET/CT findings can provide prognostic information that complements clinical and histopathologic factors. This information is important as currently combined PET/MRI equipment is becoming an available tool for staging patients with gynecological malignancies. In this setting, PET/MRI protocols will need to be balanced between high-resolution imaging of the pelvis that provides crucial information on local disease extent (especially presence of parametrial invasion) and whole-body PET/CT that provides valuable information on nodal and distant metastasis. Encouraging preliminary results have been reported in multiple malignancies using whole body MRI techniques, however further work is needed to evaluate its added value in gynecologic malignancies.

Our study has several limitations. First, it is a retrospective study hence many factors could not be controlled for, including the influence of MRI findings on FIGO stage especially in the subset of patients that underwent radiotherapy and thus pathological FIGO stage could not be obtained. Second, the retrospective nature allowed for a heterogeneous population, including patients who underwent different treatment modalities. However, the aim of our study was to evaluate potential prognostic value of multiple MRI and FDG PET/CT findings, which are, by definition, independent of treatment received. Third, a single reader evaluated both MRI and FDG PET/CT scans. Although this makes evaluation or interobserver variability impossible, it matches routine clinical practice where a single experienced reader specialized in gynecology oncology imaging reports the majority (>80%) of these studies and reviews them at the weekly tumor board meeting. Finally, subsequent validation of our results is warranted and should include a per patient measure such as a nomogram; at present, the best way to interpret our results is that they support the notion that both MRI and PET/CT provide complimentary information.

Conclusion

In pre-treatment assessment of patients with cervical cancer, a prognostic model including combined MRI and FDG PET/CT findings in addition to clinical-pathological factors adds significant incremental value to clinical-pathological factors alone.

References

1. Hricak H, Gatsonis C, Chi DS, et al. Role of imaging in pretreatment evaluation of early invasive cervical cancer: results of the intergroup study American College of Radiology Imaging Network 6651-Gynecologic Oncology Group 183. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. Dec 20; 2005 23(36):9329–9337. [PubMed: 16361632]
2. Thomeer MG, Gerestein C, Spronk S, van Doorn HC, van der Ham E, Hunink MG. Clinical examination versus magnetic resonance imaging in the pretreatment staging of cervical carcinoma: systematic review and meta-analysis. *European radiology*. Jul; 2013 23(7):2005–2018. [PubMed: 23455762]
3. Tseng JY, Yen MS, Twu NF, et al. Prognostic nomogram for overall survival in stage IIB-IVA cervical cancer patients treated with concurrent chemoradiotherapy. *American journal of obstetrics and gynecology*. Feb; 2010 202(2):174, e171–177. [PubMed: 19931041]
4. Gadducci A, Teti G, Barsotti C, et al. Clinicopathological variables predictive of clinical outcome in patients with FIGO stage Ib2-IIb cervical cancer treated with cisplatin-based neoadjuvant chemotherapy followed by radical hysterectomy. *Anticancer research*. Jan; 2010 30(1):201–208. [PubMed: 20150636]
5. Kang S, Nam BH, Park JY, et al. Risk assessment tool for distant recurrence after platinum-based concurrent chemoradiation in patients with locally advanced cervical cancer: a Korean gynecologic oncology group study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. Jul 1; 2012 30(19):2369–2374. [PubMed: 22614984]
6. Kidd EA, El Naqa I, Siegel BA, Dehdashti F, Grigsby PW. FDG-PET-based prognostic nomograms for locally advanced cervical cancer. *Gynecologic oncology*. Oct; 2012 127(1):136–140. [PubMed: 22735785]
7. Micco M, Vargas HA, Burger IA, et al. Combined pre-treatment MRI and 18F-FDG PET/CT parameters as prognostic biomarkers in patients with cervical cancer. *European journal of radiology*. Jul; 2014 83(7):1169–1176. [PubMed: 24767630]
8. Polterauer S, Grimm C, Hofstetter G, et al. Nomogram prediction for overall survival of patients diagnosed with cervical cancer. *British journal of cancer*. Sep 4; 2012 107(6):918–924. [PubMed: 22871885]
9. Shim SH, Lee SW, Park JY, et al. Risk assessment model for overall survival in patients with locally advanced cervical cancer treated with definitive concurrent chemoradiotherapy. *Gynecologic oncology*. Jan; 2013 128(1):54–59. [PubMed: 23063760]
10. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *European journal of cancer (Oxford, England : 1990)*. Jan; 2009 45(2):228–247.
11. Lin DY, Wei LJ, Ying Z. Model-checking techniques based on cumulative residuals. *Biometrics*. Mar; 2002 58(1):1–12. [PubMed: 11890304]
12. Peduzzi P, Concato J, Feinstein AR, Holford TR. Importance of events per independent variable in proportional hazards regression analysis.2. Accuracy and precision of regression estimates. *J Clin Epidemiol*. Dec; 1995 48(12):1503–1510. [PubMed: 8543964]
13. Peduzzi P, Concato J, Feinstein AR, Holford TR. Importance of events per independent variable in proportional hazards regression analysis. II. Accuracy and precision of regression estimates. *Journal of clinical epidemiology*. Dec; 1995 48(12):1503–1510. [PubMed: 8543964]
14. Narayan K, Fisher RJ, Bernshaw D, Shakher R, Hicks RJ. Patterns of failure and prognostic factor analyses in locally advanced cervical cancer patients staged by positron emission tomography and treated with curative intent. *International journal of gynecological cancer : official journal of the International Gynecological Cancer Society*. Jul; 2009 19(5):912–918. [PubMed: 19574784]

15. Crivellaro C, Signorelli M, Guerra L, et al. 18F-FDG PET/CT can predict nodal metastases but not recurrence in early stage uterine cervical cancer. *Gynecologic oncology*. Oct; 2012 127(1):131–135. [PubMed: 22772064]
16. Delgado G, Bundy B, Zaino R, Sevin BU, Creasman WT, Major F. Prospective surgical-pathological study of disease-free interval in patients with stage IB squamous cell carcinoma of the cervix: a Gynecologic Oncology Group study. *Gynecologic oncology*. Sep; 1990 38(3):352–357. [PubMed: 2227547]
17. Pecorelli S. Revised FIGO staging for carcinoma of the vulva, cervix, and endometrium. *International journal of gynaecology and obstetrics: the official organ of the International Federation of Gynaecology and Obstetrics*. May; 2009 105(2):103–104.
18. Frumovitz M, Sun CC, Schmeler KM, et al. Parametrial involvement in radical hysterectomy specimens for women with early-stage cervical cancer. *Obstetrics and gynecology*. Jul; 2009 114(1):93–99. [PubMed: 19546764]
19. Takeda N, Sakuragi N, Takeda M, et al. Multivariate analysis of histopathologic prognostic factors for invasive cervical cancer treated with radical hysterectomy and systematic retroperitoneal lymphadenectomy. *Acta obstetrica et gynecologica Scandinavica*. Dec; 2002 81(12):1144–1151. [PubMed: 12519111]
20. Kamimori T, Sakamoto K, Fujiwara K, et al. Parametrial involvement in FIGO stage IB1 cervical carcinoma diagnostic impact of tumor diameter in preoperative magnetic resonance imaging. *International journal of gynecological cancer : official journal of the International Gynecological Cancer Society*. Feb; 2011 21(2):349–354. [PubMed: 21721193]
21. Pandharipande PV, Choy G, del Carmen MG, Gazelle GS, Russell AH, Lee SI. MRI and PET/CT for triaging stage IB clinically operable cervical cancer to appropriate therapy: decision analysis to assess patient outcomes. *AJR. American journal of roentgenology*. Mar; 2009 192(3):802–814. [PubMed: 19234280]

Table 1

Clinical and Pathologic Patient Characteristics

Continuous Variables	Median	(Min-Max)
<i>Age at Diagnosis</i>	40.64	(18.22–88.93)
<i>Time b/w Scans (days)</i>	3	(0.00–50.00)
<i>Time to Progression(months) (N=40)</i>	10.4	(0.36–40.34)
Categorical Variables	N	%
<i>Histology</i>		
<i>nonSCC</i>	49	42.98
<i>SCC</i>	65	57.02
<i>Grading</i>		
<i>Grade 1/2</i>	67	58.77
<i>Grade 3</i>	47	41.23
<i>FIGO Stage</i>		
<i>I–IIa</i>	60	52.63
<i>IIb–IVB</i>	54	47.37
<i>Death</i>		
<i>Alive</i>	90	78.95
<i>Dead</i>	24	21.05
<i>Progression</i>		
<i>No</i>	74	64.91
<i>Yes</i>	40	35.09
<i>Treatment</i>		
<i>CRT Only</i>	50	43.86
<i>Surgery Only</i>	37	32.46
<i>Surgery+CRT</i>	27	23.68

Table 2

Univariate Cox Proportional Hazards Regression for Progression Free Survival

Progression free Survival		N (Prog)	HR [95% CI]	p-value
Modality	Variable	Level		
N/A	Age at Diagnosis		114 (40)	1.04 [1.02–1.06]
	Histology	SCC	65 (31)	3.28 [1.56–6.90]
		nonSCC	49 (9)	REF
	Grading	Grade 3	47 (20)	1.89 [1.01–3.52]
		Grade 1/2	67 (20)	REF
		IIb–IVB	54 (35)	12.07 [4.70–30.98]
	FIGO Stage	I–IIa	60 (5)	REF
	Tumor to ICOS Distance		113 (40)	0.94 [0.88–1.00]
	Maximum Diameter		114 (40)	1.04 [1.03–1.05]
	Tumor Volume		114 (40)	1.01 [1.00–1.01]
MRI	Spiculated Tumor	Yes	63 (35)	8.04 [3.14–20.59]
		No	51 (5)	REF
	Vessel Encasement	Yes	26 (21)	6.13 [3.26–11.52]
		No	88 (19)	REF
	Parametrial Invasion	Positive	66 (38)	19.54 [4.71–81.13]
		Negative	48 (2)	REF
	Vaginal Invasion Location	Lower 1/3	5 (5)	25.92 [7.34–91.59]
		Upper 2/3	55 (30)	8.79 [3.40–22.72]
	Pelvic Sidewall Invasion	None	54 (5)	REF
		Positive	38 (29)	8.22 [4.07–16.57]
PET	Bladder Invasion	Negative	76 (11)	REF
		Positive	13 (12)	5.93 [2.97–11.80]
	Pelvic LN Involvement	Negative	101 (28)	REF
		Positive	60 (32)	5.5 [2.51–12.06]
	Pelvic Sidewall Invasion	Negative	54 (8)	REF
		Positive	17 (14)	4.21 [2.18–8.14]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Progression free Survival	N (Prog)	HR [95% CI]	p-value
Pelvic LN Involvement	Positive	5.21 [2.64–10.30]	<.0001
	Negative	REF	
Paraaortic LN/Distant Mets	Positive	5.12 [2.59–10.11]	<.0001
	Negative	REF	

Table 3

Multivariate Cox Modeling for Progression Free Survival

Progression free Survival		Multivariate Models		
Model	Variable	HR [95%CI]	p-value	C Index [95% CI]
FIGO	Level			0.75 [0.69–0.82]
	FIGO Stage	Ib–IVB 12.07 [4.70–31.01]	<.0001	
	I–IIA	REF		
Clinical				0.68 [0.59–0.78]
	Age at Diagnosis	1.04 [1.02–1.07]	0.0001	
	Grade	Grade 3 1.99 [1.06–3.73]	0.0315	
Full	Grade 1/2	REF		
				0.81 [0.75–0.87]
	Age at Diagnosis	1.02 [1.00–1.05]	0.08	
	Grade	Grade 3 1.98 [1.04–3.75]	0.0375	
	Grade 1/2	REF		
Parametrial Invasion on MRI	Positive	13.1 [3.06–56.16]	0.0005	
	Negative	REF		
Paraaortic LN/Distant Mets on PET	Positive	2.08 [1.04–4.18]	0.0395	
	Negative	REF		