composition of the acellular BAL microbiome. The increase in *Prevotella* spp. and *Veillonella* spp. during HAART-induced immune reconstitution is provocative. When previously observed in the lung, these taxa (both common oral commensals) have had consistent associations with lung health (10, 13), and their relative abundance has been positively associated with enhanced alveolar inflammation (7, 8). Although it is tempting to speculate that the enrichment with these microorganisms observed during HAART-induced immune reconstitution may contribute to a pro-inflammatory state involved in chronic lung pathology, we do not yet know whether this change in immune tone is protective, pathogenic, or both.

The next step in our understanding of the lung microbiome will be determining how it intersects with the host immune response and disease pathogenesis. The only way to accomplish this will be by summoning a variety of complementary approaches: metagenomics, transcriptomics, and metabolomics to determine function; animal models to identify mechanism; longitudinal human studies to understand stability and natural history; and clinical trials to identify therapeutic targets. The current study is an important step in this direction. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Leopoldo N. Segal, M.D., M.S.
Division of Pulmonary and Critical Care Medicine
New York University School of Medicine
New York, New York

Robert P. Dickson, M.D.
Division of Pulmonary and Critical Care Medicine
University of Michigan Medical School
Ann Arbor, Michigan

ORCID IDs: 0000-0003-3559-9431 (L.N.S.); 0000-0002-6875-4277 (R.P.D.).

References

1. Crothers K, Huang L, Goulet JL, Goetz MB, Brown ST, Rodriguez-Barradas MC, Oursler KK, Rimland D, Gibert CL, Butt AA, *et al.* HIV infection and risk for incident pulmonary diseases in the combination antiretroviral therapy era. *Am J Respir Crit Care Med* 2011;183:388–395.
2. Hirschtick RE, Glassroth J, Jordan MC, Wilcosky TC, Wallace JM, Kvale PA, Markowitz N, Rosen MJ, Mangura BT, Hopewell PC; Pulmonary Complications of HIV Infection Study Group. Bacterial pneumonia in persons infected with the human immunodeficiency virus. *N Engl J Med* 1995;333:845–851.
3. Lozupone C, Cota-Gomez A, Palmer BE, Linderman DJ, Charlson ES, Sodergren E, Mitreva M, Abubucker S, Martin J, Yao G, *et al.*; Lung HIV Microbiome Project. Widespread colonization of the lung by *Tropheryma whippelii* in HIV infection. *Am J Respir Crit Care Med* 2013;187:1110–1117.
4. Beck JM, Schloss PD, Venkataraman A, Twigg H III, Jablonski KA, Bushman FD, Campbell TB, Charlson ES, Collman RG, Crothers K, *et al.*; Lung HIV Microbiome Project. Multicenter comparison of lung and oral microbiomes of HIV-infected and HIV-uninfected individuals. *Am J Respir Crit Care Med* 2015;192:1335–1344.
5. Twigg HL III, Knox KS, Zhou J, Crothers KA, Nelson DE, Toh E, Day RB, Lin H, Gao X, Dong Q, *et al.* Effect of advanced HIV infection on the respiratory microbiome. *Am J Respir Crit Care Med* 2016;194:226–235.
6. Dickson RP, Erb-Downward JR, Prescott HC, Martinez FJ, Curtis JL, Lama VN, Huffnagle GB. Cell-associated bacteria in the human lung microbiome. *Microbiome* 2014;2:28.
7. Segal LN, Alekseyenko AV, Clemente JC, Kulkarni R, Wu B, Gao Z, Chen H, Berger KI, Goldring RM, Rom WN, *et al.* Enrichment of lung microbiome with supraglottic taxa is associated with increased pulmonary inflammation. *Microbiome* 2013;1:19.
8. Segal LN, Clemente JC, Tsay J-CJ, Koralov SB, Keller BC, Wu BG, Li Y, Shen N, Ghedin E, Morris A, *et al.* Enrichment of the lung microbiome with oral taxa is associated with lung inflammation of a Th17 phenotype. *Nat Microbiol* 2016;1:16031.
9. Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, Turner P, Parkhill J, Loman NJ, Walker AW. Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. *BMC Biol* 2014;12:87.
10. Dickson RP, Erb-Downward JR, Martinez FJ, Huffnagle GB. The microbiome and the respiratory tract. *Annu Rev Physiol* 2016;78:481–504.
11. Gleeson K, Egli DF, Maxwell SL. Quantitative aspiration during sleep in normal subjects. *Chest* 1997;111:1266–1272.
12. Huxley EJ, Viroslav J, Gray WR, Pierce AK. Pharyngeal aspiration in normal adults and patients with depressed consciousness. *Am J Med* 1978;64:564–568.
13. Molyneux PL, Mallia P, Cox MJ, Footitt J, Willis-Owen SAG, Homola D, Trujillo-Torralbo M-B, Elkin S, Kon OM, Cookson WOC, *et al.* Outgrowth of the bacterial airway microbiome after rhinovirus exacerbation of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2013;188:1224–1231.

Copyright © 2016 by the American Thoracic Society

OPENING THE DEBATE ON THE NEW SEPSIS DEFINITION

Precision Medicine: An Opportunity to Improve Outcomes of Patients with Sepsis

Clinical outcomes from sepsis have markedly improved over the last 2 decades, facilitated in part by the development in 1992 (and refinement in 2001) of consensus definitions (1–3). Mortality has decreased, and earlier recognition and treatment of sepsis are now standard of care. Nevertheless, major challenges remain. Sepsis contributes to at least 1 of every 3 deaths in hospitalized patients in the United States (4), in addition to its major global impact on

mortality (5), and many survivors are plagued by functional and cognitive limitations with long-lasting repercussions (6). Furthermore, we still lack any specific pharmacotherapies for sepsis, pointing to a major unmet need for new approaches.

The new consensus definitions published February 2016 in *JAMA* will certainly have an important impact on epidemiologic studies of sepsis, as did their predecessors (7). In addition, as the new definitions incorporate a requirement for organ dysfunction, their application in clinical trials may result in prognostic enrichment—that is, enrollment of more severely ill patients who

Supported by National Institutes of Health grant HL131621.

may be more likely to demonstrate benefit from sepsis-focused therapies (8). Whether the new guidelines will have a commensurate impact on the identification of novel treatments for sepsis remains less certain. As our ability to quantify the clinical and biological heterogeneity in human sepsis expands, data increasingly point to divergent subtypes within the heterogeneous syndrome of sepsis, suggesting precision medicine may hold the key to therapeutic breakthroughs.

Although the National Institutes of Health has recently identified precision medicine as a major priority, highlighted by President Obama's Precision Medicine Initiative, it is hardly a novel concept. From the earliest days of the medical profession, physicians have been attempting to identify distinct subtypes of disease by analyzing variability in presenting signs and symptoms. In recent years, with advances in molecular diagnostics and "omics" approaches, the impact of precision medicine has been rapidly expanding. In oncology, for instance, the treatment approach to breast cancer now fundamentally depends on hormone receptor status (9); likewise, treatment for many patients with melanoma has been redefined by the identification of BRAF mutations (and others) and the development of specific targeted inhibitors (10). In comparison, precision medicine approaches remain in their infancy in critical care medicine.

What signals point to the potential value of precision medicine in sepsis? The first step toward precision therapeutics is identification of clinical and biological heterogeneity within a syndrome, and we now have burgeoning evidence of this heterogeneity from clinical and translational studies of human sepsis. As one example, the immune response in patients with sepsis is now widely recognized to range from an exuberant pro-inflammatory cascade (thought typically to occur early in sepsis) to a profoundly immunosuppressed phenotype (thought typically to follow the pro-inflammatory phase) (11). Despite growing recognition of this diversity, we lack accurate bedside approaches to determining where any one patient lies on the immune response spectrum at the time of emergency department or intensive care unit admission. The development of novel methods to rapidly immunophenotype patients could enable the targeted application of immunomodulatory therapies (12, 13).

Similarly, gene expression analyses have identified distinct transcriptional signatures of bacterial versus viral infection, indicating differences in the biology of the host response, as well as the potential for discrimination between infectious and noninfectious causes of systemic inflammatory response syndrome (14, 15). Such analyses could lead to identification of novel and/or more targeted therapeutics for specific subgroups of patients defined by their host response to specific infections. The advent of molecular pathogen diagnostics will further accelerate this type of research by enabling accurate identification of causative pathogens earlier in the clinical course and with greater sensitivity than conventional diagnostics permit.

While these approaches build on clinically evident distinctions (e.g., between bacterial and viral illness), data-driven analytics have the potential to identify latent subgroups that are not readily apparent at the bedside. In patients with pediatric septic shock, data-driven analyses of whole-blood transcriptomics have identified two distinct subclasses of patients, defined by differences in expression of genes related to adaptive immunity and glucocorticoid receptor signaling; interestingly, there were clinical differences between the patient

groups as well, with respect to age, white blood cell count, and mortality (16, 17). Similar approaches have been used in diseases ranging from asthma to the acute respiratory distress syndrome to identify novel subgroups that may respond differently to treatment.

Of course, precision medicine need not be confined to using just "omics"-based data. Important insights can and will be drawn from analyses of readily available clinical data, such as recent studies indicating the importance of site of infection in sepsis outcomes (e.g., lung, peritoneum, urinary tract) (18) and from the thoughtful application of available biomarker-based strategies driven by experimental insights, such as a recent study targeting patients with sepsis with detectable plasma cell-free hemoglobin (19). The key common ingredient across these studies is the recognition that disease heterogeneity may have important implications for clinical outcomes and for the effectiveness of novel therapies.

What steps are needed to move precision medicine closer to reality in sepsis? First, we need a deeper understanding of the clinical and biological heterogeneity within this syndrome. Although some aspects of this heterogeneity have been studied in some depth, including, but not limited to, those mentioned earlier, many dimensions remain to be explored or require deeper exploration, including variation in genetics/genomics, physiology, immunophenotype, comorbidities, environmental exposures, microbiota, and others. These analyses will require large, well-characterized patient cohorts with thoughtfully planned biosample repositories. Some of these analyses should be performed within the framework of randomized controlled trials so that differential response to treatment can be systematically tested (20). Integration of these human studies with experimental studies in laboratory models would accelerate their therapeutic relevance. Second, we need investment in the development of point-of-care diagnostics and other rapid assays for quickly assessing the molecular phenotype of patients at the bedside. In some cases, the technology for conducting these types of assays exists, as it does, for example, both for comparatively simple protein-based enzyme-linked immunoassays and more complex multiplex quantification of gene expression, but lack of commercial availability or high cost are prohibitive (16, 21). Third, we need the application of data-driven analytic approaches to identify novel patient subgroups that may not necessarily correspond to our presuppositions about sepsis heterogeneity. Finally, we should plan at the outset to extend studies of precision medicine in sepsis to real-world settings, including both developing countries and diverse populations, to ensure that we capture all aspects of disease heterogeneity that may be relevant to patient treatment and outcomes.

Precision medicine will not be a panacea. Issues of generalizability and cost-effectiveness will need to be carefully considered. Moreover, efforts to improve targeted treatment of patients with sepsis should not detract from efforts focused on prevention, including vaccine development and distribution, reduction of deleterious environmental exposures, provision of safe drinking water, and other similar public health measures. But we need a new approach: Simply continuing the same strategies for clinical trials in sepsis that have repeatedly failed for the last several decades is neither appealing nor logical. Given the technologies now available, the weight of the evidence supporting patient heterogeneity, and the impressive impact personalized approaches have already had in other fields, it is time for us to intensify efforts to bring the promise of precision medicine to patients with sepsis. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Carolyn S. Calfee, M.D., M.A.S.
 Department of Medicine
 Department of Anesthesia
 and
 Center for Tobacco Control Research and Education
 University of California, San Francisco
 San Francisco, California
 and
 Cardiovascular Research Institute
 San Francisco, California

References

- Kaukonen KM, Bailey M, Suzuki S, Pilcher D, Bellomo R. Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand, 2000-2012. *JAMA* 2014;311:1308-1316.
- Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, Schein RM, Sibbald WJ; The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Chest* 1992;101:1644-1655.
- Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, Cohen J, Opal SM, Vincent JL, Ramsay G; SCCM/ESICM/ACCP/ATS/SIS. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003;31:1250-1256.
- Liu V, Escobar GJ, Greene JD, Soule J, Whippy A, Angus DC, Iwashyna TJ. Hospital deaths in patients with sepsis from 2 independent cohorts. *JAMA* 2014;312:90-92.
- Fleischmann C, Scherag A, Adhikari NK, Hartog CS, Tsaganos T, Schlattmann P, Angus DC, Reinhart K; International Forum of Acute Care Trialists. Assessment of global incidence and mortality of hospital-treated sepsis: current estimates and limitations. *Am J Respir Crit Care Med* 2016;193:259-272.
- Iwashyna TJ, Ely EW, Smith DM, Langa KM. Long-term cognitive impairment and functional disability among survivors of severe sepsis. *JAMA* 2010;304:1787-1794.
- Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, Bellomo R, Bernard GR, Chiche JD, Coopersmith CM, et al. The third international consensus definitions for sepsis and septic shock (sepsis-3). *JAMA* 2016;315:801-810.
- U.S. Food and Drug Administration. Guidance for industry: enrichment strategies for clinical trials to support approval of human drugs and biological products (draft). 2012.
- Davies C, Godwin J, Gray R, Clarke M, Cutter D, Darby S, McGale P, Pan HC, Taylor C, Wang YC, et al.; Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. *Lancet* 2011;378:771-784.
- Kaufman HL. Melanoma as a model for precision medicine in oncology. *Lancet Oncol* 2014;15:251-253.
- Boomer JS, To K, Chang KC, Takasu O, Osborne DF, Walton AH, Bricker TL, Jarman SD II, Kreisel D, Krupnick AS, et al. Immunosuppression in patients who die of sepsis and multiple organ failure. *JAMA* 2011;306:2594-2605.
- Davenport EE, Burnham KL, Radhakrishnan J, Humburg P, Hutton P, Mills TC, Rautanen A, Gordon AC, Garrard C, Hill AV, et al. Genomic landscape of the individual host response and outcomes in sepsis: a prospective cohort study. *Lancet Respir Med* [online ahead of print] 22 Feb 2016; DOI: 10.1016/S2213-2600(16)00046-1.
- Hotchkiss RS, Sherwood ER. Immunology: getting sepsis therapy right. *Science* 2015;347:1201-1202.
- Hu X, Yu J, Crosby SD, Storch GA. Gene expression profiles in febrile children with defined viral and bacterial infection. *Proc Natl Acad Sci USA* 2013;110:12792-12797.
- Sweeney TE, Shidham A, Wong HR, Khatri P. A comprehensive time-course-based multicohort analysis of sepsis and sterile inflammation reveals a robust diagnostic gene set. *Sci Transl Med* 2015; 7:287ra71.
- Wong HR, Cvijanovich NZ, Anas N, Allen GL, Thomas NJ, Bigham MT, Weiss SL, Fitzgerald J, Checchia PA, Meyer K, et al. Developing a clinically feasible personalized medicine approach to pediatric septic shock. *Am J Respir Crit Care Med* 2015;191:309-315.
- Wong HR, Cvijanovich N, Lin R, Allen GL, Thomas NJ, Willson DF, Freishtat RJ, Anas N, Meyer K, Checchia PA, et al. Identification of pediatric septic shock subclasses based on genome-wide expression profiling. *BMC Med* 2009;7:34.
- Leligowicz A, Dodek PM, Norena M, Wong H, Kumar A, Kumar A; Co-operative Antimicrobial Therapy of Septic Shock Database Research Group. Association between source of infection and hospital mortality in patients who have septic shock. *Am J Respir Crit Care Med* 2014;189:1204-1213.
- Janz DR, Bastarache JA, Rice TW, Bernard GR, Warren MA, Wickersham N, Sills G, Oates JA, Roberts LJ II, Ware LB; Acetaminophen for the Reduction of Oxidative Injury in Severe Sepsis Study Group. Randomized, placebo-controlled trial of acetaminophen for the reduction of oxidative injury in severe sepsis: the Acetaminophen for the Reduction of Oxidative Injury in Severe Sepsis trial. *Crit Care Med* 2015;43:534-541.
- Calfee CS, Delucchi K, Parsons PE, Thompson BT, Ware LB, Matthay MA; NHLBI ARDS Network. Subphenotypes in acute respiratory distress syndrome: latent class analysis of data from two randomised controlled trials. *Lancet Respir Med* 2014;2: 611-620.
- Schefold JC, Hasper D, von Haehling S, Meisel C, Reinke P, Schlosser H-G. Interleukin-6 serum level assessment using a new qualitative point-of-care test in sepsis: a comparison with ELISA measurements. *Clin Biochem* 2008; 41:893-898.

Copyright © 2016 by the American Thoracic Society

OPENING THE DEBATE ON THE NEW SEPSIS DEFINITION

Medicare's Sepsis Reporting Program: Two Steps Forward, One Step Back

The recently revised definitions of sepsis and septic shock represent an important advance toward reducing the burden of this devastating syndrome (1-3). These definitions coincide with an increasing number of efforts to measure and improve the quality of sepsis care, including a recent initiative by the U.S. Centers for

Medicare & Medicaid Services requiring all U.S. hospitals to report data on the quality of care for patients with sepsis and septic shock (4). The program, which began in October 2015, requires hospitals to submit to the federal government patient-level data on adherence to the Surviving Sepsis Campaign resuscitation