Published in final edited form as:

J Parkinsons Dis. 2016 June 3; 6(3): 625–630. doi:10.3233/JPD-160782.

The Relationship between the Experience of Hypomimia and Social Wellbeing in People with Parkinson's Disease and their Care Partners

Sarah D. Gunnery^a, Barbara Habermann^b, Marie Saint-Hilaire^c, Cathi A. Thomas^c, and Linda Tickle-Degnen^a

^aDepartment of Occupational Therapy, Tufts University, Medford, MA, USA

^bSchool of Nursing, University of Delaware, Newark, DE, USA

^cDepartment of Neurology, Boston University Medical Center, Boston, MA, USA

Abstract

Background—Though hypomimia, also called *facial masking*, is experienced by many people with Parkinson's disease (PD), little is known about how the experience of this motor impairment relates to their own and their care partners' (CP) social life and relationship quality.

Objective—To test if the experience of facial masking relates to social wellbeing in people with PD and their CPs.

Method—Forty individuals with PD and their CPs rated PD's difficulty showing facial expression (facial masking), and completed questionnaires about their own social wellbeing and depression.

Results—PD-reported and CP-reported facial masking of PD were positively correlated with experience of social rejection in both partners, though this relationship was diminished when controlling for depression. CPs' rating of their partner's facial masking was negatively associated with enjoyment interacting with their partner. This relationship remained when controlling for CP and PD depression.

Conclusions—The findings suggest that the experience of facial masking is negatively associated with social wellbeing particularly for the CPs, and especially so for the quality of CPs interpersonal relationship with their partner with PD.

Keywords

Parkinson's disease; caregivers; facial expression; quality of life

Corresponding Author: Sarah D. Gunnery, Department of Occupational Therapy, 574 Boston Avenue, Medford, MA, 02155, 617-627-1099, sarah.gunnery@tufts.edu.

Conflict of Interest

The authors have no conflict of interest to report.

Introduction

Hypomimia is a symptom of PD that limits the accurate expression of emotion in the face due to a decrease in the speed and coordination with which the facial musculature is activated [1]. Practitioners and same-aged peers view people with hypomimia as being more depressed and less sociable, less socially supportive, and less cognitively capable, even when the severity of the hypomimia is quite low [2, 3]. Little is known about how self-perceptions of difficulty showing facial expression, also known as *masked face* or *facial masking* in the lay PD community [4], correspond to social wellbeing in people with PD or their care partners (CPs). Though clinician and observer rated measures of facial masking are the current clinical standard, they provide little information about how people with PD experience their symptoms and how their experience of symptoms relates to quality of life and interpersonal relationships [5, 6].

There has been some investigation of convergence between PD and CP report of hypomimia [7] and qualitative research reporting that facial masking can have detrimental effects on marital relationships [8], but to our knowledge no previous study has investigated the relationship between CP perceptions of their partners' facial expressivity and how they report their own social wellbeing. Caregiving for a person with PD can lead to increased stress, depression, and decreased health in CPs [9–11], and these effects could be compounded if the person one is caring for is less able to communicate positive emotions like gratitude and happiness [12]. Understanding how CPs' social lives are associated with their perceptions of their partners' facial expression could give rise to potential ways to improve quality of life and relationships in care partners and people with PD.

The present study measured the relationship between PD and CP experience of PD facial masking and each of their own social wellbeing. Depression was also measured as it can lead to both decreased facial expressivity and decreased social wellbeing [9, 13]. We hypothesized that report of PD facial masking by both the person with PD and their CP is related to their own social wellbeing independent of depression.

Method

Design

This is a cross-sectional correlational analysis of preliminary data of a subsample from the six months post-baseline assessment of the developing *Emergence and Evolution of Social Self-Management of Parkinson's Disease* database, a three-year longitudinal study (SocM-PD) [14].

Participants

Thirteen women and 27 men with PD ($M_{\text{age}} = 66$, $SD = 7.43$) participated with their care partner (27 women and 13 men $M_{\text{age}} = 65$, $SD = 7.06$). These 40 dyads were the complete sample of dyads who had finished the sixth month assessment at the time of data analysis. Participants were recruited through the Boston University Medical Center's Movement Disorders Clinic and through community outreach throughout the greater Boston area. There were no restrictions on who could participate as a CP, but all CPs in the sample were

spouses. PD inclusion criteria were a diagnosis of idiopathic PD using the UK Brain Bank Criteria [15], Hoehn and Yahr stage between 1 and 4 [16], and an MMSE score greater than or equal to 26. CP inclusion criterion was an MMSE score greater than or equal to 26. Participants with PD had a median Hoehn and Yahr stage of 2 (range: 1 – 4) and a mean MMSE score of 29.18 ($SD = 1.11$). CPs had a mean MMSE score of 29.20 ($SD = 1.02$) at the time of screening.

Procedure and Measures

Participants with PD and their CPs separately completed questionnaires, as described below, as part of the larger SocM-PD protocol [14]. All procedures and measures were approved by the Tufts University and Boston University Medical Center Institutional Review Boards.

Experienced hypomimia was measured by asking the person with PD, “In general, how much difficulty do you have showing expression (emotion) in your face.” Separately, care partners were asked, “In general, how much difficulty does your partner have showing expression (emotion) in their face.” Both groups answered on a five point Likert scale from 1 (no difficulty) to 5 (very severe difficulty).

Social wellbeing was measured with three different scales: social isolation, experienced stigma, and enjoyment of interactions with one’s partner (partner enjoyment). The Social Isolation domain of the Nottingham Health Profile [17] measured agreement (1 = strongly disagree to 5 = strongly agree) with 5 items assessing social isolation and loneliness. The Stigma Scale for Chronic Illness [18] included 24 items that measured experienced frequency (1 = never to 5 = always) of both enacted stigma, feeling unfavorable attitudes expressed by others, and felt stigma, feeling shame, fear, or self-discrimination as a result of internalizing negative stereotypes about oneself. Both of these scales showed convergent validity with other measures of self-reported health [18, 19]. Because the social isolation and stigma scales were correlated, $r_{PD} = .57, p < .01$; $r_{CP} = .58, p < .01$, Cronbach’s alpha = .72 (PD) and .64 (CP), we created one Social Rejection composite score of these two measures for the person with PD and one for the CP. The composite score was an average of both the social isolation mean score and social stigma mean score and varied from 1 (low) to 5 (high).

We measured partner enjoyment with 3 items from the Medical Outcomes Study: Social Support Survey [20]. These items were (1) Your partner is someone to have a good time with, (2) Your partner is someone to get together with for relaxation, and (3) Your partner is someone to do something enjoyable with. Questions were answered on a Likert scale from 1 (None of the time) to 5 (All of the time). This scale was found to be related to other self-report measures of physical health and quality of life [20].

We measured depression using the 15-item version of the Geriatric Depression Scale (GDS) [21] which employs a dichotomous yes/no response format. As is standard practice, to take into account participants who may not have answered one of the 15 items (there was only one instance of missing data from the GDS in this sample) we created average scores for the GDS (dividing the total by the number of non-missing items). These average scores were then converted back to their original 15 point scale for ease of presentation.

Data Analysis

Data were managed using Research Electronic Data Capture (REDCap) hosted at Tufts University [22]. Pearson correlations were calculated to investigate the relationships among facial masking, social rejection, partner enjoyment, and depression. For the interpretation of the size of the magnitude of the correlations, we used Cohen's criteria for small ($r = .10$), medium ($r = .30$), and large effects ($r = .50$) [23]. Multiple linear regression tested the hypothesis that report of facial masking predicts social rejection and partner enjoyment independent of depression in people with PD and their CPs. Regression analyses were only conducted for social wellbeing variables that significantly correlated with report of facial expressivity for the person with PD or the CP. Predictor variables were mean centered to account for differences in their scales. Tolerance statistics and variance inflation factors were of suitable levels to indicate a lack of multicollinearity in all analyses.

Results

Overall participants reported low levels of difficulty showing expression in their face. Table 1 shows the cross tabulation of PD-reported facial masking and their partners' CP-report of facial masking. Totals for PD and CP report are also presented in Table 1. For ease of presentation we have aggregated ratings of no difficulty and mild for both partners, and moderate and severe for CPs. No PDs self-reported having severe difficulty showing expression in their face.

Replicating Mikos et al. [7], we found no mean difference between PD and CP reported facial masking, $t(39) = 1.00$, $p = .32$. There was a correlation of $r(38) = .24$, $p = .14$ between the PD and CP indicating a small, though non-significant, size of association according to Cohen's criteria.

Participants with PD who reported less facial expressivity also reported more social rejection at a medium to large and significant level ($r = .41$) but facial expressivity was unrelated to PDs' enjoyment of their CP ($r = -.05$). PDs' own depression was positively correlated with both facial expressivity ($r = .56$, large effect) and social rejection ($r = .70$, large effect). CPs who rated their partners as having less facial expressivity reported significantly more social rejection ($r = .35$, medium effect) and significantly less partner enjoyment ($r = -.55$, large effect). CPs' ratings of their partners' facial expression were not significantly correlated with their partners' depression ($r = .27$, medium effect), but were significantly correlated their own depression ($r = .40$, medium to large effect). PD and CP depression were positively correlated with CP social rejection ($r_{pd} = .48$, large effect; $r_{cp} = .70$, large effect) and negatively correlated with CP partner enjoyment ($r_{pd} = -.35$, medium effect; $r_{cp} = -.55$, large effect). See Table 2 for descriptive statistics for all measures, correlation coefficients, and corresponding p values.

Multiple linear regression showed facial masking was not a significant predictor of social rejection when controlling for their own self-reported depression for both the people with PD and the CPs ($ps > .05$). However, for CPs, ratings of their partners' facial masking predicted how much CPs enjoyed interacting with their partners even when PD and CP

depression were held constant ($p = .01$). It should be noted that CP depression is still a significant predictor in the model as well ($p = .02$). See Table 3 for all regression statistics.

Discussion

As predicted there was a positive correlation between experienced facial masking and social rejection in people with PD, but the relationship was stronger between depression and the experience of facial masking, and between depression and social rejection. When both facial masking and depression were added into the regression analysis, depression significantly predicted social rejection but the person with PD's facial masking did not. This finding suggests an overlap in how people with PD report the difficulty that they have showing expression in their face and their own depression, perhaps because people with PD may not experience their physical symptoms and their emotional symptoms as separate [6].

There was no relationship between experienced facial masking and how much people with PD enjoyed interacting with their partners. Theories of mimicry suggest that people often mimic the nonverbal expressions of those they are interacting with [24], and thus CPs might respond to their partners with facial masking with less emotional expression or more negative emotional expression. While this could then lead people with PD to enjoy interacting with their CPs less, data from the current study do not support this. It could also be possible that people with PD are not reading the negative expressions of their partners accurately [25], and this is in turn protecting their perceptions of their relationship.

These data do show a link between CPs experience of their partners' facial masking and how much they enjoy interacting with their partners with PD independent of depression. Although CPs may be familiar with their partners' changing expressive style, these descriptive and correlational data indicate that CPs are also potentially influenced by their partners' loss of facial expression. While CPs reported more partner enjoyment when their partners reported experiencing less depression, the link between CPs' perception of their partners facial expression and their enjoyment of their partner does not appear to be driven by the CPs' or their partners' depression.

These correlational findings, though not providing clear evidence for intervention, suggest that education about hypomimia and strategies for improving communication in the dyad may be beneficial for improving quality of life in people with PD and their CPs. These strategies could be as simple as having CPs focus their attention on their PD partners' verbal rather than nonverbal cues as people with PD do verbalize their positive affect [26].

With some research showing that voice treatment may lead to short term improvement in showing expression in the face [27], future research will investigate ways to reduce hypomimia as well as the relation of reduced hypomimia to PD's and CP's experience of hypomimia. Future research will also aim to replicate these findings in a larger sample size and at a second time point to further investigate the link between self-reported hypomimia, depression, and social wellbeing in people with PD.

Acknowledgments

The authors would like to thank Michael Stevenson and the Health Quality of Life Lab for their work in data collection and data management. The study was supported in part by the National Institute of Nursing Research, United States National Institutes of Health, Grant Number R01NR013522 (PI Tickle-Degnen) and National Institute on Aging, United States National Institutes of Health, Grant Number F32AG046034 (PI Gunnery). The use of REDCap electronic data capture tools is supported by grant UL1TR000073. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Bologna M, Fabbrini G, Marsili L, Defazio G, Thompson PD, Berardelli A. Facial bradykinesia. *J Neurol Neurosurg Psychiatry*. 2013; 84:681–685. [PubMed: 23236012]
2. Hemmesch AR, Tickle-Degnen L, Zebrowitz LA. The influence of facial masking and sex on older adults' impressions of individuals with Parkinson's disease. *Psychol Aging*. 2009; 24:542–549. [PubMed: 19739910]
3. Tickle-Degnen L, Zebrowitz LA, Ma HI. Culture, gender and health care stigma: Practitioners' response to facial masking experienced by people with Parkinson's disease. *Soc Sci Med*. 2011; 73:95–102. [PubMed: 21664737]
4. American Parkinson's Disease Association. [Accessed on December 28, 2015] Motor Symptoms of Parkinson's Disease. <http://www.apdaparkinson.org/motor-symptoms-of-parkinsons-disease>
5. Frank L, Basch E, Selby JV. The PCORI perspective on patient-centered outcomes research. *JAMA*. 2014; 312:1513–1514. [PubMed: 25167382]
6. Peto V, Jenkinson C, Fitzpatrick R, Greenhall R. The development and validation of a short measure of functioning and wellbeing for individuals with Parkinson's disease. *Qual Life Res*. 1995; 4:241–248. [PubMed: 7613534]
7. Mikos AE, Springer US, Nisenzon AN, Kellison IL, Fernandez HH, Okun MS, Bowers D. Awareness of expressivity deficits in non-demented Parkinson disease. *The Clinical Neuropsychologist*. 2009; 23:805–817. [PubMed: 19169938]
8. Chiong-Rivero H, Ryan GW, Flippen C, Bordelon Y, Szumski NR, Zesiewicz TA, Vassar S, Weidmer B, Garcia RE, Bradley M, Vickrey BG. Patients' and caregivers' experiences of the impact of Parkinson's disease on health status. *Patient Relat Outcome Meas*. 2011; 2:57–70. [PubMed: 21691459]
9. Martínez-Martín P, Forjaz MJ, Frades-Payo B, Rusino AB, Fernandez-Garcia JM, Benito-Leon J, Arillo VC, Barberá MA, Sordo MP, Catalán MJ. Caregiver burden in Parkinson's disease. *Movement Disord*. 2007; 22:924–931. [PubMed: 17238193]
10. Carter JH, Stewart BJ, Archbold PG, Inoue I, Jaglin J, Lannon M, Rost-Ruffner E, Tennis M, McDermott MP, Amyot D, Barter R. Living with a person who has Parkinson's disease: the spouse's perspective by stage of disease. *Mov Disord*. 1998; 13:20–28. [PubMed: 9452321]
11. Habermann B. Spousal perspective of Parkinson's disease in middle life. *J Adv Nurs*. 2000; 31:1409–1415. [PubMed: 10849153]
12. Hemmesch AR. The detrimental effects of atypical nonverbal behavior on older adults' first impressions of individuals with Parkinson's disease. *Psychol Aging*. 2014; 29:521–527. [PubMed: 25244472]
13. Berenbaum H, Oltmanns TF. Emotional experience and expression in schizophrenia and depression. *J Abnorm Psychol*. 1992; 101:37–44. [PubMed: 1537971]
14. Tickle-Degnen L, Saint-Hilaire M, Thomas CA, Habermann B, Sprague Martinez L, Terrin N, Noubary F, Tickle-Degnen L. Emergence and evolution of social self-management of Parkinson's disease: Study protocol for a 3-year prospective cohort study. *BMC Neurol*. 2014; 14:95. [PubMed: 24885181]
15. Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease. A clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry*. 1992; 55:181–184. [PubMed: 1564476]
16. Goetz CG, Tilley BC, Shaftman SR, Stebbins GT, Fahn S, Martinez-Martin P, Poewe W, Sampaio C, Stern MB, Dodel R, Dubois B. Movement Disorder Society-sponsored revision of the Unified

Parkinson's Disease Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results. *Mov Disord*. 2008; 23:2129–2170. [PubMed: 19025984]

17. Hunt SM, McEwan J. The development of a subjective health indicator. *Sociol Health Ill*. 1980; 2:231–246.
18. Rao D, Choi SW, Victorson D, Bode R, Peterman A, Heinemann A, Cella D. Measuring stigma across neurological conditions: The development of the Stigma Scale for Chronic Illness (SSCI). *Qual Life Res*. 2009; 18:585–595. [PubMed: 19396572]
19. Hunt SM, McKenna SP, McEwen J, Williams J, Papp E. The Nottingham Health Profile: subjective health status and medical consultations. *Soc Sci Med*. 1981; 15:221–229.
20. Sherbourne CD, Stewart AL. The MOS social support survey. *Soc Sci Med*. 1991; 32:705–714. [PubMed: 2035047]
21. Sheikh, J.; Yesavage, J. Geriatric Depression Scale (GDS): Recent evidence and development of a shorter version. In: Brink, TL., editor. *Clinical Gerontology: A Guide to Assessment and Intervention*. 1986. p. 165-173.
22. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009; 42:377–381. [PubMed: 18929686]
23. Cohen, J. *Statistical Power Analysis for the Behavioral Sciences*. 2. Lawrence Erlbaum Associates; New Jersey: 1988.
24. Hess U, Blairy S. Facial mimicry and emotional contagion to dynamic emotional facial expressions and their influence on decoding accuracy. *Int J Psychophysiol*. 2001; 40:129–141. [PubMed: 11165351]
25. Gray HM, Tickle-Degnen L. A meta-analysis of performance on emotion recognition tasks in Parkinson's disease. *Neuropsychology*. 2010; 24:176–191. [PubMed: 20230112]
26. Takahashi K, Tickle-Degnen L, Coster WJ, Latham NK. Expressive behavior in Parkinson's disease as a function of interview context. *Am J Occup Ther*. 2010; 64:484–495. [PubMed: 20608279]
27. Dumer AI, Oster H, McCabe D, Rabin LA, Spielman JL, Ramig LO, Borod JC. Effects of the Lee Silverman Voice Treatment (LSVT® LOUD) on hypomimia in Parkinson's disease. *JINS*. 2014; 20:302–312. [PubMed: 24524211]

Table 1

Cross tabulation table of PD and CP experienced facial masking ratings

PD Reported	CP Reported		PD Total
	No Difficulty - Mild	Moderate -Severe	
No Difficulty - Mild	22	9	31
Moderate	4	5	9
CP Total	26	14	

Table 2

Descriptive statistics and correlations between facial masking, depression, and social wellbeing for people with PD and CPs

	α	<i>M</i>	<i>SD</i>	1.	2.	3.	4.	5.	6.	7.
1. PD Facial Masking	-	1.85	.77							
2. PD Depression	.80	3.12	2.88	.56***						
3. PD Social Rejection	.72	1.86	.48	.41**	.70***					
4. PD Partner Enjoyment	.94	4.27	.88	-.05	-.11	-.13				
5. CP Facial Masking	-	2.03	1.00	.24	.27 ⁺	.26	.25			
6. CP Depression	.89	1.83	2.97	.29 ⁺	.39*	.16	.25	.40*		
7. CP Social Rejection	.64	1.46	.45	.15	.48**	.24	-.14	.35*	.70***	
8. CP Partner Enjoyment	.95	3.89	.83	-.11	-.35*	-.11	-.02	-.55***	-.55***	-.73***

Note:

⁺ $p < .10$

* $p < .05$

** $p < .01$

*** $p < .001$.

α is Chronbach's alpha. There is no α for the facial expressivity self-report measures because there was only one item in the scale. *M* is the mean score across individuals. *SD* is the standard deviation.

Table 3

Summary of regression analysis predicting social rejection and partner enjoyment in people with PD and CPs

Dependent Variable		Beta	Standard Error	95 % Confidence Interval
PD Social Rejection	R ² = .49			
	PD Facial Masking	.03	.09	-.16 – .20
	PD Depression	.68 **	.36	.99 – 2.43
CP Social Rejection	R ² = .54			
	CP Facial Masking	.05	.06	-.09 – .14
	PD Depression	.24	.29	-.02 – 1.17
	CP Depression	.58 **	.30	.73 – 1.94
CP Partner Enjoyment	R ² = .45			
	CP Facial Masking	-.38 **	.11	-.54 – -.09
	PD Depression	-.11	.58	-1.66 – .70
	CP Depression	-.36 *	.60	-2.71 – -.29

Note:

*
 $p < .05$ **
 $p < .01$