Published in final edited form as:

Neuroreport. 2016 September 28; 27(14): 1068–1071. doi:10.1097/WNR.0000000000000658.

State-dependent alterations in CSF Aβ42 levels in cognitively intact elderly with late life major depression

Nunzio Pomara, M.D.^{1,2,3}, Davide Bruno, Ph.D.⁴, Ricardo S. Osorio, M.D.⁶, Chelsea Reichert, M.S.^{1,5}, Jay Nierenberg, M.D., Ph.D.^{1,2}, Antero S. Sarreal, M.D.¹, Raymundo T. Hernando, M.D.¹, Charles R. Marmar, M.D.^{2,3}, Thomas Wisniewski, M.D.^{2,7}, Henrik Zetterberg, M.D., Ph.D.^{8,9,10}, and Kaj Blennow, M.D., Ph.D.^{8,9}

¹Nathan S. Kline Institute, Geriatric Psychiatry Division, Orangeburg, NY, USA

²Department of Psychiatry, NYU Langone Medical Center, New York, NY, USA

³Steve and Alexandra Cohen Veterans Center, Department of Psychiatry, NYU Langone Medical Center, New York, NY, USA

⁴Liverpool John Moores University, School of Natural Sciences and Psychology, Liverpool, UK

⁵Program in Behavioral and Cognitive Neuroscience, The Graduate Center of the City University of New York, New York City, NY

⁶Center for Brain Health, NYU Langone Medical Center, New York, NY, USA

⁷Departments of Neurology, Pathology and Psychiatry, NYU Langone Medical Center, New York, NY, USA

⁸Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

⁹Institute of Neuroscience and Physiology, Department of Psychiatry and Neurochemistry, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

¹⁰Department of Molecular Neuroscience, UCL Institute of Neurology, Queen Square, London, UK

Abstract

Depression has been linked to Alzheimer's disease (AD) as either an increased risk factor for its development or as a prodromal symptom. The neurobiological basis for such association, however, remains poorly understood. Numerous studies have examined whether changes in amyloid beta (Aβ) metabolism, which are implicated in AD pathogenesis, are also found in depression. In this paper, we investigated the relationship between depressive symptoms and cerebrospinal fluid (CSF) Aβ indices, in otherwise healthy, cognitively normal elderly with late-life major depression (LLMD) and controls, using a longitudinal approach, which is a novel contribution toward the literature. Significantly lower levels of CSF Aβ42 were observed in the LLMD group at baseline and were associated with more severe depressive symptoms. During longitudinal follow up, the

Correspondence: Nunzio Pomara, MD, Nathan S. Kline Institute, 140 Old Orangeburg Rd., Orangeburg, NY 10962, Pomara@nki.rfmh.org.

Statement of Conflicts: Dr. Pomara has a potential conflict of interest related to this work. Dr. Pomara has a joint patent application with the NYU Langone Medical Center related to some of the material described in this report.

depressed group remained cognitively unchanged, but was significantly less depressed than at baseline. A greater improvement in depressive symptoms was associated with increases in CSF A β 42 levels in both groups. Increases in CSF A β 42 and A β 40 were also associated with increased CSF total tau levels. Our results suggest that LLMD may be associated with state-dependent effects of CSF A β 42 levels. Future studies should determine whether the associations reflects state-dependent changes in neuronal activity and/or brain amyloid burden in depression.

Keywords

Late-life Major Depression; Abeta42; Elderly; Alzheimer's disease

Introduction

Several lines of evidence from epidemiological, case-control and longitudinal studies provide support for an association between depression (or depressive symptoms) and an increased risk for dementia and Alzheimer's disease (AD), or for depression as a prodromal state of AD [1,2]. This relationship has been described not only for late onset depression, but also for depression starting earlier in life [3]. In a longitudinal study conducted by Wilson and colleagues [4] in cognitively normal elderly, a higher number of depressive symptoms at baseline were associated with a 19% increased risk of AD, on average, during a 7-year longitudinal follow-up period. Yet, puzzlingly there have also been results that do not support such an association [5,6], thereby highlighting the possible etiological heterogeneity of depression with respect to its association with AD, and the need for further study.

Although the neurobiological mechanisms underlying the association between AD and depression are not yet clear, it is possible that there may be a common disturbance in amyloid- β (A β) metabolism [7] in both conditions. Studies conducted by our group and others [8,9] have highlighted abnormalities in A β 40 or A β 42 levels or their ratios, in plasma or serum, in individuals with depression. Analogously, a relatively smaller number of investigations have also reported changes in cerebrospinal fluid (CSF) A β concentration or brain amyloid burden using PET imaging in individuals with depression or depressive symptoms, albeit with conflicting results [e.g. 10–13,9].

Methodological differences, however, may be the cause of these differences in results, including heterogeneity in the studied populations, such as inclusion of individuals with mild cognitive impairment in the cohorts [2,14–16]. A separate issue pertains to the use of different approaches for detecting depression, with most relying on patients' self-ratings, which may lack diagnostic specificity, and only a few studies using structured interviews on the basis of Diagnostic and Statistical Manual of Mental Disorders diagnostic criteria. Finally, and critically, standardized pre-analytical and laboratory procedures for quantifying A β across centers were not used[14]. All of the existing studies have been limited to cross sectional comparisons on the basis of a single A β determination; thus, it is not known whether these abnormalities persist over time.

To address these limitations, we carried out a longitudinal prospective study in depressed elderly and age-matched controls, all of whom were cognitively normal at baseline. All

patients were diagnosed using a structured interview on the basis of Diagnostic and Statistical Manual of Mental Disorders 4th ed. criteria and all samples were analyzed at the same lab using the same immunoassay method with demonstrated sensitivity and reliability for A β determination (see Methods section). Our aim was to determine first whether late-life major depression (LLMD) and time (baseline to follow-up) had an effect on the A β levels; and second to determine whether any time-related change in A β was associated with changes in the severity of depressive symptoms. In addition, analogous analyses were also carried out on CSF total-tau(t-tau) and phosphorylated tau (p-tau) to gauge the possible emergence of neurodegeneration and tau pathology, respectively, during the course of the longitudinal study.

Methods

This study was conducted in accordance with the Declaration of Helsinki. Approval for this study was received from the Nathan S. Kline Institute/Rockland Psychiatry Center Institutional Review Board (NKI/RPC IRB) and the NYU Langone Medical Center Institutional Review Board. All participants provided written informed consent before their participation. Ninety-one participants, aged 60 years and older, with an MMSE score of at least 28, completed a 3-year longitudinal study. At baseline, 51 of these individuals agreed to an optional lumbar puncture. Three of these individuals were excluded because of MRI findings, and an additional individual was excluded for an MMSE score below 28 (Table 1). CSF was obtained from 47 individuals (Table 2), with LLMD (N=28) and age-matched and sex-matched control group (N=19), and again at the 3-year follow-up visit (LLMD group, N=19; control group, N=17). The analyses are limited to the follow-up group. CSF levels of A β 42, A β 40, t-tau and p-tau were measured using previously established methods by board-certified laboratory technicians who were blinded to clinical data [9]. Participants underwent a comprehensive neuropsychological evaluation as well as a clinical evaluation that included the Hamilton Depression scale (HAM-D), at baseline and at follow-up. No participants were considered to have AD or other Neurodegenerative conditions including Lewy Body Disease, as determined both via interview by a geriatric psychiatrist, and by examination of neuropsychological indices, either at baseline or follow-up. Pearson's correlations were computed between A β indices and HAM-D scores. All statistical analyses were carried out using the IBM SPSS statistical software package, version 22.0 for Windows (IBM Corp., Armonk, NY, USA).

Results

To evaluate whether the clinical group (LLMD and control) and time had an effect on the A β levels, we carried out two 2 $\times$ 2 repeated measures analyses of variances (GROUP, between-patients; and TIME, within-patients) on A β 40 and A β 42. A main effect of time was detected on A β 40, P less than 0.001, showing a decrease in levels between baseline (5882.94, SD=2631.67) and follow-up (3866.17, SD=1264.98); no main effect of LLMD (P =0.202) or an interaction was observed (P =0.146). When we examined A β 42, in contrast, we found a significant interaction (P =0.050), suggesting that although depressed individuals had lower levels at baseline, this difference was not present at follow-up (see Fig. 1). To evaluate whether changes in A β were linked to changes in the severity of depressive symptoms, we

carried out Pearson's bivariate correlations between A β and Ham-D levels, using change in Ham-D scores and changes in CSF A β concentration (baseline to follow-up). The reductions in depressive symptoms observed over time were significantly correlated with increases in CSF A β 42 levels, both in the entire cohort ($r = -0.451$, $P = 0.006$) and within the LLMD group ($r = -0.547$, $P = 0.015$), specifically, but not in the control group ($P = 0.809$). The same relationship was not significant with A β 40 (P 's > 0.200).

To examine whether changes in A β 42 were related to t-tau and p-tau, Pearson's bivariate correlations were carried out between change scores in the LLMD group, as referenced above. Comparisons of the baseline to follow-up levels showed significant correlation between CSF A β 42 levels and t-tau ($r = 0.557$, $P = 0.016$). A significant correlation was found between CSF A β 40 and t-tau levels as well ($r = 0.586$, $P = 0.011$). Thus, increases in t-tau in the LLMD group, over time, were associated with increases in both CSF A β 42 and CSF A β 40. The same significant correlations were not found between p-tau and CSF A β 42 or CSF A β 40 (P 's > 0.700).

Discussion

This is the first prospective longitudinal study to have examined the relationship between different phases of depression and CSF A β indices in cognitively intact elderly. Participants were examined at baseline who either had LLMD or were controls, and CSF A β 42 and A β 40 levels were found to be lower in this depressed group compared to controls [9]. Over the 3-year longitudinal study, we observed that the depressed group became significantly less depressed than at baseline and concomitantly, we also noted that levels of A β 42 increased. CSF A β 42 has been shown consistently to correlate inversely with brain amyloid because the tendency of its soluble forms to form fibrils and plaques. This pattern of results, therefore, suggests that there may be a state-dependent association between CSF A β 42 and depressive symptoms, whereby as depressive symptoms become more severe, brain amyloid deposition intensifies, and vice versa. Critically, this would indicate that the metabolic disturbances leading to A β abnormalities in LLMD individuals may be reversible rather than fixed, and thus possibly treatable.

Unlike A β 42, CSF A β 40 was not found to change as a function of LLMD, but only to decline with age. CSF A β 40 levels do not tend to correlate with brain amyloid burden, have been reported to show inconsistent changes during longitudinal follow up, and are not used as AD biomarkers. In addition, amyloid deposits in cerebral blood vessels are known to have a greater proportion of A β 40 than A β 42. It is therefore possible that the significant longitudinal reductions in CSF A β 40 may be because of age-related increases in blood vessel deposition. Relatedly, our findings do not point to a strong role of A β 40 in relationship with LLMD and depressive symptoms.

We also found that increases in CSF A β 42 and A β 40 from baseline during the 3-year longitudinal follow-up were associated with increases in t-tau. Increases in tau have been associated with progressive cognitive decline and AD, and have been ascribed to increase neuronal and axonal degeneration. However, the correlations with tau in this study were not associated with progressive cognitive decline or the emergence of AD and the increases

remained within the normal range of CSF t-tau concentrations. This raises the possibility that other factors may have contributed toward this relationship. Several lines of evidence from preclinical studies suggest that increased neuronal activity can result in increased release of A β peptides as well as tau [17]. Thus, these results are consistent with the hypothesis that state-dependent changes in neuronal activity may underlie the aforementioned association.

Results from recent investigations of resting functional MRI connectivity in depression suggest a complex pattern of neuronal activity in LLMD with reductions in brain functional connectivity in the cognitive network as well as increases in the default mode network [18]. However, resting functional MRI connectivity studies in individuals with late-life depression, which are most pertinent to this report, have consistently described reductions in the default mode network connectivity [19,20]. Human studies using cerebral blood flow and fluorine-18-fluorodeoxyglucose-PET report reductions in cortical neuronal activity in the depressive phase of unipolar depression and improvement with remission [21]. These results are consistent with our hypothesis that state-dependent effects on neuronal activity may underlie the changes in CSF A β 42 across different phases of depression.

However, sole reliance on changes in neuronal activity is not consistent with the observation that CSF A β 40 declined longitudinally in both groups. If increased neuronal activity were the basis for the correlation between increases in CSF A β 42 and reductions in depressive symptoms in the major depressive disorder group, then CSF A β 40 should similarly be expected to increase, not decrease. Therefore, alternative hypotheses should also be considered, including the possibility of state-dependent changes in oligomeric forms of A β in depression. These forms might have escaped detection by the electrochemiluminescence technology assay that we used, as was reported previously for the ELISA method [22,23], and they may have also masked epitopes of A β 42 and A β 40, resulting in their low levels. Thus increases in oligomeric forms of A β might have contributed to the low levels of CSF plasma A β 42 and A β 40 observed at baseline and to their association with more depressive symptoms. Conversely, their reduction during the longitudinal period was associated with higher CSF A β 42 levels and reduced depression. Thus, future studies should also examine oligomeric forms of CSF A β in patients with depression. In addition, as none of the existing investigations simultaneously determined brain amyloid burden by PET or CSF A β and tau levels, future studies should therefore also examine the relationship between these AD biomarkers and measures of neuronal and functional connectivity in patients with depression, both in the depressive phase of the illness and following remission.

References

1. Jorm AF. History of depression as a risk factor for dementia: an updated review. *Australian and New Zealand Journal of Psychiatry*. 2001; 35(6):776–781. [PubMed: 11990888]
2. Osorio RS, Gumb T, Pomara N. Soluble Amyloid- β Levels and Late-Life Depression. *Current pharmaceutical design*. 2014; 20(15):2547. [PubMed: 23859552]
3. Byers AL, Yaffe K. Depression and risk of developing dementia. *Nature Reviews Neurology*. 2011; 7(6):323–331. [PubMed: 21537355]

4. Wilson RS, Barnes LL, De Leon CM, Aggarwal NT, Schneider JS, Bach J, et al. Depressive symptoms, cognitive decline, and risk of AD in older persons. *Neurology*. 2002; 59(3):364–370. [PubMed: 12177369]
5. Becker JT, Chang YF, Lopez OL, Dew MA, Sweet RA, Barnes D, et al. Depressed mood is not a risk factor for incident dementia in a community-based cohort. *The American Journal of Geriatric Psychiatry*. 2009; 17(8):653–663. [PubMed: 19634208]
6. Richard E, Reitz C, Honig LH, Schupf N, Tang MX, Manly JJ, et al. Late-life depression, mild cognitive impairment, and dementia. *JAMA neurology*. 2013; 70(3):383–389.
7. Pomara N, Doraiswamy PM. Does increased platelet release of A β peptide contribute to brain abnormalities in individuals with depression? *Medical hypotheses*. 2003; 60(5):640–643. [PubMed: 12710895]
8. Pomara N, Doraiswamy PM, Willoughby LM, Roth AE, Mulsant BH, Siddis JJ, et al. Elevation in plasma A β 42 in geriatric depression: a pilot study. *Neurochemical research*. 2006; 31(3):341–349. [PubMed: 16583267]
9. Pomara N, Bruno D, Sarreal AS, Hernando RT, Nierenberg J, Petkova E, et al. Lower CSF amyloid beta peptides and higher F2-isoprostanines in cognitively intact elderly individuals with major depressive disorder. *American Journal of Psychiatry*. 2012
10. Diniz BS, Teixeira AL, Machado-Vieira R, Talib LL, Radanovic M, Gattaz WF, Forlenza OV. Reduced cerebrospinal fluid levels of brain-derived neurotrophic factor is associated with cognitive impairment in late-life major depression. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2014; 69(6):845–851.
11. Gudmundsson P, Skoog I, Waern M, Blennow K, Pálsson S, Rosengren L, Gustafson D. The relationship between cerebrospinal fluid biomarkers and depression in elderly women. *The American journal of geriatric psychiatry*. 2007; 15(10):832–838. [PubMed: 17911361]
12. Madsen K, Hasselbalch BJ, Frederiksen KS, Haahr ME, Gade A, Law I, et al. Lack of association between prior depressive episodes and cerebral [11 C] PiB binding. *Neurobiology of aging*. 2012; 33(10):2334–2342. [PubMed: 22192243]
13. Yasuno F, Kazui H, Morita N, Kajimoto K, Ihara M, Taguchi A, et al. High amyloid- β deposition related to depressive symptoms in older individuals with normal cognition: a pilot study. *International journal of geriatric psychiatry*. 2016
14. Abbasowa L, Heegaard NH. A systematic review of amyloid- β peptides as putative mediators of the association between affective disorders and Alzheimer's disease. *Journal of affective disorders*. 2014; 168:167–183. [PubMed: 25058309]
15. do Nascimento KKF, Silva KP, Malloy-Diniz LF, Butters MA, Diniz BS. Plasma and cerebrospinal fluid amyloid- β levels in late-life depression: A systematic review and meta-analysis. *Journal of psychiatric research*. 2015; 69:35–41. [PubMed: 26343592]
16. Harrington KD, Lim YY, Gould E, Maruff P. Amyloid-beta and depression in healthy older adults: A systematic review. *Australian and New Zealand Journal of Psychiatry*. 2015; 49(1):36–46. [PubMed: 25414381]
17. Yamada K, Holth JK, Liao F, Stewart FR, Mahan TE, Jiang H, et al. Neuronal activity regulates extracellular tau in vivo. *The Journal of experimental medicine*. 211(3):387–393. [PubMed: 24534188]
18. Kenny ER, O'Brien JT, Cousins DA, Richardson J, Thomas AJ, Firbank MJ, Blamire AM. Functional connectivity in late-life depression using resting-state functional magnetic resonance imaging. *The American Journal of Geriatric Psychiatry*. 2010; 18(7):643–651. [PubMed: 20220591]
19. Wu M, Andreescu C, Butters MA, Tamburo R, Reynolds CF, Aizenstein H. Default-mode network connectivity and white matter burden in late-life depression. *Psychiatry Research: Neuroimaging*. 2011; 194(1):39–46. [PubMed: 21824753]
20. Alexopoulos GS, Hoptman MJ, Kanellopoulos D, Murphy CF, Lim KO, Gunning FM. Functional connectivity in the cognitive control network and the default mode network in late-life depression. *Journal of affective disorders*. 2012; 139(1):56–65. [PubMed: 22425432]

21. Nikolaus S, Larisch R, Beu M, Vosberg H, MÜLLER-GÄRTNER HW. Diffuse cortical reduction of neuronal activity in unipolar major depression: a retrospective analysis of 337 patients and 321 controls. *Nuclear medicine communications*. 2000; 21(12):1119–1125. [PubMed: 11200016]
22. Englund H, Degerman Gunnarsson M, Brundin RM, Hedlund M, Kilander L, Lannfelt L, Ekholm Pettersson F. Oligomerization partially explains the lowering of A β 42 in alzheimer's disease cerebrospinal fluid. *Neurodegenerative Diseases*. 2009; 6(4):139–147. [PubMed: 19521063]
23. Sten C, Englund H, Lord A, Johansson AS, Almeida CG, Gellerfors P, et al. Amyloid- β oligomers are inefficiently measured by enzyme-linked immunosorbent assay. *Annals of neurology*. 2005; 58(1):147–150. [PubMed: 15984012]
24. Hashimoto K, Bruno D, Nierenberg J, Marmar CR, Zetterberg H, Blennow K, Pomara N. Abnormality in glutamine-glutamate cycle in the cerebrospinal fluid of cognitively intact elderly individuals with major depressive disorder: a 3-year follow-up study. *Transl Psychiatry*. 2016 Mar 1.6:e744. [PubMed: 26926880]

Figure 1.

Three year follow-up of cognitively intact individuals with LLMD and control subjects. a) HAM-D scores in the LLMD group and control subjects at baseline and 3-year follow-up. There was a significant decrease in HAM-D score in the LLMD group at the 3-year follow-up as previously reported in Hashimoto et al. (2016). b) CSF Aβ42 levels in the LLMD group and control subjects at baseline and 3-year follow-up. There was a significant interaction between Aβ42 levels and time.

Baseline demographics of cognitively intact individuals with LLMD and aged-matched control subjects.

Table 1

	Baseline (Mean(SD))		Statistical Analysis		
	Control Group (N =19)	LLMD Group (N=28)	t	df	p
Age	68.1 (7.3)	66.5 (5.4)	0.84	45	0.41
Education	16.7 (2.7)	16.5 (2.7)	0.27	44	0.79
HAM-D Score	1.2 (1.9)	14.9 (8.8)	8.02	45	<.0001
MMSE Score	29.5 (0.5)	29.8 (0.6)	1.56	45	0.13

Table 2
HAM-D and CSF levels of A β 42, A β 40, total-tau (t-Tau) and p-tau at baseline and follow-up.

	LLMD		Controls	
	Baseline Mean (SD)	Follow-Up Mean (SD)	Baseline Mean (SD)	Follow-up Mean (SD)
HAM-D	14.9 (8.8)	8.74 (8.1)	1.2 (1.9)	2.24 (6.01)
CSF Aβ42	231.42 (117.64)	261.05 (148.01)	335.94 (187.71)	279.29 (118.17)
CSF Aβ40	5285.84 (2408.60)	3728.47 (1379.17)	6550.29 (2799.71)	4020.06 (1145.78)
T-tau	254.33 (122.39)	277.33 (111.98)	343.59 (152.59)	365.71 (136.17)
P-tau	48.68 (30.76)	49.63 (34.86)	48.93 (25.87)	51.12 (18.12)