Published in final edited form as:

Am J Transplant. 2016 September ; 16(9): 2624–2638. doi:10.1111/ajt.13805.

Selective Sparing of Human Tregs by Pharmacologic Inhibitors of the Phosphatidylinositol 3-Kinase and MEK Pathways

N. A. Zwang^{1,2}, R. Zhang¹, S. Germana¹, M. Y. Fan¹, W. D. Hastings³, A. Cao³, and L. A. Turka^{1,*}

¹Center for Transplantation Sciences, Massachusetts General Hospital, Boston, MA

²Massachusetts General Hospital/Brigham and Women's Hospital Nephrology Joint Fellowship Program, Boston, MA

³Novartis Pharmaceuticals, Cambridge, MA

Abstract

Phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase/extracellular signal-regulated (MEK) signaling are central to the survival and proliferation of many cell types. Multiple lines of investigation in murine models have shown that control of the PI3K pathway is particularly important for regulatory T cell (Treg) stability and function. PI3K and MEK inhibitors are being introduced into the clinic, and we hypothesized that pharmacologic inhibition of PI3K, and possibly MEK, in mixed cultures of human mononuclear cells would preferentially affect CD4⁺ and CD8⁺ lymphocytes compared with Tregs. We tested this hypothesis using four readouts: proliferation, activation, functional suppression, and signaling. Results showed that Tregs were less susceptible to inhibition by both δ and α isoform-specific PI3K inhibitors and by an MEK inhibitor compared with their conventional CD4⁺ and CD8⁺ counterparts. These studies suggest less functional reliance on PI3K and MEK signaling in Tregs compared with conventional CD4⁺ and CD8⁺ lymphocytes. Therefore, the PI3K and MEK pathways are attractive pharmacologic targets for transplantation and treatment of autoimmunity.

Introduction

Regulatory T cells (Tregs), characterized by Foxp3 expression, are central mediators of tolerance in transplantation and autoimmunity (1–4). Originally identified based on their ability to inhibit immune responses on adoptive transfer into secondary hosts (5), these cells are critical for inhibiting effector T cells (Teffs) that mediate graft rejection. Abundant

*Corresponding author: Laurence A. Turka, lturka@partners.org.

Author Contributions

N.A.Z., S.G., R.Z., M.Y.F., and W.D.H. designed and performed research. L.T. and A.C. designed research. N.A.Z. and L.T. wrote the manuscript.

Disclosure

The authors of this manuscript have conflicts of interest to disclose as described by the *American Journal of Transplantation*. W.D.H. and A.C. are employees of Novartis. The other authors have no conflicts of interest to disclose.

Supporting Information

Additional Supporting Information may be found in the online version of this article.

experimental and clinical data link Tregs to suppression of inflammation, absence of rejection, and excellent long-term outcomes (6).

The phosphatidylinositol 3-kinase (PI3K) pathway is a key mediator of multiple cell growth and survival signals in mammalian cells (7). PI3K catalyzes the conversion from inactive 2- to biologically active 3-phosphorylated phosphoinositide (8). In lymphocytes, engagement of the T cell receptor (TCR), CD28, and the interleukin (IL)-2 receptor (IL-2R) (among several cytokine receptors) activates downstream PI3K signaling, which is integral to T cell function (9,10). Loss of PI3K in T cells leads to impaired thymocyte development, lymphopenia, and autoimmunity (11,12).

There are three main PI3K classes (8). Class I PI3Ks represent the class activated by TCR, CD28, IL-2R, and related signals in T cells, and they consist of a catalytic subunit and a regulatory subunit. There are four primary class I PI3K catalytic subunit isoforms: α , β , γ , and δ . The first two are widely expressed, while the second two are relatively restricted to immune cells (13). The PI3K δ isoform has been proposed to be critical for T cell function (14), as inhibition of this isoform blocks production of inflammatory cytokines in human effector T lymphocytes (15).

PI3K signaling in Tregs is normally constrained by constitutive expression of the lipid phosphatase termed phosphatase and tensin homolog on chromosome 10 (PTEN) (16). This is important for Tregs as PI3K signals negatively regulate Foxp3 expression and Treg function (17,18). Consistent with this, conditional deletion of PTEN promotes PI3K signaling in Tregs, leading to Treg dysfunction and lineage instability with accompanying autoimmune disease (19,20). It is likely, however, that at least some minimal degree of PI3K activity is required for Treg function, as PI3K inhibition can attenuate IL-2-mediated regulatory capacity (21). Whether PI3K is an appropriate target to promote Treg survival, expansion, and function remains unresolved (22).

In addition to PI3K, mitogen-activated protein kinase/extracellular signal-regulated (MEK) signaling is important for T cell activation (23,24). Network signaling studies suggest that while Akt signaling in Tregs may be minimal, MEK signaling is highly prevalent (25). Thus, MEK signaling, mediated by transforming growth factor- β stimulation, may promote Treg differentiation (26–28). Contrasting data suggest MEK signaling may antagonize Treg function (29,30). For example, MEK inhibition attenuated graft-versus-host disease (31) and augmented resolution of autoimmune colitis in murine models (32).

The development of specific PI3K (33–35) and MEK (36) inhibitors for clinical oncology affords an opportunity to dissect these pathways. We hypothesized that PI3K, and possibly MEK, inhibition might selectively inhibit conventional CD4⁺ and CD8⁺ lymphocytes compared with Tregs. Here, we have compared pharmacologic inhibition of PI3K and MEK on normal human conventional T cell and Treg responses in mixed cultures. We find that PI3K and MEK inhibition preferentially affects CD4⁺ and CD8⁺ lymphocytes compared with Tregs while sparing Treg suppressive function. These findings correlate with decreased overall PI3K and MEK expression in Tregs compared with their conventional CD4⁺ counterparts.

Materials and Methods

Cell isolation

Normal donor buffy coat and apheresis units were obtained with institutional review board approval (Protocol 2013P002335) from the Massachusetts General Hospital Blood Bank. Samples were enriched for peripheral blood mononuclear cells (PBMCs) over a Ficoll-Paque gradient. Some samples were further enriched for CD4⁺ lymphocytes by negative selection with magnetic beads (Stemcell Technologies, Vancouver, BC, Canada). Cells were frozen in aliquots. When ready to use, cells were thawed, washed with phosphate-buffered saline (PBS), and resuspended in media. Viability was typically 95–98%.

Antibodies and flow cytometry

Antibodies were titrated for test volumes of 2×10^5 PBMCs/test. Antibodies were obtained from BioLegend unless otherwise noted and included: CD4-Phycoerythrin (PE)-Cyanine 7 (Cy7) (RPA-T4), CD25-PerCP-Cy5.5 (M-A251), Foxp3 eFluor 450 (236A/E7, eBioscience, San Diego, CA), or isotype IgG1 κ (P3.6.2.8.1, eBioscience) and Live/Dead Aqua (L34957, Life Technologies, Beverly, MA). Cells were additionally labeled with antibody to CD8-fluorescein isothiocyanate (FITC) (RPA-T8) for activation marker experiments or CD8 PE (RPA-T8) for proliferation studies. Anti-CD3-allophycocyanin (APC) (UCHT1) was used to count T lymphocytes in preparatory studies. Activation marker panels were (i) CTLA-4-PE (L3D10) and programmed cell death 1 (PD-1)-APC (EH12.2H7); (ii) inducible T cell costimulator (ICOS)-PE (C398.4A) and CD27-APC (O323); (iii) CD62L-PE (DREG-56) and HLA-DR-APC (L243); and (iv) CD39-PE (A1) and Ki67-APC. For suppression assays, PBMCs were stained with CD4-FITC (RPA-T4), CD25-PerCP-Cy5.5 (M-A251), CD69-PE (FN50), and Live/Dead Aqua. Cytometry was performed on a Navios (BD Biosciences, San Jose, CA) flow cytometer. Cytometry for preparatory cell counts was performed on an Accuri C6 (BD Biosciences) flow cytometer.

Sorting

Enriched samples of human CD4⁺ cells were labeled with antibodies to CD4-PECy7 (RPA-T4), CD25-PE (BD, 2A3), and CD127-BV421 (A019D5). They were washed, resuspended in X-VIVO 15 (Lonza, Portsmouth, NH) containing 10% human serum and sorted on a SORP FACS Aria II (BD Biosciences). Tregs were defined as CD4⁺CD25^{hi}CD127⁻.

Signaling inhibitors

IC-87114 (PI3K δ inhibitor) was obtained from Selleckchem (Houston, TX; S1268) and reconstituted to maximal solubility of 2.52 mmol/L in DMSO, and aliquots were stored at -80°C . AS-605240 (PI3K γ inhibitor) was obtained from Sigma-Aldrich (Natick, MA; A0233) and reconstituted to 9.7 mmol/L in DMSO, and aliquots were stored at -80°C . The following inhibitors were obtained from Novartis (Cambridge, MA): BKM-120 (pan PI3K), BYL-719 (PI3K α inhibitor), and MEK-162 (MEK inhibitor). These were reconstituted in DMSO 10 mmol/L, and aliquots were stored at -20°C .

Proliferation analyses

Thawed PBMCs were labeled with 0.05 $\mu\text{mol/L}$ carboxyfluorescein succinimidyl ester (CFSE, Life Technologies, C34554) in PBS for 2 min at 37°C. Cells were quenched with 100% fetal bovine serum (FBS) and washed three times in PBS supplemented with 5% FBS. Then, 2×10^5 PBMCs/well were plated in 96-well flat-bottom plates. A 2x solution containing soluble mouse antibodies to human CD3 and CD28 (final 0.25 $\mu\text{g/mL}$ each) and vehicle (final 0.2% DMSO) or inhibitors was added for a final volume of 200 μL . (For preparatory studies in Figure 1, 0.3% DMSO vehicle was used, except for IC-87114, where 1.67 $\mu\text{mol/L}$ = 0.1% DMSO and 15 $\mu\text{mol/L}$ = 0.9% DMSO.) Separate wells for Foxp3 isotype controls were plated for vehicle control and the higher concentration of each inhibitor. Samples were harvested after 72 h. Supernatant samples were saved and frozen at -80°C . Before flow cytometric analysis, a known number (20 404) of counting beads (ACFP-70-10, Spherotech, Lake Forest, IL) was added to each sample. The mitotic index was calculated as the number of mitoses among the cells that were put in the culture well. For example, if 100 cells were put in a well, 20 remained undivided, and 80 divided 4 times each, the mitotic index would be $(80 \times 4)/100 = 3.2$.

Signaling studies

Frozen PBMCs were thawed and resuspended in X-VIVO 15 containing 10% human serum containing vehicle (0.2% DMSO) or inhibitors. Aliquots of 5×10^5 PBMCs in $12 \times 75\text{-mm}$ test tubes were rested at 37°C in 5–7% CO_2 for 2–4 h. Samples were placed on ice for 15 min before labeling with 10 μL of a 1:10 solution (each) of mouse antibodies to human CD3 and CD28 (clones UCHT1 and CD28.2, respectively, BioLegend, San Diego, CA) for an additional 15 min. Cells were washed in prechilled PBS and resuspended in 100 μL of chilled PBS containing vehicle or inhibitor. Samples were coated with 10 μL of a 1:10 solution of goat anti-mouse immunoglobulin (553998, BD) on ice for an additional 15 min. Samples were stimulated in a 37°C water bath for 15 min and fixed with Cytotfix. An unstimulated negative control and a phorbol myristate acetate (up to 30 ng/mL) plus ionomycin (up to 1.5 $\mu\text{mol/L}$)-stimulated positive control was included (Figure S1). Before staining, samples were blocked with 5 μL of mouse immunoglobulin (10400C, Life Technologies) and incubated at room temperature for 15 min.

Activation marker analyses

Thawed PBMCs were resuspended in X-VIVO 15 containing 10% human serum. Aliquots of 2×10^5 PBMCs were plated in 96-well flat-bottom plates. A 2x solution containing soluble antibodies to human CD3 and CD28 (final 0.25 $\mu\text{g/mL}$ each) and vehicle (final 0.2% DMSO) or inhibitors was added for a final volume of 200 μL . Cells were harvested after 24 h. Supernatants were frozen at -80°C for Luminex analysis. For each condition, four or five technical replicates were pooled, washed, and redistributed for flow cytometric analysis.

Suppression assays

Sorted $\text{CD4}^+\text{CD25}^{\text{hi}}\text{CD127}^-$ cells (Tregs, Figure S2) were added at increasing concentrations to autologous PBMCs. Cells were suspended in X-VIVO 15 containing 10% human serum. For stimulation, anti-CD3/CD28-coated beads (Miltenyi Biotec, Cambridge,

MA; 130-095-353) in 4x vehicle (final DMSO concentration of 0.2%) or inhibitor were added at a bead:PBMC ratio of 1:6 for a final volume of 200 μ L per well in 96-well U-bottom plates. Antibodies to CD154-APC (24–31, typically 5 μ L of a 1:20 dilution per 1×10^5 PBMCs) were added as a capture assay. After 7 h of stimulation at 37°C in 5–7% CO₂, plates were removed to 4°C overnight and analyzed the next day.

Phospho flow cytometry

Samples were fixed with Cytofix (554655, BD) for 12 min at 37°C, spun, decanted, resuspended in Permeabilization Buffer III (558050, BD), and kept at 4°C for 30 min or at –80°C for longer periods. They were then washed thrice in Stain Buffer (554656, BD) and split into two panels before simultaneous labeling for surface, Foxp3, and phosphorylated signaling molecules.

Phospho antibody panels were (i) pERK1/2-PE (BD, 20A) and pAkt S473-AlexaFluor 647 (BD, M89-61) or (ii) pS6-PE (N7-548, BD) and pPDK1-AlexaFluor 647 (J66-653.44.17, BD). Phospho antibody test volumes were 4 μ L per 2.5×10^5 cells.

Luminex analysis

IL-2, IL-6, IL-8, IL-10, IL-13, interferon- γ , tumor necrosis factor- α , granulocyte-macrophage colony-stimulating factor, macrophage inflammatory protein (MIP)-1 α , and MIP-1 β were measured by Luminex.

Immunoblotting

Equal numbers of fluorescence-activated cell sorted Tregs (CD4⁺CD25^{hi}CD127[–]) and conventional CD4⁺ T lymphocytes were lysed directly in Laemmli buffer and boiled. Lysates were separated by sodium dodecyl sulfate–polyacrylamide gel electrophoresis, transferred to a polyvinylidene difluoride membrane, and probed by using immunoblotting. Primary antibodies used were PI3K δ (7176, Santa Cruz, Dallas, TX), PI3K α (4249, Cell Signaling), MEK1/2 (47E6, Cell Signaling), glyceraldehyde 3-phosphate dehydrogenase (GAPDH; G8795, Sigma), and β -tubulin (2146, Cell Signaling). Horseradish peroxidase–conjugated goat anti-rabbit (170-6515, Bio-Rad Labs, Hercules, CA) or anti-mouse (170-6516, Bio-Rad) secondary antibodies were added. Autoradiography was detected by chemiluminescence (WBLUC0500, Millipore, Billerica, MA). Densitometry analysis was performed with ImageJ (<http://imagej.nih.gov/ij/>) and normalized to GAPDH or β -tubulin loading controls.

Data analysis

Flow cytometric data were analyzed using FlowJo (Tree Star, Ashland, OR). Cells were gated for lymphocytes, singlets, and live cells. Numerical data were analyzed with Microsoft Excel, and statistical analyses were performed using heteroscedastic, one-tailed Student t-tests. Graphs were constructed using GraphPad (La Jolla, CA). In all cases, * $p < 0.05$ and ** $p < 0.01$ are indicated.

Results

Selection of inhibitors and dose titration

In preparatory studies, we assessed if there was a dose of each inhibitor that, when used in mixed PBMC cultures, (i) suppressed T lymphocyte proliferation and (ii) might differentially affect proliferation of CD4⁺ and CD8⁺ lymphocytes compared with Tregs. We tested five inhibitors: IC-87114 (PI3K δ), AS-605240 (PI3K γ), BKM-120 (pan-PI3K), BYL-719 (PI3K α), and MEK-162 (MEK). Figure 1A shows a sample titration. Representative CD4⁺ and Treg gating is shown in Figure S3. Note that we could not use CD127 for gating of Tregs from stimulated PBMC cultures, as CD127 is downregulated by *in vitro* activation of non-Tregs (37,38 and data not shown). Based on these findings, we narrowed the *in vitro* concentration range for each inhibitor and assessed their effects on mitotic indices (mitoses/precursor) of T lymphocyte subsets as described (39). To adjust for variability among donors, we normalized results of inhibitor to vehicle treatment. This allowed us to compare the effects of inhibitor with vehicle *within* each T lymphocyte subset and to compare the differential effects of each inhibitor *between* T lymphocyte subsets. As shown in Figure 1B, we observed only marginal effects of AS-605240 (PI3K γ) on proliferation at concentrations up to 3.2 μ mol/L, which is much higher than the compound's reported IC₅₀ of 8 nmol/L (40). Additionally, we observed that effects with the pan-PI3K isotype inhibitor closely resembled those using the PI3K α isoform inhibitor. Hence, in subsequent studies, we focused on the PI3K δ , PI3K α , and MEK inhibitors. The doses of IC-87114 and BYL-719 that clearly inhibited CD4⁺ and CD8⁺ lymphocyte proliferation were approximately 5 and 3 μ mol/L, respectively, and were used in subsequent experiments. Because the MEK inhibitor concentration was not in a dose-response range between 370 and 120 nmol/L (Figure 1B), we retitered (shown in Figure 1A) and selected a concentration of 40 nmol/L (which provided partial, but not complete, inhibition of proliferation).

Effect of PI3K and MEK inhibition on proliferation and naïve versus effector/memory phenotypic differentiation

We compared the effects of each inhibitor on the proliferation of T lymphocyte subsets (in mixed PBMC cultures) using the concentrations established here earlier. As shown in Figure 2A, both the PI3K and MEK inhibitors suppressed proliferation in all T lymphocyte subsets compared with vehicle. PI3K inhibition showed modest but statistically significant differences in suppressing the proliferation of Teff versus Treg populations. The combination of the PI3K δ inhibitor and MEK inhibitor had additive effects on proliferation (Figure S4) with a continued trend toward relative sparing of Tregs. A half-log-fold lower dose of each inhibitor significantly but minimally decreased T lymphocyte proliferation and did not preferentially affect any subset (Figure S5). There was minimal loss of cell viability among any cell type due to inhibitor treatment (Figure 2B). As expected, the effects of the inhibitors on proliferation were reflected in cell recovery (Figure 2B), with the exception that due to high variability of the number of Tregs remaining in culture after 72 h between experiments, the differences seen were not statistically significant. Expression of Foxp3 among Tregs was unaffected by inhibitor treatment (Figure 2C).

PI3K or MEK inhibition also blocked the conversion of CD4⁺ T cells but not Tregs, from a CD45RA⁺ (naïve) to a CD45RO⁺ (effector/memory) phenotype (Figure 2D). We considered that this might be explained by a stronger inhibition of effector CD4⁺CD45RO⁺ compared with naïve CD4⁺CD45RA⁺ lymphocytes. We found, however, that proliferation of both effector and naïve CD4⁺ lymphocytes were equivalently equally suppressed by the inhibitors (Figure 2E). Thus, the blockade of CD45RA → CD45RO conversion by PI3K or MEK inhibition appeared to be independent of their effects on proliferation.

PI3K and MEK inhibition preferentially inhibit expression of activation markers in CD4⁺ and CD8⁺ lymphocytes compared with Tregs

We next assessed the effects of PI3K or MEK inhibition on the induction of various activation markers in stimulated T cells (Figure 3). These markers included PD-1, cytotoxic T lymphocyte associate protein 4 (CTLA-4), ICOS, HLA-DR, CD27, CD25, CD62L, CD39, and Foxp3. We found that both PI3K and MEK inhibition partially blocked the normal activation-induced increase in most of these markers among CD4⁺ and CD8⁺ lymphocytes but not Tregs (Figure 3; representative gating and histograms are shown in Figures S6–S8). There were only trivial effects of PI3K and MEK inhibition on expression of CD62L and CD39 in T lymphocyte subsets (Figure S9 and data not shown, respectively).

The fold-increase in expression of some markers, such as HLA-DR, following stimulation was greater in CD4⁺ and CD8⁺ lymphocytes compared with Tregs. However, PI3K and MEK inhibition had a greater effect on induction of these molecules in CD4⁺ and CD8⁺ lymphocytes compared with Tregs as the expression for each cell subset is normalized to its own vehicle control. Among other notable findings, ICOS expression among CD8⁺ lymphocytes was especially sensitive to both PI3K and MEK inhibition, as was CD25 expression in nonregulatory CD4 T cells. Finally, although we did not see any effect of PI3K or MEK inhibition of Foxp3 upregulation at 72 h (not shown), we did observe a transient inhibitory effect at 24 h (Figure 3). The basis for this is not known at present. Taken together, these data illustrate preferential blockade of CD4⁺ lymphocyte activation compared with Tregs with both PI3K and MEK inhibitor treatment.

Effects of PI3K and MEK inhibition on cytokine production

In addition to examining differential effects of PI3K and MEK inhibitors on T cell subsets, we assessed their effects on cytokine production. There was reduced overall cytokine elaboration in the inhibitor-treated cultures compared with those treated with vehicle alone (Figure 4). The degree and pattern of cytokine suppression by inhibitors were consistent after 24 h of stimulation compared with 72 h of stimulation. This finding suggests that the cytokine profiles observed represent specific effects of the inhibitors at the *in vitro* doses selected.

We observed notable differences in the degrees of IL-6, IL-8, and IL-13 suppression in mixed cultures treated with MEK rather than PI3K inhibitors. We found that MEK inhibition suppressed IL-6 and IL-8 production in mixed culture to a greater degree than PI3K inhibition after 72 h of stimulation. MEK inhibition spared elaboration of IL-13 and, to a

lesser extent, interferon γ , IL-10, and MIP-1 α/β relative to PI3K inhibition. We note that the effects of MEK and PI3K inhibition on cytokine production do not precisely mirror effects on T cell proliferation and activation. This may be due to selectivity for particular cellular functions or the fact that many of the cytokines measured are not T cell specific but are produced by macrophages (present in our mixed PBMC cultures) as well.

PI3K and MEK inhibition minimally affect Treg suppressive function

Since PI3K and MEK inhibition spared Treg proliferation and activation compared with CD4⁺ and CD8⁺ lymphocytes, we hypothesized that PI3K and MEK inhibition would minimally inhibit Treg suppressive function. To test this, we analyzed the ability of Tregs to suppress CD69 and CD154 expression among autologous CD4⁺CD25⁻ responder lymphocytes in mixed PBMC culture after stimulation for 7 h (41,42). In this assay, decreased CD69 and CD154 expression among CD4⁺CD25⁻ responder T lymphocytes corresponds to enhanced suppression by Tregs. We found that increasing ratios of Tregs:PBMCs resulted in decreased CD69 and CD154 expression among responder T lymphocytes (Figure 5A). Neither PI3K nor MEK inhibition affected regulatory T cell suppressive function, except that MEK inhibition with the lowest Treg:PBMC ratio slightly impaired Treg suppressive function. We further suspected that both PI3K and MEK inhibition alone would suppress expression of CD69 and CD154 in Teffs (CD4⁺CD25⁻) but not Tregs (CD4⁺CD25^{hi}). To exclude any effect of exogenous Treg addition, we assessed the PBMC samples to which no autologous Tregs were added (gating illustrated in Figure S10). We found that both PI3K and MEK inhibition decreased CD69 and CD154 expression among Teffs but not Tregs after this short stimulation (Figure 5B). These findings were consistent with our studies after longer periods of stimulation: PI3K and MEK inhibition preferentially inhibited activation of conventional CD4⁺ lymphocytes versus Tregs.

PI3K and MEK inhibitors strongly inhibit PI3K signaling

Based on these findings, we asked if PI3K and MEK inhibitors selectively affected TCR-mediated signaling in CD4⁺ and CD8⁺ lymphocytes compared with Tregs. Thus, we analyzed phosphorylation at Akt (S473), ERK (1/2), S6, and PDK1 after a 5-min stimulus following preincubation with vehicle or inhibitors. Representative gating and histograms are shown in Figure 6(A), and changes in phosphorylation (percentage positive) of signaling molecules normalized to vehicle-treated samples are shown in Figure 6(B). Both PI3K δ and PI3K α inhibitors strongly inhibited Akt phosphorylation in all T lymphocyte subsets. Additionally, we found that both PI3K and MEK inhibitors blocked PDK1 (associated with PI3K signaling) and S6 [downstream of both PI3K and MEK pathways (43)] signaling equally among different lymphocyte subsets.

PI3K and MEK expression are decreased among Tregs compared with CD4⁺ lymphocytes

The preceding experiments showed that PI3K and MEK inhibition preferentially blocked both proliferation and activation among conventional CD4⁺ and CD8⁺ lymphocytes without affecting Treg suppressive function. These differences did not seem to be explained by a preferential effect on signaling. Instead, this raised the possibility of cell-intrinsic differences between conventional lymphocytes and Tregs in the roles of PI3K and MEK signaling. Therefore, we analyzed expression of these signaling molecules by immunoblotting (Figure

7). We sorted Tregs (CD4⁺CD25^{hi}CD127⁻) alongside equal numbers of conventional CD4⁺ lymphocytes and probed for expression of PI3K δ , PI3K α , and MEK1/2 by immunoblotting. The PI3K α antibody used was rigorously validated in other studies (44–46). We found decreased expression of both PI3K δ and PI3K α isoforms and of MEK in Tregs compared with conventional CD4⁺ lymphocytes. Thus, PI3K and MEK signaling networks differ between CD4⁺ lymphocytes and Tregs, at least at the level of protein expression.

Discussion

We asked whether PI3K or MEK inhibition preferentially affects conventional CD4⁺ and/or CD8⁺ lymphocytes compared with Tregs. To address this question, we studied proliferation, activation, and signaling among these lymphocyte subsets in mixed cultures in the presence of two isoform-specific PI3K inhibitors and one MEK inhibitor. We found that Tregs were less susceptible than their CD4⁺ and CD8⁺ counterparts to PI3K and MEK inhibition. This difference correlates with overall expression of PI3K and MEK in Tregs compared with CD4⁺ lymphocytes.

We found that PI3K and MEK inhibition preferentially blocked expression of a number of important activation and functional markers in CD4⁺ lymphocytes compared with Tregs, including PD-1 and CTLA-4. As well, PI3K and MEK inhibition spared c-expression of CD69 and CD154 in Tregs compared with Teff after TCR stimulation, and this corresponded with minimal impairment of Treg functional capacity in suppression assays. While the MEK and PI3K pathways are distinct, we noted that MEK inhibitors partially blocked Akt phosphorylation and, conversely, PI3K inhibitors partially blocked ERK phosphorylation. We suspect these findings are explained by cross-talk between PI3K and ERK signaling pathways (47) rather than nonspecificity of the compounds, which were tested at relatively low doses. PI3K signaling has been shown to enhance ERK activation (48), and, conversely, PI3K inhibition may as well inhibit ERK activity (49).

Defining potential isoform-specific expression of PI3K in different cell types is important for precision immunotherapy and understanding drug effects and side effects. Our literature search did not identify studies that examined qualitative expression of different PI3K isoforms in T cell subsets. Moreover, to our knowledge, PI3K α isoform expression in T lymphocytes has not been shown. It has heretofore been thought that the δ and γ isoforms are predominantly expressed in T lymphocytes, while the α isoform is primarily expressed in B but not T lymphocytes (8). Nonetheless, specific PI3K α isoform inhibition did not disrupt beta cell receptor–dependent activation in human *in vitro* or mouse *in vivo* models. Thus, expression of PI3K α does not by itself demonstrate functional importance. While we found that T cells were sensitive to a PI3K α inhibitor, we cannot exclude the possibility that effects observed were due to off-target inhibition of other PI3K isoforms. The role of PI3K γ remains controversial. PI3K γ is clearly critical for neutrophil function (50–52). Despite the fact that genetic inactivation of PI3K γ impairs differentiation and activation in murine T lymphocytes (53), we found no effect of relatively high concentrations of AS-605240 on proliferation.

Our data show a reduced role for PI3K signaling in Tregs compared with conventional CD4⁺ cells. In contrast, Abu-Eid et al found stronger inhibition of proliferation among Tregs compared with CD4⁺ lymphocytes with PI3K and Akt inhibitors (54). These results were derived using isolated cell populations rather than by analyzing cell-specific effects among mixed lymphocyte populations. One potential limitation of this approach is the fact that activation of naïve conventional T cells may induce Foxp3 expression, and thus cells examined after 72 h can be a combination of thymic Tregs and peripheral Tregs (55). However, these induced Tregs have potent functional suppressive activity and so should contribute to immune modulation. While we acknowledge the value of studying pure cell populations, we aimed to mimic the effects of inhibitors that might be seen *in vivo*, where all cell types in secondary lymphoid tissue will be affected.

Determination of cytokine profiles from supernatants of mixed cultures illustrated the pleiotropic effects of PI3K and MEK inhibitors. Some cytokines have been particularly associated with MEK or PI3K signaling. For example, PI3K but not MEK inhibition can suppress IL-10 production in peritoneal macrophages (56), while MEK but not PI3K inhibition inhibits IL-6 production (57), and IL-6 production among macrophages correlates with MEK signaling in other systems (58–60). MEK signaling may also stabilize IL-8 RNA transcripts (61), and Janus kinase 2/signal transducer and activator of transcription 5–mediated IL-8 production may attenuate the effects of PI3K/mammalian target of rapamycin inhibition (62). Thus, different signaling pathways contribute to elaboration of different cytokines.

A growing literature has demonstrated key differences in signaling (25) and metabolism (63,64) between conventional CD4⁺ lymphocytes and Tregs. Our *in vitro* studies confirmed these findings in human lymphocytes. We showed that regulatory T cells may be partially spared of the suppressive effects of PI3K inhibitors compared with CD4⁺ lymphocytes. Therefore, PI3K and MEK inhibitors might be useful primarily, or as adjunct to costimulatory blockade (65) in *ex vivo* Treg manufacturing to enhance Treg yields from a mixed starting population. Moreover, they may preferentially target Teff responses *in vivo*, while relatively preserving Treg activation and function. These findings provide a rationale for use of PI3K and possibly MEK inhibition to temper both autoimmunity and alloimmunity in transplantation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Christopher M. Borges for assistance with flow cytometry, and the Massachusetts General Hospital/Brigham and Women's Hospital Nephrology Joint Fellowship Program for its support. N.Z. was supported by the American Society of Nephrology Donald E. Wesson Research Fellowship.

Abbreviations

CFSE carboxyfluorescein succinimidyl ester

CTLA-4	cytotoxic T lymphocyte associate protein 4
FBS	fetal bovine serum
GAPDH	glyceraldehyde 3-phosphate dehydrogenase
ICOS	inducible T cell costimulator
IL-2R	interleukin-2 receptor
IL	interleukin
MEK	mitogen-activated protein kinase/extracellular signal-regulated
MIP	macrophage inflammatory protein
PBMC	peripheral blood mononuclear cell
PBS	phosphate-buffered saline
PD-1	programmed cell death 1
PI3K	phosphatidylinositol 3-kinase
PTEN	phosphatase and tensin homolog
TCR	T cell receptor
Teff	effector T cell
Treg	regulatory T cell

References

1. Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science*. 2003; 299:1057–1061. [PubMed: 12522256]
2. Godfrey VL, Wilkinson JE, Russell LB. X-linked lymphoreticular disease in the scurfy (sf) mutant mouse. *Am J Pathol*. 1991; 138:1379–1387. [PubMed: 2053595]
3. Kim JM, Rasmussen JP, Rudensky AY. Regulatory T cells prevent catastrophic autoimmunity throughout the lifespan of mice. *Nat Immunol*. 2007; 8:191–197. [PubMed: 17136045]
4. Wildin RS, Smyk-Pearson S, Filipovich AH. Clinical and molecular features of the immunodysregulation, polyendocrinopathy, enteropathy, X linked (IPEX) syndrome. *J Med Genet*. 2002; 39:537–545. [PubMed: 12161590]
5. Gershon RK. A disquisition on suppressor T cells. *Transplant Rev*. 1975; 26:170–185. [PubMed: 1101469]
6. Taflin C, Nochy D, Hill G, et al. Regulatory T cells in kidney allograft infiltrates correlate with initial inflammation and graft function. *Transplantation*. 2010; 89:194–199. [PubMed: 20098282]
7. Fruman DA, Cantley LC. Phosphoinositide 3-kinase in immunological systems. *Semin Immunol*. 2002; 14:7–18. [PubMed: 11884226]
8. Okkenhaug K. Signaling by the phosphoinositide 3-kinase family in immune cells. *Annu Rev Immunol*. 2013; 31:675–704. [PubMed: 23330955]
9. Janas ML, Varano G, Gudmundsson K, Noda M, Nagasawa T, Turner M. Thymic development beyond beta-selection requires phosphatidylinositol 3-kinase activation by CXCR4. *J Exp Med*. 2010; 207:247–261. [PubMed: 20038597]

10. Webb LM, Vigorito E, Wymann MP, Hirsch E, Turner M. Cutting edge: T cell development requires the combined activities of the p110gamma and p110delta catalytic isoforms of phosphatidylinositol 3-kinase. *J Immunol.* 2005; 175:2783–2787. [PubMed: 16116162]
11. Ji H, Rintelen F, Waltzinger C, et al. Inactivation of PI3Kgamma and PI3Kdelta distorts T-cell development and causes multiple organ inflammation. *Blood.* 2007; 110:2940–2947. [PubMed: 17626838]
12. Swat W, Montgrain V, Doggett TA, et al. Essential role of PI3K-delta and PI3Kgamma in thymocyte survival. *Blood.* 2006; 107:2415–2422. [PubMed: 16304053]
13. Okkenhaug K. Rules of engagement: distinct functions for the four class I PI3K catalytic isoforms in immunity. *Ann N Y Acad Sci.* 2013; 1280:24–26. [PubMed: 23551098]
14. Okkenhaug K, Bilancio A, Farjot G, et al. Impaired B and T cell antigen receptor signaling in p110delta PI 3-kinase mutant mice. *Science.* 2002; 297:1031–1034. [PubMed: 12130661]
15. Soond DR, Bjorgo E, Moltu K, et al. PI3K p110delta regulates T-cell cytokine production during primary and secondary immune responses in mice and humans. *Blood.* 2010; 115:2203–2213. [PubMed: 20081091]
16. Han JM, Patterson SJ, Levings MK. The role of the PI3K signaling pathway in CD4(+) T cell differentiation and function. *Front Immunol.* 2012; 3:245. [PubMed: 22905034]
17. Sauer S, Bruno L, Hertweck A, et al. T cell receptor signaling controls Foxp3 expression via PI3K, Akt, and mTOR. *Proc Natl Acad Sci U S A.* 2008; 105:7797–7802. [PubMed: 18509048]
18. Crellin NK, Garcia RV, Levings MK. Altered activation of AKT is required for the suppressive function of human CD4 + CD25 + T regulatory cells. *Blood.* 2007; 109:2014–2022. [PubMed: 17062729]
19. Shrestha S, Yang K, Guy C, Vogel P, Neale G, Chi H. Treg cells require the phosphatase PTEN to restrain TH1 and TFH cell responses. *Nat Immunol.* 2015; 16:178–187. [PubMed: 25559258]
20. Huynh A, DuPage M, Priyadharshini B, et al. Control of PI(3) kinase in Treg cells maintains homeostasis and lineage stability. *Nat Immunol.* 2015; 16:188–196. [PubMed: 25559257]
21. Yates J, Rovis F, Mitchell P, et al. The maintenance of human CD4 + CD25 + regulatory T cell function: IL-2, IL-4, IL-7 and IL-15 preserve optimal suppressive potency *in vitro*. *Int Immunol.* 2007; 19:785–799. [PubMed: 17545278]
22. Soond DR, Slack EC, Garden OA, Patton DT, Okkenhaug K. Does the PI3K pathway promote or antagonize regulatory T cell development and function? *Front Immunol.* 2012; 3:244. [PubMed: 22912633]
23. Burotto M, Chiou VL, Lee JM, Kohn EC. The MAPK pathway across different malignancies: A new perspective. *Cancer.* 2014; 120:3446–3456. [PubMed: 24948110]
24. Vella LJ, Pasam A, Dimopoulos N, et al. MEK inhibition, alone or in combination with BRAF inhibition, affects multiple functions of isolated normal human lymphocytes and dendritic cells. *Cancer Immunol Res.* 2014; 2:351–360. [PubMed: 24764582]
25. Kalland ME, Oberprieler NG, Vang T, Tasken K, Torgersen KM. T cell-signaling network analysis reveals distinct differences between CD28 and CD2 costimulation responses in various subsets and in the MAPK pathway between resting and activated regulatory T cells. *J Immunol.* 2011; 187:5233–5245. [PubMed: 22013130]
26. Cao J, Zhang X, Wang Q, et al. Cyclic AMP suppresses TGF-beta-mediated adaptive Tregs differentiation through inhibiting the activation of ERK and JNK. *Cell Immunol.* 2013; 285(1–2): 42–48. [PubMed: 24055734]
27. Lu L, Wang J, Zhang F, et al. Role of SMAD and non-SMAD signals in the development of Th17 and regulatory T cells. *J Immunol.* 2010; 184:4295–4306. [PubMed: 20304828]
28. Xu L, Kitani A, Stuelten C, McGrady G, Fuss I, Strober W. Positive and negative transcriptional regulation of the Foxp3 gene is mediated by access and binding of the Smad3 protein to enhancer I. *Immunity.* 2010; 33:313–325. [PubMed: 20870174]
29. Guo J, Zhang J, Zhang X, Zhang Z, Wei X, Zhou X. Constitutive activation of MEK1 promotes Treg cell instability *in vivo*. *J Biol Chem.* 2014; 289:35139–35148. [PubMed: 25361764]
30. Chang CF, D'Souza WN, Ch'en IL, Pages G, Pouyssegur J, Hedrick SM. Polar opposites: Erk direction of CD4 T cell subsets. *J Immunol.* 2012; 189:721–731. [PubMed: 22675204]

31. Shindo T, Kim TK, Benjamin CL, Wieder ED, Levy RB, Komanduri KV. MEK inhibitors selectively suppress alloreactivity and graft-versus-host disease in a memory stage-dependent manner. *Blood*. 2013; 121:4617–4626. [PubMed: 23575444]
32. Liu H, Yao S, Dann SM, Qin H, Elson CO, Cong Y. ERK differentially regulates Th17- and Treg-cell development and contributes to the pathogenesis of colitis. *Eur J Immunol*. 2013; 43:1716–1726. [PubMed: 23620016]
33. Colamonici M, Blyth G, Saleiro D, et al. Dual targeting of acute myeloid leukemia progenitors by catalytic mTOR inhibition and blockade of the p110alpha subunit of PI3 kinase. *Oncotarget*. 2015; 6:8062–8070. [PubMed: 25823922]
34. Bradford LS, Rauh-Hain A, Clark RM, et al. Assessing the efficacy of targeting the phosphatidylinositol 3-kinase/AKT/mTOR signaling pathway in endometrial cancer. *Gynecol Oncol*. 2014; 133:346–352. [PubMed: 24561032]
35. Bendell JC, Rodon J, Burris HA, et al. Phase I, dose-escalation study of BKM120, an oral pan-Class I PI3K inhibitor, in patients with advanced solid tumors. *J Clin Oncol*. 2012; 30:282–290. [PubMed: 22162589]
36. Zhang J, Hong Y, Shen J. Combination treatment with perifosine and MEK-162 demonstrates synergism against lung cancer cells *in vitro* and *in vivo*. *Tumour Biol*. 2015; 36:5699–5706. [PubMed: 25697899]
37. Alves NL, van Leeuwen EM, Derks IA, van Lier RA. Differential regulation of human IL-7 receptor alpha expression by IL-7 and TCR signaling. *J Immunol*. 2008; 180:5201–5210. [PubMed: 18390701]
38. Liu W, Putnam AL, Xu-Yu Z, et al. CD127 expression inversely correlates with FoxP3 and suppressive function of human CD4⁺ T reg cells. *J Exp Med*. 2006; 203:1701–1711. [PubMed: 16818678]
39. Wells AD, Gudmundsdottir H, Turka LA. Following the fate of individual T cells throughout activation and clonal expansion. Signals from T cell receptor and CD28 differentially regulate the induction and duration of a proliferative response. *J Clin Invest*. 1997; 100:3173–3183.
40. Camps M, Ruckle T, Ji H, et al. Blockade of PI3Kgamma suppresses joint inflammation and damage in mouse models of rheumatoid arthritis. *Nat Med*. 2005; 11:936–943. [PubMed: 16127437]
41. Rutenberg JJ, Boyce C, Hingorani R, Putnam A, Ghanekar SA. Rapid assessment of *in vitro* expanded human regulatory T cell function. *J Immunol Methods*. 2011; 372(1–2):95–106. [PubMed: 21781972]
42. Canavan JB, Afzali B, Scotta C, et al. A rapid diagnostic test for human regulatory T-cell function to enable regulatory T-cell therapy. *Blood*. 2012; 119:e57–e66. [PubMed: 22219224]
43. Salmond RJ, Emery J, Okkenhaug K, Zamoyska R. MAPK, phosphatidylinositol 3-kinase, and mammalian target of rapamycin pathways converge at the level of ribosomal protein S6 phosphorylation to control metabolic signaling in CD8 T cells. *J Immunol*. 2009; 183:7388–7397. [PubMed: 19917692]
44. Herzog A, Bian Y, Vander Broek R, et al. PI3K/mTOR inhibitor PF-04691502 antitumor activity is enhanced with induction of wild-type TP53 in human xenograft and murine knockout models of head and neck cancer. *Clin Cancer Res*. 2013; 19:3808–3819. [PubMed: 23640975]
45. Yoshioka K, Yoshida K, Cui H, et al. Endothelial PI3K-C2alpha, a class II PI3K, has an essential role in angiogenesis and vascular barrier function. *Nat Med*. 2012; 18:1560–1569. [PubMed: 22983395]
46. Zhao JJ, Cheng H, Jia S, et al. The p110alpha isoform of PI3K is essential for proper growth factor signaling and oncogenic transformation. *Proc Natl Acad Sci U S A*. 2006; 103:16296–16300. [PubMed: 17060635]
47. Cirit M, Wang CC, Haugh JM. Systematic quantification of negative feedback mechanisms in the extracellular signal-regulated kinase (ERK) signaling network. *J Biol Chem*. 2010; 285:36736–36744. [PubMed: 20847054]
48. Wang CC, Cirit M, Haugh JM. PI3K-dependent cross-talk interactions converge with Ras as quantifiable inputs integrated by Erk. *Mol Syst Biol*. 2009; 5:246. [PubMed: 19225459]

49. Moelling K, Schad K, Bosse M, Zimmermann S, Schweneker M. Regulation of Raf-Akt cross-talk. *J Biol Chem*. 2002; 277:31099–31106. [PubMed: 12048182]
50. Sasaki T, Irie-Sasaki J, Jones RG, et al. Function of PI3Kgamma in thymocyte development, T cell activation, and neutrophil migration. *Science*. 2000; 287:1040–1046. [PubMed: 10669416]
51. Hirsch E, Katanaev VL, Garlanda C, et al. Central role for G protein-coupled phosphoinositide 3-kinase gamma in inflammation. *Science*. 2000; 287:1049–1053. [PubMed: 10669418]
52. Ferguson GJ, Milne L, Kulkarni S, et al. PI(3)Kgamma has an important context-dependent role in neutrophil chemokinesis. *Nat Cell Biol*. 2007; 9:86–91. [PubMed: 17173040]
53. Ladygina N, Gottipati S, Ngo K, et al. PI3Kgamma kinase activity is required for optimal T-cell activation and differentiation. *Eur J Immunol*. 2013; 43:3183–3196. [PubMed: 24030559]
54. Abu-Eid R, Samara RN, Ozbun L, et al. Selective inhibition of regulatory T cells by targeting the PI3K-Akt pathway. *Cancer Immunol Res*. 2014; 2:1080–1089. [PubMed: 25080445]
55. Walker MR, Kasprovicz DJ, Gersuk VH, et al. Induction of FoxP3 and acquisition of T regulatory activity by stimulated human CD4+CD25-T cells. *J Clin Invest*. 2003; 112:1437–1443. [PubMed: 14597769]
56. Tapia-Abellan A, Ruiz-Alcaraz AJ, Hernandez-Caselles T, et al. Role of MAP kinases and PI3K-Akt on the cytokine inflammatory profile of peritoneal macrophages from the ascites of cirrhotic patients. *Liver Int*. 2013; 33:552–560. [PubMed: 23331611]
57. Ruiz-Alcaraz AJ, Martinez-Esparza M, Cano R, et al. Peritoneal macrophage priming in cirrhosis is related to ERK phosphorylation and IL-6 secretion. *Eur J Clin Invest*. 2011; 41:8–15. [PubMed: 20731703]
58. Nakamura A, Johns EJ, Imaizumi A, Yanagawa Y, Kohsaka T. Modulation of interleukin-6 by beta2-adrenoceptor in endotoxin-stimulated renal macrophage cells. *Kidney Int*. 1999; 56:839–849. [PubMed: 10469353]
59. Craig R, Larkin A, Mingo AM, et al. p38 MAPK and NF-kappa B collaborate to induce interleukin-6 gene expression and release. Evidence for a cytoprotective autocrine signaling pathway in a cardiac myocyte model system. *J Biol Chem*. 2000; 275:23814–23824. [PubMed: 10781614]
60. Chio CC, Chang YH, Hsu YW, Chi KH, Lin WW. PKA-dependent activation of PKC, p38 MAPK and IKK in macrophage: Implication in the induction of inducible nitric oxide synthase and interleukin-6 by dibutyl cAMP. *Cell Signal*. 2004; 16:565–575. [PubMed: 14751542]
61. Hoffmann E, Dittrich-Breiholz O, Holtmann H, Kracht M. Multiple control of interleukin-8 gene expression. *J Leukoc Biol*. 2002; 72:847–855. [PubMed: 12429706]
62. Britschgi A, Andraos R, Brinkhaus H, et al. JAK2/STAT5 inhibition circumvents resistance to PI3K/mTOR blockade: A rationale for cotargeting these pathways in metastatic breast cancer. *Cancer Cell*. 2012; 22:796–811. [PubMed: 23238015]
63. Basu S, Hubbard B, Shevach EM. Foxp3-mediated inhibition of Akt inhibits Glut1 (glucose transporter 1) expression in human T regulatory cells. *J Leukoc Biol*. 2015; 97:279–283. [PubMed: 25492937]
64. Priyadarshini B, Turka LA. T-cell energy metabolism as a controller of cell fate in transplantation. *Curr Opin Organ Transplant*. 2015; 20:21–28. [PubMed: 25563988]
65. Guinan EC, Cole GA, Wylie WH, et al. Ex vivo costimulatory blockade to generate regulatory T cells from individuals awaiting kidney transplantation. *Am J Transplant*. 2016 doi: 10.1111/ajt.13725 [Epub ahead of print].

Figure 1. Selection of inhibitors and dose titration *in vitro*

Human peripheral blood mononuclear cells (PBMCs) were labeled with carboxyfluorescein succinimidyl ester (CFSE) and stimulated for 72 h with soluble anti-CD3 and anti-CD28 in the presence of vehicle or inhibitors at decreasing doses. (A) Initial dose titrations for IC-81114 (δ), AS-605240 (γ), BYL-719 (α), BKM-120 (pan), and mitogen-activated protein kinase/extracellular signal-regulated (MEK)-162. Concentrations marked in red are those determined to be “optimal” and used for subsequent studies – see text for details. (B) Counting beads were used to determine mitotic indices of T lymphocyte subsets following treatment with inhibitors at indicated doses. The mitotic index (see Materials and Methods) of each inhibitor-treated T lymphocyte subset was compared with vehicle treatment to assess for significant suppression. Summary data are shown from three separate experiments. Data are mean $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$.

Figure 2. Effects of phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase/extracellular signal-regulated (MEK) inhibition on T cell proliferation

Human peripheral blood mononuclear cell (PBMCs) were labeled with carboxyfluorescein succinimidyl ester (CFSE) and stimulated for 72 h with soluble anti-CD3 and anti-CD28 in the presence of vehicle or inhibitors at the following concentrations: 5 μ mol/L (IC-87114; PI3K δ inhibitor), 3 μ mol/L (BYL-719; PI3K α inhibitor), and 40 nmol/L (MEK-162). Counting beads were used to determine output cell numbers following stimulation, and mitotic indices of T lymphocyte subsets were assessed. Summary data are shown from six separate experiments. (A) Representative CFSE dilution comparing CD4⁺ lymphocytes and regulatory T cells (Tregs; top). Summary data showing change in mitotic indices of CD4⁺, Treg, and CD8⁺ lymphocytes (bottom). (B) Viability staining and change in cell number of output conventional CD4⁺, Tregs, and CD8⁺ lymphocytes. (C) Preservation of Foxp3 expression in Tregs, summary data (left) and representative histogram (right). Data are mean $\pm$ SEM. * p < 0.05, ** p < 0.01. (D) Change in CD45RA expression in T lymphocyte subsets (note, “vehicle” is set to 1.0 to show that loss of CD45RA compared with “unstimulated”). (E) Representative CFSE dilutions for CD4⁺CD45RA⁺ and CD4⁺CD45RO⁺ lymphocytes and summary data for change in mitotic indices. Data are mean $\pm$ SEM. * p < 0.05, ** p < 0.01.

Figure 3. Effects of phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase/extracellular signal-regulated (MEK) inhibition on expression of activation markers in T cells Human peripheral blood mononuclear cells (PBMCs) were stimulated for 24 h with soluble anti-CD3 and anti-CD28 in the presence of vehicle or inhibitors at the same concentrations used in Figure 2. T cell subsets were analyzed for expression of a panel of activation markers. Change in expression with inhibitor was normalized to vehicle treatment. Shown are summary data for seven separate experiments assessing change in expression of the following markers: programmed cell death 1 (PD-1), cytotoxic T lymphocyte associate protein 4 (CTLA-4), PD-1 and CTLA-4 coexpression, HLA-DR, inducible T cell costimulator (ICOS), CD27, CD25 mean fluorescence intensities (MFIs), and Foxp3. Data are mean $\pm$ SEM. *p < 0.05, **p < 0.01.

Figure 4. Effects of phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase/extracellular signal-regulated (MEK) inhibition on cytokine production in stimulated peripheral blood mononuclear cells (PBMCs)

Supernatants from experiments depicted in Figure 3 (stimulated for 24 h) were assessed for cytokine production by Luminex. The following were analyzed: interleukin (IL)-2, IL-6, IL-8, IL-10, IL-13, interferon (IFN) γ , tumor necrosis factor (TNF) α , granulocyte-macrophage colony-stimulating factor (GM-CSF), macrophage inflammatory protein (MIP)-1 α , and MIP-1 β . The panels depict the results from seven independent cultures, and cytokine concentrations from four or five technical replicates within each separate experiment were averaged. Data are mean $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$.

Figure 5. Phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase/extracellular signal-regulated (MEK) inhibition minimally affect regulatory T cell (Treg) suppressive function

Peripheral blood mononuclear cells (PBMCs) were stimulated with anti-CD3 and anti-CD28-coated beads for 7 h in the presence of increasing ratios of Tregs:PBMCs and vehicle or inhibitors at the same concentrations used in previous experiments. Suppression of CD69 and CD154 coexpression among CD4⁺CD25⁻ lymphocytes was a readout for Treg-mediated suppression. (A) Representative flow cytometry plots showing CD69 and CD154 coexpression (top) and summary data from four separate experiments. (B) Representative flow cytometry plots showing CD69 and CD154 coexpression in CD4⁺CD25⁻ and CD4⁺CD25^{hi} lymphocytes from PBMCs without additional Tregs (left). Summary data showing change in activation marker coexpression with vehicle or inhibitor treatment (right) from four separate experiments. Data are mean $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$.

Figure 6. Phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase/extracellular signal-regulated (MEK) inhibitors inhibit PI3K signaling in CD4⁺ and CD8⁺ lymphocytes and Tregs

Human peripheral blood mononuclear cells (PBMCs) were pre-incubated with vehicle (0.2% DMSO) or inhibitor (Delta = IC-87114, 5 μ M; Alpha = BYL-719, 3 μ M; MEK = MEK-162, 40 nM). Cells were labeled with soluble anti-CD3 and CD28 and stimulated by crosslinking for 5 min. Percentage of cells positive for phosphorylation at Akt (S273), ERK, S6, and PDK1 among T lymphocyte subsets was measured in stimulated samples and normalized to vehicle. (A) Representative flow cytometry plots and histograms illustrating phosphorylation and inhibition. (B) Summary data for five separate experiments assessing inhibition of phosphorylation at Akt, ERK, and S6 following treatment with PI3K and MEK inhibitors. Four experiments are shown for phosphorylation at PDK1. Data are mean $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$. PBMC, peripheral blood mononuclear cell.

Figure 7. Phosphatidylinositol 3-kinase (PI3K) isoform and mitogen-activated protein kinase/extracellular signal-regulated (MEK) expression are decreased among regulatory T cells (Tregs) compared with CD4⁺ lymphocytes

Equal numbers of sorted conventional CD4⁺ and Tregs (CD4⁺CD25^{hi}CD127^{lo}) were lysed and subjected to immunoblotting for PI3K δ and PI3K α isoforms and MEK 1/2 (top).

Densitometry from six separate experiments was performed to quantify differential expression of proteins between CD4⁺ lymphocytes and Tregs (bottom). Data are mean $\pm$ SEM. *p < 0.05, **p < 0.01.