Histone Deacetylases with Antagonistic Roles in *Saccharomyces cerevisiae* Heterochromatin Formation

Deborah M. Thurtle-Schmidt, Anne E. Dodson, and Jasper Rine¹

Department of Molecular and Cell Biology and California Institute for Quantitative Biosciences, University of California, Berkeley, California 94720

ORCID IDs: 0000-0002-9747-9585 (D.M.T.-S.); 0000-0001-7719-0496 (A.E.D.)

ABSTRACT As the only catalytic member of the Sir-protein gene-silencing complex, Sir2's catalytic activity is necessary for silencing. The only known role for Sir2's catalytic activity in *Saccharomyces cerevisiae* silencing is to deacetylate N-terminal tails of histones H3 and H4, creating high-affinity binding sites for the Sir-protein complex, resulting in association of Sir proteins across the silenced domain. This histone deacetylation model makes the simple prediction that preemptively removing Sir2's H3 and H4 acetyl substrates, by mutating these lysines to unacetyltable arginines, or removing the acetyl transferase responsible for their acetylation, should restore silencing in the Sir2 catalytic mutant. However, this was not the case. We conducted a genetic screen to explore what aspect of Sir2's catalytic activity has not been accounted for in silencing. Mutation of a nonsirtuin histone deacetylase, Rpd3, restored Sir-protein-based silencing in the absence of Sir2's catalytic activity. Moreover, this antagonism could be mediated by either the large or the small Rpd3-containing complex. Interestingly, this restoration of silencing appeared independent of any known histone H3 or H4 substrates of Rpd3. Investigation of Sir-protein association in the Rpd3 mutant revealed that the restoration of silencing was correlated with an increased association of Sir proteins at the silencers, suggesting that Rpd3 was an antagonist of Sir2's function in nucleation of Sir proteins to the silencer. Additionally, restoration of silencing by Rpd3 was dependent on another sirtuin family member, Hst3, indicating multiple antagonistic roles for deacetylases in *S. cerevisiae* silencing.

KEYWORDS Sir2; heterochromatin; Rpd3; Hst3; H4K16

SIR2 is the founding member of the sirtuin enzyme family, which deacetylates acetylated lysines on a variety of proteins. Unlike other deacetylase enzymes that catalyze the hydrolysis of acetyl-lysine, sirtuins couple lysine deacetylation to NAD⁺ hydrolysis (Imai *et al.* 2000; Tanner *et al.* 2000; Sauve *et al.* 2006). Utilizing this high-energy cofactor for deacetylation allows sirtuins to be potentially regulated by the energy balance and/or redox balance within the cell. Sirtuins regulate diverse processes across all biological kingdoms—ranging from life-span regulation and fat metabolism to chromatin modifications (Sauve *et al.* 2006). *Saccharomy-*

ces cerevisiae has five sirtuins: Sir2 and its four paralogs named Homologs of Sir2, Hst 1–4, which are involved in cell-cycle regulation, chromatin modification, and perhaps other functions as well (Brachmann *et al.* 1995; Celic *et al.* 2006; Sauve *et al.* 2006; Downey *et al.* 2013).

Of the five yeast sirtuins, Sir2, the catalytic member of the Sir-protein silencing complex, is the most thoroughly studied. Sir2's catalytic activity is necessary for silencing *HML*, *HMR*, and some genes near telomeres. A structurally intact, but catalytically inactive Sir2 protein, encoded by the *sir2N345A* allele, cannot support gene silencing (Imai *et al.* 2000) but can assemble into a Sir-protein complex along with Sir3 and Sir4 and bind to silencers (Rusché *et al.* 2002; Thurtle and Rine 2014). The catalytic activity of Sir2 is necessary for the deacetylation of histone H4 N-terminal lysine residue 16 (H4K16) in nucleosomes at silenced loci. Point mutations changing H4K16 to glutamine, which mimics the acetylated state, disrupt silencing at *HML* (Park and Szostak 1990). This

Copyright © 2016 by the Genetics Society of America

doi: 10.1534/genetics.116.190835

Manuscript received April 26, 2016; accepted for publication July 14, 2016; published Early Online August 1, 2016.

Supplemental material is available online at <http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.116.190835/-DC1>.

¹Corresponding author: 374A Stanley Hall, University of California, Berkeley, CA 94720-3220. E-mail: jrine@berkeley.edu

H4K16-to-glutamine silencing defect can be suppressed by a point mutation in *Sir3*, indicating that acetylation of H4K16 likely disrupts the *Sir3*–nucleosome association (Johnson *et al.* 1990), as confirmed by structural studies of the *Sir3* BAH domain cocrystallized with the nucleosome core (Armache *et al.* 2011; Wang *et al.* 2013). Additionally, although *Sir2* can deacetylate H3K9-acetyl and H3K14-acetyl, it shows preference for H4K16-acetyl over other acetylated positions on histones *in vitro* (Imai *et al.* 2000), and deacetylation of H4K16 at *HML* and *HMR* *in vivo* is *Sir2* dependent (Kimura *et al.* 2002; Suka *et al.* 2002). *Sas2*, a member of the MYST family of histone acetyltransferases, functions in a complex with *Sas4* and *Sas5* to acetylate H4K16, thus functioning in opposition to *Sir2* (Meijsing and Ehrenhofer-Murray 2001; Osada *et al.* 2001; Kimura *et al.* 2002; Suka *et al.* 2002).

This evidence for *Sir2* deacetylation of H4K16-acetyl combined with other biochemical and genetic studies led to the nucleation and spreading model for yeast gene silencing: *Sir2*, *Sir3*, and *Sir4* are recruited to the silent mating loci through interactions with the ORC, *Abf1*, and *Rap1* proteins, which bind the silencers in addition to their other individual roles in the cell. In the standard model for silencing, once nucleated at the silencers, *Sir2* deacetylates the H3 and H4 tails on the neighboring nucleosome, allowing the Sir-protein complex to bind to the hypoacetylated nucleosome. Iterative rounds of deacetylation lead to the spreading of Sir protein complexes across the silenced loci, resulting in silencing at *HML* and *HMR* (Hoppe *et al.* 2002; Rusché *et al.* 2002) largely by occluding transcription factors (Loo and Rine 1994; Steakley and Rine 2015; Wang *et al.* 2015). The nucleation and spreading model makes the simple prediction that, although *Sir2*'s catalytic activity is necessary, it should be bypassed by the absence of H4K16 acetylation. However, genetically removing H4K16 acetylation either by deleting *SAS2*, the gene encoding the H4K16 acetylase, or by making the H4K16R mutant, mimicking the hypoacetylated state, does not restore the silencing defect of the *Sir2* catalytic mutant, *sir2N345A* (Yang and Kirchmaier 2006). Thus, the standard model is incomplete and missing a key role(s) for *Sir2*'s catalytic activity in silencing.

There are several possible unaccounted roles that *Sir2*'s catalytic activity might play in silencing. For example, *Sir2* may be critical for the deacetylation of other histone residues that are not targets of *Sas2*. In addition to H3 and H4 N-terminal tail lysine residues, a core lysine acetylation, H3K56-acetyl, has been implicated in silencing based upon the ability of H3K56 point mutations to disrupt telomeric silencing (Xu *et al.* 2007; Yang *et al.* 2008). H3K56-acetyl is more abundant at silenced loci in cells lacking *Sir2* function, suggesting a possible role for *Sir2* in deacetylating this core position (Xu *et al.* 2007). However, there is contradictory evidence as to whether *Sir2* can deacetylate H3K56 *in vitro* (Xu *et al.* 2007; Yang *et al.* 2008; Oppikofer *et al.* 2011). *In vivo*, there is strong evidence that H3K56-acetyl is deacetylated after deposition into the chromatin by either of two other sirtuins, *Hst3* and *Hst4* (Celic *et al.* 2006; Yang *et al.*

2008). Alternatively, *Sir2* may deacetylate an undiscovered nonhistone protein with a role in silencing. *Sir2*-dependent deacetylation of the nonhistone proteins *Ihf1* and *Pck1* certainly expands the possibilities (Lin *et al.* 2009; Downey *et al.* 2013).

The sirtuin catalytic reaction itself, rather than a specific deacetylation *per se*, could conceivably be necessary for silencing. Sirtuins couple lysine deacetylation to the hydrolysis of NAD⁺, producing the metabolites, 2-O-acetyl-ADP-ribose (OAAR) and nicotinamide (NAM) (Tanner *et al.* 2000). In mammalian cells, the histone variant macroH2A of the inactive X binds OAAR, generated from the sirtuin SIRT1, with a *K_d* of the same order of magnitude as the concentration of OAAR in a cell (Kustatscher *et al.* 2005; Lee *et al.* 2008). One study with yeast proteins reported that *in vitro* OAAR can promote structural changes in the Sir protein complex, as assayed by electron microscopy (Liou *et al.* 2005). Additionally, OAAR increases the affinity of the Sir protein complex for reconstituted chromatin (Martino *et al.* 2009), suggesting that OAAR could enhance silencing by promoting a critical level of chromatin compaction. However, an elegant genetic study has shown that a sirtuin reaction is not necessary for gene silencing in *Saccharomyces* (Chou *et al.* 2008). Thus there is no evidence for an *in vivo* role of OAAR in silencing or anything else in yeast, but the possibility remains open.

To shed light on what additional proteins or processes require *Sir2*'s catalytic function for silencing, we performed a genetic screen for mutants that could restore silencing of *HML* and *HMR* in cells with a catalytically inactive form of *Sir2*, encoded by *sir2N345A* as the only source of *Sir2*. In addition, in the parental strain, H4K16 acetylation was preemptively blocked by deletion of *SAS2*, which encodes the relevant H4K16 acetylase. *A priori*, there was one known gene, *SUM1*, in which a particular mutation (*SUM1-1*) could, in principle, be recoverable from this screen. The *SUM1-1* mutation can restore silencing in cells lacking any of the four SIR genes by causing the assembly of a different type of silenced chromatin (Klar *et al.* 1985; Laurenson and Rine 1991; Chi and Shore 1996; Rusché and Rine 2001). Although the mechanism of *SUM1-1*'s function was not germane to our goal, the existence of this very rare mutation provided a positive control for the effectiveness of the screen. The mutants recovered from the screen revealed unappreciated roles for multiple histone deacetylases with antagonistic functions at *HMR* and *HML* and *Sir2*'s catalytic function in the nucleation of the Sir proteins at the silencers.

Materials and Methods

Yeast strains, plasmids, and culture

All yeast strains were derived from the W303 background and genotypes are indicated in Supplemental Material, Table S1. Deletions were constructed through one-step integration of knockout cassettes (Longtine *et al.* 1998). The *hmr1Δ::URA3* reporter was previously described (Zill *et al.* 2010). All plasmids used are indicated in Table S2. Plasmids bearing

the histone point mutations were introduced into strains containing a pRS316 plasmid bearing wild-type H3 and H4 (pJR2657). Plasmid swap using 5-FOA counterselection against the pRS316 plasmid was used to recover yeast with only the mutant histone genes. The *rp3P264L* point mutant (JRY9568) was determined through cloning by complementation with a previously described yeast plasmid library containing both *TRP1* and G418-resistance selectable markers (Jauert *et al.* 2005). To determine silencing phenotype, five-fold serial dilutions of overnight cultures grown in rich medium were spotted onto appropriate media at an initial approximate density of 4×10^7 /ml as in Zill *et al.* (2010). To test for mating ability, cells of the strain of interest were manually mixed on a YPD plate with an α -mating tester (JRY2728) or the α -mating tester (JRY2726), grown overnight at 30°, and then replica plated onto minimal medium plates (YM), to select for successful mating events.

EMS mutagenesis

EMS mutagenesis was conducted as described (Amberg *et al.* 2005). Briefly, 10 OD units (Absorbance = 600 nm) were pelleted, washed with distilled water, and resuspended in 0.1 M sodium phosphate buffer (pH 7). A total of 30 μ l of EMS was added to the sample and incubated at 30° for 1 hr. Cells were washed with 5% sodium thiosulfate and resuspended in PBS and plated on YPD. Plates were grown overnight on YPD, then replica plated to 5-FOA. After mutagenesis, yeast was grown at room temperature to allow for the possibility of conditional alleles of essential genes.

Whole-genome-sequencing library preparation and data analysis

DNA for whole-genome sequencing analysis was isolated following the yeast DNA extraction protocol in (Hoffman and Winston 1987). For the JRY9567 (*sir2N345A*, *sas2* Δ , *hmra1* Δ ::*URA3*, *sum1-1*) libraries, three double-backcrossed segregants exhibiting the mutant phenotype were grown to saturation in a liquid culture and DNA extracted as described. The three segregants, JRY9614, JRY9615, and JRY9616, were barcoded separately and sequenced with the unmutagenized parent, JRY9097, and the original mutant, JRY9567. For the JRY9569 (*sir2N345A*, *sas2* Δ , *hmra1* Δ ::*URA3*, *rp3P397H*) and JRY9570 (*sir2N345A*, *sas2* Δ , *hmra1* Δ ::*URA3*, *rp3G290N*, *orc3A333V*) libraries, bulk segregant analysis as in Birkeland *et al.* (2010) was performed to determine causative mutations. Ten segregants with a silencing restoration phenotype and 10 segregants with a nonsilencing phenotype from the JRY9617 and JRY9618 diploids were pooled equally according to the OD₆₀₀ of each culture. DNA was then isolated from these pooled cultures. A total of 4 μ g of DNA was sheared with the S220 Focused-ultrasonicator (Covaris, Woburn, MA) and the libraries were constructed using the Illumina Tru-Seq library preparation kit. Samples were indexed and sequenced in two lanes on the Illumina HiSeq2500 as 100 bp single end reads. Reads were mapped using Burrows-Wheeler Alignment tool (BWA) (Li and

Durbin 2009) to a modified SacCer 2 genome. SNPs were detected using VCFtools (Danecek *et al.* 2011) and visualized on the Integrative Genomics Viewer (Thorvaldsdóttir *et al.* 2013). Number of reads mapped for each sample is recorded in Table S3. Reads have been deposited in the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) at <http://www.ncbi.nlm.nih.gov/sra> under accession no. SRP077398.

RNA isolation and analysis

RNA was prepared as in Ozyaydin and Rine (2010). Complementary DNA (cDNA) was synthesized using Superscript III First-Strand Synthesis (Invitrogen, Carlsbad, CA), utilizing oligo-dT primers. Amplification values for all RNA primer sets were normalized to *ACT1* cDNA amplification values for that sample. RNA analysis was performed on four independent RNA preparations and error bars represent standard error.

Chromatin isolation and immunoprecipitation

A total of 70 OD units (Absorbance = 600 nm) of logarithmically growing cells were cross-linked with 1% formaldehyde for 30 min at room temperature. Cells were lysed with 0.5-mm zirconia beads in 1 ml FA lysis buffer using the MP Fastprep-24. Chromatin was prepared as previously described (Aparicio *et al.* 2005). Chromatin was immunoprecipitated with 25 μ l of anti-c-Myc agarose (Sigma, St. Louis, MO; cat. no. 7470) overnight at 4°.

Washes, elutions, and isolation of DNA were performed as previously described (Aparicio *et al.* 2005). After the reversal of cross-links with Proteinase K, DNA was isolated with QIA-GEN PCR purification kit (Valencia, CA).

Quantitative PCR

Quantitative PCR (qRT-PCR) analysis was conducted on the MX3000P machine from Stratagene (La Jolla, CA) using DyNAmo HS SYBR Green qPCR kit (Thermo Fisher Scientific, Waltham, MA). All primer sequences for RNA and chromatin immunoprecipitation (ChIP) analysis are listed in Table S4.

Data availability

Strains and plasmids are available upon request. Table S1 contains detailed genotypes of all strains used in this study.

Results

Mutations in a few genes could restore silencing independently of Sir2's catalytic activity

To investigate what function of Sir2's catalytic activity is not accounted for in the gene silencing model, we screened for mutations able to suppress the silencing defect of a *sas2* Δ *sir2N345A* strain. As a silencing reporter, the *a1* open reading frame at *HMR* was replaced with *URA3* (Figure 1A). *URA3* expression causes sensitivity to the drug 5-FOA, allowing for selection of cells that have silenced *URA3* at *HMR* by their ability to grow on 5-FOA (Boeke *et al.* 1984). The *sir2N345A* *sas2* Δ *MATa* strain (JRY9097) was mutagenized with EMS

and mutants were initially identified by growth on 5-FOA medium. To distinguish mutants able to grow on 5-FOA because of silencing restoration from mutants with lesions in the *URA3* reporter gene, a secondary screen was conducted for putative mutant's restoration of mating ability, which required silencing at *HML α* as well as at *HMRa1 Δ ::URA3* (Figure 1A).

From 1500 5-FOA resistant colonies, four mutants were recovered that also exhibited some restoration of mating ability (Figure 1B). All four mutants were able to grow both on solid medium lacking uracil in addition to medium containing 5-FOA, indicating that silencing at *HMRa1 Δ ::URA3* was only partially restored. Molecular quantitation of the extent of restored silencing is discussed below. Silencing was also not restored to wild-type levels at *HML* as evidenced by the inefficient mating of the mutants (see also Figure 2B). Mutants were mated to an isogenic strain of the opposite mating type, and the corresponding diploids were sensitive to 5-FOA. Thus the silencing phenotype of the mutants was recessive (data not shown).

Several different strategies were used to identify the causative mutation. For one of the mutants, the causative mutation was determined by individually sequencing multiple backcrossed segregants. The causative mutation was determined as the SNP that all of the segregants with restored silencing shared in common, but was absent in the unmutagenized parent strain. This mutant had exactly the same point mutation as *SUM1-1* (Klar *et al.* 1985; Chi and Shore 1996). As noted before, this mutation appears dominant by some assays and recessive by others (Laurenson and Rine 1991).

The other three mutants had different point mutations in the gene encoding the nonsirtuin histone deacetylase, *RPD3*. One of these mutants (JRY9568) was amenable to cloning by complementation and a plasmid with a genomic fragment containing *RPD3* complemented the silencing phenotype of this mutant. Sanger sequencing of the *RPD3* gene in this mutant identified a transition mutation causing a proline-to-leucine change at amino acid 264 (hereafter referred to as *rpdp3P264L*). The causative mutation of the two additional mutants (JRY9569 and JRY9570) was determined through pooled linkage analysis (Birkeland *et al.* 2010). Both mutants analyzed by pooled linkage analysis revealed a mutation in *RPD3* specific to the mutant pool (Figure S1, A and B). Thus, three of four mutants recovered from this screen identified nonsynonymous mutations in *RPD3* (Figure 1B).

We focused most of the following analyses on understanding how *rpdp3* mutations could partially bypass the need for Sir2's catalytic activity. However, interestingly, one of the two mutants (JRY9570) analyzed by pooled linkage analysis, in addition to the *rpdp3G290N* mutation, also contained an alanine-to-valine substitution at amino acid 333 in *ORC3*, which encodes a subunit of ORC, that segregated at 94% frequency with the mutant pool (46 of 49 reads) (Figure S2). All other mutations showed a heterozygous distribution in both the mutant and wild-type pools between 35 and 65% frequency and hence were discarded from consideration. *ORC3* and *RPD3* are present on different chromosomes so the associa-

tion of the *ORC3* mutation was not due to physical linkage to *RPD3*. The tetrad analysis used to isolate single segregants for pooling showed that the restoration-of-silencing phenotype showed a 2:2 segregation, indicating a single Mendelian locus for the causative mutation. However, the segregants that were pooled were those with the most vigorous growth on 5-FOA. Thus the cosegregation of the *rpdp3* and *orc3* mutations implied that the *orc3A333V* mutation enhanced the *rpdp3* mutant's ability to restore silencing.

RPD3 as an antagonist of Sir-protein-based silencing

Rpd3 has been implicated in silencing in several ways. *Rpd3* restricts the spread of Sir proteins at the boundary of silent chromatin domains (Sun and Hampsey 1999; Zhou *et al.* 2009). An *rpdp3* mutation suppresses the silencing defect of an *HMR-E* silencer with a defective *Rap1* binding site (Vannier *et al.* 1996). Additionally, mutations in the *Drosophila* *Rpd3* ortholog, as well as deletion of yeast *RPD3*, result in increased position-effect variegation and telomere position effect, respectively (De Rubertis *et al.* 1996). To determine whether the point mutations isolated here had some novel property leading to the silencing phenotype, *RPD3* was deleted in the original *sir2N345A sas2 Δ* parent strain and a strain of similar genotype, but opposite mating type. As with the point mutant, the deletion also suppressed the silencing defect of the *sir2N345A* mutation (Figure 2, A and B). Hence the point mutation likely caused loss of *Rpd3* function. Interestingly, as judged by resistance to 5-FOA, silencing at *HMR* was slightly better in the *rpdp3P264L* mutant than the *rpdp3 Δ* . However, this point mutant was derived from the original EMS mutagenesis and may have had background mutations improving silencing slightly beyond the silencing achieved by the complete loss of *Rpd3* function.

Rpd3 is a histone deacetylase found in two distinct complexes: the *Rpd3* large complex (*Rpd3L*) and *Rpd3* small complex (*Rpd3S*). *Rpd3S* deacetylates histone H4 in nucleosomes in the coding regions of actively transcribed genes to prevent intragenic transcription (Carrozza *et al.* 2005). The large complex is involved in general transcriptional repression of many genes (Kasten *et al.* 1997; Lechner *et al.* 2000). To determine if *Rpd3L* or *Rpd3S* was specifically involved in the restoration of silencing, we deleted components specific to the small complex, *EAF3* and *RCO1*, or the large complex, *PHO23*, *SAP30*, *ASH1*, and *SDS3*, in the *sir2N345A sas2 Δ* strain to determine if, as in *rpdp3 Δ* , these deletions could also restore silencing. Neither deletions specifically affecting the small complex nor those specifically affecting the large complex restored silencing in the *sir2N345A sas2 Δ* strain (Figure 2C). However, deletion of *SIN3*, the scaffolding platform for both the *Rpd3S* and *Rpd3L* complexes, phenocopied the *rpdp3 Δ* . Thus, neither the *Rpd3L* nor *Rpd3S* specifically blocked silencing in the *sir2N345A sas2 Δ* strain.

Sin3 contains four paired amphipathic helix (PAH) domains that facilitate protein-protein interactions. To test which of these domains were required for the restoration of silencing when *SIN3* is deleted, we transformed plasmids

Figure 1 A genetic screen identified suppressors of Sir2's catalytic activity. (A) Schematic of the screen to identify mutations that restored silencing in the absence of Sir2 catalytic activity. A *sir2N345A sas2Δ* strain had the *URA3* reporter integrated in place of the *a1* ORF at *HMR*. Expression of the *URA3* gene resulted in cells that grow on complete synthetic media lacking uracil but are sensitive to 5-FOA. Expression of the α -genes from *HML* and *a*-genes from *MAT* result in a pseudodiploid cell identity that cannot mate. This strain was mutagenized with EMS and mutants were picked that re-established silencing as assayed by resistance to 5-FOA and the ability to mate. (B) The phenotypes of a *SIR2* wild-type strain (JRY10676) exhibiting complete silencing, the parent strain (JRY9097), and the four mutants that were isolated from the suppressor screen. Fivefold

serial dilutions assayed for growth phenotypes on the indicated media. Growth in the patches on the right indicates successful matings as either an *a*- or α -cell identity.

expressing *SIN3* with each of the PAH domains deleted (Wang and Stillman 1993). Only the *pah3* mutant was still able to silence *HMR* (Figure 2D), indicating that the deletion of PAH3 was required for the restoration of silencing by *sin3Δ* and *rp3Δ*.

Both *Rpd3* complexes are involved in transcriptional repression, with mutation of *RPD3* resulting in increased expression of genes (Kurdistani *et al.* 2004; Carrozza *et al.* 2005; Chen *et al.* 2012). Therefore it was odd that silencing should be partially restored in the absence of *Rpd3*. To test directly whether the effect of the *rp3Δ* mutation on silencing was transcriptional, and if so, specific to *HML* and *HMR*, or a more general effect on *MAT* gene expression, we measured RNA expression from the original unmutagenized parent strain (*RPD3*), the *rp3P264L* point mutant, and the *rp3Δ* and compared the expression from the *MAT* locus to the expression from *HML* and *HMR*. If the restoration of silencing were due to general down regulation of the *a1* promoter, then both the *a1* gene at *MAT* and the *URA3* gene at *HMR* would show decreased expression in the mutant. In contrast, if the apparent suppression of *sir2N345A* were due to silencing restoration, then the decrease in expression would be specific to *HMR* and not apparent at the *MAT* locus. Consistent with a silencing-specific role for *rp3Δ* mutants rather than a transcriptional down-regulation, decreased expression was observed only at *HML* and *HMR* and not the *MAT* locus for both the recovered *rp3P264L* point mutant and the *rp3Δ* (Figure 3A). Interestingly, as was observed by growth on 5-FOA, the *rp3P264L* point mutant showed more com-

plete silencing of *URA3* at *HMR* than the *rp3Δ*, showing concordance between the assays. Therefore, the phenotypes described here of the *rp3Δ* mutations were due to a restoration of silencing.

As described above, the *SUM1-1* mutation restores silencing independently of Sir proteins. Since the *SUM1-1* mutation was recovered in this very screen, it was possible that the *rp3Δ* mutants also restored silencing through a Sir-protein-independent mechanism. Since we were interested in understanding unaccounted roles for Sir2's catalytic activity in silencing, we tested whether the *rp3Δ* mutants' restoration of silencing was Sir-protein dependent, or whether they functioned by some independent means, like *SUM1-1*. Silencing restoration by the *rp3Δ* mutants was dependent on both *Sir3* and *Sir4*, indicating that Sir protein-dependent silencing had been restored (Figure 3B). To determine if restoration of Sir silencing by the *rp3Δ* was due to misregulation of *Sir3* and *Sir4*, we also measured messenger RNAs (mRNAs) from *SIR3* and *SIR4* in *sir2N345A sas2Δ* strains, with and without the *rp3P264L* point mutant or *rp3Δ*. *SIR3* showed a modest 1.5-fold increase in expression in both the *rp3Δ* and *rp3P264L* mutants (Figure S3); no increase was noted for *SIR4*. Previous overexpression of *SIR3* to ~16-fold wild-type levels had only marginal effects on silencing (Yang and Kirchmaier 2006), thus this modest increase in *SIR3* expression in the *rp3Δ* mutant was unlikely to be responsible for the silencing restoration. Additionally, *sir2N345A* itself was also required to restore silencing by *rp3Δ*. A *sir2Δ rp3Δ* strain that was not expressing the catalytically dead *sir2N345A* allele

Figure 2 Mutation of *RPD3* and *SIN3* restored silencing. (A) Fivefold serial dilutions of yeast were spotted on plates to determine their growth phenotype. Growth on 5-FOA indicates silencing of the *hmr1Δ::URA3* reporter. All strains are in the *sir2N345A sas2Δ* background. The first row is the unmutagenized, *RPD3* parent (JRY9097). The next two rows are replicates of the *rpd3P264L* point mutant (JRY9568). The fourth row is isogenic to the unmutagenized parent except for the opposite mating type, *MATα*, and *ade2-1* (JRY9572). In the fifth row, *RPD3* is deleted in the parent strain (JRY9573). In the last row, *RPD3* is deleted in the *MATα ade2-1* strain isogenic to the parent (JRY9574). (B) The parental strain (JRY9097) and the recovered *rpd3P264L* mutant (JRY9568), and the *rpd3Δ* (JRY9573), were crossed with the *MATα* mating tester (JRY2728) to test for silencing at *HML*. Growth on the media indicates mating and thus silencing. (C) The *Rpd3* subunit gene indicated was deleted in JRY9097 to determine if deletion of that subunit could also restore silencing. Green labels indicate *Rpd3L* subunits and red labels indicate *Rpd3S* subunits. (D) The *sin3Δ* strain that exhibited silencing restoration was transformed with plasmids expressing full-length *SIN3* or one of the four *SIN3* PAH domain deletions (M1635–M1639). Silencing was assayed on media lacking histidine to select for the plasmid.

remained unable to silence (Figure 3B). Thus the entire Sir-protein complex, but not Sir2's catalytic activity, was required for *rpd3Δ* to restore silencing.

Sir proteins have been implicated in silencing of synthetic telomeric constructs (Gottschling *et al.* 1990; Renauld *et al.* 1993; Fourel *et al.* 1999). However, more recent studies have shown that Sir-protein-based silencing at native telomeres is less pervasive than originally thought (Loney *et al.* 2009; Ellahi *et al.* 2015). To test if the restoration of silencing by the *rpd3* mutant also occurred at native telomeres, we measured RNA expression of a native telomeric gene, *COS8*,

shown to have Sir2-dependent silencing. We observed no significant change in expression of *COS8* in the *rpd3* point mutant or *rpd3Δ* (Figure S4), indicating that this mutant does not broadly affect telomeric silencing.

Additionally, the restoration of silencing by the *rpd3Δ* mutant was observed only in the absence of wild-type *SAS2* (Figure 3C). Thus *rpd3Δ* was unable to compensate for the loss of Sir2 catalytic activity when H4K16 was acetylated. These results illustrate that the phenotype revealed by *rpd3Δ* implied the existence of an unaccounted function(s) for Sir2's catalytic activity in silencing.

Figure 4 Silencing by *rpd3Δ* was independent of known Rpd3 histone H4 targets. Endogenous copies of H3 and H4 were deleted and wild-type or mutant H3 and H4 were expressed from a CEN-ARS plasmid. Complete genotypes for all strains tested are in Table S1 (JRY9605, JRY9607–JRY9613, and JRY10644–JRY10655). Silencing was tested by mating to a *MATa* tester strain (JRY2726) and replica plated to minimal media. Growth indicates successful mating, which reflected extent of silencing.

protein revealed that association of *Sir3* and *Sir4* is reduced at the silencers relative to cells with wild-type *Sir2*, though the reason for this decreased association is unknown (Thurtle and Rine 2014). The origin recognition complex (ORC) binds to all four silencers and interacts with *Sir1* to nucleate the Sir-protein complex (Bell *et al.* 1993; Zhang *et al.* 2002; Hou *et al.* 2005). Since a mutation in *ORC3*, *orc3A333V* (Figure 1B and Figure S2) also cosegregated with the strongest restoration-of-silencing phenotype in a cross with one of the *rpd3* point mutants, we considered the possibility that *Rpd3* somehow antagonized Sir protein nucleation at silencers. If so, in the absence of *Rpd3* function, Sir protein association at the silencers would be expected to increase. To test this hypothesis, we ChIPed myc-tagged *Sir4* in chromatin from *SIR2 sas2Δ* (JRY10635), *sir2N345A sas2Δ* (JRY10632), and the *sir2N345A rpd3Δ sas2Δ* (JRY10633) strains to test the association of Sir proteins with silencers. As previously reported, *Sir4* association was greatly reduced at the silencers in *sir2N345A* as compared to the enrichment in *SIR2* cells (Figure 5, comparing purple and blue bars) (Rusché *et al.* 2002; Thurtle and Rine 2014). *rpd3Δ* restored *Sir4* association at the *HML-E* and *HMR-I* silencers to near wild-type levels, whereas there was a significant, but more limited increase of *Sir4* association within *HMR* and *HML*. Association of *Sir4* at *HMR-E* was not restored to wild-type levels, which is consistent with the incomplete restoration of silencing observed in the *rpd3Δ sir2N345A* strain. Thus *rpd3Δ*

restored silencing in the *sir2N345A* strain by increasing Sir-protein association at the silencers and to a lesser degree within the silenced cassette.

Restoration of silencing was dependent on *Hst3*, but independent of H3K56

Silencing is lost when yeast is grown in the presence of NAM, a competitive inhibitor of sirtuins (Landry *et al.* 2000; Bitterman *et al.* 2002). This silencing loss has been attributed to inhibition of *Sir2*. However, NAM inhibits the four yeast Hst proteins in addition to *Sir2*. *Hst3* and *Hst4* have been implicated in silencing (Yang *et al.* 2008), suggesting the possibility that NAM's effect on silencing could be due to more than just inhibition of *Sir2*. Silencing in the *rpd3* mutant provided a unique opportunity to determine if other yeast sirtuins contributed to silencing in this mutant, which was independent of *Sir2*'s catalytic activity. Surprisingly, silencing restoration by *rpd3P264L* was sensitive to NAM, thus implicating another sirtuin in silencing (data not shown). Silencing restoration by the *rpd3* mutant depended specifically on *HST3* (Figure 6A). The *hst3Δ* mutant showed a strong phenotype, whereas the *hst4Δ* mutant had no discernible phenotype, even though both deacetylate the same substrate, H3K56-acetyl, although *HST4* is the minor partner in this role (Celis *et al.* 2006). To test whether the lack of growth on 5-FOA in the *rpd3P264L hst3Δ* was due to loss of silencing, or an unrelated sensitivity to the 5-FOA drug, as has been previously reported to interfere with reporter assays (Rossmann *et al.* 2011; Takahashi *et al.* 2011), we measured *HML* and *HMR* RNA levels in the *rpd3P264L hst3Δ sirN345A* strain. As expected if *Hst3* were required for the restoration of silencing, expression at *HML* and *HMR* in this mutant was significantly elevated as compared to the *rpd3P264L sir2N345A* strain (Figure 6B). Additionally, loss of silencing in *hst3Δ* was not due to misregulation of *SIR3* or *SIR4* as their mRNA levels were similar to their levels in the *rpd3P264L* and *rpd3Δ* mutants (Figure S3). These results identified a specific role for *Hst3* in *HMR* silencing that was independent of *Hst1*, *Hst2*, and *Hst4*.

Hst3 deacetylates histone H3K56 (Celis *et al.* 2006). To determine if the dependence of silencing by the *rpd3* mutant on *Hst3* was due to a requirement for deacetylation of H3K56-acetyl, we tested silencing in appropriately marked strains expressing either the hyperacetylated mimic (Q) or hypoacetylated mimic (R) of H3K56. As reported previously, expression of the H3K56R could not rescue the silencing defect in *sir2N345A* (Yang *et al.* 2008) (Figure 6C). Additionally, neither H3K56 acetylation-state mimic interfered with the silencing restoration in the *sir2N345A sas2Δ* strain provided by *rpd3Δ* (Figure 6C). In some contexts, the arginine and glutamine mutations may not be adequate mimics of the histone acetylation states. Thus, to provide an independent test of whether H3K56-acetyl blocked the ability of the *rpd3* mutant to restore silencing when *HST3* was deleted, we tested whether deletion of *RTT109*, which encodes the histone acetyltransferase that acetylates H3K56, could restore

Figure 5 Deletion of *RPD3* increased Sir4 association with the silencers. Sir4-myc ChIP was performed in the *MATa sas2Δ hmra1Δ::URA3* with the following *SIR2* and *RPD3* genotypes: *SIR2 RPD3* (JRY10635), *sir2N345A RPD3* (JRY10632), and *sir2N345A rpd3Δ* (JRY10633). No enrichment was observed in a strain not expressing the Sir4-myc (JRY10634). Enrichment is shown as the Immunoprecipitation (IP)/input values relative to the *SEN1* IP/input values as a negative control. Error bars represent standard error of four biological replicates. The asterisks indicate a *P*-value of <0.05 (Student's *t*-test).

silencing in the *rpd3P264L hst3Δ* strain. The *rtt109Δ* mutation did not restore silencing in cells lacking *HST3* (Figure 6D). Thus, there appeared to be another substrate of *Hst3* whose hypoacetylation promoted silencing, at least in the absence of *SIR2* and *RPD3* catalytic function.

Discussion

There were two ways to restore silencing independent of *Sir2* catalytic activity

The genetic screen identified four mutants that restored silencing independent of *Sir2*'s catalytic activity. One caused the very same amino acid substitution of *SUM1* previously reported, and the other three were all in *RPD3*. Thus few genes can be mutated to restore silencing independent of *Sir2*'s catalytic activity. The silencing restoration observed by *SUM1-1* in the *sir2N345A sas2Δ* double mutant was dependent on *Hst1* (data not shown), indicating that it restored silencing through the previously described mechanism (Rusché and Rine 2001). *Hst1* was not required for the silencing restored in *rpd3* mutants, described here. The *Rpd3* protein is part of two different complexes, suggesting that mutations in genes for other members of the two complexes could potentially have been recovered. However, of the genes tested, only those encoding subunits of both complexes, *RPD3* and *SIN3*, when deleted, restored silencing. Mutations in the genes encoding subunits specific to either Rpd3L or Rpd3S, had no impact on silencing. Hence, it appeared that both complexes prevented silencing in the *sir2N345A sas2Δ* double mutant.

We note that *rpd3Δ* is reported to be synthetically lethal in combination with *sas2Δ* in two previous reports (Collins *et al.* 2007; Ehrentraut *et al.* 2010). This synthetic lethality appears to result from the silencing of some essential gene(s), presumably near telomeres, given that the lethality is suppressed by mutations in *SIR* genes. Conceivably, such synthetic lethality would be a problem in our strains in which

silencing was restored in the *sas2Δ sir2N345A rpd3* mutant, yet we did not see any growth defect associated with the triple mutants. Presumably the level of silencing restored in these strains was not sufficient to interfere with the function of a telomere proximal essential gene(s). Additionally, we did not observe restoration of telomeric silencing in *rpd3* mutants.

Rpd3 antagonized *Sir*-protein-based silencing

Rpd3 is part of two complexes in *S. cerevisiae*: the large complex (Rpd3L) and the small complex (Rpd3S), each with distinct activities. Previous studies in yeast indicate that *Rpd3* antagonizes silencing by restricting *Sir*-protein spreading beyond the boundary of silent loci, into adjacent euchromatic domains flanking *HML* and *HMR* and adjacent to telomeres (Zhou *et al.* 2009; Ehrentraut *et al.* 2010). However, this function of *Rpd3* seems to be mediated exclusively by the Rpd3L complex (Sun and Hampsey 1999; Loewith *et al.* 2001; Zhou *et al.* 2009). In contrast, the block to silencing in the *sas2Δ sir2N345A* mutant showed no dependence on any of the Rpd3L-specific complex members tested. Although we tested those members previously shown to have the largest silencing effects in our assay, there is the possibility that other Rpd3L complex members are necessary for the silencing restoration. Additionally, there is the possibility that multiple large-complex-specific members must be deleted in combination to restore silencing in the *sir2N345A sas2Δ* strain. The Rpd3S complex functions to deacetylate nucleosomes within coding regions of actively transcribed genes to prevent aberrant transcription initiation (Carrozza *et al.* 2005), which is dependent on the small complex-specific subunit, *Eaf3*. Our data clearly showed that *eaf3Δ* provided no silencing restoration (Figure 2). We tested all members specific to the RpdS complex and none were able to restore silencing in the *sir2N345A sas2Δ* strain; thus it is unlikely that the Rpd3S complex is involved in the restoration of silencing. Additionally, we also tested which of the four PAH domains on *Sin3* were necessary for the silencing restoration by the *sin3Δ*. Only PAH3 was required to block silencing in the *sir2N345A sas2Δ* mutant (Figure 2D). Unlike PAH1, PAH2, and PAH4, which are necessary for interactions with other *Sin3* complex members (Wang and Stillman 1993; Kasten *et al.* 1996; Washburn and Esposito 2001; Mallory *et al.* 2010), PAH3 has been shown in other organisms to specifically mediate interactions with *Rpd3* (Laherty *et al.* 1997). Since none of the large or small complex-specific members had an effect on the silencing restoration, either by deletion of the subunit or deletion of the specific interacting surface on *Sin3*, the role of *RPD3* in antagonizing silencing at *HMR* and *HML* appeared mechanistically distinct from the roles of these complexes restricting the spread of heterochromatic silencing and suppression of aberrant transcription initiation.

Rpd3 has also been shown to control origin firing in budding yeast. Deletion of *RPD3* or *SIN3* causes early activation of late origins and, similar to our observations, this phenotype is

Figure 6 Silencing by the *rpd3* mutant was dependent on Hst3. (A) Fivefold serial dilutions were spotted on the indicated media to determine growth phenotype. All strains are *sir2Δ*, *sir2N345A*, *sas2Δ*, and have the *hmra1Δ::URA3* reporter to measure silencing as growth on 5-FOA. *RPD3* and *HST* genotypes are as indicated (JRY9590–JRY9597). (B) $\alpha 2$, $a 1$, and *URA3* transcripts were measured to determine expression levels at *HML*, *MAT*, and *HMR*, respectively in *RPD3* (JRY9097), *rpd3P264L* (JRY9568), and *rpd3P264L hst3Δ* (JRY9596). RNA was normalized to *ACT1* for each sample and reported as relative to the *RPD3* sample. Error bars are the standard error of four biological replicates. The asterisks indicate a *P*-value of <0.05 (Student's *t*-test). (C) Endogenous copies of H3 and H4 were deleted and mutant H3 was expressed from a CEN-ARS plasmid (JRY9604 and JRY9606). Silencing was tested by mating to a *MATa* tester strain (JRY2726) and replica plated to minimal media. Growth indicates successful mating and silencing. (D) Fivefold serial dilutions of the indicated *RPD3*, *HST3*, and *RTT109* genotypes were spotted to determine the growth phenotype. The strains assayed are (from top to bottom) JRY9097, JRY9567, JRY9573, JRY10636, JRY10637, and JRY10638. All strains are *sir2Δ*, *sas2Δ*, *hmra1Δ::URA3*, and *sir2N345A*.

independent of any of the other complex members tested (Aparicio *et al.* 2004). Additionally, *Rpd3*'s repression of late origins is in opposition to *Sir2*, which promotes early origin firing (Yoshida *et al.* 2014). The silencers at *HMR* are late firing origins and the *HML* silencers can act as origins of replication on plasmids, but typically do not do so in the chromosome (Rusché *et al.* 2003). Additionally, establishment of Sir-based silencing is dependent on passage through S phase; however, this S-phase requirement is independent of replication (Miller and Nasmyth 1984; Kirchmaier and Rine 2001; Li *et al.* 2001). Nonetheless, our observations that Sir

proteins show increased association at the silencers in the *rpd3Δ* (Figure 5) and that a mutation in *ORC3* cosegregated with one of the *Rpd3* mutations recovered (Figure S2) may indicate that *Rpd3*'s effect on origins is antagonistic to Sir-based silencing.

***Sir2* catalytic activity promoted Sir-protein nucleation at silencers**

Previously, it had been assumed that *Sir2*'s catalytic activity was necessary for the spreading of Sir proteins across the silenced cassette and that nucleation of Sir proteins was

independent of Sir2's catalytic activity. This view emerged from nonquantitative ChIP studies that showed Sir-protein association at the silencers, but not across *HMR*, in the absence of Sir2's catalytic activity (Rusché *et al.* 2002). However, a more recent, quantitative study showed that Sir-protein association is reduced at the silencers of *HMR* and *HML* in *sir2N345A* mutants (Thurtle and Rine 2014). The results reported here confirmed that Sir-protein nucleation at silencers is defective in *sir2N345A* mutants and increased in *rpd3* mutants commensurate with the partial restoration of silencing.

What is the protein target of Rpd3 whose acetylation promotes Sir-protein binding at the silencers? We investigated all known histone H3 and H4 targets of Rpd3, individually and in combinations, which revealed no involvement in the restoration of silencing observed here (Figure 4). Additionally, the silencers are nucleosome-free regions (Weiss and Simpson 1998; Ravindra *et al.* 1999). These data suggest an alternative acetylation as the target of Sir2 and Rpd3. Although historically named histone deacetylases, Sir2 and Rpd3 deacetylate proteins other than histones (Lin *et al.* 2009; Henriksen *et al.* 2012; Kaluarachchi Duffy *et al.* 2012; Downey *et al.* 2013). For example, Rpd3 deacetylates Snf2, a subunit of the SWI/SNF chromatin remodeling complex (Kim *et al.* 2010). Another possible candidate of a non-histone Rpd3 target is the ORC. ORC binds to each of the silencers and the *HMR* silencers can act as origins of replication. A mutation in *ORC3* cosegregated with the stronger restoration-of-silencing phenotypes of one of the *RPD3* mutations. The Orc2 subunit is acetylated in human cells (Iizuka *et al.* 2006), the Hat1 acetyl transferase associates with *Saccharomyces* ORC, and mutations in *HAT1* exacerbate the defect of temperature-sensitive *orc* mutations (Suter *et al.* 2007). Moreover, N-terminal acetylation of Orc1 in *Saccharomyces* contributes to its roles in replication and silencing (Geissenhöner *et al.* 2004). Perhaps the catalytically inactive Sir2 can still recognize and bind to acetylated Orc subunits, especially if their level is elevated in the *rpd3* mutant, and contribute to the recruitment of the Sir-protein complex to the silencer.

Hst3 activity independent of H3K56 deacetylation was critical for silencing

Previous studies have implicated two other sirtuins, Hst3 and Hst4, as important for silencing at telomeres through their role in deacetylation of H3K56-acetyl (Yang *et al.* 2008). Additionally, H3K56-acetyl and Hst3 have a role in the stability of silencing at *HML* (Dodson and Rine 2015). The work reported here clearly highlighted a role for Hst3 in silencing at *HML* and *HMR* that was independent of H3K56-acetyl and perhaps of Hst4. The lack of an effect of an *hst4* mutation may simply reflect a lower activity level of Hst4 relative to Hst3, or possibly a nonshared substrate. Nevertheless, it seems inescapable that Hst3's role in silencing in the *rpd3* mutant reflects a heretofore-overlooked substrate.

How could two sirtuins (Sir2 and Hst3), the broadly acting Rpd3 deacetylase, and the H4K16 acetyltransferase (Sas2)

collaborate to produce the phenotypes described in this study? If we restrict speculation to models involving direct effects at *HML* and *HMR*, the models converge on a paradox. Mutations in Rpd3 result in elevated acetylation of most acetylated positions in core histones, with the exception of H4K16 (Robyr *et al.* 2002). Nevertheless, all such positions are hypoacetylated in heterochromatin. So how can the hyperacetylation resulting from the *rpd3* mutation, at all positions except H4K16 (due to the *sas2Δ* mutation), restore silencing in a way that depends upon the sirtuin Hst3? The function of sirtuins is NAD⁺ dependent and results in the production of two cleavage products of NAD⁺: NAM and 2-OAAR (Jackson and Denu 2002). The 2-OAAR has been proposed to be a small molecule effector of silencing. To date there is only *in vitro* evidence that this molecule has the potential for affecting chromatin structure (Liou *et al.* 2005; Martino *et al.* 2009; Ehrentraut *et al.* 2010), and one *in vivo* study shows that silencing can be established without 2-OAAR in the presence of cleverly designed fusion proteins (Chou *et al.* 2008). Nevertheless there is still room to consider the possibility that 2-OAAR might be a critical ligand to promote silencing. In this view, the *rpd3* mutation would result in elevated levels of acetylation of a substrate for Hst3, which would then produce 2-OAAR. To test this hypothesis, we mutated combinations of H3 and H4 histone N terminus residues individually and in combination (Figure 4) to test if their acetylation was necessary for the restoration of silencing by *rpd3* mutants, and none were. However our study did not evaluate the entire acetylome, so an as yet unidentified Hst3 substrate could be deacetylated by the Hst3 sirtuin to generate 2-OAAR locally and promote silencing.

Additionally, there is the possibility that global, rather than local, increases in 2-OAAR production in the cell are necessary for silencing. In the absence of Rpd3, many of its targets are now acetylated and may become substrates for Hst3, generating 2-OAAR, which may promote silencing through some allosteric mechanism. As a partial test of this hypothesis, we deleted the Nudix hydrolase *YSA1* in the *sir2N345A sas2Δ* strain. This deletion has been shown to increase intracellular 2-OAAR (Tong *et al.* 2009), but did not restore silencing (data not shown). However, Ysa1 localizes to the mitochondria, thus its activity may not alter nuclear 2-OAAR levels. Hence, for the time being, the 2-OAAR hypothesis remains a possible explanation for some or all of the phenotypes described here.

Acknowledgments

We thank Erin Osborne Nishimura for initial conception of the screen; Gavin Schlissel and Kelsey Van Dalfsen for strain contributions; David Stillman for the plasmids containing the paired amphipathic helix domain mutations in *SIN3*; members of the Rine lab for critical discussions, advice, and support; and Minyong Chung and the Vincent J. Coates Genomics Sequencing Laboratory at the University of California, Berkeley, which is supported by National Institutes of

Health (NIH) S10 Instrumentation grants S10-RR029668 and S10-RR027303. This work has been supported by National Science Foundation predoctoral fellowships (to D.M.T.-S. and A.E.D.); a University of California, Berkeley's training grant from the NIH (T32 GM 007232 34); and by a grant from the NIH National Institute of General Medical Sciences (NIGMS-31105 to J.R.).

Literature Cited

- Aparicio, J. G., C. J. Viggiani, D. G. Gibson, and O. M. Aparicio, 2004 The Rpd3-Sin3 histone deacetylase regulates replication timing and enables intra-S origin control in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 24(11): 4769–4780.
- Aparicio, O., J. V. Geisberg, E. Sekinger, A. Yang, Z. Moqtaderi *et al.*, 2005 Chromatin immunoprecipitation for determining the association of proteins with specific genomic sequences in vivo. *Curr. Protoc. Mol. Biol.* Chapter 21: Unit 21.3.
- Armache, K.-J., J. D. Garlick, D. Canzio, G. J. Narlikar, and R. E. Kingston, 2011 Structural basis of silencing: Sir3 BAH domain in complex with a nucleosome at 3.0 Å resolution. *Science* 334(6058): 977–982.
- Bell, S. P., R. Kobayashi, and B. Stillman, 1993 Yeast origin recognition complex functions in transcription silencing and DNA replication. *Science* 262(5141): 1844–1849.
- Birkeland, S. R., N. Jin, A. C. Ozdemir, R. H. Lyons, Jr., L. S. Weisman *et al.*, 2010 Discovery of mutations in *Saccharomyces cerevisiae* by pooled linkage analysis and whole-genome sequencing. *Genetics* 186: 1127–1137.
- Bitterman, K. J., R. M. Anderson, H. Y. Cohen, M. Latorre-Esteves, and D. A. Sinclair, 2002 Inhibition of silencing and accelerated aging by nicotinamide, a putative negative regulator of yeast sir2 and human SIRT1. *J. Biol. Chem.* 277(47): 45099–45107.
- Boeke, J. D., F. LaCroute, and G. R. Fink, 1984 A positive selection for mutants lacking orotidine-5'-phosphate decarboxylase activity in yeast: 5-fluoro-orotic acid resistance. *Mol. Gen. Genet.* 197(2): 345–346.
- Brachmann, C. B., J. M. Sherman, S. E. Devine, E. E. Cameron, L. Pillus *et al.*, 1995 The SIR2 gene family, conserved from bacteria to humans, functions in silencing, cell cycle progression, and chromosome stability. *Genes Dev.* 9(23): 2888–2902.
- Burke, D. J., D. C. Amberg, and J. N. Strathern (Editors), 2005 *Methods in Yeast Genetics: A Cold Spring Harbor Laboratory Course Manual*. Cold Spring Laboratory Press, Cold Spring Harbor, NY.
- Carrozza, M. J., B. Li, L. Florens, T. Suganuma, S. K. Swanson *et al.*, 2005 Histone H3 methylation by Set2 directs deacetylation of coding regions by Rpd3S to suppress spurious intragenic transcription. *Cell* 123(4): 581–592.
- Celic, I., H. Masumoto, W. P. Griffith, P. Meluh, R. J. Cotter *et al.*, 2006 The sirtuins hst3 and Hst4p preserve genome integrity by controlling histone h3 lysine 56 deacetylation. *Curr. Biol.* 16(13): 1280–1289.
- Chen, X.-F., B. Kuryan, T. Kitada, N. Tran, J.-Y. Li *et al.*, 2012 The Rpd3 core complex is a chromatin stabilization module. *Curr. Biol.* 22(1): 56–63.
- Chi, M. H., and D. Shore, 1996 SUM1-1, a dominant suppressor of SIR mutations in *Saccharomyces cerevisiae*, increases transcriptional silencing at telomeres and HM mating-type loci and decreases chromosome stability. *Mol. Cell. Biol.* 16(8): 4281–4294.
- Chou, C.-C., Y.-C. Li, and M. R. Gartenberg, 2008 Bypassing Sir2 and O-acetyl-ADP-ribose in transcriptional silencing. *Mol. Cell* 31(5): 650–659.
- Collins, S. R., K. M. Miller, N. L. Maas, A. Roguev, J. Fillingham *et al.*, 2007 Functional dissection of protein complexes involved in yeast chromosome biology using a genetic interaction map. *Nature* 446(7137): 806–810.
- Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks *et al.* 1000 Genomes Project Analysis Group, 2011 The variant call format and VCFtools. *Bioinformatics* 27(15): 2156–2158.
- De Rubertis, F., D. Kadosh, S. Henchoz, D. Pauli, G. Reuter *et al.*, 1996 The histone deacetylase RPD3 counteracts genomic silencing in *Drosophila* and yeast. *Nature* 384(6609): 589–591.
- Dodson, A. E., and J. Rine, 2015 Heritable capture of heterochromatin dynamics in *Saccharomyces cerevisiae*. *eLife* 4: e05007.
- Downey, M., B. Knight, A. A. Vashisht, C. A. Seller, J. A. Wohlschlegel *et al.*, 2013 Gcn5 and sirtuins regulate acetylation of the ribosomal protein transcription factor Ifh1. *Curr. Biol.* 23(17): 1638–1648.
- Ehrentauf, S., J. M. Weber, J. N. Dybowski, D. Hoffmann, and A. E. Ehrenhofer-Murray, 2010 Rpd3-dependent boundary formation at telomeres by removal of Sir2 substrate. *Proc. Natl. Acad. Sci. USA* 107(12): 5522–5527.
- Ellahi, A., D. M. Thurtle, and J. Rine, 2015 The chromatin and transcriptional landscape of native *Saccharomyces cerevisiae* telomeres and subtelomeric domains. *Genetics* 200: 505–521.
- Fourel, G., E. Revardel, C. E. Koering, and E. Gilson, 1999 Cohabitation of insulators and silencing elements in yeast subtelomeric regions. *EMBO J.* 18(9): 2522–2537.
- Geissenhöner A., C. Weise, and A. E. Ehrenhofer-Murray, 2004 Dependence of ORC Silencing Function on NatA-Mediated N α Acetylation in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* 24: 10300–10312.
- Gottschling, D. E., O. M. Aparicio, B. L. Billington, and V. A. Zakian, 1990 Position effect at *S. cerevisiae* telomeres: reversible repression of Pol II transcription. *Cell* 63(4): 751–762.
- Henriksen, P., S. A. Wagner, B. T. Weinert, S. Sharma, G. Bacinskaja *et al.*, 2012 Proteome-wide analysis of lysine acetylation suggests its broad regulatory scope in *Saccharomyces cerevisiae*. *Mol. Cell. Proteomics* 11(11): 1510–1522.
- Hoffman, C. S., and F. Winston, 1987 A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene* 57(2-3): 267–272.
- Hoppe, G. J., J. C. Tanny, A. D. Rudner, S. A. Gerber, S. Danaie *et al.*, 2002 Steps in assembly of silent chromatin in yeast: Sir3-independent binding of a Sir2/Sir4 complex to silencers and role for Sir2-dependent deacetylation. *Mol. Cell. Biol.* 22(12): 4167–4180.
- Hou, Z., D. A. Bernstein, C. A. Fox, and J. L. Keck, 2005 Structural basis of the Sir1-origin recognition complex interaction in transcriptional silencing. *Proc. Natl. Acad. Sci. USA* 102(24): 8489–8494.
- Iizuka, M., T. Matsui, H. Takisawa, and M. M. Smith, 2006 Regulation of replication licensing by acetyltransferase Hbo1. *Mol. Cell. Biol.* 26(3): 1098–1108.
- Imai, S., C. M. Armstrong, M. Kaerberlein, and L. Guarente, 2000 Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature* 403(6771): 795–800.
- Jackson, M. D., and J. M. Denu, 2002 Structural identification of 2'- and 3'-O-acetyl-ADP-ribose as novel metabolites derived from the Sir2 family of beta -NAD $^{+}$ -dependent histone/protein deacetylases. *J. Biol. Chem.* 277(21): 18535–18544.
- Jauert, P. A., L. E. Jensen, and D. T. Kirkpatrick, 2005 A novel yeast genomic DNA library on a geneticin-resistance vector. *Yeast* 22(8): 653–657.
- Johnson, L. M., P. S. Kayne, E. S. Kahn, and M. Grunstein, 1990 Genetic evidence for an interaction between SIR3 and histone H4 in the repression of the silent mating loci in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 87(16): 6286–6290.

- Kaluvarachchi Duffy, S., H. Friesen, A. Baryshnikova, J.-P. Lambert, Y. T. Chong *et al.*, 2012 Exploring the yeast acetylome using functional genomics. *Cell* 149(4): 936–948.
- Kasten, M. M., D. E. Ayer, and D. J. Stillman, 1996 SIN3-dependent transcriptional repression by interaction with the Mad1 DNA-binding protein. *Mol. Cell. Biol.* 16(8): 4215–4221.
- Kasten, M. M., S. Dorland, and D. J. Stillman, 1997 A large protein complex containing the yeast Sin3p and Rpd3p transcriptional regulators. *Mol. Cell. Biol.* 17(8): 4852–4858.
- Kelly, T. J., S. Qin, D. E. Gottschling, and M. R. Parthun, 2000 Type B histone acetyltransferase Hat1p participates in telomeric silencing. *Mol. Cell. Biol.* 20(19): 7051–7058.
- Kim, J.-H., A. Saraf, L. Florens, M. Washburn, and J. L. Workman, 2010 Gcn5 regulates the dissociation of SWI/SNF from chromatin by acetylation of Swi2/Snf2. *Genes Dev.* 24(24): 2766–2771.
- Kimura, A., T. Umehara, and M. Horikoshi, 2002 Chromosomal gradient of histone acetylation established by Sas2p and Sir2p functions as a shield against gene silencing. *Nat. Genet.* 32(3): 370–377.
- Kirchmaier, A. L., and J. Rine, 2001 DNA replication-independent silencing in *S. cerevisiae*. *Science* 291(5504): 646–650.
- Klar, A. J., S. N. Kakar, J. M. Ivy, J. B. Hicks, G. P. Livi *et al.*, 1985 SUM1, an apparent positive regulator of the cryptic mating-type loci in *Saccharomyces cerevisiae*. *Genetics* 111: 745–758.
- Kurdistani, S. K., S. Tavazoie, and M. Grunstein, 2004 Mapping global histone acetylation patterns to gene expression. *Cell* 117(6): 721–733.
- Kustatscher, G., M. Hothorn, C. Pugieux, K. Scheffzek, and A. G. Ladurner, 2005 Splicing regulates NAD metabolite binding to histone macroH2A. *Nat. Struct. Mol. Biol.* 12(7): 624–625.
- Laherty, C. D., W. M. Yang, J. M. Sun, J. R. Davie, E. Seto *et al.*, 1997 Histone deacetylases associated with the mSin3 corepressor mediate mad transcriptional repression. *Cell* 89(3): 349–356.
- Landry, J., A. Sutton, S. T. Tafrov, R. C. Heller, J. Stebbins *et al.*, 2000 The silencing protein SIR2 and its homologs are NAD-dependent protein deacetylases. *Proc. Natl. Acad. Sci. USA* 97(11): 5807–5811.
- Laurenson, P., and J. Rine, 1991 SUM1-1: a suppressor of silencing defects in *Saccharomyces cerevisiae*. *Genetics* 129: 685–696.
- Lechner, T., M. J. Carrozza, Y. Yu, P. A. Grant, A. Eberharter *et al.*, 2000 Sds3 (suppressor of defective silencing 3) is an integral component of the yeast Sin3[middle dot]Rpd3 histone deacetylase complex and is required for histone deacetylase activity. *J. Biol. Chem.* 275(52): 40961–40966.
- Lee, S., L. Tong, and J. M. Denu, 2008 Quantification of endogenous sirtuin metabolite O-acetyl-ADP-ribose. *Anal. Biochem.* 383(2): 174–179.
- Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25(14): 1754–1760.
- Li, Y. C., T. H. Cheng, and M. R. Gartenberg, 2001 Establishment of transcriptional silencing in the absence of DNA replication. *Science* 291(5504): 650–653.
- Lin, Y. Y., J. Y. Lu, J. Zhang, W. Walter, W. Dang *et al.*, 2009 Protein acetylation microarray reveals that NuA4 controls key metabolic target regulating gluconeogenesis. *Cell* 136(6): 1073–1084.
- Liou, G. G., J. C. Tanny, R. G. Kruger, T. Walz, and D. Moazed, 2005 Assembly of the SIR complex and its regulation by O-acetyl-ADP-ribose, a product of NAD-dependent histone deacetylation. *Cell* 121(4): 515–527.
- Loewith, R., J. S. Smith, M. Meijer, T. J. Williams, N. Bachman *et al.*, 2001 Pho23 is associated with the Rpd3 histone deacetylase and is required for its normal function in regulation of gene expression and silencing in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 276(26): 24068–24074.
- Loney, E. R., P. W. Inglis, S. Sharp, F. E. Pryde, N. A. Kent *et al.*, 2009 Repressive and non-repressive chromatin at native telomeres in *Saccharomyces cerevisiae*. *Epigenetics Chromatin* 2(1): 18.
- Longtine, M. S., A. McKenzie, 3rd, D. J. Demarini, N. G. Shah, A. Wach *et al.*, 1998 Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* 14(10): 953–961.
- Loo, S., and J. Rine, 1994 Silencers and domains of generalized repression. *Science* 264(5166): 1768–1771.
- Mallory, M. J., M. J. Law, L. E. Buckingham, and R. Strich, 2010 The Sin3p PAH domains provide separate functions repressing meiotic gene transcription in *Saccharomyces cerevisiae*. *Eukaryot. Cell* 9(12): 1835–1844.
- Martino, F., S. Kueng, P. Robinson, M. Tsai-Pflugfelder, F. van Leeuwen *et al.*, 2009 Reconstitution of yeast silent chromatin: multiple contact sites and O-AADPR binding load SIR complexes onto nucleosomes in vitro. *Mol. Cell* 33(3): 323–334.
- Meijsing, S. H., and A. E. Ehrenhofer-Murray, 2001 The silencing complex SAS-I links histone acetylation to the assembly of repressed chromatin by CAF-I and Asf1 in *Saccharomyces cerevisiae*. *Genes Dev.* 15(23): 3169–3182.
- Miller, A. M., and K. A. Nasmyth, 1984 Role of DNA replication in the repression of silent mating type loci in yeast. *Nature* 312(5991): 247–251.
- Oppikofer, M., S. Kueng, F. Martino, S. Soeroes, S. M. Hancock *et al.*, 2011 A dual role of H4K16 acetylation in the establishment of yeast silent chromatin. *EMBO J.* 30(13): 2610–2621.
- Osada, S., A. Sutton, N. Muster, C. E. Brown, J. R. Yates, 3rd *et al.*, 2001 The yeast SAS (something about silencing) protein complex contains a MYST-type putative acetyltransferase and functions with chromatin assembly factor ASF1. *Genes Dev.* 15(23): 3155–3168.
- Ozaydin, B., and J. Rine, 2010 Expanded roles of the origin recognition complex in the architecture and function of silenced chromatin in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 30(3): 626–639.
- Park, E. C., and J. W. Szostak, 1990 Point mutations in the yeast histone H4 gene prevent silencing of the silent mating type locus HML. *Mol. Cell. Biol.* 10(9): 4932–4934.
- Ravindra, A., K. Weiss, and R. T. Simpson, 1999 High-resolution structural analysis of chromatin at specific loci: *Saccharomyces cerevisiae* silent mating-type locus HMRA. *Mol. Cell. Biol.* 19(12): 7944–7950.
- Renauld, H., O. M. Aparicio, P. D. Zierath, B. L. Billington, S. K. Chhablani *et al.*, 1993 Silent domains are assembled continuously from the telomere and are defined by promoter distance and strength, and by SIR3 dosage. *Genes Dev.* 7(7A): 1133–1145.
- Robyr, D., Y. Suka, I. Xenarios, S. K. Kurdistani, A. Wang *et al.*, 2002 Microarray deacetylation maps determine genome-wide functions for yeast histone deacetylases. *Cell* 109(4): 437–446.
- Rossmann, M. P., W. Luo, O. Tsaponina, A. Chabes, and B. Stillman, 2011 A common telomeric gene silencing assay is affected by nucleotide metabolism. *Mol. Cell* 42(1): 127–136.
- Rundlett, S. E., A. A. Carmen, R. Kobayashi, S. Bavykin, B. M. Turner *et al.*, 1996 HDA1 and RPD3 are members of distinct yeast histone deacetylase complexes that regulate silencing and transcription. *Proc. Natl. Acad. Sci. USA* 93(25): 14503–14508.
- Rusché, L. N., and J. Rine, 2001 Conversion of a gene-specific repressor to a regional silencer. *Genes Dev.* 15(8): 955–967.
- Rusché, L. N., A. L. Kirchmaier, and J. Rine, 2002 Ordered nucleation and spreading of silenced chromatin in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 13(7): 2207–2222.
- Rusché, L. N., A. L. Kirchmaier, and J. Rine, 2003 The establishment, inheritance, and function of silenced chromatin in

- Saccharomyces cerevisiae*. Annu. Rev. Biochem. 72: 481–516.
- Sauve, A. A., C. Wolberger, V. L. Schramm, and J. D. Boeke, 2006 The biochemistry of sirtuins. Annu. Rev. Biochem. 75: 435–465.
- Steakley, D. L., and J. Rine, 2015 On the mechanism of gene silencing in *Saccharomyces cerevisiae*. G3 (Bethesda) 5(8): 1751–1763.
- Suka, N., Y. Suka, A. A. Carmen, J. Wu, and M. Grunstein, 2001 Highly specific antibodies determine histone acetylation site usage in yeast heterochromatin and euchromatin. Mol. Cell 8(2): 473–479.
- Suka, N., K. Luo, and M. Grunstein, 2002 Sir2p and Sas2p oppositely regulate acetylation of yeast histone H4 lysine16 and spreading of heterochromatin. Nat. Genet. 32(3): 378–383.
- Sun, Z. W., and M. Hampsey, 1999 A general requirement for the Sin3-Rpd3 histone deacetylase complex in regulating silencing in *Saccharomyces cerevisiae*. Genetics 152: 921–932.
- Suter, B., O. Pogoutse, X. Guo, N. Krogan, P. Lewis *et al.*, 2007 Association with the origin recognition complex suggests a novel role for histone acetyltransferase Hat1p/Hat2p. BMC Biol. 5: 38.
- Takahashi, Y.-H., J. M. Schulze, J. Jackson, T. Hentrich, C. Seidel *et al.*, 2011 Dot1 and histone H3K79 methylation in natural telomeric and HM silencing. Mol. Cell 42(1): 118–126.
- Tanner, K. G., J. Landry, R. Sternglanz, and J. M. Denu, 2000 Silent information regulator 2 family of NAD- dependent histone/protein deacetylases generates a unique product, 1-O-acetyl-ADP-ribose. Proc. Natl. Acad. Sci. USA 97(26): 14178–14182.
- Thorvaldsdóttir, H., J. T. Robinson, and J. P. Mesirov, 2013 Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Brief. Bioinform. 14(2): 178–192.
- Thurtle, D. M., and J. Rine, 2014 The molecular topography of silenced chromatin in *Saccharomyces cerevisiae*. Genes Dev. 28(3): 245–258.
- Tong, L., S. Lee, and J. M. Denu, 2009 Hydrolase regulates NAD+ metabolites and modulates cellular redox. J. Biol. Chem. 284(17): 11256–11266.
- Vannier, D., D. Balderes, and D. Shore, 1996 Evidence that the transcriptional regulators SIN3 and RPD3, and a novel gene (SDS3) with similar functions, are involved in transcriptional silencing in *S. cerevisiae*. Genetics 144: 1343–1353.
- Wang, F., G. Li, M. Altaf, C. Lu, M. A. Currie *et al.*, 2013 Heterochromatin protein Sir3 induces contacts between the amino terminus of histone H4 and nucleosomal DNA. Proc. Natl. Acad. Sci. USA 110(21): 8495–8500.
- Wang, H., and D. J. Stillman, 1993 Transcriptional repression in *Saccharomyces cerevisiae* by a SIN3-LexA fusion protein. Mol. Cell. Biol. 13(3): 1805–1814.
- Wang, X., G. Bryant, A. Zhao, and M. Ptashne, 2015 Nucleosome avidities and transcriptional silencing in yeast. Curr. Biol. 25(9): 1215–1220.
- Washburn, B. K., and R. E. Esposito, 2001 Identification of the Sin3-binding site in Ume6 defines a two-step process for conversion of Ume6 from a transcriptional repressor to an activator in yeast. Mol. Cell. Biol. 21(6): 2057–2069.
- Weiss, K., and R. T. Simpson, 1998 High-resolution structural analysis of chromatin at specific loci: *Saccharomyces cerevisiae* silent mating type locus HMLalpha. Mol. Cell. Biol. 18(9): 5392–5403.
- Xu, F., Q. Zhang, K. Zhang, W. Xie, and M. Grunstein, 2007 Sir2 deacetylates histone H3 lysine 56 to regulate telomeric heterochromatin structure in yeast. Mol. Cell 27(6): 890–900.
- Yang, B., and A. L. Kirchmaier, 2006 Bypassing the catalytic activity of SIR2 for SIR protein spreading in *Saccharomyces cerevisiae*. Mol. Biol. Cell 17(12): 5287–5297.
- Yang, B., A. Miller, and A. L. Kirchmaier, 2008 HST3/HST4-dependent deacetylation of lysine 56 of histone H3 in silent chromatin. Mol. Biol. Cell 19(11): 4993–5005.
- Yoshida, K., J. Bacal, D. Desmarais, I. Padioleau, O. Tsaponina *et al.*, 2014 The histone deacetylases sir2 and rpd3 act on ribosomal DNA to control the replication program in budding yeast. Mol. Cell 54(4): 691–697.
- Zhang, Z., M. K. Hayashi, O. Merkel, B. Stillman, and R.-M. Xu, 2002 Structure and function of the BAH-containing domain of Orc1p in epigenetic silencing. EMBO J. 21(17): 4600–4611.
- Zhou, J., B. O. Zhou, B. A. Lenzmeier, and J.-Q. Zhou, 2009 Histone deacetylase Rpd3 antagonizes Sir2-dependent silent chromatin propagation. Nucleic Acids Res. 37(11): 3699–3713.
- Zill, O. A., D. Scannell, L. Teytelman, and J. Rine, 2010 Co-evolution of transcriptional silencing proteins and the DNA elements specifying their assembly. PLoS Biol. 8(11): e1000550.

Communicating editor: M. Hampsey

GENETICS

Supporting Information

<http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.116.190835/-/DC1>

Histone Deacetylases with Antagonistic Roles in *Saccharomyces cerevisiae* Heterochromatin Formation

Deborah M. Thurtle-Schmidt, Anne E. Dodson, and Jasper Rine

A. **B.**

Figure S1. *RPD3* point mutations as identified by pooled linkage analysis. Sequencing reads from the JRY9569 mutant pool and JRY9569 wildtype pools (A) and JRY9570 mutant and wildtype pools (B) were visualized with IGV (THORVALDSDÓTTIR et al. 2013) at *RPD3* (YNL330C). Red indicates forward reads and blue indicates reverse reads and the grey bar above each base pair position indicates coverage at that position. Below is the wild-type sequence for the region of *RPD3* with the relevant reading frame indicated in blue below. (A) 14/14 reads in the JRY9569 mutant pool revealed a C to T mutation resulting in a proline to histidine mutation. (B) 18/18 reads in the JRY9570 mutant pool showed a C to T mutation resulting in a Glycine to Asparagine coding mutation. Neither wild-type pools showed any reads with mutations in *RPD3*.

Figure S2. *ORC3* point mutation in JRY9570 as identified by pooled linkage analysis. Sequencing reads from the JRY9570 mutant and wild-type pools were visualized with IGV (THORVALDSDÓTTIR et al. 2013) at *ORC3* (YLL004W). Red indicates forward reads and blue indicates reverse reads and the grey bar above each base pair position indicates coverage at that position. Below is the wild-type *SacCer2* sequence for the region of *ORC3* with the relevant reading frame indicated in blue below. 46/49 reads (not all shown) in the JRY9570 mutant pool showed a C to T mutation resulting in an Alanine to Valine coding mutation. The wild-type pool showed only 4 reads of 24 with the same C to T mutation. Additionally, the sequencing revealed a synonymous G to A mutation 7 basepairs downstream of the identified mutation present in our lab W303 strain as compared to the *SacCer2* genome.

Figure S3. Expression of *SIR3* and *Sir4* in *rpd3* mutants and *rpd3P264L hst3Δ* double mutant. RNA was normalized to *ACT1* for each sample and reported as relative to the *RPD3* sample. Error bars are the standard error of 4 biological replicates. The asterisks indicate a p-value less than .05 (student's t-test).

Figure S4. Expression of *COS8* in *rpd3* mutants. RNA was normalized to *ACT1* for each sample and reported as relative to the *RPD3* sample. Error bars are the standard error of 4 biological replicates. No significant differences were found (student's t-test).

Table S1. Strains used in this study.

Name	Genotype	Source
JRY2726	<i>MATa his4</i>	Rine Lab Collection
JRY2728	<i>MATα his4</i>	Rine Lab Collection
JRY9097	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A</i>	This Study
JRY9567	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, SUM1-1</i> *derived from EMS mutagenesis	This Study
JRY9568	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L</i> *derived from EMS mutagenesis	This Study
JRY9569	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P397H</i> *derived from EMS mutagenesis	This Study
JRY9570	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3G290N, orc3A333V</i> *derived from EMS mutagenesis	This Study
JRY9572	<i>MATα, ade2-1, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A</i>	This Study
JRY9573	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3Δ::KanMX</i>	This Study
JRY9574	<i>MATα, ade2-1, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, , rpd3Δ::KanMX</i>	This Study
JRY9575	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, pho23Δ::NatMX</i>	This Study
JRY10630	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, sds3Δ::KanMX</i>	This Study
JRY10631	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, ash1Δ::KanMX</i>	This Study
JRY9576	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, sap30Δ::NatMX</i>	This Study
JRY9577	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, eaf3Δ::NatMX</i>	This Study
JRY9578	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rco1Δ::NatMX</i>	This Study
JRY9579	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, sin3Δ::NatMX</i>	This Study
JRY9580	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L, sir3Δ::NatMX</i>	This Study
JRY9581	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L, sir3Δ::NatMX</i>	This Study
JRY9582	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3Δ::KanMX, sir3Δ::NatMX</i>	This Study
JRY9583	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3Δ::KanMX, sir3Δ::NatMX</i>	This Study

JRY9584	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>sir4Δ::NatMX</i>	This Study
JRY9585	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>Sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>sir4Δ::NatMX</i>	This Study
JRY9586	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>Sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i> , <i>sir4Δ::NatMX</i>	This Study
JRY9587	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i> , <i>sir4Δ::NatMX</i>	This Study
JRY9588	<i>MATα</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>leu2-3,112</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9589	<i>MATα</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>leu2-3,112</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9590	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hst1Δ::KanMX</i>	This Study
JRY9591	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hst2Δ::KanMX</i>	This Study
JRY9592	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hst3Δ::KanMX</i>	This Study
JRY9593	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hst4Δ::KanMX</i>	This Study
JRY9594	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>hst1Δ::KanMX</i>	This Study
JRY9595	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>hst2Δ::KanMX</i>	This Study
JRY9596	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>hst3Δ::KanMX</i>	This Study
JRY9597	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>hst4Δ::KanMX</i>	This Study
JRY9598	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i>	This Study
JRY9599	<i>MATa</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i>	This Study
JRY9600	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::Hyg</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9601	<i>MATα</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9602	<i>MATa</i> , <i>lys2-</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9603	<i>MATα</i> , <i>lys2-</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9604	<i>MATα</i> , <i>sas2Δ::HygMX</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hht1-bhf1Δ::HygMX</i> , <i>hht2-bhf2Δ::NatMX</i> , <i>pJR3212 [H3K56R-H4]</i>	This Study
JRY9605	<i>MATα</i> , <i>sas2Δ::HygMX</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hht1-bhf1Δ::HygMX</i> , <i>hht2-bhf2Δ::NatMX</i> , <i>pJR2759[H3-H4]</i>	This Study
JRY9606	<i>MATα</i> , <i>sas2Δ::HygMX</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hht1-bhf1Δ::HygMX</i> , <i>hht2-bhf2Δ::NatMX</i> , <i>pJR3213[H3K56Q-H4]</i>	This Study
JRY9607	<i>MATα</i> , <i>sas2Δ::HygMX</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hht1-</i>	This Study

	<i>hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR2760[H3-H4K5,8,12R]</i>	
JRY9608	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR2761[H3-H4K5,8,12,16R]</i>	This Study
JRY9609	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3212[H3K56R-H4]</i>	This Study
JRY9610	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR2759[H3-H4]</i>	This Study
JRY9611	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3213[H3K56Q-H4],</i>	This Study
JRY9612	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR2760[H3-H4K5,8,12R]</i>	This Study
JRY9613	<i>MATα, sas2Δ::Hyg, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR2761[H3-H4K5,8,12,16R]</i>	This Study
JRY9614	<i>MATα, ADE2, lys2-, sas2Δ::Hyg, hmrα1Δ::URA3, Sir2Δ::HIS3, LEU2::Sir2N345A, sum1-1 *double backcrossed</i>	This Study
JRY9615	<i>MATα, ADE2, lys2-, sas2Δ::Hyg, hmrα1Δ::URA3, Sir2Δ::HIS3, LEU2::Sir2N345A, sum1-1 *double backcrossed</i>	This Study
JRY9616	<i>MATα, ADE2, lys2-, sas2Δ::Hyg, hmrα1Δ::URA3, Sir2Δ::HIS3, LEU2::Sir2N345A, sum1-1 *double backcrossed</i>	This Study
JRY9617	<i>MATα/ MATα, ADE/ade2-1, LYS/lys2-, sas2Δ::Hyg/sas2Δ::Hyg, hmr-α1Δ::URA3/ hmr-α1Δ::URA3, sir2Δ::HIS3/sir2Δ::HIS3, Leu2::Sir2N345A/Leu2::Sir2N345A, rpd3G290N/RPD3, orc3A333V/ORC3</i>	This Study
JRY9618	<i>MATα/ MATα, ADE/ade2-1, LYS/lys2-, sas2Δ::Hyg/sas2Δ::Hyg, hmr-α1Δ::URA3/ hmr-α1Δ::URA3, sir2Δ::HIS3/sir2Δ::HIS3, Leu2::Sir2N345A/Leu2::Sir2N345A, rpd3P397H/RPD3</i>	This Study
JRY10632	<i>MATα ade2-1, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, SIR4-myc::KanMX</i>	This Study
JRY10633	<i>MATα ade2-1, sas2Δ::Hyg, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, SIR4-myc::KanMX, rpd3Δ::NatMX</i>	This Study
JRY10634	<i>MATα ade2-1, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3Δ::NatMX</i>	This Study
JRY10635	<i>MATα sas2Δ::HygMX, hmrα1Δ::URA3, LEU2::sir2N345A, SIR4-myc::KanMX</i>	This Study
JRY10636	<i>MATα lys2-, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rtt109Δ::NatMX</i>	This Study
JRY10637	<i>MATα lys2-, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L, rtt109Δ::Nat</i>	This Study
JRY10638	<i>MATα lys2-, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L, hst3Δ::KanMX, rtt109Δ::NatMX</i>	This Study
JRY10644	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3383 [H3K9R-H4K5,8,12,16R]</i>	This Study

JRY10645	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3384 [H3K9Q-H4K5,8,12,16R]</i>	This Study
JRY10646	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3385 [H3K14R-H4K5,8,12,16R]</i>	This Study
JRY10647	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3386 [H3K14Q-H4K5,8,12,16R]</i>	This Study
JRY10648	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3387 [H3K27R-H4K5,8,12,16R]</i>	This Study
JRY10649	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3388 [H3K27Q-H4K5,8,12,16R]</i>	This Study
JRY10650	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR3383 [H3K9R-H4K5,8,12,16R]</i>	This Study
JRY10651	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR3384 [H3K9Q-H4K5,8,12,16R]</i>	This Study
JRY10652	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR3385 [H3K14R-H4K5,8,12,16R]</i>	This Study
JRY10653	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR3386 [H3K14Q-H4K5,8,12,16R]</i>	This Study
JRY10654	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR3387 [H3K27R-H4K5,8,12,16R]</i>	This Study
JRY10655	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR3388 [H3K27Q-H4K5,8,12,16R]</i>	This Study
JRY10676	<i>MATα, ade2-1, sas2Δ::HygMX, hmra1Δ::URA3, LEU2::sir2N345A</i>	This Study
JRY10698	<i>MATα, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::KanMX, LEU2::sir2N345A, sin3Δ::NatMX</i>	This Study

Table S2. Plasmids used in this study

Plasmid	Backbone	Yeast marker	Bacterial marker	Insert	Source
pJR2759	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>HHT2-HHF2</i>	Rine Lab Collection
pJR2760	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2-hhf2(K5,8,12R)</i>	Rine Lab Collection
pJR2761	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2-hhf2(K5,8,12,16R)</i>	Rine Lab Collection
pJR3212	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K56R)-HHF2</i>	Rine Lab Collection
pJR3213	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K56Q)-HHF2</i>	Rine Lab Collection
pJR3383	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K9R)- hhf2(K5,8,12,16R)</i>	This Study
pJR3384	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K9Q)- hhf2(K5,8,12,16R)</i>	This Study
pJR3385	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K14R)- hhf2(K5,8,12,16R)</i>	This Study
pJR3386	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K14Q)- hhf2(K5,8,12,16R)</i>	This Study
pJR3387	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K27R)- hhf2(K5,8,12,16R)</i>	This Study
pJR3388	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K27Q)- hhf2(K5,8,12,16R)</i>	This Study
pJR734	pRS313	<i>HIS3</i>	<i>ampR</i>	<i>None</i>	Rine Lab Collection
M1635	YCp	<i>HIS3</i>	<i>ampR</i>	<i>SIN3</i>	(WANG and STILLMAN 1993)
M1636	YCp	<i>HIS3</i>	<i>ampR</i>	<i>sin3ΔPAH1</i>	(WANG and STILLMAN 1993)
M1637	YCp	<i>HIS3</i>	<i>ampR</i>	<i>sin3ΔPAH2</i>	(WANG and STILLMAN 1993)
M1638	YCp	<i>HIS3</i>	<i>ampR</i>	<i>sin3ΔPAH3</i>	(WANG and STILLMAN 1993)
M1639	YCp	<i>HIS3</i>	<i>ampR</i>	<i>sin3ΔPAH4</i>	(WANG and STILLMAN 1993)

Table S3. Number of sequencing reads mapped.

Experiment	Aligned Reads
JRY9097 Sequencing	24324128
JRY9614 Sequencing	24773456
JRY96915 Sequencing	29407529
JRY9616 Sequencing	22330041
JRY9567 Sequencing	31451777
JRY9569 Wildtype Pool	8405843
JRY9569 Mutant Pool	7714958
JRY9570 Wildtype Pool	10454114
JRY9570 Mutant Pool	12837055

Table S4. Primers used in this study.

Primer Name	Sequence	Use
<i>ACT1</i> Forward	TGTCCTTGTACTCTTCCGGT	RNA
<i>ACT1</i> Reverse	CCGGCCAAATCGATTCTCAA	RNA
<i>alpha2</i> Forward	TCCACAAATCACAGATGAGT	RNA /ChIP
<i>alpha2</i> Reverse	GTTGGCCCTAGATAAGAATCC	RNA /ChIP
a1 Forward	TGGATGATATTTGTAGTATGGCGGA	RNA
a1 Reverse	TCCCTTTGGGCTCTTCTCTT	RNA
<i>URA3</i> Forward	CTTCCAAGGGTTCTCTAGCACACG	RNA /ChIP
<i>URA3</i> Reverse	CTGTACTGCTGACCCAATGCATCG	RNA /ChIP
<i>SIR3</i> Forward	TCATATGCGGTGCCTGGGA	RNA
<i>SIR3</i> Reverse	TGTTGGTGGTCAAATGCAGTCC	RNA
<i>SIR4</i> Forward	TATCGTTTTGTCGAAAAGCAG	RNA
<i>SIR4</i> Reverse	CTTCGTTACTGGTCTTTTGT	RNA
<i>HML-E</i> Forward	AAAGGACGAGACACCAGAAGA	ChIP
<i>HML-E</i> Reverse	TCCAAATAAGTCCGTGCCGA	ChIP
<i>HMR-E</i> Forward	CGAACGATCCCCGTCCAAGTTATG	ChIP
<i>HMR-E</i> Reverse	TCGGAATCGAGAATCTTCGTAATGC	ChIP
<i>HMR-I</i> Forward	TCCTTTAGTTGGAATTTGCACAAG	ChIP
<i>HMR-I</i> Reverse	CAATCCTTGCATCCAGCTTTCTCT	ChIP

Table S1. Strains used in this study.

Name	Genotype	Source
JRY2726	<i>MATa his4</i>	Rine Lab Collection
JRY2728	<i>MATα his4</i>	Rine Lab Collection
JRY9097	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A</i>	This Study
JRY9567	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, SUM1-1</i> *derived from EMS mutagenesis	This Study
JRY9568	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L</i> *derived from EMS mutagenesis	This Study
JRY9569	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P397H</i> *derived from EMS mutagenesis	This Study
JRY9570	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3G290N, orc3A333V</i> *derived from EMS mutagenesis	This Study
JRY9572	<i>MATα, ade2-1, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A</i>	This Study
JRY9573	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3Δ::KanMX</i>	This Study
JRY9574	<i>MATα, ade2-1, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, , rpd3Δ::KanMX</i>	This Study
JRY9575	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, pho23Δ::NatMX</i>	This Study
JRY10630	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, sds3Δ::KanMX</i>	This Study
JRY10631	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, ash1Δ::KanMX</i>	This Study
JRY9576	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, sap30Δ::NatMX</i>	This Study
JRY9577	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, eaf3Δ::NatMX</i>	This Study
JRY9578	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rco1Δ::NatMX</i>	This Study
JRY9579	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, sin3Δ::NatMX</i>	This Study
JRY9580	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L, sir3Δ::NatMX</i>	This Study
JRY9581	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L, sir3Δ::NatMX</i>	This Study
JRY9582	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3Δ::KanMX, sir3Δ::NatMX</i>	This Study
JRY9583	<i>MATa, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3Δ::KanMX, sir3Δ::NatMX</i>	This Study

JRY9584	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>sir4Δ::NatMX</i>	This Study
JRY9585	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>Sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>sir4Δ::NatMX</i>	This Study
JRY9586	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>Sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i> , <i>sir4Δ::NatMX</i>	This Study
JRY9587	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i> , <i>sir4Δ::NatMX</i>	This Study
JRY9588	<i>MATα</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>leu2-3,112</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9589	<i>MATα</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>leu2-3,112</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9590	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hst1Δ::KanMX</i>	This Study
JRY9591	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hst2Δ::KanMX</i>	This Study
JRY9592	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hst3Δ::KanMX</i>	This Study
JRY9593	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hst4Δ::KanMX</i>	This Study
JRY9594	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>hst1Δ::KanMX</i>	This Study
JRY9595	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>hst2Δ::KanMX</i>	This Study
JRY9596	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>hst3Δ::KanMX</i>	This Study
JRY9597	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3P264L</i> , <i>hst4Δ::KanMX</i>	This Study
JRY9598	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i>	This Study
JRY9599	<i>MATa</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i>	This Study
JRY9600	<i>MATa</i> , <i>lys2-</i> , <i>sas2Δ::Hyg</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9601	<i>MATα</i> , <i>lys2-</i> , <i>sas2Δ::HygMX</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9602	<i>MATa</i> , <i>lys2-</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9603	<i>MATα</i> , <i>lys2-</i> , <i>hmra1Δ::URA3</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>rp3Δ::KanMX</i>	This Study
JRY9604	<i>MATα</i> , <i>sas2Δ::HygMX</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hht1-bhf1Δ::HygMX</i> , <i>hht2-bhf2Δ::NatMX</i> , <i>pJR3212 [H3K56R-H4]</i>	This Study
JRY9605	<i>MATα</i> , <i>sas2Δ::HygMX</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hht1-bhf1Δ::HygMX</i> , <i>hht2-bhf2Δ::NatMX</i> , <i>pJR2759[H3-H4]</i>	This Study
JRY9606	<i>MATα</i> , <i>sas2Δ::HygMX</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hht1-bhf1Δ::HygMX</i> , <i>hht2-bhf2Δ::NatMX</i> , <i>pJR3213[H3K56Q-H4]</i>	This Study
JRY9607	<i>MATα</i> , <i>sas2Δ::HygMX</i> , <i>sir2Δ::HIS3MX</i> , <i>LEU2::sir2N345A</i> , <i>hht1-</i>	This Study

	<i>hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR2760[H3-H4K5,8,12R]</i>	
JRY9608	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, pJR2761[H3-H4K5,8,12,16R]</i>	This Study
JRY9609	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3212[H3K56R-H4]</i>	This Study
JRY9610	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR2759[H3-H4]</i>	This Study
JRY9611	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3213[H3K56Q-H4],</i>	This Study
JRY9612	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR2760[H3-H4K5,8,12R]</i>	This Study
JRY9613	<i>MATα, sas2Δ::Hyg, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR2761[H3-H4K5,8,12,16R]</i>	This Study
JRY9614	<i>MATα, ADE2, lys2-, sas2Δ::Hyg, hmrα1Δ::URA3, Sir2Δ::HIS3, LEU2::Sir2N345A, sum1-1 *double backcrossed</i>	This Study
JRY9615	<i>MATα, ADE2, lys2-, sas2Δ::Hyg, hmrα1Δ::URA3, Sir2Δ::HIS3, LEU2::Sir2N345A, sum1-1 *double backcrossed</i>	This Study
JRY9616	<i>MATα, ADE2, lys2-, sas2Δ::Hyg, hmrα1Δ::URA3, Sir2Δ::HIS3, LEU2::Sir2N345A, sum1-1 *double backcrossed</i>	This Study
JRY9617	<i>MATα/ MATα, ADE/ade2-1, LYS/lys2-, sas2Δ::Hyg/sas2Δ::Hyg, hmr-α1Δ::URA3/ hmr-α1Δ::URA3, sir2Δ::HIS3/sir2Δ::HIS3, Leu2::Sir2N345A/Leu2::Sir2N345A, rpd3G290N/RPD3, orc3A333V/ORC3</i>	This Study
JRY9618	<i>MATα/ MATα, ADE/ade2-1, LYS/lys2-, sas2Δ::Hyg/sas2Δ::Hyg, hmr-α1Δ::URA3/ hmr-α1Δ::URA3, sir2Δ::HIS3/sir2Δ::HIS3, Leu2::Sir2N345A/Leu2::Sir2N345A, rpd3P397H/RPD3</i>	This Study
JRY10632	<i>MATα ade2-1, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, SIR4-myc::KanMX</i>	This Study
JRY10633	<i>MATα ade2-1, sas2Δ::Hyg, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, SIR4-myc::KanMX, rpd3Δ::NatMX</i>	This Study
JRY10634	<i>MATα ade2-1, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3Δ::NatMX</i>	This Study
JRY10635	<i>MATα sas2Δ::HygMX, hmrα1Δ::URA3, LEU2::sir2N345A, SIR4-myc::KanMX</i>	This Study
JRY10636	<i>MATα lys2-, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rtt109Δ::NatMX</i>	This Study
JRY10637	<i>MATα lys2-, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L, rtt109Δ::Nat</i>	This Study
JRY10638	<i>MATα lys2-, sas2Δ::HygMX, hmrα1Δ::URA3, sir2Δ::HIS3MX, LEU2::sir2N345A, rpd3P264L, hst3Δ::KanMX, rtt109Δ::NatMX</i>	This Study
JRY10644	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-hhf1Δ::HygMX, hht2-hhf2Δ::NatMX, rpd3Δ::KanMX, pJR3383 [H3K9R-H4K5,8,12,16R]</i>	This Study

JRY10645	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, rpd3Δ::KanMX, pJR3384 [H3K9Q-H4K5,8,12,16R]</i>	This Study
JRY10646	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, rpd3Δ::KanMX, pJR3385 [H3K14R-H4K5,8,12,16R]</i>	This Study
JRY10647	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, rpd3Δ::KanMX, pJR3386 [H3K14Q-H4K5,8,12,16R]</i>	This Study
JRY10648	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, rpd3Δ::KanMX, pJR3387 [H3K27R-H4K5,8,12,16R]</i>	This Study
JRY10649	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, rpd3Δ::KanMX, pJR3388 [H3K27Q-H4K5,8,12,16R]</i>	This Study
JRY10650	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, pJR3383 [H3K9R-H4K5,8,12,16R]</i>	This Study
JRY10651	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, pJR3384 [H3K9Q-H4K5,8,12,16R]</i>	This Study
JRY10652	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, pJR3385 [H3K14R-H4K5,8,12,16R]</i>	This Study
JRY10653	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, pJR3386 [H3K14Q-H4K5,8,12,16R]</i>	This Study
JRY10654	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, pJR3387 [H3K27R-H4K5,8,12,16R]</i>	This Study
JRY10655	<i>MATα, sas2Δ::HygMX, sir2Δ::HIS3MX, LEU2::sir2N345A, hht1-bhf1Δ::HygMX, hht2-bhf2Δ::NatMX, pJR3388 [H3K27Q-H4K5,8,12,16R]</i>	This Study
JRY10676	<i>MATα, ade2-1, sas2Δ::HygMX, hmra1Δ::URA3, LEU2::sir2N345A</i>	This Study
JRY10698	<i>MATα, lys2-, sas2Δ::HygMX, hmra1Δ::URA3, sir2Δ::KanMX, LEU2::sir2N345A, sin3Δ::NatMX</i>	This Study

Table S2. Plasmids used in this study

Plasmid	Backbone	Yeast marker	Bacterial marker	Insert	Source
pJR2759	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>HHT2-HHF2</i>	Rine Lab Collection
pJR2760	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2-hhf2(K5,8,12R)</i>	Rine Lab Collection
pJR2761	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2-hhf2(K5,8,12,16R)</i>	Rine Lab Collection
pJR3212	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K56R)-HHF2</i>	Rine Lab Collection
pJR3213	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K56Q)-HHF2</i>	Rine Lab Collection
pJR3383	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K9R)- hhf2(K5,8,12,16R)</i>	This Study
pJR3384	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K9Q)- hhf2(K5,8,12,16R)</i>	This Study
pJR3385	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K14R)- hhf2(K5,8,12,16R)</i>	This Study
pJR3386	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K14Q)- hhf2(K5,8,12,16R)</i>	This Study
pJR3387	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K27R)- hhf2(K5,8,12,16R)</i>	This Study
pJR3388	pRS314	<i>TRP1</i>	<i>ampR</i>	<i>hht2(K27Q)- hhf2(K5,8,12,16R)</i>	This Study
pJR734	pRS313	<i>HIS3</i>	<i>ampR</i>	<i>None</i>	Rine Lab Collection
M1635	YCp	<i>HIS3</i>	<i>ampR</i>	<i>SIN3</i>	(WANG and STILLMAN 1993)
M1636	YCp	<i>HIS3</i>	<i>ampR</i>	<i>sin3ΔPAH1</i>	(WANG and STILLMAN 1993)
M1637	YCp	<i>HIS3</i>	<i>ampR</i>	<i>sin3ΔPAH2</i>	(WANG and STILLMAN 1993)
M1638	YCp	<i>HIS3</i>	<i>ampR</i>	<i>sin3ΔPAH3</i>	(WANG and STILLMAN 1993)
M1639	YCp	<i>HIS3</i>	<i>ampR</i>	<i>sin3ΔPAH4</i>	(WANG and STILLMAN 1993)

Table S3. Number of sequencing reads mapped.

Experiment	Aligned Reads
JRY9097 Sequencing	24324128
JRY9614 Sequencing	24773456
JRY96915 Sequencing	29407529
JRY9616 Sequencing	22330041
JRY9567 Sequencing	31451777
JRY9569 Wildtype Pool	8405843
JRY9569 Mutant Pool	7714958
JRY9570 Wildtype Pool	10454114
JRY9570 Mutant Pool	12837055

Table S4. Primers used in this study.

Primer Name	Sequence	Use
<i>ACT1</i> Forward	TGTCCTTGTACTCTTCCGGT	RNA
<i>ACT1</i> Reverse	CCGGCCAAATCGATTCTCAA	RNA
<i>alpha2</i> Forward	TCCACAAATCACAGATGAGT	RNA /ChIP
<i>alpha2</i> Reverse	GTTGGCCCTAGATAAGAATCC	RNA /ChIP
a1 Forward	TGGATGATATTTGTAGTATGGCGGA	RNA
a1 Reverse	TCCCTTTGGGCTCTTCTCTT	RNA
<i>URA3</i> Forward	CTTCCAAGGGTTCTCTAGCACACG	RNA /ChIP
<i>URA3</i> Reverse	CTGTACTGCTGACCCAATGCATCG	RNA /ChIP
<i>SIR3</i> Forward	TCATATGCGGTGCCTGGGA	RNA
<i>SIR3</i> Reverse	TGTTGGTGGTCAAATGCAGTCC	RNA
<i>SIR4</i> Forward	TATCGTTTTGTCGAAAAGCAG	RNA
<i>SIR4</i> Reverse	CTTCGTTACTGGTCTTTTGT	RNA
<i>HML-E</i> Forward	AAAGGACGAGACACCAGAAGA	ChIP
<i>HML-E</i> Reverse	TCCAAATAAGTCCGTGCCGA	ChIP
<i>HMR-E</i> Forward	CGAACGATCCCCGTCCAAGTTATG	ChIP
<i>HMR-E</i> Reverse	TCGGAATCGAGAATCTTCGTAATGC	ChIP
<i>HMR-I</i> Forward	TCCTTTAGTTGGAATTTGCACAAG	ChIP
<i>HMR-I</i> Reverse	CAATCCTTGCATCCAGCTTTCTCT	ChIP