Published in final edited form as:

Epilepsia. 2016 September ; 57(9): 1406–1415. doi:10.1111/epi.13480.

Midazolam-Ketamine Dual Therapy Stops Cholinergic Status Epilepticus And Reduces Morris Water Maze Deficits

Jerome Niquet^{1,2,*}, Roger Baldwin², Keith Norman², Lucie Suchomelova², Lucille Lumley³, and Claude G. Wasterlain^{1,2,4}

¹Department of Neurology, David Geffen School of Medicine at UCLA, Los Angeles, CA

²Epilepsy Research Laboratory (151), Veterans Affairs Greater Los Angeles Healthcare System, Los Angeles, CA

³US Army Medical Research Institute of Chemical Defense (USAMRICD), 2900 Ricketts Point Rd., Aberdeen Proving Ground, MD 21010, USA

⁴Brain Research Institute, David Geffen School of Medicine at UCLA, Los Angeles, CA.

SUMMARY

Objective—Pharmacoresistance remains an unsolved therapeutic challenge in status epilepticus (SE) and in cholinergic SE induced by nerve agent intoxication. SE triggers a rapid internalization of synaptic GABA_A receptors and externalization of NMDA receptors that may explain the loss of potency of standard anti-epileptic drugs (AED). We hypothesized that a drug combination aimed at correcting the consequences of receptor trafficking would reduce SE severity and its long-term consequences.

Methods—A severe model of SE was induced in adult Sprague-Dawley rats with a high dose of lithium and pilocarpine. The GABA_A receptor agonist midazolam, the NMDA receptor antagonist ketamine and/or the AED valproate were injected 40 min after SE onset in combination or as monotherapy. Measures of SE severity were the primary outcome. Secondary outcomes were acute neuronal injury, spontaneous recurrent seizures (SRS), and Morris water maze (MWM) deficits.

Results—Midazolam-ketamine dual therapy was more efficient than double-dose midazolam or ketamine monotherapy or than valproate-midazolam or valproate-ketamine dual therapy in reducing several parameters of SE severity, suggesting a synergistic mechanism. In addition, midazolam-ketamine dual therapy reduced SE-induced acute neuronal injury, epileptogenesis and MWM deficits.

Significance—This study showed that a treatment aimed at correcting maladaptive GABA_A receptor and NMDA receptor trafficking can stop SE and reduce its long-term consequences. Early

*Corresponding author: Jerome Niquet, PhD, Epilepsy Research Laboratory, VA Greater Los Angeles Healthcare System, 11301 Wilshire Boulevard, Building 114, Room 139, West Los Angeles, CA 90073. Phone number: 310-268-3595; jniquet@ucla.edu.

Sentence line describing the portrait (J. Niquet): Associate Researcher at the Department of Neurology, David Geffen School of Medicine at UCLA

Disclosure of Conflicts of Interest

None of the authors has any conflict of interest to disclose.

Midazolam-ketamine dual therapy may be superior to monotherapy in the treatment of benzodiazepine-refractory SE.

Keywords

refractory status epilepticus; cholinergic seizures; neuronal injury; hippocampal sclerosis; spatial memory; epileptogenesis

INTRODUCTION

The development of pharmacoresistance remains an unsolved therapeutic challenge in status epilepticus (SE), a serious neurological condition affecting 102-152,000 cases/year in the USA, with an estimated 22,000-42,000 deaths yearly,¹ and in SE induced by nerve agent intoxication.²⁻⁴ Benzodiazepine monotherapy, which is the standard treatment failed to stop seizures in 35–69% of cases of SE.⁵⁻⁸ Similarly, diazepam or midazolam monotherapy is the current antidote for the treatment of nerve agent-induced seizures.^{4; 9-11}

Studies in rodent models of SE may have elucidated the mechanism of pharmacoresistance. Seizures quickly become self-sustaining, independent of their initial cholinergic trigger, and become refractory to anticonvulsants.^{12; 13} Studies from our laboratory and others showed that SE causes internalization (and inactivation) of synaptic GABA_A receptors (GABA_AR),¹⁴⁻¹⁶ and promotes movement of NMDA receptor (NMDAR) subunits to synapses where they form additional functional receptors.¹⁷ These maladaptive changes due to receptor trafficking may explain the mechanism of pharmacoresistance to benzodiazepines. These drugs work well early in cholinergic SE because they have lots of GABA_AR to bind to, later their potency is reduced because of the loss of GABA_AR, and because they cannot counteract enhanced glutamatergic excitation. This might suggest that dual or triple therapy aimed at correcting the consequences of receptor trafficking may be more effective than monotherapy targeting only the remaining synaptic GABA_AR.² A combination of a GABA_AR agonist (diazepam) and an NMDAR antagonist (dizocilpine or ketamine) stopped SE in moderate models of cholinergic SE.^{18; 19} However, these studies did not investigate the long-term effects of SE (acute injury, epileptogenesis and behavior deficit). In the present study, we used a severe model of SE, induced by a high dose of lithium and pilocarpine, which was designed to mimic the effect of $1.6-2.0 \times LD_{50}$ soman exposure.²⁰ We examined the effect of midazolam-ketamine, valproate-ketamine or valproate-midazolam dual therapy on SE severity and compared them to double-dose midazolam, ketamine or valproate monotherapy. Consistent with the receptor trafficking hypothesis, we found that midazolam-ketamine dual therapy was the most potent combination and reduced several parameters of SE severity. In addition, this combination reduced SE-induced acute injury, prevented epileptogenesis and reduced spatial memory deficits.

METHODS

Animals

Male Sprague-Dawley rat (200-300g, mean 249g; Charles River, MA) were used. Rats were housed in a temperature- and humidity- controlled room with 12 h light-dark cycles (7 am-7 pm) and had free access to food and water. All experiments were conducted with the approval and in accordance with the regulations of the Institutional Animal care and Use Committee of West Los Angeles VA Medical Center.

Induction of SE, Monotherapies and Dual Therapies

Rats were administered lithium chloride (5 mEq/kg; #L-0505 Sigma, St. Louis MO, USA) subcutaneously and, 16 h later, SE was induced with i.p. pilocarpine hydrochloride (320 mg/kg; #P6503 Sigma). Only lithium/pilocarpine-treated rats displaying behavioral /EEG seizures were used. All rats received scopolamine methyl bromide (1mg/kg; i.p., #S8502; Sigma), a muscarinic antagonist that does not cross the blood-brain barrier, at the same time as pilocarpine, to decrease peripheral cholinergic effects such as pulmonary secretions. Seizures occurred 7.6 $\pm$ 2.7 min. after pilocarpine injection, so that time from pilocarpine injection to mono or dual therapy was approximately 48 min. All animals subsequently received scopolamine (10 mg/kg i.p.; #S1013; Sigma) to remove the original seizure trigger without stopping SE, and sham injection (control SE group), one drug (monotherapy) or a combination of two drugs (dual therapy) i.p. 40 min after EEG seizure onset to make sure that pharmacoresistance and self-sustaining seizures were well established. Drugs for monotherapy groups included midazolam (9 mg/kg; Caraco Pharmaceutical Laboratories Ltd), ketamine (90 mg/kg; #RL3760 Hospira) and sodium valproate (270 mg/kg; #P4543 Sigma). The valproate group was added as an additional “control” group because mortality rate in the control group was 100% at one week post-SE and valproate monotherapy did not measurably alter seizure activity. Dual therapy groups included combination of 4.5 mg/kg midazolam with 45 mg/kg ketamine, or 4.5 mg/kg midazolam with 135 mg/kg valproate, or 45 mg/kg ketamine with 135 mg/kg valproate. Note that monotherapy groups have double dose of drugs compared to dual therapy groups to compensate for the amount of drugs. For long-term behavioral studies, a sham group, which did not receive drug treatment and was not exposed to SE, was added.

Rats not capable of coordinated walking and movement 16 h after SE were injected SC (10ml/kg) with 5% glucose twice per day until capable of coordinated movement or until euthanasia at three days. Water moistened food pellets and or gelatin cubes were placed in the cage in Petri dishes. Euthanasia criteria consisted of failure to achieve coordinated movement three days after SE. Animals were euthanized if showing a weight loss of 5% sustained over two days after the coordination criterion had been achieved.

Implantation of electrodes

Under isoflurane anesthesia, the animals were implanted with stainless steel skull screws to serve as recording electrodes. Two electrodes were used for bipolar recording and were located 3mm anterior to lambda and 4 mm left and right of the medial suture. The third electrode served as reference and was located 1mm anterior to bregma and 1mm to the right

of the mid-line defined by the medial suture. The screw electrodes were connected to a tri-polar connector (Plastics One, VA) and dental cement was used to cover the electrodes so that only the connector was exposed. Following surgical implantation of electrodes, lidocaine was applied to the sutured area near the head cap and 0.05 mg/kg of buprenorphine administered subcutaneously. Buprenorphine treatment (0.05mg/kg S.C.) was continued twice a day for three days post surgery. Animals were used one to two weeks after electrode implantation. The BioPac Systems MP150 was used to record digital EEG using a BioPac UM100A preamplifier. Sampling rate was 200Hz.

Acute video-EEG monitoring

Recording for controls (10 mg/kg scopolamine alone; n=10), midazolam (9 mg/kg; n=10), ketamine (90 mg/kg; n=8), valproate (270 mg/kg; n=6), 4.5 mg/kg midazolam with 45 mg/kg ketamine (n=9), 4.5 mg/kg midazolam with 135 mg/kg valproate (n=10), or 45 mg/kg ketamine with 135 mg/kg valproate (n=11) groups was started before pilocarpine injection and was continuous for 24 h which included an initial pre-pilocarpine segment of EEG, the development of SE, drug treatment, and the overnight recovery period (Fig. 1 A). The EEGs were processed offline to detect seizures and spikes using Stellate Systems Harmony software (Natus) with default parameters: amplitude threshold 2.7, minimum frequency 3 Hz, maximum coefficient of variation 40% for seizure detection and a spike amplitude threshold of 6 for spike detection.

The outcome measures were the ratio of EEG power at T time divided by the average baseline EEG power (before pilocarpine); the number of seizures per 24 h, the cumulative seizure time per 24 h (time spent seizing, subtracting post- and inter-ictal time); the number of spikes per 24 h; the time spent in high amplitude ($> 2\times$ pre-pilocarpine baseline) EEG discharge per 24 h; and the time needed for EEG amplitude to fall for the first time below 2 times the prepilocarpine EEG amplitude and be free of semi-periodic spikes or sharp waves for at least one minute, which in this experimental paradigm is close to the time of termination of SE.

Tissue preparation for detection of acute neuronal injury

The animals were anesthetized with an overdose of pentobarbital (100 mg/kg i.p.) 48 h after induction of SE. Then, the animals underwent transcardiac perfusion with 4% phosphate-buffered formaldehyde (#P-6148 Sigma). Brains were kept in situ at 4 °C overnight, after which they were removed and postfixed in the same perfusate for 2-3 h. Subsequently, brains were kept in PB 0.1 M containing 30% sucrose for 48-72 h. Floating sections (30 μ M thickness) were obtained using a sliding microtome. Coronal sections were mounted, dried, incubated in potassium permanganate solution (0.06%; w/v) for 15 min, washed and incubated in fluoro-Jade B staining solution for 30 min. After 3 rinses, slides were dried overnight at room temperature, cleared three times in xylene and coverslipped with Permount medium. In the hilus of the dentate gyrus, the number of injured cells was counted by unbiased stereology using the optical disector method. The first series of one in five sections were stained with fluoro-Jade B, and the analysis was performed using a microscope (Olympus AX70) with a motorized stage connected to a computer running the Stereo Investigator software (MBF Bioscience). A counting frame of $45 \times 45 \mu$ m was

randomly positioned in a sampling grid of $70 \times 120 \mu\text{m}$. In the other areas (CA1, CA3, frontoparietal, entorhinal, and piriform cortices, thalamus, and amygdala), distribution of fluoro-jade B-positive cells was scored as follows: 0, no injury; 1: 1-30 positive cells per field; 2: 31-60 positive cells per field; 3: 61-100 positive cells per field; 4: more than 100 positive cells per field.

Recording of spontaneous recurrent seizures (SRS)

For chronic recordings, control (valproate 270 mg/kg; n=7), midazolam (9 mg/kg; n=10), ketamine (90 mg/kg, n=10) and 4.5 mg midazolam with 45 mg/kg ketamine groups (n=10) were implanted 4 weeks after SE and monitored 2 weeks later for 2 weeks (24/7) with a digital video-EEG system which saves 1 min of recording before and after each seizure (Figure 1A). Electrographic seizures were analyzed off-line and seizures were confirmed by manual review of the tracing morphology and of the digital videos.²¹ Outcome measure was the average number of SRS per week.

Morris water maze paradigm

Spatial learning and memory were evaluated one week after SRS recording with a modified Morris water maze paradigm, by requiring the rats to swim in a pool 170 cm in diameter, with the water kept at 20°C, to find a 12-cm diameter circular platform submerged 2 cm beneath the surface of the water, which was opacified by the addition of black non-toxic tempera paint. The platform was in a constant position during training, as there were a number of visual cues in the testing room. Experiments were monitored with a Sony CCD-IRIS high-resolution camera mounted above the pool and using indirect lighting from a 25 W bulb. A video-tracking system (Ethovision; Noldus, Inc. Wageningen, The Netherlands) was used for data acquisition. The rats were brought to the experimental room at least 30 min prior to an experiment. Each rat was trained to find the hidden platform kept in the same location for one session of eight trials per day, for 5 consecutive days. The start sequence was randomly selected and was different for each day. For training, the rat was released in the water from one of the four starting positions, facing the wall of the pool. It was given 60s to locate and climb onto the platform, where it stayed for 30s. If a rat did not find the platform within 60s, it was gently guided to it by experimenter. After the end of the last session, the rat was dried with absorbent paper and kept in a warm cage. Ten days later, acquisition test for long-term memory retention was performed.

Behavioral battery tests

Animals were submitted to a battery of behavioral tests slightly modified from others²² one day after the MWM. Four tests were used to discriminate animal hyperexcitability. In the *approach-response test*, a pen held vertically is moved toward the face of the animal. Responses were scored as follows: 1. No reaction, 2. The rat sniffs and turns toward the object, 3. The rat moves away from the object, 4. The rat jumps away from the object, 5. The rat jumps at or attacks the object. In the *touch-response test*, the animal is gently prodded in the rump with the blunt end of a pen. Responses were scored as follows: 1. No reaction, 2. The rat turns toward the object, 3. The rat moves away from the object, 4. The rat tries to avoid the object, running away, 5. The rat jumps with or without vocalization. In the *finger-snap test*, a finger snaps several inches above the head of the animal. Responses were

scored as follows: 1. No reaction, 2. Slight reaction (jerk), 3. Slight jump, 4. Violent jump. *In the pick-up test*, the animal is picked up by grasping him around the body. Responses were scored as follows: 1. Very easy, 2. Easy with vocalization, 3. Some difficulty, the rat rears and faces the hand, 4. Difficult, the rat avoids the hand and runs away, 5. Very difficult, the rat attacks the hand.

Statistical analyses

EEG and cell injury data showed a non-Gaussian distribution and were analyzed with nonparametric statistical methods: Kruskal-Wallis test followed by Dunn's multiple comparison test (GraphPad version 6). MWM data was analyzed by 2-way ANOVA (GraphPad version 6). Statistical significance was defined as $p < 0.05$. In all graphs, data are presented as median values with the interquartile range, which is the difference between the 75th and 25th percentile.

RESULTS

Dual therapy with midazolam-ketamine synergistically reduces the severity of SE

In the acute studies, EEGs were recorded for 24 h following pilocarpine injection (Figure 1B) and we examined the effect of mono and dual therapy on several parameters of SE severity. We examined the effect of monotherapies and dual therapies on the EEG power integral over the first hour posttreatment. Kruskal-Wallis analysis showed that ketamine monotherapy ($n = 8$) decreased EEG power compared to SE control ($n = 10$) or valproate monotherapy ($n = 10$; Fig. 1C). Among all groups, the combination midazolam-ketamine ($n = 9$) is the only treatment that decreased EEG power below pre-pilocarpine baseline. Midazolam-ketamine dual therapy showed a significant reduction compared to SE control and to double-dose midazolam ($n = 10$), ketamine, or valproate monotherapy. While the other two dual therapies were significantly different from SE control and valproate monotherapy, they were not significantly more potent than midazolam or ketamine monotherapy (Fig. 1C).

We examined the time needed for EEG amplitude to decline to twice the pre-seizure baseline. The midazolam-ketamine combination ($n = 9$) had the lowest time measured among all groups, which was significantly different from SE control and valproate monotherapy (Fig. 1D), suggesting that the midazolam-ketamine was the most potent combination tested.

High mortality was found in the first 24 h following pilocarpine in the SE control (5 of 10) and valproate (4 of 10) groups. Survival rate at 24 h post-SE was 100% in the midazolam ($n = 10$) and ketamine ($n = 8$) monotherapy groups and the three dual therapy groups ($n = 9-11$). Several other parameters of seizure severity were measured during the first 24 h following pilocarpine (table 1). The three dual therapies significantly reduced the EEG power integral over 6 h post-treatment and the number of computer-detected spikes/24 h when compared to SE control. Midazolam-ketamine was the only combination significantly reducing the time in spike/burst activity when compared to SE controls or to the valproate group (table 1).

Dual therapy with midazolam-ketamine reduces SE-induced acute neuronal injury—We examined the distribution of neuronal injury by fluoro-jade B staining in

animals perfused 48 h after pilocarpine injection. CA1 neuronal injury was found in 3 out of 5 animals from the SE control group, 4 out of 5 animals in the valproate group, 3 of 10 animals in the midazolam group, 1 of 7 animals in the ketamine group, and 1 of 9 animals from the midazolamketamine group. Semi-quantitative analysis showed that the dual therapy was the only treatment that significantly reduced CA1 neuronal injury compared to the SE control group. The ketamine and the dual therapy groups significantly reduced injury compared to the valproate group. CA3 neuronal injury was observed in 3 out of 5 animals from the SE control group, 4 out of 5 animals in the valproate group, and in 3 out of 10 animals in the midazolam group. The ketamine and dual therapy groups, which did not show neuronal injury in CA3, showed significant neuroprotection compared to the valproate group. Neuronal injury in the hilus was found in all animals studied. Unbiased stereological counting showed that ketamine monotherapy significantly reduced hilar injury compared to the SE control group. Monotherapy and dual therapy significantly reduced hilar injury compared to the valproate group (Fig. 2O). Semi-quantitative analysis showed that ketamine monotherapy and dual therapy significantly reduced neuronal injury in the frontoparietal, entorhinal and piriform cortices, and in thalamus and amygdala (table 2).

Dual therapy with midazolam-ketamine prevented SE-induced epileptogenesis

—We then examined whether midazolam-ketamine dual therapy has a long-term effect on the development of SRS. No rat in the SE control group survived long enough to be tested (most of them died in the first few days after SE). Rats which received 270 mg/kg of valproate, which increased long-term survival but did not affect the severity of SE (Fig. 1 and table 1), had a high incidence of SRS. Figure 3 shows that the midazolam + ketamine combination prevented epileptogenesis (Valproate: 4.6 ± 1 SRS per week, $n=7$; dual therapy: 0, $n=10$, $p<0.0001$). It did not differ significantly from the midazolam group, in spite of the fact that some midazolam-treated animals did develop SRS.

Dual therapy with midazolam-ketamine reduced MWM deficits—Performance in the MWM (Fig. 4A) was impaired in the valproate group ($n=7$) compared to sham controls ($n=8$). Again, no SE control survived long enough to be tested. The midazolam + ketamine group ($n=10$) performed better than the valproate group ($p<0.0001$), the midazolam group ($n=10$; $p<0.05$) and the ketamine group ($n=10$; $p<0.01$). In the retention test, the valproate group was significantly different from the dual therapy and sham groups (Fig. 4 B). Altogether, these observations showed that dual therapy reduced the MWM deficits.

In three other behavioral tests (finger snap, approach response and pick up), the dual therapy group was better than the valproate group but did not differ significantly from the sham, ketamine or midazolam groups (Fig. 4C).

DISCUSSION

The simultaneous trafficking of GABA_AR and NMDAR in opposite directions during repetitive seizures is a maladaptive change which increases excitation¹⁷ at the same time that it reduces inhibition.^{14; 16} Current standard therapy (benzodiazepine monotherapy) addresses only half the problem, and leaves runaway excitation untouched. Combined treatment with midazolam and ketamine tackles both types of receptor trafficking: it reduces

NMDAR-mediated excitation, and allosterically stimulates the remaining synaptic GABA_AR. This would be expected to reduce seizure activity, providing that enough GABA_AR are left in synapses at the time of treatment. The present study showed that the midazolam-ketamine combination indeed stops cholinergic SE and is more potent than double-dose midazolam or double-dose ketamine, suggesting that the effect of the two drugs is synergistic and not just additive, and confirming a previous study showing synergism between diazepam and ketamine.¹⁸ Furthermore, the midazolam-ketamine combination was more potent than the valproate-ketamine or valproate-midazolam combinations, showing that not all dual therapies show synergism, and suggesting that simultaneously targeting GABA_AR and NMDAR is a valid therapeutic strategy. This is in agreement with our hypothesis that correcting the consequences of GABA_AR and NMDAR trafficking would stop benzodiazepine-refractory SE, but does not prove that hypothesis since ketamine has several mechanisms of action in addition to NMDAR channel blockage.^{23; 24} Clinical and experimental studies suggest that ketamine administration can control refractory SE, and its combination with propofol is effective in controlling super-refractory SE in patients.²⁵⁻²⁷ A combination of diazepam and phenobarbital (with the anticholinergic scopolamine) has also been reported to stop refractory SE and to prevent CA1 damage in a lower dose lithiumpilocarpine model of SE.^{26; 28} The difference in models makes comparison with the current results difficult.

Midazolam-ketamine dual therapy eliminated SE-induced acute neuronal injury in CA1. This effect may be due to a reduction of SE severity and/or to the neuroprotective properties of midazolam and/or ketamine.²⁹ In fact, double-dose or even single-dose ketamine (data not shown) was as neuroprotective as dual therapy without stopping SE, confirming the well-known neuroprotective properties of ketamine, and suggesting an NMDAR-dependent mechanism of injury in that sector. The fact that neuroprotection was only partial in the hilus of the dentate gyrus may suggest that neuronal injury in that area is less dependent on NMDAR activation than in CA1. Previous studies have shown that the same seizures can cause neuronal injury by different mechanisms in CA1 and in the hilus.^{30; 31}

Dual therapy completely blocked epileptogenesis (Fig. 3), since no animal showed spontaneous recurrent seizures (SRSs). It was significantly better than ketamine or valproate treatment, but did not differ statistically from double-dose midazolam monotherapy. Midazolam or ketamine monotherapy reduced the frequency of SRSs compared to valproate-treated rats. Of course, this could simply reflect a reduction in SE severity. Several studies have shown that chronic treatment with valproate, gabapentin or lamotrigine reduced epileptogenesis and/or neuronal loss induced by SE.³²⁻³⁵ In our model, one treatment with dual therapy could reduce both acute injury and epileptogenesis.

Studies of spatial memory in the MWM showed major deficits in valproate-treated rats, which had little decrease in SE severity compared to vehicle-treated controls. Midazolamketamine dual therapy was superior to all three types of monotherapy in reducing MWM deficits, showing that combining a GABA_AR agonist and an NMDAR antagonist reduces not only the severity of SE but that of its long-term consequences as well.

Acknowledgements

This work was supported in part by Merit Review Award # I01 BX000273-07 from the United States (U.S.) Department of Veterans Affairs (Biomedical Laboratory Research and Development Service) and NINDS (grant U01 NS074926; CW). We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

REFERENCES

- DeLorenzo RJ, Hauser WA, Towne AR, et al. A prospective, population-based epidemiologic study of status epilepticus in Richmond, Virginia. *Neurology*. 1996; 46:1029–1035. [PubMed: 8780085]
- Chen JW, Wasterlain CG. Status epilepticus: pathophysiology and management in adults. *Lancet Neurol*. 2006; 5:246–256. [PubMed: 16488380]
- McDonough JH, McMonagle JD, Shih TM. Time-dependent reduction in the anticonvulsant effectiveness of diazepam against soman-induced seizures in guinea pigs. *Drug Chem Toxicol*. 2010; 33:279–283. [PubMed: 20429808]
- Shih T, McDonough JH Jr, Koplovitz I. Anticonvulsants for soman-induced seizure activity. *J Biomed Sci*. 1999; 6:86–96. [PubMed: 10087439]
- Treiman DM, Meyers PD, Walton NY, et al. A comparison of four treatments for generalized convulsive status epilepticus. Veterans Affairs Status Epilepticus Cooperative Study Group. *N Engl J Med*. 1998; 339:792–798. [PubMed: 9738086]
- Holtkamp M, Othman J, Buchheim K, et al. Predictors and prognosis of refractory status epilepticus treated in a neurological intensive care unit. *J Neurol Neurosurg Psychiatry*. 2005; 76:534–539. [PubMed: 15774441]
- Mayer SA, Claassen J, Lokin J, et al. Refractory status epilepticus: frequency, risk factors, and impact on outcome. *Arch Neurol*. 2002; 59:205–210. [PubMed: 11843690]
- Glauser T, Shinnar S, Gloss D, et al. Evidence-Based Guideline: Treatment of Convulsive Status Epilepticus in Children and Adults: Report of the Guideline Committee of the American Epilepsy Society. *Epilepsy Curr*. 2016; 16:48–61. [PubMed: 26900382]
- Capacio BR, Byers CE, Merk KA, et al. Pharmacokinetic studies of intramuscular midazolam in guinea pigs challenged with soman. *Drug Chem Toxicol*. 2004; 27:95–110. [PubMed: 15198070]
- Reddy SD, Reddy DS. Midazolam as an anticonvulsant antidote for organophosphate intoxication-- A pharmacotherapeutic appraisal. *Epilepsia*. 2015; 56:813–821. [PubMed: 26032507]
- McDonough JH Jr, McMonagle J, Copeland T, et al. Comparative evaluation of benzodiazepines for control of soman-induced seizures. *Arch Toxicol*. 1999; 73:473–478. [PubMed: 10650919]
- Kapur J, Macdonald RL. Rapid seizure-induced reduction of benzodiazepine and Zn²⁺ sensitivity of hippocampal dentate granule cell GABAA receptors. *J Neurosci*. 1997; 17:7532–7540. [PubMed: 9295398]
- Mazarati AM, Baldwin RA, Sankar R, et al. Time-dependent decrease in the effectiveness of antiepileptic drugs during the course of self-sustaining status epilepticus. *Brain Res*. 1998; 814:179–185. [PubMed: 9838100]
- Goodkin HP, Joshi S, Mtchedlishvili Z, et al. Subunit-specific trafficking of GABA(A) receptors during status epilepticus. *J Neurosci*. 2008; 28:2527–2538. [PubMed: 18322097]
- Goodkin HP, Yeh JL, Kapur J. Status epilepticus increases the intracellular accumulation of GABAA receptors. *J Neurosci*. 2005; 25:5511–5520. [PubMed: 15944379]
- Naylor DE, Liu H, Wasterlain CG. Trafficking of GABA(A) receptors, loss of inhibition, and a mechanism for pharmacoresistance in status epilepticus. *J Neurosci*. 2005; 25:7724–7733. [PubMed: 16120773]
- Naylor DE, Liu H, Niquet J, et al. Rapid surface accumulation of NMDA receptors increases glutamatergic excitation during status epilepticus. *Neurobiol Dis*. 2013; 54:225–238. [PubMed: 23313318]
- Martin BS, Kapur J. A combination of ketamine and diazepam synergistically controls refractory status epilepticus induced by cholinergic stimulation. *Epilepsia*. 2008; 49:248–255. [PubMed: 17941842]

19. Rice AC, DeLorenzo RJ. NMDA receptor activation during status epilepticus is required for the development of epilepsy. *Brain Res.* 1998; 782:240–247. [PubMed: 9519269]
20. Tetz LM, Rezk PE, Ratcliffe RH, et al. Development of a rat pilocarpine model of seizure/status epilepticus that mimics chemical warfare nerve agent exposure. *Toxicol Ind Health.* 2006; 22:255–266. [PubMed: 16924957]
21. Haas KZ, Sperber EF, Moshe SL. Kindling in developing animals: expression of severe seizures and enhanced development of bilateral foci. *Brain Res Dev Brain Res.* 1990; 56:275–280. [PubMed: 2261687]
22. Rice AC, Floyd CL, Lyeth BG, et al. Status epilepticus causes long-term NMDA receptor-dependent behavioral changes and cognitive deficits. *Epilepsia.* 1998; 39:1148–1157. [PubMed: 9821978]
23. Kohrs R, Durieux ME. Ketamine: teaching an old drug new tricks. *Anesth Analg.* 1998; 87:1186–1193. [PubMed: 9806706]
24. Zanos P, Moaddel R, Morris PJ, et al. NMDAR inhibition-independent antidepressant actions of ketamine metabolites. *Nature.* 2016
25. Sabharwal V, Ramsay E, Martinez R, et al. Propofol-ketamine combination therapy for effective control of super-refractory status epilepticus. *Epilepsy Behav.* 2015; 52:264–266. [PubMed: 26523340]
26. Löscher W. Single versus combinatorial therapies in status epilepticus: Novel data from preclinical models. *Epilepsy Behav.* 2015; 49:20–25. [PubMed: 25819944]
27. Gaspard N, Foreman B, Judd LM, et al. Intravenous ketamine for the treatment of refractory status epilepticus: a retrospective multicenter study. *Epilepsia.* 2013; 54:1498–1503. [PubMed: 23758557]
28. Brandt C, Töllner K, Klee R, et al. Effective termination of status epilepticus by rational polypharmacy in the lithium-pilocarpine model in rats: Window of opportunity to prevent epilepsy and prediction of epilepsy by biomarkers. *Neurobiol Dis.* 2015; 75:78–90. [PubMed: 25549873]
29. Fujikawa DG. Neuroprotective effect of ketamine administered after status epilepticus onset. *Epilepsia.* 1995; 36:186–195. [PubMed: 7821277]
30. Lopez-Meraz ML, Niquet J, Wasterlain CG. Distinct caspase pathways mediate necrosis and apoptosis in subpopulations of hippocampal neurons after status epilepticus. *Epilepsia.* 2010; 51(Suppl 3):56–60. [PubMed: 20618402]
31. Lopez-Meraz ML, Wasterlain CG, Rocha LL, et al. Vulnerability of postnatal hippocampal neurons to seizures varies regionally with their maturational stage. *Neurobiol Dis.* 2010; 37:394–402. [PubMed: 19879360]
32. Bolanos AR, Sarkisian M, Yang Y, et al. Comparison of valproate and phenobarbital treatment after status epilepticus in rats. *Neurology.* 1998; 51:41–48. [PubMed: 9674776]
33. Halonen T, Nissinen J, Pitkanen A. Effect of lamotrigine treatment on status epilepticus-induced neuronal damage and memory impairment in rat. *Epilepsy Res.* 2001; 46:205–223. [PubMed: 11518623]
34. Cilio MR, Bolanos AR, Liu Z, et al. Anticonvulsant action and long-term effects of gabapentin in the immature brain. *Neuropharmacology.* 2001; 40:139–147. [PubMed: 11077080]
35. Brandt C, Gastens AM, Sun M, et al. Treatment with valproate after status epilepticus: effect on neuronal damage, epileptogenesis, and behavioral alterations in rats. *Neuropharmacology.* 2006; 51:789–804. [PubMed: 16806297]

Key Point Box

- Pharmacoresistance is an unsolved therapeutic challenge in clinical SE and in nerve agent-induced SE.
- Several dual therapies were tested in a rat model of severe SE induced by a high amount of lithium and pilocarpine.
- Midazolam-ketamine dual therapy synergistically stopped SE and reduced several parameters of SE severity.
- This dual therapy reduced SE-induced acute neuronal injury, prevented epileptogenesis and reduced Morris water maze deficits.
- The midazolam-ketamine combination may be effective in treating benzodiazepine-refractory SE and its consequences.

A**Acute Studies** **Long-term studies** **B** **C** **D**

Figure 1. Midazolam-ketamine dual therapy is more efficient than double-dose midazolam or ketamine in reducing SE severity

A) Experimental flow: A severe form of status epilepticus (SE) was induced by administration of a high dose of lithium + scopolamine methyl bromide (scop. m. b), followed by an injection of a high dose of pilocarpine. Drug(s) or vehicle, and scopolamine (scop.) were injected 40 minutes after seizure onset. In acute studies, animals were implanted 1 week prior SE induction and sacrificed 48 h after SE onset to assess neuronal injury with fluoro-jade B staining. In long-term studies, animals were implanted 4 weeks

after SE and monitored for the detection of spontaneous seizures during 2 weeks. They were then studied in the Morris Water Maze and subjected to a battery of behavioral tests.

B) The left panels show the compressed EEG from SE control, midazolam, ketamine or midazolam-ketamine animals up to 75 min following treatment. The right panels show the magnified 6-sec EEG traces prior to SE or following SE (marked by vertical lines a-c). Vertical bar = 0.5 mV; horizontal bar = 1 sec.

C) This graph shows the ratio of EEG power integral over the first hour to initial EEG power at baseline, before pilocarpine injection. The midazolam-ketamine group (n=9), which lowered the EEG power below pre-pilocarpine baseline, is significantly different from SE control (n=10, **** p <0.0001), midazolam (n=10; # p <0.05), ketamine (n=8, α p <0.05) and valproate (n=10, *** p <0.0001) by Kruskal-Wallis, followed by Dunn's test. The ketamine group is significantly different from SE control, valproate and midazolam (α p <0.05; Kruskal-Wallis, followed by Dunn's test.).

D) This graph shows the time needed to reach an EEG amplitude of twice the pre-seizure baseline. The midazolam-ketamine group (n=9), which has the lowest time needed among all groups, is significantly different from SE control (n=10, *** p <0.001) and valproate (n=8, **** p <0.0001) by Kruskal-Wallis, followed by Dunn's test. The midazolam-valproate group (n=10) is significantly different from valproate (• p <0.05) by Kruskal-Wallis, followed by Dunn's test.

Figure 2. Reduction of neuronal injury in animals treated with midazolam-ketamine dual therapy or double-dose midazolam or ketamine

(A-L) These confocal images show fluoro-jade B staining in CA1 area (A-D), CA3 (E-H) and the hilus (I-L) 48 h following SE in SE control (A, E and I), midazolam (B, F and J), ketamine (C, G and K) and midazolam-ketamine (D, H and L) groups. Bars = 100 microns. (M-O) These graph shows the injury score (median values) in in CA1 (M) and CA3 (N), and the number of fluoro-jade B- positive cells counted by an unbiased stereological method in the hilus (O). Midazolam-ketamine dual therapy reduced neuronal injury compared to SE control in CA1. * $p < 0.05$; ** $p < 0.01$ by Kruskal-Wallis analysis followed by Dunn's test.

Figure 3. Midazolam-ketamine dual therapy prevents epileptogenesis

(A) The upper panel shows the compressed 80-sec EEG of a SRS. The lower panel shows the magnified 8-sec EEG traces marked by horizontal lines (a-f) (vertical bar = 0.5 mV; horizontal bar = 20 sec in upper panel and 2 sec in lower panel).

(B) Graph showing the number of SRS per week. Dual therapy midazolam-ketamine was more effective than double-dose ketamine in reducing the number of SRS. * $p < 0.05$, ** $p < 0.01$ and **** $p < 0.0001$ vs valproate (Kruskal-Wallis followed by Dunn's test).

Figure 4. Midazolam-ketamine dual therapy reduces behavioral deficits

(A-B) Performance in the Morris water maze. Graph A shows the latency to reach the hidden platform (y-axis) on each testing day (x-axis). Data are presented as mean $\pm$ SEM. * $p < 0.05$ vs Mz4.5+Ket45; ** $p < 0.01$ vs Mz4.5+Ket45 and **** $p < 0.0001$ vs Mz4.5+Ket45 by 2-way ANOVA. Graph B shows the latency during the retention test. ** $p < 0.01$ vs valproate 270 mg/kg (by Kruskal-Wallis followed by Dunn's test).

(C) Performance in the battery of behavioral tests. Data are presented as median values. ## $p < 0.01$ or ##### $p < 0.0001$ vs Mz4.5+Ket45 (Kruskal-Wallis followed by Dunn's test).

Table 1

Effect of monotherapies and dual therapies on several EEG measures recorded during the first 24 h following treatment.

		SE control (n=5-7)	Midazolam 9 mg/kg (n=10)	Ketamine 90 mg/kg (n=8)	Valproate 270 mg/kg (n=6-7)	Mz 4.5 + Ket 45 (n=9)	Mz 4.5 + Valp 135 (n=10)	Ket 45 + Valp 135 (n=11)
EEG power integral 6 h / pre-pilocarpine EEG power (%)	25% Percentile	986.0	834.8	830.0	1692	-60.50	367.0	2.000
	Median	2760	1440	1070	3641	479.2^{**}	638.0[•]	859.0[•]
Number of computer-detected spikes/24 hrs	75% Percentile	2998	1729	1227	3739	978.3	1136	1382
	25% Percentile	3495	1587	2152	3059	823.0	611.5	989.0
	Median	4031	2113	3305	5943	1255^{***}	1248^{**}	1378^{**}
	75% Percentile	7152	2958	3853	9958	1747	3874	1738
Time in high amplitude discharge (hrs)	25% Percentile	2.8	1.55	0.525	2.975	1.1	0.8103	0.4
	Median	3.5	4.45	1.4	5.7	1.7	1.663	0.875[#]
	75% Percentile	4.9	7.125	7.2	9.6	2.55	3.37	1.93
	25% Percentile	19.23	24.07	11.48	26.67	7.27	10.77	10.99
Cumulative seizure time (min)	Median	15.44	22.56	6.48	21.65	5.80	9.08	10.15
	75% Percentile	6.90	7.13	2.29	8.84	1.93	2.87	3.06
Number of computer-detected seizures/24 hrs	25% Percentile	24.5	20.25	19	11.75	6.5	7.5	8
	Median	53	28.5	25.5	68	12	24.5	18
	75% Percentile	77.5	92.5	45	109	28.5	39.75	32
	25% Percentile	2.85	0.95	0.425	1.6	0.5	0.9513	0.833
Time in spike/burst activity (hrs) per 24 hrs	Median	3.4	1.85	1.45	5.45	1.1^{**}	1.125	1.76
	75% Percentile	4.4	3.675	2.75	11.2	1.75	3.447	3.42

Abbreviations: Mz 4.5 = midazolam 4.5 mg/kg; Ket 45 = ketamine 45 mg/kg; Valp 135 = valproate 135 mg/kg.

* p < 0.05 or

** p < 0.01 vs SE control

p < 0.05 vs midazolam

• p < 0.05 or

proate.
vs valproate.
d
p < 0.01
**

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Effect of monotherapies and dual therapies on neuronal injury assessed by fluoro-jade B staining.

	SE control (n= 5)	Valp 270 mg/kg (n=5)	Mz 9mg/kg (n=10)	Ket 90 mg/kg (n=7)	Mz 4.5 + Ket 45 (n= 9)
Frontoparietal cortex	4 (1-4)	4 (4)	2 (1-4)	1 (1) ^{**}	1 (1-2) ^{**}
Entorhinal cortex	4 (4)	4 (4)	1 (1-4)	1 (1) ^{***}	1 (1) ^{***}
Piriform cortex	4 (4)	4 (4)	1 (1-4)	1 (1) ^{***}	1 (1-2) ^{**}
Thalamus	4 (1-4)	4 (3-4)	2 (1-4)	1 (1-2) [*]	1 (1) ^{**}
Amygdala	4 (4)	4 (4)	2 (1-4)	1 (1-2) [*]	1 (1) ^{***}

The first numbers represent the median damage score. The numbers in parentheses indicate the range of neuronal injury.

* p < 0.05 or

** p < 0.01 vs SE control

• p < 0.05 or

•• p < 0.01 vs valproate.