Published in final edited form as:

Clin Psychol Sci. 2016 July ; 4(4): 629–640. doi:10.1177/2167702615602864.

Self-distancing Buffers High Trait Anxious Pediatric Cancer Caregivers against Short- and Longer-term Distress

Louis A Penner¹, Darwin A. Guevarra², Felicity W. K. Harper¹, Jeffrey Taub³, Sean Phipps⁴,
Terrance L. Albrecht¹, and Ethan Kross²

¹Wayne State University and Karmanos Cancer Institute Detroit, Michigan

²University of Michigan, Ann Arbor, Michigan

³Children's Hospital of Michigan, Detroit, Michigan

⁴Saint Jude's Children's Research Hospital, Memphis Tennessee

Abstract

Pediatric cancer caregivers are typically present at their child's frequent, invasive treatments, and such treatments elicit substantial distress. Yet, variability exists in how even the most anxious caregivers cope. Here we examined one potential source of this variability: caregivers' tendencies to self-distance when reflecting on their feelings surrounding their child's treatments. We measured caregivers' self-distancing and trait anxiety at baseline, anticipatory anxiety during their child's treatment procedures, and psychological distress and avoidance three months later. Self-distancing buffered high (but not low) trait anxious caregivers against short- and long-term distress without promoting avoidance. These findings held when controlling for other buffers, highlighting the unique benefits of self-distancing. These results identify a coping process that buffers vulnerable caregivers against a chronic life stressor while also demonstrating the ecological validity of laboratory research on self-distancing. Future research is needed to explicate causality and the cognitive and physiological processes that mediate these results.

Keywords

self-distancing; pediatric cancer; vulnerable caregivers; anticipatory anxiety; psychological distress

Pediatric cancer is the leading cause of disease-related deaths for children and adolescents in the United States (American Cancer Society, 2014). Although the threat of losing a child to cancer and the multiple psychosocial challenges faced by caregivers clearly contribute to their distress, the difficult, invasive treatment-related procedures that their child undergoes play a major role in predicting caregivers' distress (Best, Streisand, Catania, & Kazak, 2001;

Corresponding Author Information: Louis A Penner, Department of Oncology, Wayne State University and Karmanos Cancer Institute, 4100 John R. Detroit Michigan, 48201, pennerl@karmanos.org.

Portions of this research were presented in a poster at the 12th Annual Meeting of the International Psycho-Oncology Society, Brisbane, Australia

Financial Disclosure: The authors have no financial relationships relevant to this article to disclose.

Kazak et al., 1995; Norberg & Boman, 2008, 2013; Phipps, Dunavant, Lensing, & Rai, 2005). Caregivers are almost invariably present during these procedures, which they often describe as worse than the disease itself (Hedström, Haglund, Skolin, & Von Essen, 2003; Ljungman et al., 1996).

To date, most research on pediatric cancer caregivers has focused on answering questions about the level of the modal caregiver's long-term stress reactions (e.g. Dunn et al., 2012; Phipps et al., 2015; Stuber et al., 1996) and identifying the social and environmental resources that generally endanger or protect caregivers from the psychosocial difficulties associated with their situation (e.g., Greening & Stoppelbein, 2007; Pai & Kazak, 2006; Rosenberg et al., 2014). While this research generally ignores individual differences among caregivers, other research on pediatric cancer caregivers suggest that individual differences in personal dispositions play an important role in determining which caregivers' are most likely to experience high levels of distress. Specifically, this research suggests that high levels of trait anxiety put caregivers at risk for both heightened anxiety at the time of treatments and longer-term symptoms of psychological distress (e.g., Dunn et al., 2012; Harper et al., 2014, 2015; Harper et al., 2013; Peterson et al., 2014; Sloper, 2000). Indeed, in an earlier study of pediatric cancer caregivers, Hoekstra-Weebers, Jaspers, Kamps, and Klip (1999) concluded that, "Trait anxiety was the strongest predictor of [caregivers'] future distress." (p.1526).

Yet, even highly anxious caregivers vary substantially in their immediate and long-term reactions to their child's treatments, but beyond some prior work, by the first author and his colleagues (Harper et al., 2013, 2015; Peterson et al., 2014), little is known about the sources of such variability. In the present study we aimed to fill this knowledge gap by turning to basic psychological theory and research on self-distancing. Specifically, we examined whether a caregiver's tendency to spontaneously self-distance when reflecting on their feelings surrounding their child's painful cancer treatments plays a role in explaining this variability, as the body of work on emotion regulation suggests it should.

When people encounter stressful events they often try to make sense of their feelings (Wilson & Gilbert, 2008) by visualizing them in their imagination (Kross & Ayduk, 2011). Prior research performed with unselected, subclinical, and clinical samples indicates that people who adopt a self-distanced or "fly on the wall" visual perspective (versus a self-immersed, first person visual perspective) are buffered against many of the harmful consequences associated with analyzing negative experiences (e.g., rumination, exaggerated emotional and physiological reactivity; e.g., Kross, Ayduk, & Mischel, 2005; Kross, Gard, Deldin, Clifton, & Ayduk, 2012; Wisco et al., 2015; Wisco & Nolen-Hoeksema, 2011). This has been found in studies where participants self-distanced in the context of experimental manipulations and studies where participants engaged in this process spontaneously (e.g., Ayduk & Kross, 2010; Grossmann & Kross, 2010; Kross et al., 2005; Verduyn et al., 2012). Moreover, many of these studies find that self-distancing is particularly helpful for buffering vulnerable individuals against exaggerated distress (Bruehlman-Seneca, Orvell, Kross, & Ayduk, 2015; Kross & Ayduk, 2009, Kross et al., 2012).

Collectively, these findings suggest that self-distancing may serve as a buffer against stress among high trait anxious pediatric cancer caregivers. However, all of this self-distancing work with vulnerable groups has been performed in the laboratory. No research of which we are aware has examined whether these findings would be observed in vulnerable individuals as they grapple with ongoing stressors in their daily lives. The primary goal of this work was to address this issue. In so doing, we aimed to simultaneously address two basic science questions: (a) does research on self-distancing generalize to the real world and (b) does self-distancing buffer high vulnerable pediatric cancer caregivers against short- and longer-term distress.

Guided by the self-distancing research described above, we predicted that high trait anxious caregivers who spontaneously self-distanced as they reflected on their child's painful treatments would be buffered against both heightened anticipatory anxiety just prior to their child's treatments and against longer-term distress associated with these anxiety reactions. Because prior research has generated conflicting findings on the benefits of self-distancing for people not identified as psychologically vulnerable, with some work indicating that self-distancing is useful for such individuals (Wisco & Nolen-Hoeksema, 2011) and other work indicating an absence of any effects (e.g., Kross & Ayduk, 2009; Kross et al., 2012), we were agnostic about whether these beneficial effects would be present but much weaker in low trait anxious caregivers or simply not present at all.

We also aimed to address two important secondary questions. First, does self-distancing have significant buffering effects even when "pitted" against other variables that buffer caregiver distress? Prior research has identified two significant buffers of caregiver distress: the size of the caregivers' social support network (Harper et al., 2015) and the caregivers' perceived self-efficacy for caregiving tasks at the time of treatments (Harper et al., 2013; Peterson et al., 2014). Therefore, an important test of whether self-distancing serves as a protective factor for high anxious groups is whether it moderates the effects of trait anxiety on short- and long-term distress while controlling for these other protective factors. We expected that it would.

Second, prior research has yielded conflicting findings regarding the relationship between spontaneous self-distancing and avoidance. Studies examining people's tendency to adopt a self-distanced perspective as they *analyze* their feelings has consistently failed to reveal significant relationships between self-distancing and avoidance (e.g., Ayduk & Kross, 2009, 2010; Kross et al., 2012). However, some other studies examining people's tendency to adopt a self-distanced perspective when they simply *recall* negative experiences has found positive links between self-distancing and avoidance (e.g., Kenny et al., 2009; Williams & Moulds, 2007; cf. McIsaac & Eich, 2004). In the current study, caregivers provided data on their levels of avoidance at the three-month follow-up. This provided a naturalistic opportunity to study how self-distancing and avoidance are related. Guided by prior research indicating that self-distancing does not promote avoidance when people adopt this perspective to analyze their feelings—i.e., the process we measured in the current study—we did not expect self-distancing to promote avoidance.

Method

Overview

This study was part of a larger, longitudinal study of families of children in active treatment for cancer. It was conducted at two major children's hospitals; one was in the Midwest and the other in the south central part of the United States. All families were initially contacted by medical staff to take part in the study. Research assistants obtained consent from eligible families, 87% of whom agreed to participate. Families were eligible if: (a) the child was between 3 and 12 years old; (b) parent and child were able to speak and the parent could read English; and (c) the child had been diagnosed with cancer within the last 18 months and was receiving cancer treatment-related procedures regularly. Data were collected at study entry, immediately before and after up to 3 subsequent treatment procedures, and at follow-ups 3 and 9 months after the last treatment procedure at which data were collected. Although the larger study concerns both primary caregivers and their children, in this study we focus only on the primary caregivers who provided data at more than one treatment. Further, because prior research has not yet documented a direct relationship between anticipatory anxiety before treatments and psychological distress measured at nine months (and we expected self-distancing to primarily affect anticipatory anxiety), we only examined data from the three-month follow. Primary caregivers received \$15 gift cards for initial assessments and \$20 gift cards for each subsequent assessment; their children received \$10 gift cards for the same assessments. The Institutional Review Boards at both hospitals approved the study.

Participants

Participants were 99 primary caregivers from the larger sample who had answered two questions about self-distancing questions and provided data for at least two of their children's treatment procedures. Some of these participants or their children participated in other cross-sectional or longitudinal studies on coping with pediatric cancer (Harper et al., 2013; Harper et al., 2014; Harper et al., 2014, 2015; Peterson et al., 2014)¹. Their mean age was 33.98 ($SD=6.96$) (children $M_{age}=6.59$, $SD=3.11$). Caregivers were either parents (97%) or grandparents (3%). Eighty two percent of caregivers were the mothers of the child in treatment. Seventy-three percent of caregivers self-identified as White, 20% as Black/African American, 4% as Hispanic/Latino, and 3% as other racial/ethnic identities. A majority (60%) of the children were male; 79% were diagnosed with Acute Lymphocytic Leukemia, the most frequent pediatric cancer in the United States. Children's ethnicities reflected those of their caregivers.

Procedure

Upon entry into the study, caregivers completed an interview and several questionnaires about themselves and their children. In the months that followed, we assessed how anxious caregivers felt just prior to cancer-related treatments. Data were collected for up to three cancer-related treatments. These treatment procedures involved port-starts (insertion of a needle into a port implanted in the child's chest), lumbar punctures, or bone marrow

¹None of these studies concerned self-distancing and somewhat different dependent measures were used.

aspirations. Three months after the last procedure at which data were collected, caregivers were mailed questionnaires that asked questions relevant to their current psychosocial status.

Independent Variables

Trait Anxiety—Caregivers' trait anxiety was assessed at study entry with the 20-item Trait Anxiety subscale of the *State-Trait Anxiety Inventory* (Spielberger, Gorsuch, Lushene, Vagg, & Jacobs, 1983). The trait anxiety subscale is intended to measure the predisposition to experience chronically high levels of anxiety. A representative item is "I feel nervous and restless" (Item $M=2.33$, $SD=.63$, $\alpha=.91$). Caregivers used a 5-point Likert response scale ("Not at all like me" to "Exactly like me"). The trait anxiety subscale of the STAI is a widely used measure of anxiety as a personality attribute; an extensive literature supports both the reliability and construct validity of this measure (Spielberger, 1989; Spielberger & Reheiser, 2004).

Spontaneous Self-Distancing—Caregivers' tendency to spontaneously self-distance while reflecting on their feelings surrounding their child's painful cancer treatments was also assessed at study entry as part of an interview. Prior research has assessed this tendency by asking participants to first recall and then try to make sense of a recent painful past experience. Subsequently, participants are asked to answer two questions that assess their tendency to spontaneously adopt a self-distanced or "fly on the wall" visual perspective as they try to make sense on their feelings (e.g., Kross et al., 2014; Mischkowski, Kross, & Bushman, 2012; Park et al., 2014). The self-distancing questions that accompany these instructions have been used in multiple studies and have been demonstrated to be construct valid (e.g., Ayduk & Kross, 2010; Kross et al., 2012; Kross et al., 2014; Mischkowski et al., 2012; Park et al., 2014).

The current study employed a version of this protocol and two questions that were tailored to target caregivers' experiences thinking about their child's stressful cancer procedures. Specifically, a research assistant instructed each caregiver to *"...think back to your child's most recent hospital visit, and your experience while he or she received treatment or a diagnostic procedure related to their illness. Please go back to the time and place of the experience and see the scene in your imagination. Take a few moments to recall this event. Once you have it in mind, let me know."* The assistant waits for caregiver to indicate that they recalled the event. The research assistant then instructs the caregiver that *"...to spend about a minute thinking about the causes and reasons underlying the thoughts and feelings you experienced during that experience. Try to understand the emotions you experienced during the event."*

Caregivers were given up to 60 seconds to complete this imagery task. Next, we used two items to assess spontaneous self-distancing. First, participants rated the extent to which they adopted the perspective of an immersed participant (i.e., "saw the event replay through my own eyes, as if I were right there ...") versus a distanced observer (i.e., "watched the event unfold from the perspective of an observer, in which I could see myself from afar ...") as they pondered their deepest thoughts and feelings during the task. Caregivers used a scale from 1 (predominantly immersed participant) to 7 (predominantly distanced observer). Next,

we had participants rate how far they were from the scene in their mind's eye during the task; responses were made on a scale from 1 (very close, saw it through my own eyes) to 7 (very far, saw it as if an observer). Responses to the two questions were averaged to create a single self-distancing index ($M=2.55$, $SD=1.66$).²

Although responses to the two questions were significantly related ($\alpha=.75$), preliminary analyses revealed that several participants gave quite different responses to them. Given that the two questions are intended to assess the same construct (i.e., self-distancing), we performed a residual analysis to identify outliers with regard to consistency in response to the two questions—i.e., we plotted the standardized residuals for the two self-distancing items on a boxplot, used the 1.5 interquartile range away from the 25th and 75th percentile as cut offs, and identified eight participants who were outliers (i.e., the size of their residuals exceeded these parameters); they were removed from the primary analyses. When these eight cases were removed, mean and the coefficient alpha for the remaining primary caregivers were $M=2.45$, $SD=1.69$, $\alpha=.89$. The results of analyses with and without the eight outliers were virtually identical. However, because the self-distancing scores of the latter group were more reliable, we report the findings of analyses that exclude the outliers.

Covariates

Social Support Network Size—Size of the caregivers' social support network was assessed with Sarason, Levine, Basham, & Sarason's (1983) brief measure of social support. Specifically the size of the network was operationally defined as the average of caregivers' report of the number of people who would provide support across the six different areas of need (e.g., providing dependable help) ($M=4.23$, $SD=2.46$; α across six areas $=.95$). A large research literature supports the reliability and validity of this assessment of perceived size of one's social support network (Prodicano & Smith, 2013; Sarason, B., Sarason I., Hacker, & Basham, 1985; Sarason, I., & Sarason B., 1985).

Caregiver self-efficacy—The procedures for assessing caregiver treatment self-efficacy are described in detail in Harper et al. (2013). Prior to each treatment, caregivers were presented with 6 caregiving tasks specific to their child's treatment procedure that day (e.g., keeping the child calm before the procedures began; gaining needed information from medical staff). For each task, caregivers rated its importance and their confidence in performing the task effectively. Self-efficacy was operationalized as the product of these scales. Peterson et al.'s (2014) findings indicate that this measure is construct valid. These scores were averaged across treatment sessions (all α 's across treatment sessions $>.76$) and then standardized with a z score transformation. Research using this measure of self-efficacy (Harper et al., 2013; Peterson et al., 2014) finds that self-efficacy about keeping the child calm before the procedure predicts caregivers' state anxiety prior to cancer-related procedures, their self-rated distress during the actual procedures, mood following the procedures, and symptoms of post traumatic stress at the three month follow-up. As would

²Although not the focus of this study, we also asked participants to rate the degree to which they re-experienced their emotions after reflecting on their feelings surrounding their child's cancer treatment. Spontaneous self-distancing was not significantly correlated with responses to this item.

be expected, all of these relationships are negative; that is the greater the self-efficacy the less psychological distress³.

Dependent Variables

Anticipatory Anxiety Prior to Treatment Procedures—Caregivers' anticipatory anxiety just prior to their child's treatment procedures was assessed with the 20-item State Anxiety subscale of Spielberger's State-Trait Anxiety Inventory (Spielberger et al., 1983). This subscale is intended to measure anxiety in response to a specific stimulus or situation; this kind of anxiety is considered transitory (Hedberg, 1972). A representative item is "I am worried." Caregivers used a 5-point response scale ("Not at all" to "Exactly") to indicate their level of state anxiety. A large research literature supports the reliability and construct validity of this subscale of the STAI (Spielberger, 1989; Spielberger & Reheiser, 2004).

All caregivers provided state anxiety scores for at least two treatments and 85 of them (87%) provided state anxiety scores for all three treatments. State anxiety scores were reliable across treatment procedures ($\alpha=.87$). Thus, they were averaged across treatments to form a single state anxiety index (Mean item response = 2.50, $SD=.42$).⁴

Psychological Distress at Three-month Follow-up—Three months after the last treatment procedure, (months since baseline assessment: $M=6.83$, $SD=2.62$, range 3.65–13.17 months⁵) caregivers received a mailed questionnaire that contained two scales that assessed symptoms of psychological distress—the 18-item Brief Symptom Inventory (BSI-18; Derogatis, 2001) and the 22-item Impact of Events Scale (IES-R; Horowitz, Wilner, & Alvarez, 1979; Weiss & Marmar, 1997). Both measures had a 0 (Never) to 4 (Very Often) response scales, which caregivers used to describe how frequently they were currently experiencing the symptoms presented in each measure. The BSI-18 asked caregivers about psychiatric symptoms of anxiety (6 items, $M=.82$, $SD=.71$, $\alpha=.84$), depression (6 items, $M=.76$, $SD=.66$, $\alpha=.84$), and somatization (6 items, $M=.53$, $SD=.69$, $\alpha=.88$). A large literature supports the reliability and construct validity of BSI-18 as a screening for symptoms of psychiatric distress (Boothroyd, 2003; Mariush, 2004; Zabora et al., 2001)

The IES-R asked participants about symptoms associated with posttraumatic stress: hyperarousal (6 items $M=.98$, $SD=.76$, $\alpha=.73$), intrusion (8 items, $M=1.38$, $SD=.87$, $\alpha=.90$) and avoidance (8 items $M=1.03$, $SD=.74$, $\alpha=.82$) in relation to their child's previous treatments over the past several months. The IES-R has been shown to correlate well with independent assessments of PTSD using criteria from the Diagnostic Statistical Manual. Research on the IES-R indicates it is a reliable and construct valid measure of symptoms of posttraumatic distress (Creamer, Bell, & Salvina, 2003; Weiss, 2004).

³We also used self-efficacy about keeping the child calm *during the treatment* as the covariate and the findings were essentially identical. Therefore, they are not discussed further. Also, although age was not correlated with the variables of interest in this study, it was also included as a covariate to make the present findings comparable to other studies using this sample of caregivers that included age as a covariate.

⁴Caregivers' mood (PANAS; Clark, Watson, & Tellegen, 1988) was measured after the treatment procedures. Given the focus of this study on anxiety. It is not discussed further.

⁵One family mistakenly received the follow-up questionnaires upon completion of the final treatment. Exclusion of their data does not change any of the findings.

The subscales were all significantly intercorrelated within each measure and across the two measures. We combined all of these BSI and IES subscales (except avoidance) to create a reliable single composite measure of psychological distress at the three-month follow-up. To do this we z-score transformed the anxiety, depression, somatization, hyperarousal and intrusion subscales and then averaged them to create a single composite index of global psychological distress. The measure was internally consistent, $\alpha=.85$. We did not include avoidance in this composite measure because, as noted earlier, we wanted to examine the relationship between self-distancing and avoidance independently of the other psychological distress measures.

Results

Overview of Analyses

First, zero order correlations were computed to determine if trait anxiety and spontaneous self-distancing measured at study entry correlated with caregivers' anticipatory (i.e., state) anxiety during treatments, their symptoms of psychological distress (i.e., total score on the composite measure), and avoidance at the three-month follow-up or any of the covariates.

Next, Model 1 of Hayes (2013) regression utility for SPSS ("Process") was used to determine whether trait anxiety and self-distancing interacted to affect either anticipatory anxiety (short term outcome), or the composite measure of psychological distress, and avoidance (longer-term outcomes). In these analyses all predictor variables were centered.

Prior studies with pediatric cancer caregivers (e.g., Harper et al., 2014; Harper et al., 2013) have shown that anticipatory anxiety mediates the relationship between trait anxiety and the measures of longer-term psychological distress used in the present study (specifically, the BSI-18 and the IES-R [including avoidance] measured separately). To determine whether self-distancing moderates this relationship, we used Model 8 of Hayes (2013) regression utility for SPSS ("Process"), which tests for moderated mediation. Again this was done with and without the covariates.

Because all variables in the regression equations had been converted to z scores, the correlation coefficients presented in Table 1 and the regression coefficients reported in the text and/or Table 2 represent standardized estimates of *effect size*. Specifically, the standardized regression coefficients (β) are the amount of change in the dependent variable for each change of one standard deviation in the variable of interest, while holding all other predictor variables in the equation constant. All estimates of regression effect sizes (i.e., standardized regression weights) in the equations are accompanied by 95% confidence intervals. Each regression coefficient was tested for significance using a "t" test with the alpha level set at .05.

Correlates of Caregivers' Trait Anxiety and Self-distancing

Prior research with different samples from the larger study (e.g., Harper et al., 2014, 2015; Harper et al., 2013) has shown that trait anxiety significantly affects anticipatory (i.e., state) anxiety at treatment procedures and psychological distress during the follow-up. As Table 1 shows, this was also the case with this particular sample for both anticipatory anxiety and

the composite measure of psychological distress assessed three months after the last treatment. Table 1 also shows that spontaneous self-distancing was not significantly correlated with trait anxiety, any of the covariates, state anxiety, or the composite measure of longer-term psychological distress. It was also uncorrelated with avoidance.

Moderation of Caregivers' Anticipatory Anxiety Before Treatment Procedures

We predicted that spontaneous self-distancing would have a buffering effect on high trait anxious caregivers. That is, we expected high self-distancing would minimize the impact of trait anxiety on their anticipatory anxiety. This prediction was confirmed; we found a significant interaction between trait anxiety and spontaneous self-distancing, $\beta = -.23$, 95% CI $[-.41, -.05]$, $t(89) = 2.53$, $p = .01$. The full regression equation is presented in Table 2. Figure 1 Panel A presents the regression slopes for low and high trait anxious caregivers. Among caregivers who were *low* in trait anxiety (i.e., 1 standard deviation below the sample trait anxiety mean), their tendency to spontaneously self-distance had no implications for how much anticipatory anxiety they experienced before their child's treatments. However, among *high* trait anxious caregivers (i.e., 1 standard deviation above the sample trait anxiety mean), self-distancing served a protective function. That is, among high anxious caregivers trait anxiety significantly affected anticipatory anxiety only for caregivers who were *low* in self-distancing, $\beta = -.37$, 95% CI $[-.65, -.10]$, $t(89) = -2.67$, $p = .01$. Among the *high* self-distancing caregivers, this relationship was not significant, $\beta = .07$, 95% CI $[-.14, .28]$, $t(90) = .67$, $p = .50$.

Moderation of Caregivers' Longer-term Symptoms of Psychological Distress

As expected, we observed a significant trait anxiety by spontaneous self-distancing interaction, $\beta = -.20$, 95% CI $[-.37, -.02]$, $t(85) = 2.27$, $p = .03$. As Figure 1, Panel B demonstrates, the nature of this interaction was similar to the interaction reported for anticipatory anxiety. Specifically, if a caregiver scored *low* on trait anxiety, their tendency to spontaneously self-distance had no implications for their longer-term psychological distress. However, if a caregiver scored *high* on trait anxiety, spontaneous self-distancing served a protective function. That is, among high anxious caregivers trait anxiety significantly affected longer-term psychological distress only among caregivers who were *low* in self-distancing, $\beta = -.28$, 95% CI $[-.53, -.03]$, $t(85) = 2.19$, $p = .03$. Among the *high* self-distancing caregivers, this relationship was not significant, $\beta = .08$, 95% CI $[-.11, .27]$, $t(85) = -.87$, $p = .39$.

Moderation of Caregivers' Avoidance

Self-distancing did not interact with trait anxiety to predict avoidance at the three-month follow-up, $\beta = .08$, 95% CI $[-.14, .31]$, $t(85) = .72$, $p = .47$.

⁶Moderation analyses without the covariates (i.e. social support, self-efficacy, and child age) produced a marginally significant interaction term for anticipatory anxiety, $\beta = -.18$, 95% CI $[-.36, .01]$, $t(94) = 1.89$, $p = .06$; and a significant interaction term for longer-term psychological distress $\beta = -.18$, 95% CI $[-.34, -.01]$, $t(89) = 2.12$, $p = .04$. Consistent with the results of the covariate analyses, the regressions of the outcome variables onto self-distancing were only significant among the *high* anxiety caregivers. Anticipatory anxiety: $\beta = -.31$, 95% CI $[-.59, -.02]$, $t(94) = -2.13$, $p = .04$; psychological distress: $\beta = -.26$, 95% CI $[-.50, -.02]$, $t(89) = -2.12$, $p = .04$. The results for caregivers low in trait anxiety were not significant: anticipatory anxiety, $\beta = .04$, 95% CI $[-.18, .26]$, $t(94) = .35$, $p = .73$; psychological distress, $\beta = -.07$, 95% CI $[-.12, .25]$, $t(89) = .68$, $p = .50$.

Moderation of Relation Between Caregiver Anticipatory Anxiety and Longer-Term Psychological Distress

Our final set of analyses aimed to enhance our understanding of how self-distancing buffers high trait anxious caregivers against long-term distress. Towards this end, we first examined whether participant's anticipatory anxiety mediated the relationship between trait anxiety and longer-term psychological distress—as noted earlier, this mediational pathway has been documented in prior research with another subset of the caregivers in the larger study (e.g., Harper et al., 2014). There was also significant mediation in the present subsample with the composite measure of psychological distress at the three month follow-up as the outcome, indirect effects, $\beta=.10$, 95% CI [.03, .19] (Hayes (2013) Model 4, simple mediation).

Next, we examined whether self-distancing affects this pathway. That is, we conducted an analysis of moderated mediation (Hayes [2013] Model 8 for SPSS) of the indirect path from trait anxiety to longer-term psychological distress. We expected that the indirect path with state anxiety as the mediator would only be significant among the low self-distancing caregivers.

The overall moderated mediation model was significant, $\beta=-.04$, 95% CI [-.13, -.001]. Inspection of the indirect effects of state anxiety on psychological distress among caregivers low and high in self-distancing indicated that whereas there was a significant indirect effect among *low* self-distancing caregivers, $\beta=.10$, 95% CI [.01, .25]; there was no significant indirect effect among *high* self-distancing caregivers, $\beta=.02$, 95% CI [-.03, .11]. This analysis indicates that higher levels of self-distancing may serve a buffering function by disrupting the relationship between caregivers' state anxiety in response to the treatments and their psychological distress three months following their child's final cancer treatment.

Discussion

These findings demonstrate that among high trait anxious pediatric cancer caregivers, high levels of spontaneous self-distancing served to virtually eliminate the impact of trait anxiety on (a) anticipatory anxiety during the time of children's painful cancer treatments and (b) longer term psychological distress. These effects were found when controlling for other established buffers of caregiver trait anxiety, highlighting the specificity and robustness of the effects of self-distancing. Finally, consistent with prior work (e.g., Ayduk & Kross, 2009, 2010; Kross et al., 2012), self-distancing did not promote avoidance.

These findings have three important basic science implications. First, they add to our understanding of the well-documented individual differences in how pediatric cancer caregivers cope with the psychosocial effects of their child's cancer. As the literature review presented in the introduction indicates, there is widespread agreement on the important role trait anxiety plays in the way in which pediatric cancer caregivers react to the treatments their children receive and other aspects of the disease (e.g., Hoekstra-Weebers et al., 1999; Link & Fortier, 2015). Yet there is relatively little research on the cognitive processes that mitigate the impact of trait anxiety on these caregivers. The present research begins to fill this knowledge gap by identifying spontaneous self-distancing as a moderator that buffers

high trait anxious caregivers against short- and longer-term distress related to coping with their child's painful cancer treatments.

These findings also have important implications for theory and research on self-distancing. First, and perhaps most importantly, they show that laboratory research on self-distancing is ecologically valid (Shadish, Cook & Campbell, 2002) and can be conceptually replicated in the real world. Second, they provide additional evidence suggesting that self-distancing is specifically useful for helping “high risk,” highly vulnerable individuals cope with ongoing emotional threats. Indeed, the data represented in Figures 1a and 1b indicate that on both measures of distress the high trait anxious/high self-distancing caregivers were almost indistinguishable from the low trait anxious caregivers. In this vein it is noteworthy that we did not observe any comparable beneficial effects of self-distancing for people who scored low on trait anxiety. This finding is consistent with a number of recent laboratory studies on self-distancing that have likewise found that emotional vulnerability factors moderate the benefits of this process in the same direction observed here (Kross & Ayduk, 2009; Kross et al., 2012). More broadly, they suggest that a certain level of emotional distress may be needed for self-distancing to exert its beneficial effects; if not enough distress is experienced, there may be little room for self-distancing to operate.

Finally, the fact that self-distancing led to reductions in long-term distress without increasing avoidance conceptually replicates prior work (Ayduk & Kross, 2010; Kross & Ayduk, 2009; Kross et al., 2012) and argues against the idea that self-distancing is a de facto form of avoidance. This is not to say that we expect self-distancing to invariably be beneficial. Instead, whether this process is adaptive or not likely depends critically on the context in which people engage in this process. That is, are people self-distancing to approach or avoid their feelings? In this vein it is important to emphasize that all participants were specifically asked to confront their emotions in this study – i.e., “think about *the causes and reasons underlying the thoughts and feelings [you] experienced during [your] child's last cancer treatment experience.*” We then asked participants to report on the degree to which they self-distanced as they engaged in this meaning-making process.

A number of caveats are in order before we turn to suggestions for future research. First, the real world context in which this study was conducted—caregivers coping with the cancer-related treatment procedures their children receive—provided a unique opportunity to expand our understanding of how spontaneous self-distancing operates outside the laboratory, as vulnerable individuals grapple with intense emotional experiences. Although there are clear advantages associated with this approach, there are also disadvantages. For example, as is the case in any naturalistic study of psychological processes, our ability to include the kinds of control conditions and experimental manipulations that would permit strong inferences about causality was limited in this study. Thus, although the results regarding the links between self-distancing and our outcomes are consistent with our a priori hypotheses and a large body of prior research, in the absence of random assignment to experimental conditions we are unable to state it was spontaneous self-distancing, independent of any unmeasured other variables that was responsible for the buffering effects among the high trait anxious caregivers.

In a similar vein, a caveat is in order when considering the mediation analyses we reported, which suggested that the interaction between trait anxiety and spontaneous self-distancing predicted longer-term distress because of their impact on caregivers' anticipatory anxiety at the time of the children's treatments. Although this mediational analysis was theory-predicted and utilized temporally distinct longitudinal data points to test the predicted pathway of interest—both assets for drawing inferences about likely causal sequences from non experimental data—we were not able to manipulate either of the major predictor variables (trait anxiety and spontaneous self-distancing) or the mediating variable (anticipatory anxiety). Thus, causality cannot be determined with the present data.

Finally is important to acknowledge that we did not include multiple types of measures of our constructs of interest. Thus, although the self-report tools we used to assess our core independent and dependent variables have all been validated and used extensively in the past, future research should consider measuring these constructs with other types of measures (e.g., physiological [Ebner-Priemer & Kubiak, 2007] and linguistic measures [Dirske, Hadjistavropoulos, Hesser, & Barak, 2015] of anxiety, neural measures of self-distancing [Park et al., 2014], or informant measures of a person's distress and/or psychosocial adjustment [e.g., McDade, Watson, O'Hara, & Denberg, 2008]).

Although these finding raise multiple questions for future research, three stand out to us as most pressing. First, future research is needed to test the causal relationships that are suggested, but cannot be tested or proven with the present data. While trait anxiety cannot be meaningfully manipulated among pediatric cancer caregivers (or other vulnerable groups) for ethical reasons, it would be possible to conduct small randomized field experiments that enhance caregivers' propensity to self-distance and compare their responses to an untreated comparison group. Such experiments are justifiable on the basis of the present findings and would speak directly to the causal impact of self-distancing on caregivers' short and longer-term distress.

Second, what are the pathways through which self-distancing buffers high anxious pediatric cancer caregivers or other vulnerable individuals? On the one hand, it is possible that self-distancing leads to adaptive shifts in the way caregivers mentally present their child's cancer treatments. For example, rather than focusing on the concrete, aversive features of these events (i.e., the invasive nature of the procedures, the child's displays of distress—crying, for example), self-distancing might lead caregivers to focus more broadly on the potential good that may come from the treatment (i.e., they will help my child live). It is also possible, however, that self-distancing instigates physiological changes that work to combat their anxiety. For example, prior research indicates that self-distancing buffers individuals against the autonomic consequences of ruminating over negative experiences (i.e., chronically elevated cardiovascular reactivity and shortened cardiovascular recovery; e.g., Ayduk & Kross, 2008; Brosschot, Gerin, & Thayer, 2006; Wisco et al., 2015). Thus, it is possible that there is simply less aversive physiological reactivity among high self-distancing, high trait anxious caregivers. Future research is needed to examine the role that these different pathways play in either independently or jointly accounting for the current results.

A third important question is whether there is a “tipping point” that captures when self-distancing ceases to be useful and becomes harmful? The current study was motivated by research suggesting that encouraging people to “take a step back” in the heat of the moment can be useful for allowing them to confront and approach negative experiences constructively, in ways that facilitating adaptive coping and self-reflection. However, the distribution of responses to the self-distancing questions in the current study indicates that even caregivers classified as high self-distancing (i.e., those who scored 1 SD about the group mean) only engaged in moderate levels of distancing (just above the mid-point on the seven point scale of tendency to self-distance). It is possible that a caregiver could distance themselves too much from their child’s treatment experience, which could diminish their care and concern for the child. The absence of such feelings make treatments much more difficult for children (See Cline et al., 2006). Future research is needed to address this question and is important for refining our understanding of self-distancing and the factors that promote effective pediatric cancer caregiving.

Acknowledgments

This work is part of a larger investigation of caregivers’ influence on child coping with cancer treatment and was supported by the National Cancer Institute at the National Institutes of Health (NCI#1R01CA138981; PI: L. Penner) and The Herrick Foundation (PI: L. Penner).

References

- American Cancer Society. Cancer facts & figures special section: Cancer in children and adolescents. 2014. Retrieved from <http://www.cancer.org/acs/groups/content/@research/documents/webcontent/acspc-041787.pdf>
- Ayduk O, Kross E. Enhancing the pace of recovery self-distanced analysis of negative experiences reduces blood pressure reactivity. *Psychological Science*. 2008; 19(3):229–231. [PubMed: 18315794]
- Ayduk O, Kross E. Asking ‘why’ from a distance facilitates emotional processing: A reanalysis of Wimalaweera and Moulds (2008). *Behaviour Research and Therapy*. 2009; 47(1):88–92. [PubMed: 19013553]
- Ayduk O, Kross E. From a distance: implications of spontaneous self-distancing for adaptive self-reflection. *Journal of Personality and Social Psychology*. 2010; 98(5):809–829. [PubMed: 20438226]
- Best M, Streisand R, Catania L, Kazak AE. Parental distress during pediatric leukemia and posttraumatic stress symptoms (PTSS) after treatment ends. *Journal of Pediatric Psychology*. 2001; 26(5):299–307. [PubMed: 11390572]
- Boothroyd, RA. Review of the brief symptom inventory 18. In: Plake, BS.; Impara, JC.; Spies, RA., editors. *The fifteenth mental measurements yearbook*. Lincoln, NE: Buros Institute of Mental Measurements of the University of Nebraska-Lincoln; 2003. p. 136-138.
- Brosschot JF, Gerin W, Thayer JF. The perseverative cognition hypothesis: A review of worry, prolonged stress-related physiological activation, and health. *Journal of Psychosomatic Research*. 2006; 60(2):113–124. [PubMed: 16439263]
- Bruehlman-Seneca, E.; Orvell, A.; Kross, E.; Ayduk, O. From laboratory to daily life: A brief self-distancing intervention buffers against rumination and depression. 2015. Manuscript in preparation
- Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of Personality and social Psychology*. 1988; 54(6):1063–1070. [PubMed: 3397865]
- Cline RJ, Harper FW, Penner LA, Peterson AM, Taub JW, Albrecht TL. Parent communication and child pain and distress during painful pediatric cancer treatments. *Social Science & Medicine*. 2006; 63(4):883–898. [PubMed: 16647174]

- Cremer M, Bell R, Failla S. Psychometric properties of the impact of event scale—revised. *Behaviour Research and Therapy*. 2003; 41(12):1489–1496. [PubMed: 14705607]
- Derogatis, LR. BSI 18, Brief Symptom Inventory 18: Administration, scoring and procedures manual. Minneapolis, MN: NCS Pearson Inc; 2001.
- Dirkse D, Hadjistavropoulos HD, Hesser H, Barak A. Linguistic analysis of communication in therapist-assisted internet-delivered cognitive behavior therapy for generalized anxiety disorder. *Cognitive Behaviour Therapy*. 2015; 44(1):21–32. [PubMed: 25244051]
- Dunn MJ, Rodriguez EM, Barnwell AS, Grossenbacher JC, Vannatta K, Gerhardt CA, Compas BE. Posttraumatic stress symptoms in parents of children with cancer within six months of diagnosis. *Health Psychology*. 2012; 31(2):176–185. [PubMed: 21942750]
- Ebner-Priemer UW, Kubiak T. Psychological and psychophysiological ambulatory monitoring: A review of hardware and software solutions. *European Journal of Psychological Assessment*. 2007; 23(4):214–226.
- Grossmann I, Kross E. The impact of culture on adaptive versus maladaptive self-reflection. *Psychological Science*. 2010; 21(8):1150–1157. [PubMed: 20624935]
- Greening L, Stoppelbein L. Brief report: Pediatric cancer, parental coping style, and risk for depressive, posttraumatic stress, and anxiety symptoms. *Journal of Pediatric Psychology*. 2007; 32(10):1272–1277. [PubMed: 17675304]
- Harper FWK, Goodlett BD, Trentacosta CJ, Albrecht TL, Taub J, Phipps S, Penner L. Temperament, Personality, and Quality of Life in Pediatric Cancer Patients. *Journal of Pediatric Psychology*. 2014; 39:459–468. [PubMed: 24443742]
- Harper FWK, Peterson AM, Albrecht TL, Taub JW, Phipps S, Penner LA. Posttraumatic stress symptoms in parents of pediatric cancer patients: A mediational analysis. *Journal of Traumatic Stress Disorders & Treatment*. 2014; 3(4):1000133. [PubMed: 25580441]
- Harper FWK, Peterson AM, Albrecht TL, Taub JW, Phipps S, Penner LA. Satisfaction with support versus size of network: Differential effects of social support on psychological distress in parents of pediatric cancer patients. *Psycho-Oncology*. 2015; Advance online publication. doi: 10.1002/pon.3863
- Harper FWK, Peterson AM, Uphold H, Albrecht TL, Taub JW, Orom H, Penner LA. Longitudinal study of parent caregiving self-efficacy and parent stress reactions with pediatric cancer treatment procedures. *Psycho-Oncology*. 2013; 22(7):1658–1664. [PubMed: 23034930]
- Hayes, AF. Introduction to mediation, moderation, and conditional process analysis: A regression-based approach. New York, NY: Guilford Press; 2013.
- Hedberg AG. Review of state-trait anxiety inventory. *Professional Psychology*. 1972; 3:389–390.
- Hedström M, Haglund K, Skolin I, Von Essen L. Distressing events for children and adolescents with cancer: Child, parent, and nurse perceptions. *Journal of Pediatric Oncology Nursing*. 2003; 20(3):120–132. [PubMed: 12776260]
- Hoekstra-Weebers JE, Jaspers JP, Kamps WA, Klip EC. Risk factors for psychological maladjustment of parents of children with cancer. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1999; 38(12):1526–1535. [PubMed: 10596253]
- Horowitz M, Wilner N, Alvarez W. Impact of event scale: A measure of subjective stress. *Psychosomatic Medicine*. 1979; 41(3):209–218. [PubMed: 472086]
- Kazak AE, Boyer BA, Brophy P, Johnson K, Scher CD, Covelman K, Scott S. Parental perceptions of procedure-related distress and family adaptation in childhood leukemia. *Children's Health Care*. 1995; 24(3):143–158.
- Kenny LM, Bryant RA, Silove D, Creamer M, O'Donnell M, McFarlane AC. Distant memories a prospective study of vantage point of trauma memories. *Psychological Science*. 2009; 20(9):1049–1052. [PubMed: 19594861]
- Kross E, Ayduk O. Facilitating adaptive emotional analysis: distinguishing distanced-analysis of depressive experiences from immersed-analysis and distraction. *Personality and Social Psychology Bulletin*. 2008; 34(7):924–938. [PubMed: 18469151]
- Kross E, Ayduk O. Boundary conditions and buffering effects: Does depressive symptomology moderate the effectiveness of self-distancing for facilitating adaptive emotional analysis? *Journal of Research in Personality*. 2009; 43(5):923–927. [PubMed: 20161055]

- Kross E, Ayduk O. Making meaning out of negative experiences by self-distancing. *Current Directions in Psychological Science*. 2011; 20(3):187–191.
- Kross E, Ayduk O, Mischel W. When asking “why” does not hurt: Distinguishing rumination from reflective processing of negative emotions. *Psychological Science*. 2005; 16(9):709–715. [PubMed: 16137257]
- Kross E, Bruehlman-Senecal E, Park J, Burson A, Dougherty A, Shablack H, Ayduk O. Self-talk as a regulatory mechanism: How you do it matters. *Journal of Personality and Social Psychology*. 2014; 106(2):304–324. [PubMed: 24467424]
- Kross E, Duckworth A, Ayduk O, Tsukayama E, Mischel W. The effect of self-distancing on adaptive versus maladaptive self-reflection in children. *Emotion*. 2011; 11(5):1032–1039. [PubMed: 21728415]
- Kross E, Gard D, Deldin P, Clifton J, Ayduk O. “Asking why” from a distance: Its cognitive and emotional consequences for people with major depressive disorder. *Journal of Abnormal Psychology*. 2012; 121(3):559–569. [PubMed: 22708885]
- Link CJ, Fortier MA. The Relationship Between Parent Trait Anxiety and Parent-reported Pain, Solicitous Behaviors, and Quality of Life Impairment in Children With Cancer. *Journal of Pediatric Hematology/oncology*. 2015; Advance online publication. doi: 10.1097/MPH.0000000000000376
- Ljungman L, Hovén E, Ljungman G, Cernvall M, Essen L. Does time heal all wounds? A longitudinal study of the development of posttraumatic stress symptoms in parents of survivors of childhood cancer and bereaved parents. *Psycho-Oncology*. 2015; Advance online publication. doi: 10.1002/pon.3856
- Ljungman G, Kreuger A, Gordh T, Berg T, Sörensen S, Rawal N. Treatment of pain in pediatric oncology: A Swedish nationwide survey. *Pain*. 1996; 68(2):385–394. [PubMed: 9121828]
- Maruish, ME., editor. The use of psychological testing for treatment planning and outcomes assessment: Volume 3: Instruments for adults. 3. Mahwah, NJ: Lawrence Erlbaum Associates Publishers; 2004.
- McDade-Montez EA, Watson D, O’Hara MW, Denburg NL. The effect of symptom visibility on informant reporting. *Psychology and Aging*. 2008; 23(4):940–946. [PubMed: 19140663]
- McIsaac HK, Eich E. Vantage point in traumatic memory. *Psychological Science*. 2004; 15(4):248–253. [PubMed: 15043642]
- Mischkowski D, Kross E, Bushman BJ. Flies on the wall are less aggressive: Self-distancing “in the heat of the moment” reduces aggressive thoughts, angry feelings and aggressive behavior. *Journal of Experimental Social Psychology*. 2012; 48(5):1187–1191.
- Norberg AL, Boman KK. Parent distress in childhood cancer: a comparative evaluation of posttraumatic stress symptoms, depression and anxiety. *Acta Oncologica*. 2008; 47(2):267–274. [PubMed: 17851875]
- Norberg AL, Boman KK. Mothers and fathers of children with cancer: loss of control during treatment and posttraumatic stress at later follow-up. *Psycho-Oncology*. 2013; 22(2):324–329. [PubMed: 22021113]
- Pai AL, Kazak AE. Pediatric medical traumatic stress in pediatric oncology: Family systems interventions. *Current Opinion in Pediatrics*. 2006; 18(5):558–562. [PubMed: 16969172]
- Park J, Ayduk O, O’Donnell L, Chun J, Gruber J, Kamali M, Kross E. Regulating the high cognitive and neural processes underlying positive emotion regulation in bipolar I disorder. *Clinical Psychological Science*. 2014; 2(6):661–674. [PubMed: 26719819]
- Peterson AM, Harper FWK, Albrecht TL, Taub JW, Orom H, Phipps S, Penner LA. Parent caregiver self-efficacy and child reactions to pediatric cancer treatment procedures. *Journal of Pediatric Oncology Nursing*. 2014; 31(1):18–27. [PubMed: 24378818]
- Phipps S, Dunavant M, Lensing S, Rai SN. Psychosocial predictors of distress in parents of children undergoing stem cell or bone marrow transplantation. *Journal of Pediatric Psychology*. 2005; 30(2):139–153. [PubMed: 15681309]
- Phipps S, Long A, Willard VW, Okado Y, Hudson M, Huang Q, Noll R. Parents of Children With Cancer: At-Risk or Resilient? *Journal of Pediatric Psychology*. 2015; Advance online publication. doi: 10.1093/jpepsy/jsv047

- Prodicano, N.; Smith, WW. Assessing perceived social support. The importance of context. In: Pierce, GR.; Lakey, B.; Sarason, IG.; Sarason, BR., editors. *Sourcebook of social support and personality*. New York, NY: Springer; 2013. p. 93-106.
- Rosenberg AR, Wolfe J, Bradford MC, Shaffer ML, Yi-Frazier JP, Curtis JR, Baker KS. Resilience and psychosocial outcomes in parents of children with cancer. *Pediatric Blood & Cancer*. 2014; 61(3): 552–557. [PubMed: 24249426]
- Sarason, IG.; Sarason, BR. Social support: Insights from assessment and experimentation. In: Sarason, IG.; Sarason, BR., editors. *Social support: Theory, research, and applications*. Dordrecht, The Netherlands: Martinus-Nijhoff; 1985. p. 39-50.
- Sarason BR, Sarason IG, Hacker TA, Basham RB. Concomitants of social support: Social skills, physical attractiveness, and gender. *Journal of Personality and Social Psychology*. 1985; 49(2): 469–480.
- Sarason IG, Levine HM, Basham RB, Sarason BR. Assessing social support: the social support questionnaire. *Journal of Personality and Social Psychology*. 1983; 44(1):127–139.
- Shadish, WR.; Cook, TD.; Campbell, DT. *Experimental and quasi-experimental designs for generalized causal inference*. Boston, MA: Houghton Mifflin; 2002.
- Sloper P. Predictors of distress in parents of children with cancer: A prospective study. *Journal of Pediatric Psychology*. 2000; 25(2):79–91. [PubMed: 10820946]
- Spielberger, CD. *State-Trait Anxiety Inventory: Bibliography*. 2. Palo Alto, CA: Consulting Psychologists Press; 1989.
- Spielberger, CD.; Gorsuch, RL.; Lushene, RE.; Vagg, PR.; Jacobs, GA. *Manual for the State-Trait Anxiety Inventory*. Palo Alto, CA: Consulting Psychologists Press; 1983.
- Spielberger, CD.; Reheiser, EC. Measuring anxiety, anger, depression, and curiosity as emotional states and personality traits with the STAI, STAXI and STPI. In: Hilsenroth, MJ.; Segal, DL., editors. *Comprehensive handbook of psychological assessment: Vol. 2. Personality assessment*. Hoboken, NJ: Wiley; 2004. p. 70-86.
- Stuber ML, Christakis DA, Houskamp B, Kazak AE. Posttrauma symptoms in childhood leukemia survivors and their parents. *Psychosomatics*. 1996; 37(3):254–261. [PubMed: 8849502]
- Verduyn P, Van Mechelen I, Kross E, Chezzi C, Van Bever F. The relationship between self-distancing and the duration of negative and positive emotional experiences in daily life. *Emotion*. 2012; 12(6):1248–1263. [PubMed: 22642344]
- Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*. 1988; 54(6): 1063–1070. [PubMed: 3397865]
- Weiss, DS. The Impact of Event Scale—Revised. In: Wilson, JP.; Keane, TM., editors. *Assessing psychological trauma and PTSD: A handbook for practitioners*. 2. New York: Guilford Press; 2004. p. 168-190.
- Weiss, DS.; Marmar, CR. The Impact of Event Scale—Revised. In: Wilson, JP.; Keane, TM., editors. *Assessing psychological trauma and PTSD: A handbook for practitioners*. New York, NY: Guilford Press; 1997. p. 399-411.
- Williams AD, Moulds ML. Cognitive avoidance of intrusive memories: Recall vantage perspective and associations with depression. *Behaviour Research and Therapy*. 2007; 45(6):1141–1153. [PubMed: 17067549]
- Wilson TD, Gilbert DT. Explaining away a model of affective adaptation. *Perspectives on Psychological Science*. 2008; 3(5):370–386. [PubMed: 26158955]
- Wisco BE, Marx BP, Sloan DM, Gorman KR, Kulish AL, Pineles SL. Self-distancing from trauma memories reduces physiological but not subjective emotional reactivity among veterans with posttraumatic stress disorder. *Clinical Psychological Science*. 2015; Advance online publication. doi: 10.1177/2167702614560745
- Wisco BE, Nolen-Hoeksema S. Effect of visual perspective on memory and interpretation in dysphoria. *Behaviour Research and Therapy*. 2011; 49(6):406–412. [PubMed: 21550018]
- Zabora J, Brintzenhofesoc K, Jacobsen P, Curbow B, Piantadosi S, Hooker C, Derogatis L. A new psychosocial screening instrument for use with cancer patients. *Psychosomatics*. 2001; 42(3):241–246. [PubMed: 11351113]

Figure 1. Moderation of anticipatory anxiety (Panel A) and longer term-distress (Panel B) by self-distancing and trait anxiety. The scores on the y-axis represent z-scores.

Table 1

Zero-Order correlations among primary variables

Variable	1	2	3	4	5	6	7	8
1. Trait Anxiety	—							
2. Self-Distancing	-.14	—						
3. Child's Age	.004	.13	—					
4. Social Support Size	-.02	-.01	-.07	—				
5. Caregiver Self-Efficacy	-.18	-.004	.36**	.01	—			
6. Anticipatory Anxiety	.42***	-.15	-.17	-.12	-.35***	—		
7. Longer-Term Distress	.33***	-.13	.02	-.03	-.09	.37***	—	
8. Avoidance	.29***	-.03	-.003	-.08	-.17	.46***	.55***	—

Note. $N = 92 - 99$

* $p < .05$

** $p < .01$

*** $p < .001$.

Table 2

Summary of regression analyses for anticipatory anxiety, three-month longer term distress, and three-month avoidance.

Variable	Anticipatory Anxiety			Three-month Longer Term Distress			Three-month Avoidance		
	β (SE)	<i>t</i>	95% CI	β (SE)	<i>t</i>	95% CI	β (SE)	<i>t</i>	95% CI
Trait Anxiety	.34(.09)	3.98***	[.17, .52]	.24(.08)	2.90**	[.08, .41]	.31(.11)	2.80**	[.09, .53]
Self-Distancing	-.15(.09)	-1.75	[-.33, .02]	-.10(.08)	-1.24	[-.25, .06]	.06(.10)	.59	[-.14, .26]
Trait Anxiety x Self-Distancing	-.23(.09)	-2.53*	[-.41, -.05]	-.20(.09)	-2.27*	[-.37, -.02]	.08(.11)	.72	[-.14, .31]
Child's Age	-.06(.09)	-.65	[-.24, .12]	.03(.08)	.32	[-.13, .19]	-.02(.10)	-.15	[-.22, .19]
Social Support Size	-.08(.08)	-1.01	[-.25, .08]	-.02(.08)	-.22	[-.17, .14]	-.11(.10)	-1.05	[-.31, .10]
Caregiver Self-Efficacy	-.36(.12)	-3.04***	[-.60, -.12]	-.09(.11)	-.84	[-.30, .12]	-.13(.14)	-.95	[-.41, .14]

Note. *N* = 96 for anticipatory anxiety, *N* = 92 for three-month longer-term distress, *N* = 92 for three-month avoidance.

*
p < .05

**
p < .01

p < .001