Single-cell RNAseq reveals cell adhesion molecule profiles in electrophysiologically defined neurons

Csaba Földy^{a,b,1}, Spyros Darmanis^c, Jason Aoto^{a,d}, Robert C. Malenka^e, Stephen R. Quake^{c,f}, and Thomas C. Südhof^{a,f,1}

^aDepartment of Molecular and Cellular Physiology, Stanford University, Stanford, CA 94305; ^bBrain Research Institute, University of Zürich, 8057 Zurich, Switzerland; ^cDepartment of Bioengineering, Stanford University, Stanford, CA 94305; ^dDepartment of Pharmacology, University of Colorado Denver, Aurora, CO 80045; ^eNancy Pritzker Laboratory, Stanford University, Stanford, CA 94305; and ^fHoward Hughes Medical Institute, Stanford University, Stanford, CA 94305

Contributed by Thomas C. Südhof, July 10, 2016 (sent for review May 21, 2016; reviewed by Thomas Biederer, Tamas F. Freund, and Li-Huei Tsai)

In brain, signaling mediated by cell adhesion molecules defines the identity and functional properties of synapses. The specificity of presynaptic and postsynaptic interactions that is presumably mediated by cell adhesion molecules suggests that there exists a logic that could explain neuronal connectivity at the molecular level. Despite its importance, however, the nature of such logic is poorly understood, and even basic parameters, such as the number, identity, and single-cell expression profiles of candidate synaptic cell adhesion molecules, are not known. Here, we devised a comprehensive list of genes involved in cell adhesion, and used single-cell RNA sequencing (RNAseq) to analyze their expression in electrophysiologically defined interneurons and projection neurons. We compared the cell type-specific expression of these genes with that of genes involved in transmembrane ion conductances (i.e., channels), exocytosis, and rho/rac signaling, which regulates the actin cytoskeleton. Using these data, we identified two independent, developmentally regulated networks of interacting genes encoding molecules involved in cell adhesion, exocytosis, and signal transduction. Our approach provides a framework for a presumed cell adhesion and signaling code in neurons, enables correlating electrophysiological with molecular properties of neurons, and suggests avenues toward understanding synaptic specificity.

synapse | cell adhesion | single cell | RNAseq

The brain's "connectivity code" is thought to confer exquisite specificity to brain circuits by dictating connectivity between different types of neurons. Although its existence has not yet been conclusively demonstrated, synaptic cell adhesion molecules likely comprise a large part of such a code (1–4). Cell adhesion molecules are encoded by ~2% of the genome and play central roles in all tissues. During brain development, precisely matching presynaptic and postsynaptic cell adhesion molecule interactions likely guide synapse formation and specify the properties of synapses by activating signal transduction cascades and recruiting scaffolding molecules, receptors, and active-zone proteins; in addition, such interactions could provide structural support to synapses. However, the molecular mechanisms involved are not understood. Because cell adhesion molecule-based interactions likely code for synapse specificity in a combinatorial fashion (2, 3), an important step toward gaining insight into these molecular mechanisms is to eliminate nonfunctional possibilities, which—to a great extent—can be done by examining cell type-specific expression of these molecules.

If cell adhesion molecules dictate synapse properties, then such differences must be clearly present in interneuron and pyramidal cell classes, which have diverse synaptic properties. For example, within the hippocampal circuit, CA1 pyramidal (CA1-PYR) neurons receive convergent inputs from multiple, electrophysiologically distinct inhibitory interneurons within the hippocampus (5). Inhibitory inputs include synapses formed by fast-spiking (FS) and regular-spiking (RS) interneurons (ref. 6; for reviews, see refs. 7 and 8). We previously showed using single-cell transcriptional profiling in FS parvalbumin (PV)- and RS cholecystokinin (CCK)-containing GABAergic interneurons that

neurexin (Nrxn1 and Nrxn3; presynaptic cell adhesion molecules) isoforms were expressed cell type-specifically, with remarkable consistency in respective cell types (9). We also found that genetic deletion of neuroligin-3 (Nlgn3) (postsynaptic cell adhesion molecule) in PYR cells disabled tonic, cannabinoid type 1 receptor-mediated, endocannabinoid signaling in RS CCK synapses, but had no detectable phenotype in FS PV synapses (10). Thus, although no systematic assessment of cell adhesion molecules in GABAergic interneurons is available, previous studies established that cell adhesion molecules play a central role in controlling their properties.

Similar to their inputs, outputs of CA1-PYR cells display functional dichotomy: they primarily project to the subiculum, forming synapses on two, electrophysiologically different principal cell types: regular-spiking pyramidal (RS-PYR) and burst-spiking pyramidal (BS-PYR) cells. Analysis of these neurons is particularly difficult because, although RS-PYR and BS-PYR cells exhibit distinct electrophysiological signatures as well as dramatically different forms of long-term plasticity, no molecular markers are available to distinguish these neurons (11, 12). In examining synapse-specific mechanisms in these cells, we have shown that different forms of long-term plasticity were determined presynaptically by expression of specific neurexin isoforms in CA1-PYR cells (13–15). Together, these molecular and physiological analyses revealed specific control of synaptic properties (such as forms of LTP and endocannabinoid signaling) by cell adhesion molecules in CA1-PYR cell inputs and

Significance

Synapses functionally connect neurons in the brain and mediate information processing relevant to all aspects of life. Among others, synaptic connections are enabled by cell adhesion molecules, which connect presynaptic and postsynaptic membranes by binding to each other via the synaptic cleft. Mammalian genomes express hundreds of cell adhesion molecules whose combinatorial utilization is thought to contribute to the brain's "connectivity code." Such code could explain the versatility of synapses as well as the logic of connectivity between cell types. Here, we used single-cell RNA sequencing to analyze the expression of cell adhesion molecules and other signaling proteins in defined cell types, and found developmental patterns that potentially identify relevant elements of the connectivity code.

Author contributions: C.F., S.D., J.A., R.C.M., S.R.Q., and T.C.S. designed research; C.F., S.D., and J.A. performed research; C.F. analyzed data; and C.F. and T.C.S. wrote the paper.

Reviewers: T.B., Tufts University; T.F.F., Institute of Experimental Medicine; and L.-H.T., Massachusetts Institute of Technology.

The authors declare no conflict of interest.

Data deposition: The sequence reported in this paper have been deposited in the Gene Expression Omnibus (GEO) database, www.ncbi.nlm.nih.gov/geo (accession no. GSE75386).

¹To whom correspondence may be addressed. Email: foldy@hifo.uzh.ch or tcs1@stanford.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1610155113/-DCSupplemental.

outputs, and raised the question whether other synaptic properties were also controlled by cell adhesion molecules. Because there are a large number of molecules with such potential, a logical step to this direction is the cell type-specific analysis of cell adhesion-related gene expression.

Here, we combined electrophysiology and single-cell RNA sequencing (RNAseq) to identify cells in the pathway involving hippocampal FS interneuron (FS-INT), RS-INT, CA1-PYR cells, and subiculum RS-PYR and BS-PYR cells, and to analyze their gene expression profiles. Our data represent an initial circuit-level single-cell RNAseq analysis from synaptically and electrophysiologically defined neurons. We find surprising differences in the total number of expressed genes among neuron types and show that hippocampal neurons can be characterized by the expression of two common, developmentally regulated gene networks comprising shared cell adhesion and signaling molecules. Moreover, we demonstrate that each type of electrophysiologically defined neuron expresses a separate set of candidate synaptic cell adhesion and signaling molecules. Finally, we extended these analyses to the two types of subiculum PYR neurons that feature highly similar transcriptomes but express a limited number of unique markers, including cell adhesion-related genes, which could potentially be used in future analyses. In this manner, the approaches and data developed here lay the foundation for a biologically relevant analysis of how cell type-specific neuronal gene expression may sculpt neural circuits during development and beyond.

Results

Cell Adhesion Molecules in the Hippocampus. As a starting point, we examined the transcriptome of the entire hippocampus (Fig. 1A) at five developmental stages (in triplicate at postnatal days P0, P7, P14, P21, and P28). In these samples, the total number and distribution of expressed genes were similar (Fig. 1B and C, and Fig. S1). To specifically analyze cell adhesion molecules, we created a comprehensive list of candidate genes involved in, or related to, transsynaptic cell adhesion (collectively referred to as CAMs, for candidate cell adhesion-related molecules). For this, we first considered all molecules with single transmembrane domains (a general, although not unique property of membrane surface adhesion molecules) and narrowed this list down to 406 genes based on preexisting data (Fig. S1C). This curated list of candidate cell adhesion molecules includes proteins implicated in cell–cell signaling but excludes, for example, intracellular signal transducers linked to cell adhesion signaling. We found these genes to be broadly represented in all replicates and developmental stages (Fig. 1D), highlighting diversity of CAMs in the hippocampal circuit and throughout development. Expression profiles (Fig. 1E) for each developmental stage included genes that were consistently highly expressed at all developmental stages [for example, amyloid precursor protein (App), contactins (Ctnn), neuroligins (Nrxn), and protein-tyrosine phosphatases (Ptpn); Fig. S1D], as well as genes with large changes during development (for example, Ncam1 and Sdc3). Normalization of individual gene

Fig. 1. RNAseq of the total hippocampus during development: analysis of the changing landscape of CAMs. (A) Experimental design and strategy for mRNA extraction from hippocampal tissue samples (represented in photograph) and RNAseq analyses. (B and C) Similar number of detected genes in our analyses of three sample replicates at five developmental stages (from P0 to P28) suggested replicability, and similar normalized gene count distributions (i.e., histogram of gene numbers with given normalized gene count) suggested comparability between samples. (D) Number of detected CAMs was consistent across developmental stages. (E) Averaged expression of CAMs in five developmental stages (in the plot, each gene is spaced equidistant and represented by mean $\pm$ SEM; expression values are connected with lines for visibility; for numerical values, see Dataset S1). (F) Heat plot of normalized gene expression shows developmental regulation of CAMs, which were ranked by linear fit of developmental expression values. (G) Average gene expression levels grouped by peak expression throughout development. Label Insets detail number of genes that belong to each group. Averaged data represent mean $\pm$ SEM.

expression levels also revealed an average change of 60% between P0 and P28 (Fig. 1 *F* and *G*).

Single-Cell RNAseq from Electrophysiologically Identified Cells. To analyze gene expression in specific cell types in a defined circuit, we developed a method—which was also recently introduced by others (16, 17)—with which we first characterized neurons electrophysiologically, and subsequently analyzed their transcriptome by aspiration of cytosol followed by single-cell RNAseq. Using this approach, we examined FS and RS interneurons that directly inhibit CA1-PYR cells, as well as the target CA1-PYR cells for these interneurons. We patched FS and regular-firing interneurons located within, or in close proximity to, the pyramidal cell layer in wild-type mice, and then used paired recordings by simultaneously patching PYR cells to test whether presynaptic action potentials (APs) evoked inhibitory postsynaptic responses, characterizing these cells as FS or RS inhibitory interneurons (Fig. 2*A*). Note that such distinction did not necessarily identify individual cell types. For instance, FS cells could include PV+ basket, bistratified, and axoaxonic cells, whereas RS cells could include CCK+ basket or bistratified cells as well. At the end of the recordings, we collected the neuronal cytosol via pipette aspiration for RNAseq (Fig. 2*A*). Upon alignment of sequencing reads and assignment of gene counts for each gene, we applied stringent quality control metrics to each cell (Fig. S2 *A* and *B*). RNAseq data obtained in this manner from 18 RS-INT, 9 FS-INT, and 14 CA1-PYR cells passed quality control.

To gain insight into the molecular identity of recorded cells, we examined expression of genes that had been previously associated with RS-INT, FS-INT, or PYR cells (Fig. 2*B*). As expected, *Gad1* and *Gad2* were highly expressed in interneurons but not in PYR cells, consistent with their respective neurotransmitters. Although CCK peptide is a unique marker for RS CCK cells, we found that the CCK gene was nonspecifically expressed in all three cell types. Similarly, although cannabinoid type-1 receptor (*Cnr1*) is most highly expressed in RS CCK cells (for specific examples, see refs. 10, 18, and 19; for review, see ref. 7), it was also produced in FS-PV and PYR cells. In contrast, *Htr3a* and PV (parvalbumin) were more specific to RS-INT and FS-INT cells, respectively. Although the latter is a unique marker for FS subtypes at the protein level, the corresponding mRNA was detected in two RS-INT and three PYR cells as well. To identify PYR cells, we examined *Camk2a* and *Neurod6* expression: although the former's expression was rather nonspecific, *Neurod6* was a reliable predictor of PYR cell identity. We also found that *Grik3* (kainate receptor *GluR7*) uniquely, but not exclusively, identified FS-INT cells.

Examining overall gene expression, we found that interneurons consistently expressed almost twice as many genes as CA1-PYR cells (Fig. 2*C*; see Fig. S2*F* for controls), and that detection of individual genes was also more consistent in RS- and FS-INT cells than in CA1-PYR cells (Fig. 2*D*). A potential explanation for the latter observation involves spatial gene expression gradients in the hippocampus that likely underlie heterogeneity of CA1 pyramidal neurons (20).

Molecular Correlates of Electrophysiological Properties. To link single-cell transcriptional profiles to functional properties, we examined expression of 140 voltage-gated ion channels (see *Materials and Methods* for gene set assembly; Fig. 3*A*, Fig. S3*A*, and Dataset S1). Because RS-, FS-, and CA1-PYR cells have distinct electrophysiological properties (Fig. 3*A*), we asked whether ion channel expression could account for their physiological differences. On average, we detected more ion channels and more consistent gene expression in interneurons than in PYR cells (Fig. 3*B* and *C*), following total transcript levels. As a pivotal example of cell type-specific gene expression, we found enrichment of Na^+ and K^+ channels in FS cells (Fig. 3*D* and *E*, Upper).

Next, we quantified electrophysiological parameters ($n = 11$; Fig. 3*A* and *E*, Left) because their correlation with gene expression patterns could reflect characteristic, cell type-specific properties in these cells. We used unsupervised clustering and plotted genes/electrophysiology parameters with at least one correlation value of greater than 0.35 or less than -0.35 (arbitrary threshold to eliminate no or minimal correlations; Fig. 3*E*). Here, we found that AP firing rate, threshold, symmetry, and afterhyperpolarization, parameters that typically exhibit high values in FS cells, correlated with high expression of Na^+ and K^+ channels (Fig. 3*E*), including *Kcnc1* (Kv3.1) and *Kcnc2* (Kv3.2). Furthermore, we found that expression of these genes inversely correlated with AP peak-to-trough and width times (i.e., larger gene expression values correspond to lower electrophysiological property values, and vice versa), which as properties also correspond to the fast signaling characteristics of FS-INT cells. These results are not unexpected as expression of *Kcnc1* and *Kcnc2* is a hallmark of FS PV interneurons (refs. 21 and 22; for review, see ref. 23), validating the single-cell RNAseq analysis.

Surprisingly, unbiased correlation analysis can also deliver seemingly contradictory results. For example, the observation of high *Hcn1* expression but a low hyperpolarization sag in FS-INT is puzzling because a primary readout for *Hcn*-mediated I_h currents is the sag amplitude. However, the correlation between hyperpolarization sag and I_h current is not absolute, nor the involvement of *Hcn1* in modulating active membrane properties (24). In addition, we found significantly higher expression of *Kcnc3*, *Scn1a*, and *Trpc6* (Fig. 3*E*) in FS cells (described for PV

Fig. 2. Single-cell RNAseq in situ of electrophysiologically characterized neurons in identified microcircuits. (*A*) Experimental design and strategy. Single-cell mRNA samples were collected after paired electrophysiological recordings from RS-INT, FS-INT, and CA1-PYR cells. White box indicates the CA1 region. After sequencing, transcripts were aligned to the mm10 assembly. (*B*) Representative heat map shows examples of gene expression in each cell type. For example, *Gad1* and *Gad2* expression in most presynaptic FS-INT and RS-INT cells confirmed that they were GABAergic, whereas *Neurod6* was specific to postsynaptic PYR cells. (*C*) Consistency of gene expression within and between cell types. On average, RS-INT and FS-INT cells expressed twice as many genes as CA1-PYR cells. (*D*) Plot of the percentage of genes that were detected in less (1–50%) and more (50–100%) than one-half of the cells (data points were connected for visualization) in respective cell types. RS-INT and FS-INT cells had more consistent gene expression profiles than CA1-PYR cells. Averaged data in *C* represent means $\pm$ SEM.

Fig. 3. Combined electrophysiological and transcriptional analysis. (A) Ion channels define neuronal excitability that can be quantified using active (AP-related) and passive electrophysiological properties (e.g., voltage response for hyperpolarizing and depolarizing current injections). These parameters differ between RS-INT, FS-INT, and CA1-PYR cells, and the lower panel shows differences in AP firing in the three cell types. (B) Number of detected ion channels was similar in RS-INT and FS-INT cells, but lower in CA1-PYR cells. (C) Consistency of ion channel expression was similar in all three cell types. (D) Averaged, single-cell ion channel expression in RS-INT ($n = 14$), FS-INT ($n = 9$), and CA1-PYR ($n = 14$) cells, and in aged-matched P21 tissue control (for numerical values, see [Dataset S1](#)). (E) Correlation analysis of electrophysiological parameters and ion channel expression in RS-INT, FS-INT, and CA1-PYR cells. (Upper) Normalized gene expression data and (Left) electrophysiology parameter measurements for RS-INT, FS-INT, and CA1-PYR cells. Averaged data represent mean $\pm$ SEM. Asterisks represent significant differences between cell types (*difference between two cell types; **difference between two-to-two cell types; ***difference between all three cell types).

cells in refs. 25–27, respectively), whereas specific expression of Kcng4 (Kv6.3, Kv6.4) identifies a thus far uncharacterized component of the ion channel repertoire of FS cells. For further clues about molecular architecture of these cell types, see expression of ligand-gated ion channels in [Fig. S3 C–E](#).

Together, these combined analyses extend previous single-cell studies (including refs. 22 and 26–29) that used probe-based gene expression analysis, and enable comprehensive and unbiased RNAseq-based functional inferences.

CAMs in Single Cells. Our analyses in whole-tissue preparations (Fig. 1) consistently detected transcripts encoding 383.2 ± 1.1 (mean $\pm$ SEM) different CAMs in the hippocampus, which comprises a large variety of neuronal and nonneuronal cell types. In examining expression of CAMs at the single-cell level, we focused on how many CAMs were expressed in a neuron, and whether these were expressed with cell type specificity. In single RS-INT, FS-INT, and CA1-PYR cells, we detected $N_{\text{RS-INT}} = 121.6 \pm 11.2$, $N_{\text{FS-INT}} = 128.7 \pm 12.7$, and $N_{\text{CA1-PYR}} = 82.7 \pm 4.8$ different CAM transcripts, respectively [$P_{\text{RS-INT vs. FS-INT}} = 0.73$, $P_{\text{RS-INT vs. CA1-PYR}} = 0.02$, $P_{\text{FS-INT vs. CA1-PYR}} = 0.006$; pairwise Mann–Whitney (MW) test; Fig. 4 A and B and [Dataset S1](#)]. Examining these genes revealed that expression of CAMs partially overlapped between different cell types. There were $N_{\text{RS-INT vs. FS-INT}} = 48$, $N_{\text{RS-INT vs. CA1-PYR}} = 58$, and $N_{\text{FS-INT vs. CA1-PYR}} = 62$ differentially expressed CAMs between cell types (pairwise MW test, $P < 0.05$ in all cases). Thirty CAMs were expressed higher ($n = 26$; MW test, $P < 0.05$ in all cases) or lower ($n = 4$; Eph4, Mdga1, Pcdhgc5, Sema3e; MW test, $P < 0.05$

in all cases) in both interneuron types compared with PYRs. We found that genes that were highly expressed in tissue samples were also highly expressed in the RS-INT, FS-INT, and CA1-PYR cells (note that for sequencing libraries, tissue samples were diluted and processed using same conditions and reagents as for single cells, albeit with larger amounts of starting mRNAs). Such genes included App, Chl1, Cntn1, Nptn, Nrnx1, and Ptpns that were consistently detected in all three cell types (Fig. 4C, “All cell types”).

The 26 CAMs that were expressed higher in interneurons than in CA1-PYR cells included Cln3, Cntnap4, and Lrrc4, which have been implicated in specification of inhibitory synapses (30–32), as well as Nrnx3, a presumed presynaptic organizer of synapse function (13, 14, 33). Note that, although expression of Nrnx3 was not specific for interneurons, it was consistently enriched in interneurons, which we also considered here as a cell type-specific feature. Other examples for cell type specificity included Cbln2 and Ephb2 for RS-INT cells, Nphx1 for FS-INT cells, and Eph4 for CA1-PYR cells. Although some of these molecules have been directly implicated in trans-synaptic signaling (e.g., Nrnx1, Nrnx3, Ptpns) or in synapse specialization (see examples above), others have not been studied in detail. Nevertheless, these data suggest cell type-specific expression of CAMs can potentially explain neuronal connectivity at the molecular level. Such cell type-specific features were further highlighted by principal-component axes (PCA) analysis, which revealed two main components, including overall expression as well as differential expression in interneurons vs. PYR cells (Fig. S4 A–C).

Fig. 4. Hierarchy of CAMs in single neurons. (A) Averaged, single-cell expression of CAMs in RS-INT, FS-INT, and CA1-PYR cells, and in age-matched P21 tissue control (for numerical values, see [Dataset S1](#)). (B) Number of CAMs detected in the three cell types and in age-matched tissue control. (C) Heat plot shows single-cell expression for ubiquitously ("All cell types") as well as cell type-specifically expressed CAMs. Statistical differences were determined using MW test, with $P < 0.05$. Averaged data in *B* represent mean $\pm$ SEM.

Beyond cell type-specific control of gene expression levels, diversity in CAM signaling could have been generated by other cellular mechanisms, especially by alternative splicing. Some of the most notable examples for alternative splicing include CAMs, such as *Dscam* in *Drosophila* and neuexins in mammals, which can express thousands of alternatively spliced isoforms (34–37). Therefore, we examined alternative exon use of CAMs at the single-cell level. We analyzed genes with reliable end-to-end exon junction coverage, which was applicable to $n = 139$ CAM genes. Out of these, we identified a single isoform in $n = 67$ and multiple isoforms in $n = 72$ genes (Fig. S4 D and E). Because of their potential cell type specificity, presence of different isoforms in the latter group would be especially interesting. However, currently available single-cell RNAseq platforms—including ours—require library fragmentation. As a result, precise exon use of full-length mRNAs can be inferred only indirectly, with sample sizes typically larger than those of single cells, hindering further analyses of cell type-specific splice variants. Nevertheless, we examined whether any splice variants of the highly expressed ubiquitous CAMs had the potential to be expressed with cell type specificity. We found that some of these molecules had only single isoforms that may or may not involve alternative splicing (for example, in *Ptprn*, we consistently detected exon skipping, whereas in *Ptprn* and *Cadm*, we only detected canonical isoforms), whereas others had multiple isoforms (for example, in *Nptn*, *Nrxn1*, and *Cd47*; Fig. S4F). Together, these exon junction observations suggested further diversification of cell adhesion signaling in single

cells, which—at least regarding neuexins—was also suggested in our previous findings (9).

Synaptic Vesicle Exocytosis and Intracellular Signal Transduction.

One presumed role of CAMs is to convey transsynaptic information and thereby establish a functional match between presynaptic and postsynaptic specializations. However, neither the precise molecular targets nor the underlying mechanisms have been described for most CAMs. Thus, we analyzed expression of two broadly defined groups of genes that are likely linked to cell adhesion function, namely genes related to vesicle exocytosis and to RhoGAP/RhoGEF signaling (see detailed list in Fig. S5A and [Dataset S1](#)). Exocytosis-related molecules (38) were expressed broadly in all single cells and at all developmental stages (Fig. 5 A and B, and Fig. S5B). In single cells, we detected both ubiquitous and cell type-specific features, where the former included *Snap25*, *Synaptobrevin-2* (*Vamp2*), *Snap47*, and *Syt11* (for which no function has been uncovered yet, despite its high abundance), whereas *Cplx1*, *Stx1a*, *Syt13*, and *Synaptobrevin-1* (*Vamp1*) were expressed higher in interneurons (MW test, $P < 0.05$, for all genes; expression of *Synaptobrevin-1* also appeared to be exclusive, as it was not detected in any PYRs; Fig. 5 A and B, and [Dataset S1](#)). In addition, we found exclusive expression of *Sncg* and *Syt6* genes in RS-INT cells, with unknown functional consequences.

Next, we analyzed the expression of RhoGAP- and RhoGEF-domain-containing and related proteins. These are membrane-associated proteins that are thought to transform extracellular receptor-mediated signals into an intracellular response, prominently

First, we independently examined gene correlations in tissue and single-cell data, and used unsupervised clustering to consolidate these results (Fig. 6*A* and *B*), which reflected developmental and cell type-specific characteristics, respectively (Fig. S64). We reasoned that functionally relevant correlations may be present in both sets with high correlation coefficients, and therefore selected gene pair correlations that were in the tail of distributions (>92nd percentile; note that this threshold was chosen arbitrarily but that consequences of this analysis were thoroughly tested at multiple threshold values; Fig. 6*E*) both for tissue and single-cell data (Fig. 6*C*). This step narrowed our focus to 269 genes and 692 correlations. Genes were outnumbered by correlations because some genes correlated with multiple others (i.e., they represented correlation “hubs”). We used graph analysis to test such possibility. In such graphs, each node represents a gene and each vertex represents correlation between two genes, if any, with a length inversely proportional to the respective coefficient; genes with more correlations simply had more vertices connecting them.

Next, we sought to further examine graph structures using added knowledge about identity of single-cell samples. We reasoned that robust gene expression correlations should be consistently present if cell types were analyzed independently. Therefore, we repeated analyses such that Fig. 6B only included RS-INT, or FS-INT, or CA1-PYR cells separately, or RS-INT

Fig. 6. Coregulation and coexpression of synaptic molecules. (A) Heat plot of correlations between genes representing cell adhesion-, exocytosis-, RhoGAP-, and RhoGEF-related molecules in hippocampal tissue. Coefficients were computed based on gene expression values in five developmental stages in tissue samples. Unsupervised clustering revealed 12 clusters (labeled on the *Right* and *Top*) based on 395,641 gene correlations. (B) Same analysis as in A, but for single-cell data. Coefficients were computed based on single-cell gene expression values independent of cell type identity. Unsupervised clustering revealed five clusters (labeled on the *Right* and *Top*) based on the 187,489 gene correlations. (C) Combined data of correlation coefficients from tissue and single-cell samples. Gene pairs with both coefficients larger than the 92nd percentile of the respective distributions were further analyzed (green shaded area; see text for further information). (D) Graph representation of correlation coefficients displayed in the green shaded area of C. Each vertex connects two genes according to the correlation coefficient between the two. Graph representation of correlations in C revealed two independent, uncorrelated subgraphs. (E, *Left*) Subgraph 1 and subgraph 2 remained independent when relaxing gene inclusion criteria (i.e., when green area in C was gradually increased). (*Right*) Density of vertex distribution (number of gene–gene correlations) followed power law distribution, indicating that both subgraphs were scale-free and nonrandom in nature. (F, *Left*) Mean normalized gene expression values in the two subgraphs showed diametrically opposite developmental trajectories, and suggest that genes in subgraph were developmentally coregulated. Because single-cell expression was a selection criterion in C, these genes were also coexpressed at the single-cell level. (*Right*) As suggested by the developmental differences, genes in subgraph 1 and subgraph 2 had different expression values in single cells of all three cell types, collected at ~P21. (G) Core early- and late-gene networks (subgraph 1 and subgraph 2, respectively) were identified by common motifs found in cell type-specific analyses (i.e., these correlations were present in pooled data from the three cell types, in pooled data from RS-INT and FS-INT cells, representing interneurons, and RS-INT, FS-INT, and CA1-PYR data, analyzed independently). Averaged data represent mean $\pm$ SEM.

and FS-INT cells combined, because they both represented GABAergic interneurons. We found that each of these analyses confirmed the independence of subgraphs 1 and 2 (Fig. S6B), and that the gene representation in each cell type-specific dataset was at least 80% identical to that in Fig. 6D (Fig. S6B), suggesting that coexpression of these genes occurs at the single-cell level. Moreover, we identified core parts of these subgraphs (i.e., parts that can be consistently detected in each cell type) by taking the intersection of the cell type-specific analyses (Fig. 6G). Because of their robust presence in each cell type, these correlations conceivably represented ubiquitous features, at different developmental stages of synapse maturation. This hypothesis was supported by the existence of correlations between structurally relevant RhoGEF and RhoGAP molecules in early development, when synapses are formed (Fig. 6G, *Left*, and Fig. S6D, *Upper*). Conversely, emerging correlations between cell adhesion and exocytosis in the late-gene network may reflect synapse specialization (Fig. 6G, *Right*, and Fig. S6D, *Lower*). Although our analyses did not reveal a mechanistic explanation for the observed correlations, they identified synaptic genes that were both developmentally coregulated and coexpressed at the single-cell level.

CAMs in BS-PYR and RS-PYR Cells of the Subiculum. Our findings thus far revealed ubiquitous and cell type-specific features of CAM and signaling molecule expression as well as the contextual determinants of their expression, involving molecules related to exocytosis and Rho signaling. Next, we wanted to examine the generality of our central findings, including that of (i) the identity of ubiquitously expressed CAMs and of (ii) the existence and identity of synaptic early- (subgraph 1) and late-gene interaction

networks (subgraph 2). To this end, we performed single-cell RNAseq experiments to analyze electrophysiologically defined BS-PYR and RS-PYR neurons in the subiculum (Fig. 7A), which are the major projection targets of CA1-PYR cells. We hypothesized that their gene expression profiles are different, because they exhibit distinct excitability and synaptic properties (12, 13, 15).

We detected $9,671 \pm 278$ and $9,499 \pm 405$ genes in BS-PYR and RS-PYR cells, respectively ($P = 0.85$; Fig. 7B and Fig. S7A), with more consistent gene expression patterns than among CA1-PYRs (Fig. 7C and Fig. S7B). Next, we examined the molecular profiles of these cells, especially seeking exclusive markers that would unequivocally identify these functionally different cell types. However, we found that they were largely identical and detected only a surprisingly small number of exclusively expressed genes (Fig. 7D). Such genes included, for example, the neuropeptide cortistatin (Cort) and the c-Fos-induced growth factor (Figf) (a member of the VEGF family). Apart from these genes, BS-PYR and RS-PYR neurons expressed similar patterns of voltage-gated ion channels (Fig. 7E; see differently expressed ion channels in Fig. S7C), CAMs (Fig. 7F), ligand-gated ion channels, and exocytosis- and RhoGAP-/RhoGEF-related molecules (Fig. S7 C–G and Dataset S1).

We next examined expression of ubiquitously present CAMs (as identified in CA1 interneurons and PYR cells), which we reliably detected in subiculum PYR cells similar to CA1-PYRs (Figs. 4C and 7G), further strengthening the notion of general importance of these genes in neuron and synapse function. Although 42 CAMs had different expression levels between the two subiculum PYR cell types ($P < 0.05$; pairwise MW test; Dataset S1), none of these CAMs was exclusively expressed in one or the other cell type. However, their expression differed markedly from that of

Finally, the subiculum neuron data allowed us to test the validity of synaptic gene correlation networks described in Fig. 6.

PNAS | Published online August 16, 2016 | E5229

Discussion

A molecular description of gene expression in individual, independently identified neurons in combination with knowledge of their physiological properties and synaptic connectivity is a prerequisite for insight into how synaptic connections are formed and maintained in brain. Moreover, such data are essential for understanding why disease-associated gene mutations impair brain function. RNAseq of single neurons can be routinely obtained from cells isolated from tissue lysates (44–47) and has recently been achieved from electrophysiologically recorded cells (the current study; also see refs. 16 and 17). Here, we used single-cell RNAseq of individual synaptically and electrophysiologically characterized neurons, and cross-referenced their gene expression profiles to that of total brain tissue to explore cell adhesion signaling in a well-defined hippocampal circuit. In particular, we examined inhibitory projections from RS and FS interneurons to CA1-PYR cells (6, 7, 10) and excitatory projections from CA1-PYR cells to subiculum BS-PYR and RS-PYR cells (13, 15). Our results show that capture of mRNAs by aspiration of cytosol allows production of high-quality transcriptome data (Figs. S2 and S7), opening up identification of the precise expression profiles of single neurons as a molecular portal for a detailed functional understanding.

Functional implications of gene expression can be most directly examined by relating gene expression to electrophysiological properties (16, 17, 26–29, 48). Therefore, we first tested whether the electrophysiological properties of a neuron could be explained by its ion channel expression pattern. Previous studies established that expression of Kv3 K⁺ channels and a high Na⁺-channel density enable the FS properties of PV cells (23). In line with this conclusion, we found specific enrichment of Kcnc1 (Kv3.1), Kcnc2 (Kv3.2), and Scn1a (NaV1.1) in FS-INT cells. By taking advantage of our comprehensive gene expression data, we extended these findings and correlated 11 electrophysiological properties with 104 ion channel genes. We found strong correlations between specific sets of genes and physiological properties, consistent with the different cell types. Although these correlations may mostly reflect group-level differences, an added value of these analyses is the conclusion that electrophysiological properties can be identified using single-cell transcriptome information. More detailed quantification of electrophysiological properties in combination with extended sample sets and morphological analyses should, for example, allow differentiation of interneuron subtypes (5) on a molecular basis. In addition, these results suggest that the validity of RNAseq data in electrophysiologically defined cells extends to genes whose functional readouts were not readily available, as is the case for CAMs.

We hypothesized that combinatorial expression of CAMs in single cells may define connectivity in a neuronal network. A corollary of this hypothesis is that CAMs in a presynaptic terminal must interact with complementary CAMs in the postsynaptic compartment; this interaction is—through reciprocal ligand–receptor coupling—enabled via the synaptic cleft. Abundant presence of CAMs in hippocampal tissue (Fig. 1) supports this hypothesis, because the diversity of CAMs likely permits a large number of combinations, possibly defining synaptic connectivity between the more than 20 cell types in the hippocampus and beyond (5). Our data present the most current account of CAMs in the brain, described by expression analysis of 406 genes. A limitation of our RNAseq results is that a transcriptome profile represents the abundance of mRNAs in a given cell type but does not provide information regarding protein abundance or localization. A given neuron receives heterogeneous inputs (excitatory and inhibitory synapses from multiple sources) and can innervate multiple postsynaptic cells of differing identities. Thus, it is plausible that CAMs are differentially localized and combinatorially used in a synapse-specific manner. To understand the organization of these molecules and possibly interpret them as a code for connectivity, we need to clarify which of these molecules are expressed in specific synapses and physically interact with each other. An important step to this direction is to examine cell type-specific expression of these molecules.

Our hypothesis suggests that expression of CAMs should differ among cell types, including RS-INT, FS-INT, CA1-PYR, BS-PYR,

and RS-PYR cells, in which we previously established functionally relevant differences in cell adhesion signaling (9, 10, 13, 15). Thus, we examined candidate synaptic CAMs in single cells from these types. Presynaptic and postsynaptic CAMs presumably engage in homophilic and heterophilic interactions in the synaptic cleft, but the rules that associate particular CAMs to specific synapses and their transcriptional regulation in different cell types were not known. In examining the cell type-specific expression of CAMs, we made two fundamental observations (Figs. 4 and 7).

First, CAMs are hierarchically expressed. This conclusion is based on the observation that some CAMs were consistently and highly expressed in all five cell types examined (these included Nptn, Nrnx1, App, Ptpr, Nrcam, Ppfia2, Chl1, Cntn1, Ptpn, and Bsg), whereas others were expressed cell type-specifically (these included, for example, Cntnap4, Fstl5, Igsf8, Lrrc4, Cbln2, Eph2b, Idgc4, Nxph1, Eph4, and Sema3e). The most likely explanation for such hierarchy is that commonly expressed molecules are required for core synaptic features as well as other neuronal cell interactions, whereas cell type-specific molecules may enable functional specialization. For example, it is intriguing to speculate which, if any, of these molecules are transsynaptically involved in sculpting endocannabinoid signaling in RS CCK or opioid signaling in FS PV synapses, respectively (6, 7, 10). Overall, these data support our hypothesis that CAMs are combinatorially expressed at the single-cell level.

Second, we identified synaptic genes that were both developmentally coregulated as well as coexpressed in single cells (Figs. 5–7). This was made possible by a combined analysis of tissue and single-cell data, and revealed two independent gene networks: an “early-gene” and a “late-gene” network. Beyond developmental differences, molecular composition of the two networks implicated distinct functional relevance. Specifically, interactions between intracellular signal transducer molecules were enriched in the early-gene cluster, whereas representation of exocytosis-related molecules as well as their interactions with CAMs were more apparent in the late-gene cluster. Such findings are consistent with the developmental sequels of structural formation and functional maturation of synapses and neuron–glia interactions. Because they are comprehensive in nature, experimental probing of these networks is challenging as it may require simultaneous manipulation of multiple genes. Such approach, targeting multiple heavily connected genes or hubs (e.g., Clstn1, Clstn3, or Igsf8), could reveal susceptibility to genetic conditions with especially disruptive pathophysiological consequences.

Together, our results provide a description of which electrophysiologically and synaptically characterized neurons express which specific CAMs. Some of these molecules are already known to play important synaptic roles, as their genetic disruptions have been linked to neuropsychiatric disorders in humans and have been shown to be lethal in mice (33). However, the function of the majority of these molecules as well as the logic that integrates them into assembly and specification of synapses or other intercellular junctions remain unknown. Here, we made progress toward understanding such logic by measuring developmental and single-cell expression of CAMs. When revealed, underlying principles can help us to use the brain’s cell adhesion code to formally describe connectivity in neuronal circuits. Such applications would lead to developments in molecular diagnostics allowing comprehensive analysis of brain circuits from single-cell samples.

Materials and Methods

Electrophysiology and Single-Cell Sample Collection. All animal protocols and husbandry practices were approved by the Institutional Animal Care and Use Committee at Stanford University. Hippocampal slices (300 μ m) were prepared from 3- to 4-wk-old wild-type CD1 mice, as described in ref. 10 and *SI Materials and Methods*.

cDNA Preparation and Library Preparation for Next-Generation Sequencing. Single-cell mRNA was performed using the Clontech’s SMARTer Ultra Low Input RNA Kit. As a first step, cells were collected via pipette aspiration into sample collection buffer, were spun briefly, and were snap frozen on dry ice.

Samples were stored at -80°C until further processing, which was performed according to manufacturer's protocol. Library preparation was performed using Nextera XT DNA Sample Preparation Kit (Illumina) as described in the protocol. Then, cells were pooled and sequenced in an Illumina NextSeq500 instrument using 2×75 paired end reads on a NextSeq high-output kit (Illumina; see *SI Materials and Methods* for details).

Processing of mRNA Sequencing Data. After de-multiplexing the raw reads to single-cell datasets, we used *Prinseq* to remove short reads. After trimming and removal of overrepresented sequences and adapters, remaining reads were aligned to the mm10 genome with *STAR* aligner. Aligned read were converted to gene counts using *HTSeq* (see *SI Materials and Methods* for detailed parameter descriptions).

Gene Categories. For gene expression analysis, we examined six functionally related categories: CAMs, voltage-gated ion channels, ligand-gated ion

channels, synaptic exocytosis-related molecules, as well as RhoGAP and RhoGEF signaling-related molecules. For each of these, we assembled a comprehensive list, which included all genes examined (see *SI Materials and Methods* for a detailed description of categories).

Data Analysis. All data analyses were performed using Mathematica10 (Wolfram Research). These analyses included (i) normalization of gene expression data, (ii) quality control, (iii) analysis of physiological properties, (iv) analysis of exon junctions, and (v) correlation and graph analyses of expression data (see *SI Materials and Methods* for more details).

ACKNOWLEDGMENTS. This work was supported by National Institute on Drug Abuse Grant K99 DA034029 (to C.F.), Swiss National Science Foundation Grant CRETP3_166815 (to C.F.), and the National Institute of Mental Health Grants R37 MH52804 (to T.C.S.) and K99 MH103531 (to J.A.).

- Sperry RW (1963) Chemoaffinity in the orderly growth of nerve fiber patterns and connections. *Proc Natl Acad Sci USA* 50:703–710.
- Koch C, Laurent G (1999) Complexity and the nervous system. *Science* 284(5411):96–98.
- Itzkovitz S, Baruch L, Shapiro E, Segal E (2008) Geometric constraints on neuronal connectivity facilitate a concise synaptic adhesive code. *Proc Natl Acad Sci USA* 105(27):9278–9283.
- de Wit J, Ghosh A (2016) Specification of synaptic connectivity by cell surface interactions. *Nat Rev Neurosci* 17(1):22–35.
- Klausberger T, Somogyi P (2008) Neuronal diversity and temporal dynamics: The unity of hippocampal circuit operations. *Science* 321(5885):53–57.
- Glickfeld LL, Atallah BV, Scanziani M (2008) Complementary modulation of somatic inhibition by opioids and cannabinoids. *J Neurosci* 28(8):1824–1832.
- Freund TF, Katona I (2007) Perisomatic inhibition. *Neuron* 56(1):33–42.
- Caroni P (2015) Inhibitory microcircuit modules in hippocampal learning. *Curr Opin Neurobiol* 35:66–73.
- Fucillo MV, et al. (2015) Single-cell mRNA profiling reveals cell-type specific expression of neurexin isoforms. *Neuron* 87(2):326–340.
- Földy C, Malenka RC, Südhof TC (2013) Autism-associated neurexin-3 mutations commonly disrupt tonic endocannabinoid signaling. *Neuron* 78(3):498–509.
- Staff NP, Jung HY, Thiagarajan T, Yao M, Spruston N (2000) Resting and active properties of pyramidal neurons in subiculum and CA1 of rat hippocampus. *J Neurophysiol* 84(5):2398–2408.
- Wozny C, Maier N, Schmitz D, Behr J (2008) Two different forms of long-term potentiation at CA1-subiculum synapses. *J Physiol* 586(11):2725–2734.
- Aoto J, Martinelli DC, Malenka RC, Tabuchi K, Südhof TC (2013) Presynaptic neurexin-3 alternative splicing trans-synaptically controls postsynaptic AMPA receptor trafficking. *Cell* 154(1):75–88.
- Aoto J, Földy C, Ilcus SM, Tabuchi K, Südhof TC (2015) Distinct circuit-dependent functions of presynaptic neurexin-3 at GABAergic and glutamatergic synapses. *Nat Neurosci* 18(7):997–1007.
- Anderson GR, et al. (2015) β -Neurexins control neural circuits by regulating synaptic endocannabinoid signaling. *Cell* 162(3):593–606.
- Cadwell CR, et al. (2016) Electrophysiological, transcriptomic and morphologic profiling of single neurons using Patch-seq. *Nat Biotechnol* 34(2):199–203.
- Fuzik J, et al. (2016) Integration of electrophysiological recordings with single-cell RNA-seq data identifies neuronal subtypes. *Nat Biotechnol* 34(2):175–183.
- Neu A, Földy C, Soltesz I (2007) Postsynaptic origin of CB1-dependent tonic inhibition of GABA release at cholecystokinin-positive basket cell to pyramidal cell synapses in the CA1 region of the rat hippocampus. *J Physiol* 578(Pt 1):233–247.
- Lenkey N, et al. (2015) Tonic endocannabinoid-mediated modulation of GABA release is independent of the CB1 content of axon terminals. *Nat Commun* 6:6557.
- Cembrowski MS, et al. (2016) Spatial gene-expression gradients underlie prominent heterogeneity of CA1 pyramidal neurons. *Neuron* 89(2):351–368.
- Du J, Zhang L, Weiser M, Rudy B, McBain CJ (1996) Developmental expression and functional characterization of the potassium-channel subunit Kv3.1b in parvalbumin-containing interneurons of the rat hippocampus. *J Neurosci* 16(2):506–518.
- Martina M, Schultz JH, Ehmk H, Monyer H, Jonas P (1998) Functional and molecular differences between voltage-gated K^+ channels of fast-spiking interneurons and pyramidal neurons of rat hippocampus. *J Neurosci* 18(20):8111–8125.
- Hu H, Gan J, Jonas P (2014) Interneurons. Fast-spiking, parvalbumin $^+$ GABAergic interneurons: From cellular design to microcircuit function. *Science* 345(6196):1255–1263.
- Aponte Y, Lien C-C, Reisinger E, Jonas P (2006) Hyperpolarization-activated cation channels in fast-spiking interneurons of rat hippocampus. *J Physiol* 574(Pt 1):229–243.
- Chang SY, et al. (2007) Distribution of Kv3.3 potassium channel subunits in distinct neuronal populations of mouse brain. *J Comp Neurol* 502(6):953–972.
- Ogiwara I, et al. (2007) Nav1.1 localizes to axons of parvalbumin-positive inhibitory interneurons: A circuit basis for epileptic seizures in mice carrying a Scn1a gene mutation. *J Neurosci* 27(22):5903–5914.
- Nagy GA, et al. (2013) DAG-sensitive and Ca^{2+} permeable TRPC6 channels are expressed in dentate granule cells and interneurons in the hippocampal formation. *Hippocampus* 23(3):221–232.
- Lambole B, Audinat E, Bochet P, Crépel F, Rossier J (1992) AMPA receptor subunits expressed by single Purkinje cells. *Neuron* 9(2):247–258.
- Toledo-Rodriguez M, et al. (2004) Correlation maps allow neuronal electrical properties to be predicted from single-cell gene expression profiles in rat neocortex. *Cereb Cortex* 14(12):1310–1327.
- Lu Z, et al. (2014) Calsyntenin-3 molecular architecture and interaction with neurexin 1 α . *J Biol Chem* 289(50):34530–34542.
- Pettem KL, et al. (2013) The specific α -neurexin interactor calsyntenin-3 promotes excitatory and inhibitory synapse development. *Neuron* 80(1):113–128.
- Karayannis T, et al. (2014) Cntnap4 differentially contributes to GABAergic and dopaminergic synaptic transmission. *Nature* 511(7508):236–240.
- Südhof TC (2008) Neuroligins and neurexins link synaptic function to cognitive disease. *Nature* 455(7215):903–911.
- Ullrich B, Ushkaryov YA, Südhof TC (1995) Cartography of neurexins: More than 1000 isoforms generated by alternative splicing and expressed in distinct subsets of neurons. *Neuron* 14(3):497–507.
- Schmucker D, et al. (2000) *Drosophila* Dscam is an axon guidance receptor exhibiting extraordinary molecular diversity. *Cell* 101(6):671–684.
- Treutlein B, Gokke O, Quake SR, Südhof TC (2014) Cartography of neurexin alternative splicing mapped by single-molecule long-read mRNA sequencing. *Proc Natl Acad Sci USA* 111(13):E1291–E1299.
- Schreiner D, et al. (2014) Targeted combinatorial alternative splicing generates brain region-specific repertoires of neurexins. *Neuron* 84(2):386–398.
- Südhof TC (2013) Neurotransmitter release: The last millisecond in the life of a synaptic vesicle. *Neuron* 80(3):675–690.
- Jaudon F, et al. (2015) The RhoGEF DOCK10 is essential for dendritic spine morphogenesis. *Mol Biol Cell* 26(11):2112–2127.
- Lai KO, Ip NY (2013) Structural plasticity of dendritic spines: The underlying mechanisms and its dysregulation in brain disorders. *Biochim Biophys Acta* 1832(12):2257–2263.
- Cheadle L, Biederer T (2012) The novel synaptogenic protein Farp1 links postsynaptic cytoskeletal dynamics and transsynaptic organization. *J Cell Biol* 199(6):985–1001.
- Van de Ven TJ, Van Dongen HM, Van Dongen AM (2005) The nonkinase phorbol ester receptor α 1-chimerin binds the NMDA receptor NR2A subunit and regulates dendritic spine density. *J Neurosci* 25(41):9488–9496.
- Vidal M, Cusick ME, Barabási AL (2011) Interactome networks and human disease. *Cell* 144(6):986–998.
- Usoskin D, et al. (2015) Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nat Neurosci* 18(1):145–153.
- Zeisel A, et al. (2015) Brain structure. Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science* 347(6226):1138–1142.
- Darmanis S, et al. (2015) A survey of human brain transcriptome diversity at the single cell level. *Proc Natl Acad Sci USA* 112(23):7285–7290.
- Hanchate NK, et al. (2015) Single-cell transcriptomics reveals receptor transformations during olfactory neurogenesis. *Science* 350(6265):1251–1255.
- Geiger JR, et al. (1995) Relative abundance of subunit mRNAs determines gating and Ca^{2+} permeability of AMPA receptors in principal neurons and interneurons in rat CNS. *Neuron* 15(1):193–204.
- Catania MV, et al. (1998) AMPA receptor subunits are differentially expressed in parvalbumin- and calretinin-positive neurons of the rat hippocampus. *Eur J Neurosci* 10(11):3479–3490.
- Milenkovic I, et al. (2013) The parvalbumin-positive interneurons in the mouse dentate gyrus express GABA $_A$ receptor subunits α 1, α 2, and δ along their extrasynaptic cell membrane. *Neuroscience* 254:80–96.
- Ferando I, Mody I (2015) In vitro gamma oscillations following partial and complete ablation of δ subunit-containing GABA $_A$ receptors from parvalbumin interneurons. *Neuropharmacology* 88:91–98.
- Gao B, Fritschy JM (1994) Selective allocation of GABA $_A$ receptors containing the α 1 subunit to neurochemically distinct subpopulations of rat hippocampal interneurons. *Eur J Neurosci* 6(5):837–853.
- Frerking M, Malenka RC, Nicoll RA (1998) Synaptic activation of kainate receptors on hippocampal interneurons. *Nat Neurosci* 1(6):479–486.