APOBEC3G expression correlates with T cell infiltration and improved clinical outcomes in high-grade serous ovarian carcinoma

Brandon Leonard^{1,2,*}, Gabriel J. Starrett^{1,2,*}, Matthew J. Maurer³, Ann L. Oberg³, Mieke Van Bockstal⁴, Jo Van Dorpe⁴, Olivier De Wever⁵, Jozien Helleman⁶, Anieta M. Sieuwerts⁶, Els M.J.J. Berns⁶, John W.M. Martens⁶, Brett D. Anderson^{1,2}, William L. Brown^{1,2}, Kimberly R. Kalli⁷, Scott H. Kaufmann^{8,9}, and Reuben S. Harris^{1,2,10}

¹Biochemistry, Molecular Biology and Biophysics Department, University of Minnesota, Minneapolis, MN 55455, USA ²Masonic Cancer Center, University of Minnesota, Minneapolis, MN 55455, USA ³Division of Biomedical Statistics and Informatics, Department of Health Sciences Research, Mayo Clinic, Rochester, MN 55905, USA ⁴Department of Pathology, Ghent University Hospital, Ghent, Belgium ⁵Department of Radiation Oncology and Experimental Cancer Research, Ghent University and Ghent University Hospital, Ghent, Belgium ⁶Department of Medical Oncology, Erasmus MC Cancer Institute, Rotterdam, Netherlands ⁷Women's Cancer Program, Mayo Clinic Cancer Center, Rochester, MN 55905 USA ⁸Division of Oncology Research, Department of Oncology, Mayo Clinic, Rochester, MN 55905 USA ⁹Department of Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic, Rochester, MN 55905, USA ¹⁰Howard Hughes Medical Institute, University of Minnesota, Minneapolis, MN 55455, USA

Abstract

Purpose—APOBEC3 DNA cytosine deaminase family members normally defend against viruses and transposons. However, deregulated APOBEC3 activity causes mutations in cancer. Due to broad expression profiles and varying mixtures of normal and cancer cells in tumors, including immune cell infiltration, it is difficult to determine where different APOBEC3s are expressed. Here, we ask whether correlations exist between *APOBEC3* expression and T cell infiltration in high-grade serous ovarian cancer (HGSOC), and assess whether these correlations have prognostic value.

Experimental Design—Transcripts for *APOBEC3G*, *APOBEC3B*, and the T cell markers, *CD3D*, *CD4*, *CD8A*, *GZMB*, *PRF1*, and *RNF128* were quantified by RT-qPCR for a cohort of 354 HGSOC patients. Expression values were correlated with each other and clinical parameters. Two additional cohorts were used to extend HGSOC clinical results. Immuno-imaging was used to colocalize APOBEC3G and the T cell marker CD3. TCGA data extended expression analyses to additional cancer types.

Address correspondence to: Reuben S. Harris, Howard Hughes Medical Institute, University of Minnesota, 2231 6th St. S.E., Minneapolis, MN 55455, USA. Phone: 612.624.0457; Fax: 612.625.2163; rsh@umn.edu.

*These authors contributed equally to this work

Conflict of Interest Statement: R.S.H. is a co-founder of ApoGen Biotechnologies Inc. The other authors have no conflicts of interest to disclose.

Results—A surprising positive correlation was found for expression of *APOBEC3G* and several T cell genes in HGSOC. Immunohistochemistry and immunofluorescent imaging showed protein colocalization in tumor-infiltrating T lymphocytes. High *APOBEC3G* expression correlated with improved outcomes in multiple HGSOC cohorts. TCGA data analyses revealed that expression of *APOBEC3D* and *APOBEC3H* also correlates with *CD3D* across multiple cancer types.

Conclusions—Our results identify *APOBEC3G* as a new candidate biomarker for tumor-infiltrating T lymphocytes and favorable prognoses for HGSOC. Our data also highlight the complexity of the tumor environment with respect to differential *APOBEC* family gene expression in both tumor and surrounding normal cell types.

Keywords

APOBEC3G; immune cell infiltration; ovarian cancer; ovarian tumor heterogeneity; prognostic biomarker

Introduction

Ovarian cancer is the deadliest gynecological malignancy worldwide (1). The most common type of ovarian cancer, high-grade serous ovarian carcinoma (HGSOC), accounts for over 60% of cases, and is the most aggressive reproductive track malignancy (2). Due to the lack of efficient detection methods, HGSOC generally presents at advanced stages and is associated with high rates of recurrence and mortality. Interestingly, several studies have identified T cell infiltration as a favorable prognostic factor for HGSOC (3–7). Additional markers of T cell infiltration and particularly those reflecting high-quality anti-tumor responses are needed to fully realize the clinical impact of this finding.

The APOBEC enzymes are an 11-member family of zinc-coordinating enzymes that convert cytosines to uracils (C-to-U) in ssDNA (8). The enzymatic activity of specific family members is essential for both adaptive (AID) and innate immune responses [APOBEC3 subfamily members; (9) and references therein]. AID is well-studied due to its known roles in antibody diversification through somatic hypermutation and class switch recombination in B cells (10). Other well-studied family members include APOBEC3D, APOBEC3F, APOBEC3G, and APOBEC3H because of demonstrated anti-HIV-1 activity (11). These enzymes are capable of introducing C-to-U lesions in viral cDNA intermediates that manifest as G-to-A mutations in proviral genomes (12–14). APOBEC3 subfamily members have also been implicated in restricting the replication of many other DNA-based parasites including transposable elements [(9) and references therein]. It is important to note that most APOBEC family members are expressed broadly and constitutively (12,15–17), but several also can be further upregulated by specific conditions such as *APOBEC3A* by interferon- α (18,19), *APOBEC3B* by non-canonical NF- κ B activation and HPV infection (20–22), and *APOBEC3C*, *APOBEC3D*, *APOBEC3F*, *APOBEC3G*, and *APOBEC3H* by the combination of T cell activation and HIV-1 infection (13,14).

While the APOBEC3 enzymes have been shown to have important physiologic roles in protecting cells from endogenous and exogenous DNA-based pathogens, their dysregulation has also been linked to cancer. For instance, AID has been linked to various B cell

malignancies [reviewed in (23)]. A second prominent example is the recent finding that APOBEC3B is overexpressed and a significant source of mutation in breast, ovarian, and several other cancers [(17,24–27) and reviewed in (28–30)]. APOBEC3B deaminates cytosines in genomic DNA to produce promutagenic uracil lesions, which can result in mutations if they are not repaired faithfully. APOBEC3B is overexpressed and its mutation signature is overrepresented in genomic sequences of HGSOC and other cancers. Elevated APOBEC3B expression levels have also been linked to poor prognosis in multiple cancer types (31–36).

In addition to APOBEC3B, related subfamily members have also been implicated in cancer mutagenesis to varying degrees (37–40). Because most *APOBEC* family members are expressed in many normal cell types and tissues, a major confounding factor in quantification of *APOBEC* expression levels in tumors is cellular heterogeneity due to surrounding normal tissue and immune cell infiltration. To address these issues, we quantified *APOBEC* expression in a large cohort of HGSOC patients and asked whether expression levels correlated with immune cell infiltration and could be distinguished from surrounding normal tissues. A strong positive correlation was identified between *APOBEC3G* expression and markers of T cell infiltration, and co-expression was confirmed by immunohistochemical and immunofluorescent staining of primary HGSOC specimens. Moreover, high *APOBEC3G* expression levels correlated significantly with improved prognosis. These findings were extended to multiple cancer types through an analysis of publically available RNA sequencing (RNAseq) data from The Cancer Genome Atlas (TCGA). Collectively, these results highlight the complexity of *APOBEC3* family member expression in HGSOC specimens comprised of tumor cells, surrounding normal tissues, and in many instances infiltrating immune cells. Our studies are the first to identify *APOBEC3G* as a new candidate biomarker for effective T cell responses and potentially for immunotherapies against HGSOC.

Materials and Methods

Analysis of expression correlations in Mayo Clinic cohort

Primary tumor samples from 354 HGSOC cases at the Mayo Clinic were selected based on histology, grade, stage, and availability of clinical outcome data (IRB #13-002487). Following cryosectioning of each snap frozen specimen, TRIzol based RNA extractions were performed. cDNA was synthesized in triplicate using Transcriptor Reverse Transcriptase (Roche) and reverse transcription quantitative PCR (RT-qPCR) for *APOBEC3G*, *APOBEC3B*, *CD3D*, *CD4*, *CD8A*, *GZMB*, *PRF1*, *RNF128*, and *TBP* was performed using the primer and probe combinations listed in Table S1 (validation in Fig. S1). Correlations between *APOBEC3G*, *APOBEC3B*, and the various T cell markers were determined using Spearman's correlation. Spearman's correlation coefficient (r_s) and p-values are reported.

Immunohistochemistry

IHC for CD3, CD4, CD8, and APOBEC3G was performed on 7 paraffin embedded primary HGSOC specimens obtained from patients who underwent debulking surgery at Ghent

University Hospital. This study was approved by the ethics committee of the Ghent University Hospital. The tissues were fixed in a 4% formaldehyde solution for 12–48 hours and embedded in paraffin. IHC was performed on 3.5 µm tissue sections on Superfrost slides (Menzel-Gläser) using a Benchmark XT automated slide stainer (Roche), according to the manufacturer's instructions. The following antibodies were used: mouse monoclonal anti-CD3 (clone F7.2.38, dilution 1/10, Dako), rabbit monoclonal anti-CD4 (clone SP35, dilution 1/25, Cell Marque), mouse monoclonal anti-CD8 (clone C8/144B, no dilution, NeoMarkers), and a locally developed rabbit monoclonal anti-APOBEC3G [clone 5211-110-19, dilution 1/50]. Although this rabbit monoclonal antibody recognizes APOBEC3G, APOBEC3A, and APOBEC3B due to unavoidable homology, we are confident that it is only detecting APOBEC3G in the immunohistochemistry studies described here because APOBEC3A is not expressed in T lymphocytes and, for reasons still under investigation, this monoclonal antibody does not recognize endogenous APOBEC3B by these procedures. It is important to emphasize that APOBEC3A is not expressed in most normal cell types, and it is only induced to detectable levels in interferon-activated myeloid lineage cells (12,15,17,18). Moreover, comparatively few CD68-positive macrophages were detected in the HGSOc specimens described here and positive signals from APOBEC3A or APOBEC3G expression in this cell type are minor (data not shown). Heat-induced epitope retrieval was done using Cell Conditioning 1 (Roche) for CD3, CD8, and APOBEC3G, and using Cell Conditioning 2 (Roche) for CD4. Visualization of all primary antibodies was achieved with the ultraView™ Universal DAB Detection Kit (Roche). Counterstaining with hematoxylin, dehydration of the tissue sections, and application of coverslips were carried out using an automated coverslipper (Tissue-Tek).

Immunofluorescent imaging

Fluorescence-based co-localization experiments were done using a subset of the same HGSOc specimens used above for IHC following published procedures (41). After sample preparation, permeabilization, and blocking with 10% goat serum in PBS at room temperature (DAKO X0907), each slide was stained first with rabbit polyclonal anti-APOBEC3G (Sigma Atlas Antibody HPA001812) diluted 1/50 in PBS (RT, 1 hr), second with mouse monoclonal anti-CD3 (described above) diluted 1/10 in PBS (4°C, overnight), and finally with a combination of secondary antibodies diluted 1/500 in PBS [RT, 1 hr; anti-rabbit IgG-AlexaFluor 594 (Invitrogen A11012) and anti-mouse IgG1-AlexaFluor 488 (Invitrogen A21121)]. Finally, slides were stained with DAPI diluted 1/5000 in methanol (RT, 5 min), mounted with a cover slip, and imaged using a fluorescence microscope equipped with appropriate filters (Olympus BX40, Tokyo, Japan). Multiple PBS washes were done between each step of the procedures.

Survival analysis in Mayo Clinic cohort

Kaplan-Meier curves were constructed by dividing specimens at the median expression level for each gene. OS data was available for all 354 patients, while PFS data was only available for 348 patients. P-values, hazard ratios and 95% confidence intervals were determined using Cox regression models on the continuous log2-transformed expression, adjusting for stage and debulking status.

Survival analysis in additional cohorts

Ovarian cancer data from TCGA and Gene Expression Omnibus (GEO) were compiled by the 2015 version of the Kaplan-Meier plotter database on a PostgreSQL server (42). GEO accession numbers were GSE14764, GSE15622, GSE18520, GSE19829, GSE23554, GSE26193, GSE26712, GSE27651, GSE30161, GSE3149, GSE51373, and GSE9891. All gene expression data were determined using only the HG-U133A, HG-U133 Plus 2.0, and HG-U133A 2.0 Affymetrix microarray platforms so that comparisons could be made between datasets. Specifically, *APOBEC3G* expression was determined using the 204205_at probe. Grade 3 serous ovarian cancers were the only samples used in this analysis. OS data were available for 856 patients, while PFS data was available for 753 patients. Kaplan-Meier plots were constructed and p-values, hazard ratios and 95% confidence intervals were calculated using the Mantel-Cox log-rank test.

Medical ethics approval for the Dutch cohort was obtained in part previously (43) (n=37) and in part more recently (n=36 and n=15; MEC-2008-183). *APOBEC3G* mRNA levels were measured by RTqPCR using an assay on demand (Hs00222415_m1, Applied Biosystems) and three reference genes were measured with the primers listed in Table S1 and quantification using SYBR green. Relative *APOBEC3G* expression levels were determined by normalization to the average of 3 reference genes as described (31). Kaplan-Meier curves were constructed by dividing specimens at the median expression level for each gene. P-values, hazard ratios and 95% confidence intervals were determined using the Mantel-Cox log-rank test.

TCGA analysis

Normalized RNAseqV2 data were downloaded from TCGA in July 2015. *APOBEC3*, *CD3D*, and *CD20* mRNA levels were quantified using normalized read counts. r_s and p-values for linear models of *APOBEC* versus immune-marker genes were calculated using Spearman's rank correlation coefficient with the R statistical environment. Cancer types were grouped by hierarchical clustering (hclust) of the r_s values for each *APOBEC* family member in the R statistical environment, and these results were used to generate a dendrogram of these relationships. All data were graphed using the ggplot2 R package (44). P-values were calculated from the r_s values and adjusted for multiple comparisons using the Benjamini-Hochberg method and significance was defined as a p-value less than 0.05.

Results

***APOBEC3G* expression correlates with activated T lymphocyte infiltration in HGSOC**

APOBEC3G is expressed constitutively in many different cell lines and tissue types and is also known to be upregulated by HIV-1 infection of primary T lymphocytes, which is one of many distinct mechanisms of immune cell activation [e.g., (12–14)]. Although virus infection is unlikely to be part of the etiology of ovarian cancer (45), the presence of activated immune cells (infiltrate) in HGSOC is known to correlate with better overall outcomes, most likely due to strong anti-tumor immune responses (3–7). We therefore asked whether *APOBEC3G* expression correlated with T cell infiltration and clinical outcomes

using a cohort of 354 primary HGSOC samples procured at the Mayo Clinic (clinical characteristics in Table 1).

RNA was prepared from fresh frozen HGSOC tissues, and a previously validated, highly specific RT-qPCR assay was used to assess *APOBEC3G* mRNA levels (12). In parallel, expression of the related gene *APOBEC3B*, which has been implicated in ovarian cancer genome mutagenesis (24), was assayed. In addition, the mRNA levels of several established T cell markers were quantified, including *CD3D* (total T cells), *CD4* (helper T cells), *CD8A* (cytotoxic T cells), *GZMB* (activated cytotoxic T cells), *PRF1* (activated cytotoxic T cells), and *RNF128* (anergic T cells) (primers and probes in Table S1) (46,47).

Strong positive correlations were observed between *APOBEC3G* mRNA expression levels and *CD3D* ($p<0.0001$, $r_s=0.6159$), *CD4* ($p<0.0001$, $r_s=0.5825$), *CD8A* ($p<0.0001$, $r_s=0.6168$), *GZMB* ($p<0.0001$, $r_s=0.6591$), and *PRF1* ($p<0.0001$, $r_s=0.6422$) (Fig. 1A–E). The lone exception was *RNF128* ($p=0.7665$, $r_s=0.0161$), which is a marker for T cell anergy (48), suggesting that the positive signals emanate from *bona fide* activated T lymphocytes (Fig. 1F). Interestingly, *APOBEC3G* expression showed a similar positive correlation with *CD8A* and *CD4*, suggesting that *APOBEC3G* is expressed in both the cytotoxic and helper T cell subsets (Fig. 1B vs. 1C). This result was corroborated by similarly strong positive correlations between *APOBEC3G* and two markers of CTL activation, *GZMB* and *PRF1* (Fig. 1D and 1E). In contrast, weaker and less significant correlations were found between *APOBEC3B* and the expression of any of these T cell genes (Fig. 1G–L). This result was expected, however, because prior studies have indicated that *APOBEC3B* is only expressed at very low levels in normal tissues and upregulated in tumor cells (12,17,24,25). Taken together, our data confirm prior studies documenting T cell infiltration in HGSOC (3–7) and, importantly, reveal an unanticipated association between high levels of *APOBEC3G* expression and CTL activation in HGSOC. This result for *APOBEC3G* was especially unexpected given its broad expression profile documented previously by several groups including our own (12,15–17).

APOBEC3G protein visualization in infiltrating T lymphocytes in HGSOC

To determine whether APOBEC3G protein expression co-localizes with the same T cell markers analyzed above at the mRNA level by RT-qPCR, we performed hematoxylin staining and immunohistochemistry for CD3, CD4, CD8, and APOBEC3G on seven unrelated HGSOC samples (representative image sets in Fig. 2A–T with hematoxylin staining in Fig. 2A, F, K, and P). As expected from the RT-qPCR analysis above, several of these additional tumor specimens showed clear evidence for T lymphocyte infiltration. Two HGSOC lesions contained low expression of all examined T cell markers (representative images for CD3, CD4, and CD8 in Fig. 2B–D) and correspondingly low levels of APOBEC3G expression (representative image in Fig. 2E). In contrast, two tumors showed extensive T lymphocyte infiltration (representative images for CD3, CD4 and CD8 in Fig. 2G–I, L–N, and Q–S). Interestingly, there was a strong colocalization between these markers and the expression of APOBEC3G in these tumors (representative image in Fig. 2J, O, and T). Moreover, different regions of a single HGSOC can be heterogenous, with one region showing dense clusters of APOBEC3G-high infiltrating T cells and another showing a more

dispersed distribution (compare Fig. 2F–J and K–O). The remaining three samples showed moderate expression of both infiltrating T cell markers and APOBEC3G, and colocalization was still observed (data not shown). These immunohistochemical experiments confirm the above correlations and show that APOBEC3G is indeed expressed at the protein level within tumor infiltrating T lymphocytes.

To directly test whether APOBEC3G is expressed within tumor infiltrating T lymphocytes, we performed a series of fluorescence-based co-localization experiments with CD3 and APOBEC3G. Here, we were only able to image well-isolated infiltrating T lymphocytes because groups of cells caused excessive background fluorescence. Nevertheless, these experiments enabled us to demonstrate for the first time that CD3 and APOBEC3G are indeed co-expressed in the same T lymphocyte (representative image in Fig. 2U–X). Moreover, the DAPI co-stain and 1000x total magnification combine to show that both proteins are excluded from the nuclear compartment and predominantly cytoplasmic (the additional cell surface localization of CD3 is more difficult to visualize in tissue cross sections).

***APOBEC3G* is a candidate biomarker for improved HGSOC patient outcomes**

Long-term clinical follow-up data were available for all of the Mayo Clinic HGSOC patients. The aforementioned gene expression data were therefore correlated with clinical information to determine whether *APOBEC3G* expression levels predict the length of progression free survival (PFS) and/or overall survival (OS) in HGSOC. As a positive control, the T cell markers above were also analyzed with respect to clinical information. Kaplan-Meier plots were constructed by splitting each gene expression data set at the median to create high and low expression groups (Fig. 3A and B). P-values, hazard ratios (HR), and 95% confidence intervals (CI) were determined using Cox regression models on the log2-transformed expression that were corrected for stage and debulking status (Table 2).

Higher expression levels of *CD3D* ($p=0.020$, $HR=0.94$ [95% confidence interval 0.90, 0.99]), *CD4* ($p=0.0046$, $HR=0.90$ [0.84, 0.97]), *CD8A* ($p=0.0053$, $HR=0.93$ [0.88, 0.98]), *GZMB* ($p=0.011$, $HR=0.94$ [0.90, 0.99]), and *PRF1* ($p=0.0049$, $HR=0.91$ [0.86, 0.97]) were all associated with improved PFS (Fig. 3A and Table 2). As expected, *RNF128* ($p=0.43$, $HR=0.97$ [0.91, 1.04]) did not correlate with PFS (Fig. 3A and Table 2). Interestingly, *APOBEC3G* ($p<0.0001$, $HR=0.81$ [0.73, 0.89]) surpassed all of these genes as the most indicative marker of improved PFS in HGSOC (Fig. 3A). The results compiled from an analysis of OS largely mirrored those of PFS (Fig. 3B and Table 2).

Next, we used Kaplan-Meier plotter (kmplot.com) to generate a validation cohort using HGSOC data from TCGA and GEO. In this composite analysis of HGSOC patients, high levels of *APOBEC3G* correlated with improved durations of PFS ($p=0.0057$, $HR=0.78$ [0.65, 0.93]) and OS ($p=0.063$, $HR=0.84$ [0.70, 1.01]) (Fig. S2). Although only the former correlation reached statistical significance, the latter trended toward significance and also supported the observations above with the HGSOC Mayo Clinic cohort. A larger degree of variation is to be expected in this composite cohort because clinical variables are more difficult to take into account.

Finally, we analyzed HGSOC specimens from a Dutch cohort consisting of 88 patient samples (clinical information in Table S2). *APOBEC3G* expression was quantified using an independent RT-qPCR strategy (housekeeping gene primer sequences in Table S1 and see **Material and Methods** for details). Again, *APOBEC3G* expression levels associated with improved OS and more weakly with PFS (Fig. S2).

***APOBEC3B* expression levels do not associate with HGSOC outcomes**

APOBEC3B has been implicated recently as an endogenous mutagen in several cancers [(28–30) and references therein], including ovarian cancer (24). Moreover, its overexpression has been linked to poor patient outcomes in multiple cancer types (31–36). Using the Mayo cohort gene expression data and clinical information from above, we asked whether *APOBEC3B* affects patient outcomes in HGSOC. These analyses revealed a trend toward high *APOBEC3B* and improved, rather than worsened, PFS and OS outcomes, although these relationships were less significant statistically than those for *APOBEC3G* (Fig. 3).

***APOBEC* expression correlates with immune cell markers in multiple human cancers**

To extend our findings from HGSOC to additional human cancers, publically available RNAseq data from TCGA were analyzed to determine if correlations exist between expression of *APOBEC* genes and immune cell markers. At the time of these analyses, the TCGA had RNAseq data available for 7,861 samples spanning 22 different tumor types (details in Table S3). For each tumor type, expression of each *APOBEC* family member was quantified and correlated with the T cell marker *CD3D* (Fig. 4; top heatmap). Hierarchical clustering was also performed to elucidate similar correlation patterns between cancer types (Fig. 4A top). These analyses revealed that, in addition to *APOBEC3G*, *APOBEC3D* and *APOBEC3H* also correlated significantly with *CD3D* across multiple cancer types. *APOBEC3F*, also known to be expressed broadly and in T cells (12,13,15–17), did not correlate as strongly. The same analysis was also performed with *CD20*, a well-known marker for B cells (Fig. 4 bottom heatmap). The expression of the antibody diversification gene, *AID*, was the only *APOBEC* family member that significantly correlated with *CD20* in a majority of cancer types (Fig. 4B). These analyses indicate that much of the expression of several *APOBEC* family members in cancer is likely due to T and B cell infiltration.

Discussion

Our studies have led to the surprising identification of *APOBEC3G* as a new candidate biomarker for activated T lymphocyte infiltration in HGSOC and improved patient outcomes. This result was unexpected because prior work has shown that *APOBEC3G* is expressed broadly, constitutively, and in some instances inducibly (12,13,15–17). The analysis of a cohort of 354 HGSOC patients also identified a strong correlation between *APOBEC3G* and several markers of T cell infiltration (Fig. 1A–E). These results were validated at the protein level by immunohistochemistry and immunofluorescent staining of independent HGSOC tumor samples (representative image sets in Fig. 2). Clinical data revealed that *APOBEC3G* also associates with improved outcomes in a large HGSOC cohort as well as in two additional independent ovarian cancer cohorts (Table 2, Fig. 3, and Fig.

S2). Finally, a global analysis of gene expression in 22 cancer types identified a similar correlation for two additional *APOBEC3* genes, *APOBEC3D* and *APOBEC3H*, and a marker of T cells, *CD3D* (Fig. 4). Together, these data suggest that *APOBEC3G* expression levels in tumor infiltrating T lymphocytes may be a predictive biomarker for strong anti-cancer T cell responses and improved HGSOC outcomes.

Over the past several years, substantial evidence has accumulated to indicate that *APOBEC3B* and *AID* contribute to cancer genome mutagenesis. *APOBEC3B* is thought to mutate the genome of several different cancer types, including breast, lung, bladder, cervical, head/neck, and ovarian (17,24–27). The carcinogenic effect of *AID* appears more limited, as its characteristic deamination signature has only been found in certain types of B cell leukemias and lymphomas (23,27). The idea that *AID* expression is constrained to B lineage cell types is consistent with our data showing that *AID* mRNA expression correlates with *CD20* mRNA expression in several solid tumor types (Fig. 4 bottom heatmap). In addition to *APOBEC3B* and *AID*, several other *APOBEC* family members have also been implicated in carcinogenesis (37–40). For example, it has been proposed that *APOBEC3G* drives hepatocellular carcinoma tumorigenesis (37). This idea, however, is difficult to reconcile with our observation that *APOBEC3G* mRNA expression in HGSOC and other tumor types correlates with the expression of activated T lymphocyte markers and, at least for HGSOC, this strong correlation could be validated visually at the protein level (Figs. 1, 2, and 4). Additional studies are clearly warranted to extend the *APOBEC3G* immunohistochemistry results described here to other tumor types including hepatocellular carcinoma.

We were also surprised by the analysis of *APOBEC3B* in HGSOC, which revealed no significant clinical correlations and even a slight trend in the Mayo Clinic cohort toward better overall survival (Fig. 3). This finding differs significantly from estrogen receptor (ER)-positive breast cancer, where high *APOBEC3B* expression associates with shorter periods of disease-free survival and poorer rates of overall survival (31–34). A previous deep-sequencing study highlighted the similarities between HGSOC and another type of breast cancer, triple negative breast cancer (TNBC) (49), and, also found no correlation between *APOBEC3B* expression and survival in TNBC (32). One major difference between these two cancer types and ER-positive breast cancer is therapeutic options. There are multiple targeted therapeutics available for the treatment of ER-positive breast cancer that are administered based on molecular markers. In contrast, nearly all HGSOC and many TNBC patients are treated with platinum-based therapies. Because platinum-based therapies induce DNA damage, it is possible that these drugs become synergistic with *APOBEC3B*-catalyzed cytosine deamination and create a synthetic lethal state in cancer cells. This idea is reasonable as increased mutation loads have been shown to correlate with improved clinical outcomes in HGSOC patients treated with cisplatin (50). Furthermore, a synergistic effect created by these two forms of DNA damage could explain the slight trend found here toward a positive correlation between increased *APOBEC3B* expression and improved outcomes. Another possible explanation is that the levels of *APOBEC3B* mutagenesis in ovarian cancer may be not high enough to manifest clinically. Indeed, the *APOBEC3B* mutation signature is not as strong in ovarian cancer as it is in many other cancer types despite similar mRNA and protein expression levels (17,21,24–27). The underlying causes for this apparent discordancy are unknown, but several factors could be involved, including altered DNA

repair capacities, differential protein regulation, and mutational contributions from other sources. More work is needed to determine the threshold of APOBEC3B mutagenesis needed to have a clinical impact.

Our global analysis of TCGA data revealed that the correlation between *APOBEC3F* and *CD3D* was substantially diminished as compared to *APOBEC3D*, *APOBEC3G*, and *APOBEC3H* (Fig. 4). This is surprising because APOBEC3F is thought to be broadly expressed and play an equally important role as APOBEC3D, APOBEC3G, and APOBEC3H in HIV-1 restriction in CD4-positive T lymphocytes, and prior studies have found that *APOBEC3F* is expressed at comparable levels to *APOBEC3D* and higher than *APOBEC3H* in human primary CD4+ T cells (12–14). Our data suggest that *APOBEC3F* may be under- and/or heterogeneously-expressed in T cells associated with the tumor microenvironment. These differences are potentially interesting as the overall *APOBEC* family member expression profile may be a useful property for distinguishing anti-viral from anti-tumor immune responses. It will be interesting in future studies to determine whether APOBEC3G, the overall APOBEC family expression profile, APOBEC mutation signature, and/or classical markers for T cell activation are the best predictors of successful clinical responses to immunotherapies for HGSOV as well as for other tumor types.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Emily Law and David Masopust for thoughtful discussions, and Xuan Bich Trinh, Peter van Dam, Peter Vermeulen, Steven van Laere, and Luc Dirix from the Center for Oncological Research (CORE) at the University of Antwerp (Belgium) for contributing clinical specimens to the Dutch cohort.

Grant Support

Salary support for BL was provided by a Cancer Biology Training Grant (NIH T32 CA009138), and GJS by a National Science Foundation Graduate Research Fellowship (DGE 13488264). MJM, ALO, KRK, and SHK were supported in part by the Mayo Clinic SPORE in Ovarian Cancer (P50 CA136393). Ovarian cancer research in the Harris laboratory was supported by grants from the Minnesota Partnership for Biotechnology and Medical Genomics and the Minnesota Ovarian Cancer Alliance. RSH is an Investigator of the Howard Hughes Medical Institute.

References

1. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin.* 2011; 61:69–90. [PubMed: 21296855]
2. Seidman JD, Horkayne-Szakaly I, Haiba M, Boice CR, Kurman RJ, Ronnett BM. The histologic type and stage distribution of ovarian carcinomas of surface epithelial origin. *Int J Gynecol Pathol.* 2004; 23:41–4. [PubMed: 14668549]
3. Knutson KL, Maurer MJ, Preston CC, Moysich KB, Goergen K, Hawthorne KM, et al. Regulatory T cells, inherited variation, and clinical outcome in epithelial ovarian cancer. *Cancer Immunol Immunother.* 2015; 64:1495–504. [PubMed: 26298430]
4. Nielsen JS, Sahota RA, Milne K, Kost SE, Nesslinger NJ, Watson PH, et al. CD20+ tumor-infiltrating lymphocytes have an atypical CD27– memory phenotype and together with CD8+ T cells promote favorable prognosis in ovarian cancer. *Clin Cancer Res.* 2012; 18:3281–92. [PubMed: 22553348]

5. Zhang L, Conejo-Garcia JR, Katsaros D, Gimotty PA, Massobrio M, Regnani G, et al. Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N Engl J Med*. 2003; 348:203–13. [PubMed: 12529460]
6. Sato E, Olson SH, Ahn J, Bundy B, Nishikawa H, Qian F, et al. Intraepithelial CD8+ tumor-infiltrating lymphocytes and a high CD8+/regulatory T cell ratio are associated with favorable prognosis in ovarian cancer. *Proc Natl Acad Sci USA*. 2005; 102:18538–43. [PubMed: 16344461]
7. Preston CC, Maurer MJ, Oberg AL, Visscher DW, Kalli KR, Hartmann LC, et al. The ratios of CD8+ T cells to CD4+CD25+ FOXP3+ and FOXP3– T cells correlate with poor clinical outcome in human serous ovarian cancer. *PLoS ONE*. 2013; 8:e80063. [PubMed: 24244610]
8. Harris RS, Petersen-Mahrt SK, Neuberger MS. RNA editing enzyme APOBEC1 and some of its homologs can act as DNA mutators. *Mol Cell*. 2002; 10:1247–53. [PubMed: 12453430]
9. Refsland EW, Harris RS. The APOBEC3 family of retroelement restriction factors. *Curr Top Microbiol Immunol*. 2013; 371:1–27. [PubMed: 23686230]
10. Di Noia JM, Neuberger MS. Molecular mechanisms of antibody somatic hypermutation. *Annu Rev Biochem*. 2007; 76:1–22. [PubMed: 17328676]
11. Harris RS, Hultquist JF, Evans DT. The restriction factors of human immunodeficiency virus. *Journal of Biological Chemistry*. 2012; 287:40875–83. [PubMed: 23043100]
12. Refsland EW, Stenglein MD, Shindo K, Albin JS, Brown WL, Harris RS. Quantitative profiling of the full APOBEC3 mRNA repertoire in lymphocytes and tissues: implications for HIV-1 restriction. *Nucleic Acids Res*. 2010; 38:4274–84. [PubMed: 20308164]
13. Hultquist JF, Lengyel JA, Refsland EW, LaRue RS, Lackey L, Brown WL, et al. Human and rhesus APOBEC3D, APOBEC3F, APOBEC3G, and APOBEC3H demonstrate a conserved capacity to restrict Vif-deficient HIV-1. *Journal of Virology*. 2011; 85:11220–34. [PubMed: 21835787]
14. Refsland EW, Hultquist JF, Harris RS. Endogenous origins of HIV-1 G-to-A hypermutation and restriction in the nonpermissive T cell line CEM2n. *PLoS Pathog*. 2012; 8:e1002800. [PubMed: 22807680]
15. Koning FA, Newman ENC, Kim E-Y, Kunstman KJ, Wolinsky SM, Malim MH. Defining APOBEC3 expression patterns in human tissues and hematopoietic cell subsets. *Journal of Virology*. 2009; 83:9474–85. [PubMed: 19587057]
16. Liddament MT, Brown WL, Schumacher AJ, Harris RS. APOBEC3F properties and hypermutation preferences indicate activity against HIV-1 in vivo. *Curr Biol*. 2004; 14:1385–91. [PubMed: 15296757]
17. Burns MB, Lackey L, Carpenter MA, Rathore A, Land AM, Leonard B, et al. APOBEC3B is an enzymatic source of mutation in breast cancer. *Nature*. 2013; 494:366–70. [PubMed: 23389445]
18. Stenglein MD, Burns MB, Li M, Lengyel J, Harris RS. APOBEC3 proteins mediate the clearance of foreign DNA from human cells. *Nat Struct Mol Biol*. 2010; 17:222–9. [PubMed: 20062055]
19. Thielen BK, McNevin JP, McElrath MJ, Hunt BVS, Klein KC, Lingappa JR. Innate immune signaling induces high levels of TC-specific deaminase activity in primary monocyte-derived cells through expression of APOBEC3A isoforms. *Journal of Biological Chemistry*. 2010; 285:27753–66. [PubMed: 20615867]
20. Lucifora J, Xia Y, Reisinger F, Zhang K, Stadler D, Cheng X, et al. Specific and nonhepatotoxic degradation of nuclear hepatitis B virus cccDNA. *Science*. 2014; 343:1221–8. [PubMed: 24557838]
21. Leonard B, McCann JL, Starrett GJ, Kosyakovsky L, Luengas EM, Molan AM, et al. The PKC/NF- κ B signaling pathway induces APOBEC3B expression in multiple human cancers. *Cancer Research*. 2015; 75:4538–47. [PubMed: 26420215]
22. Vieira VC, Leonard B, White EA, Starrett GJ, Temiz NA, Lorenz LD, et al. Human papillomavirus E6 triggers upregulation of the antiviral and cancer genomic DNA deaminase APOBEC3B. *MBio*. 2014; 5:e02234–14. [PubMed: 25538195]
23. Robbiani DF, Nussenzweig MC. Chromosome translocation, B cell lymphoma, and activation-induced cytidine deaminase. *Annu Rev Pathol*. 2013; 8:79–103. [PubMed: 22974238]
24. Leonard B, Hart SN, Burns MB, Carpenter MA, Temiz NA, Rathore A, et al. APOBEC3B upregulation and genomic mutation patterns in serous ovarian carcinoma. *Cancer Research*. 2013; 73:7222–31. [PubMed: 24154874]

25. Burns MB, Temiz NA, Harris RS. Evidence for APOBEC3B mutagenesis in multiple human cancers. *Nat Genet.* 2013; 45:977–83. [PubMed: 23852168]
26. Roberts SA, Lawrence MS, Klimczak LJ, Grimm SA, Fargo D, Stojanov P, et al. An APOBEC cytidine deaminase mutagenesis pattern is widespread in human cancers. *Nat Genet.* 2013; 45:970–6. [PubMed: 23852170]
27. Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SAJR, Behjati S, Biankin AV, et al. Signatures of mutational processes in human cancer. *Nature.* 2013; 500:415–21. [PubMed: 23945592]
28. Burns MB, Leonard B, Harris RS. APOBEC3B: pathological consequences of an innate immune DNA mutator. *Biomedical Journal.* 2015; 38:102–10. [PubMed: 25566802]
29. Swanton C, McGranahan N, Starrett GJ, Harris RS. APOBEC enzymes: mutagenic fuel for cancer evolution and heterogeneity. *Cancer Discovery.* 2015; 5:704–12. [PubMed: 26091828]
30. Roberts SA, Gordenin DA. Hypermutation in human cancer genomes: footprints and mechanisms. *Nat Rev Cancer.* 2014; 14:786–800. [PubMed: 25568919]
31. Sieuwerts AM, Willis S, Burns MB, Look MP, Meijer-Van Gelder ME, Schlicker A, et al. Elevated APOBEC3B correlates with poor outcomes for estrogen-receptor-positive breast cancers. *Horm Cancer.* 2014; 5:405–13. [PubMed: 25123150]
32. Cescon DW, Haibe-Kains B, Mak TW. APOBEC3B expression in breast cancer reflects cellular proliferation, while a deletion polymorphism is associated with immune activation. *Proceedings of the National Academy of Sciences.* 2015; 112:2841–6.
33. Periyasamy M, Patel H, Lai C-F, Nguyen VTM, Nevedomskaya E, Harrod A, et al. APOBEC3B-Mediated Cytidine Deamination Is Required for Estrogen Receptor Action in Breast Cancer. *Cell Rep.* 2015; 13:108–21. [PubMed: 26411678]
34. Tsuboi M, Yamane A, Horiguchi J, Yokobori T, Kawabata-Iwakawa R, Yoshiyama S, et al. APOBEC3B high expression status is associated with aggressive phenotype in Japanese breast cancers. *Breast Cancer.* 2015 in press. (Oct. 17th Epub ahead of print).
35. Xu L, Chang Y, An H, Zhu Y, Yang Y, Xu J. High APOBEC3B expression is a predictor of recurrence in patients with low-risk clear cell renal cell carcinoma. *Urol Oncol.* 2015; 33:340, e1–8. [PubMed: 26051357]
36. Zhang J, Wei W, Jin H-C, Ying R-C, Zhu A-K, Zhang F-J. The roles of APOBEC3B in gastric cancer. *Int J Clin Exp Pathol.* 2015; 8:5089–96. [PubMed: 26191203]
37. Ding Q, Chang C-J, Xie X, Xia W, Yang J-Y, Wang S-C, et al. APOBEC3G promotes liver metastasis in an orthotopic mouse model of colorectal cancer and predicts human hepatic metastasis. *J Clin Invest.* 2011; 121:4526–36. [PubMed: 21985787]
38. Caval V, Suspène R, Shapira M, Vartanian J-P, Wain-Hobson S. A prevalent cancer susceptibility APOBEC3A hybrid allele bearing APOBEC3B 3'UTR enhances chromosomal DNA damage. *Nat Commun.* 2014; 5:5129. [PubMed: 25298230]
39. Nik-Zainal S, Wedge DC, Alexandrov LB, Petljak M, Butler AP, Bolli N, et al. Association of a germline copy number polymorphism of APOBEC3A and APOBEC3B with burden of putative APOBEC-dependent mutations in breast cancer. *Nat Genet.* 2014; 46:487–91. [PubMed: 24728294]
40. Chan K, Roberts SA, Klimczak LJ, Sterling JF, Saini N, Malc EP, et al. An APOBEC3A hypermutation signature is distinguishable from the signature of background mutagenesis by APOBEC3B in human cancers. *Nat Genet.* 2015; 47:1067–72. [PubMed: 26258849]
41. Verschuere S, Bracke KR, Demoor T, Plantinga M, Verbrugghe P, Ferdinande L, et al. Cigarette smoking alters epithelial apoptosis and immune composition in murine GALT. *Lab Invest.* 2011; 91:1056–67. [PubMed: 21537330]
42. Györfy B, Lánckzy A, Szallasi Z. Implementing an online tool for genome-wide validation of survival-associated biomarkers in ovarian-cancer using microarray data from 1287 patients. *Endocr Relat Cancer.* 2012; 19:197–208. [PubMed: 22277193]
43. Schuyer M, van der Burg ME, Henzen-Logmans SC, Fieret JH, Klijn JG, Look MP, et al. Reduced expression of BAX is associated with poor prognosis in patients with epithelial ovarian cancer: a multifactorial analysis of TP53, p21, BAX and BCL-2. *Br J Cancer.* 2001; 85:1359–67. [PubMed: 11720475]

44. Wickham, H. ggplot2: elegant graphics for data analysis. New York: Springer Publishing Company; 2016.
45. Bast RC, Hennessy B, Mills GB. The biology of ovarian cancer: new opportunities for translation. *Nat Rev Cancer*. 2009; 9:415–28. [PubMed: 19461667]
46. Sabek O, Dorak MT, Kotb M, Gaber AO, Gaber L. Quantitative detection of T-cell activation markers by real-time PCR in renal transplant rejection and correlation with histopathologic evaluation. *Transplantation*. 2002; 74:701–7. [PubMed: 12352889]
47. Zheng Y, Zha Y, Gajewski TF. Molecular regulation of T-cell anergy. *EMBO Rep*. 2008; 9:50–5. [PubMed: 18174897]
48. Schwartz RH. T cell anergy. *Annu Rev Immunol*. 2003; 21:305–34. [PubMed: 12471050]
49. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature*. 2011; 474:609–15. [PubMed: 21720365]
50. Sohn I, Jung WY, Sung CO. Somatic hypermutation and outcomes of platinum based chemotherapy in patients with high grade serous ovarian cancer. *Gynecol Oncol*. 2012; 126:103–8. [PubMed: 22484402]

Translational Relevance

Ovarian cancer is the deadliest cancer of the female reproductive tract, partly due to a lack of biomarkers for early clinical detection and assessment of prognosis. Several members of the APOBEC family of antiviral DNA cytosine deaminases, including *APOBEC3G*, are expressed broadly in human tissues, and *APOBEC3B* is overexpressed in many tumor types including ovarian cancer. Here, we show that *APOBEC3G* is expressed at surprisingly high levels in T lymphocytes within high-grade serous ovarian tumors. *APOBEC3G* expression correlates positively with improved patient outcomes. The latter result is consistent with prior work demonstrating that T cell infiltration correlates with improved outcomes. Our results are the first to demonstrate that high *APOBEC3G* levels in ovarian tumors are attributable to T cell infiltration and not, for instance, expression in other normal cells or tumor cells. Thus, *APOBEC3G* is a new candidate biomarker for T cell infiltration and positive anti-tumor immune responses.

Figure 1. Correlations between *APOBEC3* expression and T cell markers in HGSOc
 Dot plots illustrating correlations between *APOBEC3G* (A–F) or *APOBEC3B* (G–L) expression and the indicated T cell marker (n=354). mRNA expression was determined using RT-qPCR and normalized to the housekeeping gene *TBP*. Spearman's correlation coefficients (r_s) and p-values are shown. Best-fit lines are shown for qualitative comparison, and were calculated using linear regression models.

Figure 2. Immunohistochemistry and immunofluorescence of T cell markers in HGSOC
Photomicrographs of immunohistochemistry and immunofluorescence staining performed on HGSOC specimens illustrating the association between levels of T lymphocyte infiltration and the intensity of APOBEC3G expression. Representative staining of one HGSOC specimen with low (patient 6) and three staining sets from two HGSOC specimens with high (patients 3 and 2) levels of T cell infiltration are shown. The images depict hematoxylin (A, F, K, and P), CD3 (B, G, L, and Q), CD4 (C, H, M, and R), CD8 (D, I, N, and S), and APOBEC3G (E, J, O, and T). The dotted box in the 40x hematoxylin images indicates the approximate location of the subsequent panels at 100x magnification. The bottom row shows representative 1000x magnification images of colocalization of CD3 and APOBEC3G by immunofluorescent staining. DAPI-stained nuclei are blue.

Figure 3. Clinical correlates of T cell marker expression in HGSOc

Kaplan-Meier plots illustrating associations between progression free survival (PFS) (**A**, n=354) or overall survival (OS) (**B**, n=348) and either one of the conventional T cell markers or *APOBEC3G* or *APOBEC3B* expression in the Mayo cohort of patients. Samples were split at the median expression level for each gene with red representing tumors with high and blue representing tumors with low mRNA levels.

Figure 4. Correlations between APOBEC expression and immune cell markers across 22 cancer types

Heatmap of Spearman's correlation coefficients calculated from the comparison of the T cell marker *CD3D* (A) or the B cell marker *CD20* (B) with the indicated APOBEC family member. Expression levels were determined using TCGA RNAseq data (see Table S3 for the long form for each tumor abbreviation). Dark red squares indicate strong positive correlations, dark blue squares indicate strong negative correlations and white squares indicate a lack of correlation.

Table 1

Clinical information for Mayo cohort (n=354)

Tumor characteristic	Number
Morphology	
Serous	354 (100%)
Grade	
2	11 (3.1%)
3	343 (96.9%)
Stage	
1	14 (3.9%)
2	8 (2.3%)
3	253 (71.5%)
4	79 (22.3%)
Debulking status	
No residual disease	162 (45.8%)
<=1 cm remaining	144 (40.7%)
<=1 cm remaining, possibly 0	48 (13.5%)

Table 2

Mayo cohort Cox regression analysis

Gene	Parameter ¹	p-value	HR ²	95% CI ³
<i>CD3D</i>	PFS	0.020	0.94	[0.90,0.99]
<i>CD3D</i>	OS	0.087	0.95	[0.90,1.01]
<i>CD4</i>	PFS	0.0046	0.90	[0.84,0.97]
<i>CD4</i>	OS	0.018	0.91	[0.84,0.98]
<i>CD8A</i>	PFS	0.0053	0.93	[0.88,0.98]
<i>CD8A</i>	OS	0.015	0.93	[0.87,0.99]
<i>GZMB</i>	PFS	0.011	0.94	[0.90,0.99]
<i>GZMB</i>	OS	0.047	0.95	[0.90,1.00]
<i>PRF1</i>	PFS	0.0049	0.91	[0.86,0.97]
<i>PRF1</i>	OS	0.018	0.92	[0.86,0.99]
<i>RNF128</i>	PFS	0.43	0.97	[0.91,1.04]
<i>RNF128</i>	OS	0.44	0.97	[0.91,1.04]
<i>APOBEC3G</i>	PFS	<0.0001	0.81	[0.73,0.89]
<i>APOBEC3G</i>	OS	0.0003	0.82	[0.73,0.91]
<i>APOBEC3B</i>	PFS	0.034	0.92	[0.85, 0.99]
<i>APOBEC3B</i>	OS	0.06	0.93	[0.87, 1.00]

¹OS= overall survival (n= 354); PFS= progression free survival (n= 348)²HR= hazard ratio³CI= confidence interval