Published in final edited form as:

Pharmacol Res. 2016 September ; 111: 357–373. doi:10.1016/j.phrs.2016.06.024.

mTOR-Independent Autophagy Inducer Trehalose Rescues against Insulin Resistance-Induced Myocardial Contractile Anomalies: Role of p38 MAPK and Foxo1

Qiurong Wang¹ and Jun Ren^{1,2}

¹Center for Cardiovascular Research and Alternative Medicine, University of Wyoming College of Health Sciences, Laramie, WY 82071, USA

²Shanghai Institute of Cardiovascular Diseases, Zhongshan Hospital, Fudan University, Shanghai 200032, China

Abstract

Insulin resistance is associated with cardiovascular diseases although the precise mechanisms remain elusive. Akt2, a critical member of the Akt family, plays an essential role in insulin signaling. This study was designed to examine the effect of trehalose, an mTOR-independent autophagy inducer, on myocardial function in an Akt2 knockout-induced insulin resistance model. Adult WT and Akt2 knockout (Akt2^{-/-}) mice were administered trehalose (1 mg/g/day, i.p.) for two days and were then given 2% trehalose in drinking water for two more months. Echocardiographic and myocardial mechanics, intracellular Ca²⁺ properties, glucose tolerance, and autophagy were assessed. Apoptosis and ER stress were evaluated using TUNEL staining, Caspase 3 assay and Western blot. Autophagy and autophagy flux were examined with a focus on p38 mitogen activated protein kinase (MAPK), Forkhead box O (Foxo1) and Akt. Akt2 ablation impaired glucose tolerance, myocardial geometry and function accompanied with pronounced apoptosis, ER stress and dampened autophagy, the effects of which were ameliorated by trehalose treatment. Inhibition of lysosomal activity using bafilomycin A1 negated trehalose-induced induction of autophagy (LC3B-II and p62). Moreover, phosphorylation of p38 MAPK and Foxo1 were upregulated in Akt2^{-/-} mice, the effect of which was attenuated by trehalose. Phosphorylation of Akt was suppressed in Akt2^{-/-} mice and was unaffected by trehalose. In vitro findings revealed that the p38 MAPK activator anisomycin and the Foxo1 inhibitor (through phosphorylation) AS1842856 effectively masked trehalose-offered beneficial cardiomyocyte contractile response against Akt2 ablation. These data suggest that trehalose may rescue against insulin resistance-induced myocardial contractile defect and apoptosis, via autophagy associated with dephosphorylation of p38 MAPK and Foxo1 without affecting phosphorylation of Akt.

Correspondence should be addressed to: Dr. Jun Ren, University of Wyoming College of Health Sciences, Laramie, WY 82071, USA, Tel: (307)766-6131; Fax: (307) 766-2953; jren@uwyo.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Trehalose; autophagy; cardiac; contractile function; Akt2

1. INTRODUCTION

Insulin resistance, or impaired sensitivity to insulin, is a hallmark of metabolic morbidities including type 2 diabetes mellitus, obesity and hypertension, clustered together as the “metabolic syndrome” (1, 2). Incidence of cardiovascular disease is much higher in individuals with insulin resistance, as evidenced by dampened cardiac energy utilization efficiency (glucose oxidation versus fatty acid oxidation), compromised ventricular function and coronary heart diseases (1). Insulin resistance-associated myocardial anomalies are reminiscent of those found in diabetic individuals with compromised heart function, characterized by impaired ventricular compliance, prolonged action potential duration and ventricular diastole, and delayed cytosolic Ca^{2+} clearance (3–5). It is suggested that compromised insulin signaling in the heart plays an important role in insulin resistance-induced myocardial contractile and geometric anomalies (6, 7). Within the complex insulin regulatory signaling cascade, the phosphatidylinositol 3-kinase (PI3K)-Akt pathway governs the metabolic properties of insulin and serves as an essential gate-keeper for post-insulin receptor signaling transduction (8, 9). Akt, a serine/threonine kinase downstream of PI3K, comprises three isoforms Akt1, Akt2 and Akt3. Although the structure and function of these three isoforms are highly conserved, ample studies have depicted somewhat distinct yet overlapping roles for these isoforms in physiological and pathophysiological conditions (10, 11). Dysregulation of Akt is known to trigger a number of diseases including cardiovascular diseases, metabolic disorders and cancer (12–14). A dominant negative Akt2 mutation (R274H) has been demonstrated to be linked with severe hyperinsulinemia and diabetes mellitus in human (15). In mouse models, knockout of Akt2 triggers overt global pre-diabetic insulin resistance (11, 16) and myocardial contractile dysfunction (17) although the precise mechanism(s) underscoring Akt2 ablation-associated myocardial contractile and geometric derangements remains elusive.

Autophagy is a highly conserved intracellular lysosomal catabolic process to degrade aged, damaged or dysfunctional proteins, intracellular organelles and cytoplasmic components to maintain cellular homeostasis (18–20). Basal level of autophagy plays a unique housekeeping role in the regulation of cardiac geometry and function (21–23). Impaired autophagy may contribute to various end organ complications in insulin resistance and diabetes, including cardiomyopathy and nephropathy (24, 25). Autophagy is believed to be downregulated in insulin resistance while tissue-specific autophagy knockout mice displayed overt insulin resistance (25–27). Autophagy is usually regulated by both mTOR (mammalian target of rapamycin)-dependent and -independent mechanisms (28). mTOR pathway is deemed the classical regulation route of autophagy, which negatively regulates autophagy involving two functional complexes: mTORC1 and mTORC2, with a much more predominant role for mTORC1. Inhibition of mTORC1 can induce autophagy which is associated with reduced phosphorylation of two downstream targets p70S6K (ribosomal protein S6 kinase-1) and 4E-BP1 (translation initiation factor 4E-binding protein-1) (18,

29). Various signaling molecules/pathways may function as upstream regulators for mTORC1 to control autophagy, including Rag (Ras-related GTP-binding protein) (30, 31), PI3K/Akt/TSC (tuberous sclerosis complex) (29, 32, 33) and AMPK/TSC (34–37). Despite the regulation of autophagy by mTORC1 and its upstream signaling components, several mTOR-independent pathways may also participate in the regulation of autophagy. Up-to-date, a number of cellular machineries including inositol (38), Ca^{2+} /calpain (39), cAMP/Epac/Ins(1,4,5) $\text{p}3$ (39), c-Jun N terminal kinase (JNK)/Beclin1/PI3K (40), and p38/Atg9 (41) have been reported to regulate autophagy in a mTOR-independent manner.

Trehalose, a natural occurring α -linked disaccharide widely distributed in non-mammalian species such as fungi, yeast, bacteria, invertebrates, insects and plants, functions to provide energy sources and protects the integrity of cells against various environmental stresses (42–45). Trehalose has also been demonstrated to protect against apoptosis in an autophagy-dependent manner (42, 43). Trehalose induces autophagy by facilitating the recruitment of LC3B to the autophagosomal membranes in an mTOR-independent manner (42). Nonetheless, the precise signaling mechanism underneath trehalose-regulated autophagy still remains unclear. To this end, this study was designed to examine the effect of Akt2 knockout-induced insulin resistance on myocardial function and geometry as well as the impact of trehalose on Akt2 knockout-induced myocardial anomalies, if any. To better elucidate the underlying mechanism of trehalose-induced autophagy in the regulation of cardiac geometry and function, potential mTOR-independent cellular regulatory pathways were examined. Our data suggested that Akt2 deletion may directly contribute to the down-regulation of autophagy and up-regulation of apoptosis and ER stress due to the loss of transmission of insulin signaling to mTORC1 pathway. More importantly, trehalose treatment was capable of restoring the level of autophagy (likely through improved autophagy flux) and inhibiting apoptosis in a p38 MAPK-dependent albeit mTOR-independent manner.

2. MATERIALS AND METHODS

2.1 Experimental animals and trehalose treatment

All animal experimental procedures carried out here were approved by the Animal Use and Care Committees at the University of Wyoming (Laramie, WY, USA). Adult Akt2 knockout (Akt2^{-/-}) mice were obtained from Prof. Morris Birnbaum at the University of Pennsylvania (Philadelphia, PA, USA) and their wild-type (WT) littermates were used as wild-type controls (11). All mice were housed in a temperature-controlled room ($22.8 \pm 2.0^\circ\text{C}$, 45–50% humidity) under a 12 hr/12 hr light/dark and allowed access to food and water ad libitum. WT and Akt2 knockout mice were divided into two groups: one group with trehalose treatment, mice were first administered with trehalose (1 mg/g/day) intraperitoneal injection for two days and then given 2% (w/v) trehalose (Sigma-Aldrich Ltd, St. Louis, MO, USA) in drinking water for two months, while the other group were offered regular drinking water (46).

2.2 Intraperitoneal glucose tolerance test (IPGTT)

Following trehalose treatment, mice were fasted for 12 hrs and were then given an intraperitoneal injection of glucose (2 g/kg body weight). Blood samples were drawn from tail veins and serum glucose levels were determined immediately before glucose challenge, as well as 15, 30, 60 and 120 min thereafter using the CONTOUR™ NEXT EZ blood glucose monitoring system (Bayer Diabetes Care, Tarrytown, NY, USA) (47).

2.3 Echocardiographic assessment

Cardiac geometry and function were evaluated in anesthetized (ketamine 80 mg/kg and xylazine 12 mg/kg, i.p.) mice using the two-dimensional guided M-mode echocardiography (Philips SONOS 5500) equipped with a 15–6 MHz linear transducer (Phillips Medical Systems, Andover, MD, USA). Left ventricular (LV) anterior and posterior wall dimensions during diastole and systole were recorded from three consecutive cycles in M-mode using the method adopted by the American Society of Echocardiography. Fractional shortening was calculated from LV end-diastolic (LVEDD) and end-systolic (LVESD) diameters using the equation $(LVEDD - LVESD)/LVEDD \times 100$. Echocardiographic LV mass was estimated using the following equation $[(LVEDD + \text{septal wall thickness} + \text{posterior wall thickness})^3 - LVEDD^3] \times 1.055$, where 1.055 (mg/mm³) is the density of myocardium. Heart rates were averaged over 10 consecutive cycles (37, 48).

2.4 Isolation of murine cardiomyocytes

After ketamine/xylazine sedation, hearts were removed and were mounted onto a temperature-controlled (37°C) Langendorff system. Following perfusing with a modified Tyrode solution (Ca²⁺ free) for 2 min, the heart was digested for 16–20 min with a Ca²⁺-free KHB buffer containing Liberase Blendzyme 4 (Hoffmann-La Roche Inc., Indianapolis, IN, USA). The modified Tyrode's solution (pH 7.4) contained (in mM): NaCl 135, KCl 4.0, MgCl₂ 1.0, HEPES 10, NaH₂PO₄ 0.33, glucose 10, and butanedione monoxime 10, and the solution was gassed with 5% CO₂–95% O₂. Digested hearts were then removed from the cannula and left ventricles were cut into small pieces in the modified Tyrode's solution. Tissue pieces were gently agitated and pellet of cells was resuspended. Extracellular Ca²⁺ was added incrementally back to 1.20 mM over a period of 30 min. A yield of at least 60–70% viable rod-shaped cardiomyocytes with clear sarcomere striations was achieved (which was unaffected by Akt2 ablation or trehalose treatment). Only rod-shaped cardiomyocytes with clear edges were selected for mechanical and intracellular Ca²⁺ studies (49).

2.5 Cell shortening/relengthening

Mechanical properties of myocytes were assessed using a SoftEdge Myocam system (IonOptix Corporation, Milton, MA, USA). IonOptix SoftEdge software was used to capture changes in cardiomyocyte length during shortening and relengthening. In brief, cardiomyocytes were placed in a Warner chamber mounted on the stage of an inverted microscope (Olympus IX-70) and superfused (~1 ml/min at 25°C) with a buffer containing (in mM) 131 NaCl, 4 KCl, 1 CaCl₂, 1 MgCl₂, 10 glucose, and 10 HEPES at pH 7.4. Myocytes were field stimulated with supra-threshold voltage at a frequency of 0.5 Hz (unless otherwise stated), 3-ms duration, using a pair of platinum wires placed on opposite

sides of the chamber connected to a FHC stimulator (Brunswick, NE, USA). The myocyte being studied was displayed on the computer monitor using an IonOptix MyoCam camera. IonOptix SoftEdge software was used to capture changes in cell length during shortening and relengthening. Cell shortening and relengthening were assessed using the following indices: peak shortening (PS)-the amplitude myocytes shortened on electrical stimulation, which is an indicative of peak ventricular contractility; time-to-PS (TPS) - the duration of myocyte shortening, which is an indicative of systolic duration; time-to-90% relengthening (TR₉₀) - the duration to reach 90% relengthening, which represents cardiomyocyte diastolic duration (90% rather than 100% relengthening was used to avoid noisy signal at baseline concentration); and maximal velocities of shortening (+ dL/dt) and relengthening (– dL/dt) - maximal slope (derivative) of shortening and relengthening phases, which are indicatives of maximal velocities of ventricular pressure rise/fall (37, 48).

To further assess the role of p38 MAPK and Foxo1 in Akt2 knockout-induced cardiomyocyte contractile responses, cardiomyocytes from adult WT and Akt2^{-/-} mice were treated with the p38 MAPK activator anisomycin (10 µg/ml) for 1 hr (50) or the Foxo1 inhibitor AS1842856 (10 µM) for 1 hr (51) in the presence or absence of trehalose (100 mM) for 2 additional hrs (45). To assess the impact of Akt2 knockout and trehalose on autophagic flux, cardiomyocytes from adult WT and Akt2^{-/-} mice were treated with the lysosomal inhibitor bafilomycin A1 (100 nM) for 2 hrs (52) in the presence or absence of trehalose (100 mM) for 1 additional hr prior to the assessment of autophagy protein markers.

2.6 Intracellular Ca²⁺ handling

A cohort of myocytes was loaded with fura-2/AM (0.5 µM) for 10 min and fluorescence intensity was recorded with a dual-excitation fluorescence photomultiplier system (Ionoptix). Myocytes were placed onto an Olympus IX-70 inverted microscope and imaged through a Fluor × 40 oil objective. Cells were exposed to light emitted by a 75W lamp and passed through either a 360 or a 380 nm filter, while being stimulated to contract at 0.5 Hz. Fluorescence emissions were detected between 480 and 520 nm and qualitative change in fura-2 fluorescence intensity (FFI) was inferred from FFI ratio at the two wavelengths (360/380). Fluorescence decay time (single exponential decay rate) was measured as an indication of intracellular Ca²⁺ clearing rate (37).

2.7 Caspase 3 assay

Tissue was homogenized and centrifuged (10,000 g at 4°C, 10 min) and pellets were lysed in an ice-cold cell lysis buffer (50 mM HEPES, 0.1% CHAPS, 1 mM dithiothreitol, 0.1 mM EDTA, 0.1% NP40). The assay was carried out in a 96-well plate with each well containing 30 µl cell lysate, 70 µl of assay buffer (50 mM HEPES, 0.1% CHAPS, 100 mM NaCl, 10 mM dithiothreitol, and 1 mM EDTA) and 20 µl of caspase 3 colorimetric substrate Ac-DEVD-pNA. The 96-well plate was then incubated at 37°C for 1 hr, when the caspase in the sample was allowed to cleave the chromophore p-NA from the substrate molecule. The absorbency of cleaved p-NA was detected at 405 nm with caspase 3 activity being proportional to color reaction. Protein content was determined by Bradford method. Caspase 3 was calculated as the picomoles of p-NA released per microgram of protein per minute (53).

2.8 Wheat germ agglutinin (WGA) staining

Ventricular tissues were stained with FITC-conjugated wheat germ agglutinin (Sigma, St. Louis, MO, USA) and cardiomyocyte cross-sectional areas were calculated from 400 randomly selected cells on a digital microscope (x400) using the Image J (version 1.34S) software (54).

2.9 TUNEL staining

Terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL) staining of DNA strand breaks was performed using the In Situ Death Detection Kit (Roche, Branchburg, NJ, USA), as previously described (55). Fresh frozen heart sections were cut using a Leica cryostat (Model CM3050S, Leica Microsystems) to produce 7- μ m tissue sections. Tissue sections were fixed with 4% paraformaldehyde for 20 min and permeabilized in 0.1% Triton X-100 in 0.1% sodium citrate for 2 min at 4°C. To distinguish myocytes from nonmyocytes in the heart, tissue sections were incubated with anti-Desmin (Cell Signaling, 1:100) followed by incubation with anti-Rabbit secondary conjugate with Alex Fluor 568 (Invitrogen). Fifty microliters of a reaction mixture containing terminal deoxynucleotidyl transferase (TdT), fluorescein-dUTP was added to each section and incubated in a humidified chamber for 60 min at 37°C. Sections were washed three times with PBS and counterstained with 4',6'-diamidino-2-phenylindole (DAPI, 5 μ g/ml) for 1 min. Slides were mounted with the Prolong Gold mounting medium (Invitrogen, Carlsbad, CA, USA), and five images per tissue section were obtained using a Zeiss confocal microscope equipped with an Olympus MagnaFire SP digital camera and ImagePro image analysis software as previously described (56).

2.10 Western blot analysis

Heart tissues from study mice were homogenized and sonicated in a lysis buffer containing 20 mM Tris (pH 7.4), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton, 0.1% sodium dodecyl sulfate, and a protease inhibitor cocktail. For in vitro study, isolated cardiomyocytes from WT mice treated with paraquat or rapamycin were sonicated in the lysis buffer as described above. Myocardial protein samples were incubated with anti-Akt, anti-phosphorylated Akt (Ser⁴⁷³), anti-Atg5, anti-LC3B, anti-Bec1, anti-p38 MAPK, anti-phosphorylated p38 MAPK (Thr¹⁸⁰/Tyr¹⁸²), anti-Bad, anti-Bax, anti-Bcl2, anti-cleaved-caspase 3, anti-cleaved-caspase 9, anti-GRP78, anti-JNK, anti-phosphorylated JNK (Thr¹⁸³/Tyr¹⁸⁵), anti-glyceraldehyde-3-phosphate dehydrogenase (GAPDH; loading control) (1:1000; Cell Signaling Technology, Danvers, MA, USA), anti-p62 (1:1000; Guinea Pig; Enzo Life Sciences, Plymouth Meeting, PA, USA), anti-IRE1 α , anti-phosphorylated IRE1 α (Ser724) (1:1000, Abcam, Cambridge, MA, USA), anti-GADD153 (1:1000, Santa Cruz Biotechnology, Inc. Dallas, TX, USA) antibodies. Horseradish peroxidase-coupled secondary antibodies were used for membrane incubation. After immunoblotting, the films were scanned and detected with a Bio-Rad calibrated densitometer and the intensity of immunoblot bands was normalized with corresponding band intensity of GAPDH (37).

2.11 Real-time quantitative PCR

Total mRNA was extracted from heart tissues using the Trizol[®] Reagent per the manufacturer's instruction (Invitrogen, Carlsbad, CA, USA). Total mRNA was treated with DNase I (RNase-free) (Invitrogen) and was purified by a RNeasy Mini Kit (Qiagen, Valencia, CA, USA). It was then reverse transcribed into cDNA by iScript[™] cDNA Synthesis Kit (Bio-Rad, Hercules, CA, USA). RT-PCR was performed on CFX96[™] RT-PCR detection system (Bio-Rad). The SsoAdvanced[™] Universal SYBR[®] Green Supermix (Bio-Rad) was used together with the primers listed in Table 1. PCR reaction cycles were as follows: 3 min at 95°C, then 10 s at 95°C, 30 s at 56°C for 40 cycles, finally for melt curve analysis by increment 0.5°C from 55°C to 95°C for 30 sec. After amplification, melting curves were used to confirm product purity. Fluorescent signals were normalized to an internal reference, and the threshold cycle (CT) was set within the exponential phase of the PCR. Results are expressed relative to GAPDH using CT method (57).

2.12 Statistical analysis

Data are mean $\pm$ SEM. Statistical significance ($p < 0.05$) for each variable was estimated by a one-way analysis of variance followed by Tukey's test for *the post hoc* analysis.

3. RESULTS

3.1 Effect of trehalose treatment on Akt2 knockout-induced cardiac hypertrophy and insulin resistance

Akt2 knockout overtly increased heart-to-body weight ratio as well as cardiomyocyte cross-sectional area, which were attenuated by trehalose treatment (Fig. 1A–C). Similarly, expression levels of hypertrophic genes (ANP and BNP), were significantly upregulated in Akt2^{−/−} mice, the effect of which was mitigated by trehalose treatment (Fig. 1D–E). However, neither Akt2 knockout nor trehalose treatment affected expression levels of both α -MHC and β -MHC, except that trehalose significantly reduced the expression level of β -MHC in Akt2^{−/−} mice (Fig. 1F–G). Following intraperitoneal glucose challenge, serum glucose levels began to drop after peaking at 15 min and returned towards near baseline after 120 min, although mice subjected to trehalose treatment displayed slightly higher serum glucose levels. Akt2^{−/−} mice exhibited higher serum glucose levels between 15 and 120 min with a somewhat delayed peak at 30 min (Fig. 1H). This is supported by significantly greater area underneath the IPGTT curve (AUC) in Akt2^{−/−} mice, indicating glucose intolerance in Akt2^{−/−} insulin resistant mice. Trehalose did not overtly affect IPGTT curve in Akt2^{−/−} mice not did it alter AUC in both WT and Akt2^{−/−} mice (Fig. 1I).

3.2 Echocardiographic characteristics of WT and Akt2 knockout mice with or without trehalose treatment

Neither Akt2 knockout nor trehalose treatment overtly affected heart rate. However, Akt2 knockout significantly reduced fractional shortening while increased LVEDD, LVESD and LV mass/body weight ratio, the effect of which was mitigated or significantly attenuated by trehalose treatment. Trehalose itself did not exert any significant effect on any echocardiographic indices tested in WT mice (Fig. 2).

3.3 Effect of trehalose and Akt2 knockout on cardiomyocyte contractile properties

Neither Akt2 knockout nor trehalose treatment affected resting cell length. However, cardiomyocytes from Akt2^{-/-} mice displayed significantly decreased PS and $\pm$ dL/dt associated with overtly prolonged TPS and TR₉₀. Trehalose treatment abrogated Akt2 knockout-induced cardiomyocyte mechanical dysfunctions without eliciting any obvious effect by itself (Fig. 3).

3.4 Cardiomyocyte intracellular Ca²⁺ transient properties

To investigate the possible mechanisms behind cardiomyocyte contractile properties in response to Akt2 knockout and trehalose treatment, intracellular Ca²⁺ handling was evaluated using Fura-2 fluorescence. Our data indicated that Akt2 knockout significantly suppressed the resting and peak intracellular Ca²⁺ levels, as well as the rise in fura-2 fluorescence intensity (FFI) associated with prolonged intracellular Ca²⁺ clearance. These changes of intracellular Ca²⁺ properties in response to Akt2 ablation were significantly attenuated or mitigated by trehalose treatment. Trehalose treatment itself failed to elicit any overt intracellular Ca²⁺ responses (Fig. 4).

3.5 Trehalose attenuates Akt2 knockout-induced apoptosis

Given that apoptosis may play a key role in Akt2 knockout-induced cardiac dysfunction (17), apoptosis was examined in murine cardiomyocytes from WT and Akt2^{-/-} mice using confocal microscopy. Our triple immunofluorescence TUNEL staining displayed that Akt2 ablation significantly promoted apoptosis in cardiomyocytes, the effect of which was abrogated by trehalose. Trehalose itself did not affect cardiomyocyte apoptosis (Fig. 5A–B). Consistent with the findings from TUNEL staining, Akt2 knockout significantly increased Caspase 3 activity, a key marker for apoptosis, the effect of which was negated by trehalose (Fig 5C).

3.6 Effect of trehalose and Akt2 knockout on autophagy protein markers

To explore the potential role of autophagy in Akt2 knockout-induced cardiac damage, protein levels of the autophagy makers Atg5, Beclin1, LC3I/II and the autophagy cargo adaptor protein p62 were evaluated using western blot analysis. Our data revealed that Akt2 knockout significantly suppressed protein levels of Atg5, Beclin1, p62, LC3II as well as the LC3B-II-to-LC3B-I ratio. Trehalose treatment itself significantly promoted protein levels of these autophagy markers (including Atg5, LC3B-II, LC3B-II-to-LC3B-I ratio) and obliterated Akt2 knockout-induced changes in autophagy protein markers (with the exception of LC3B-II). In addition, trehalose significantly reduced protein level of LC3B-I in Akt2^{-/-} mice. Protein level of p62 was significantly reduced in myocardium from both WT and Akt2^{-/-} mice following trehalose treatment (Fig 6).

Given the changes of p62 cargo adaptor protein in WT and Akt2^{-/-} mice with or without trehalose treatment, autophagic flux was examined in both WT and Akt2^{-/-} mice. Freshly isolated cardiomyocytes from WT and Akt2^{-/-} mice were treated with the lysosomal inhibitor bafilomycin A1 (100 nM for 2 hrs) in the presence or absence of trehalose treatment (100 mM). In accordance with our *in vivo* findings, trehalose treatment significantly promoted protein levels of LC3B-II and LC3B-II-to-LC3B-I ratio and

suppressed protein level of p62 in both WT and Akt2^{-/-} groups. Moreover, trehalose nullified Akt2 knockout-induced changes in autophagy makers. Although bafilomycin A1 itself failed to affect protein levels of LC3B-II, LC3B-I and LC3B-II-to-LC3B-I ratio in WT group, it significantly increased protein levels of p62. In addition, bafilomycin A1 cancelled off trehalose-induced changes in p62 and LC3B-II-to-LC3B-I ratio in both WT and Akt2^{-/-} mice. Last but not least, bafilomycin A1 promoted trehalose-induced changes in LC3B-I level in both WT and Akt2^{-/-} mice (Fig 7).

3.7 Effect of trehalose and Akt2 knockout on apoptosis and ER stress protein markers

Further assessment of apoptosis using Western blot analysis revealed upregulated protein levels of the pro-apoptotic proteins BAD, BAX, cleaved Caspase 3 and cleaved Caspase 9 along with down-regulated anti-apoptotic protein BCL2 in Akt2^{-/-} mice, the effects of which were significantly attenuated or ablated by trehalose treatment. Little effect was noted for trehalose treatment itself on these apoptotic markers in WT mice (Fig. 8). To further examine the potential role of ER stress in Akt2 knockout- and trehalose-induced cardiac responses, protein levels of the ER stress markers GRP78, IRE1 α , JNK and GADD153 were evaluated using western blot analysis in WT and Akt2^{-/-} mice. Our result showed that Akt2 knockout significantly upregulated protein expression of GRP78, phosphorylation of IRE1 α and JNK, the effects of which were mitigated by trehalose treatment. Neither Akt2 knockout nor trehalose treatment exhibited any notable effect on protein level of GADD13 as well as the pan expression of IRE1 α and JNK (Fig. 9).

3.8 Effect of trehalose and Akt2 knockout on pan and phosphorylated Akt, Foxo1 and p38 MAPK

To further explore the signaling mechanisms behind trehalose-mediated autophagy responses in Akt2 knockout murine model, Akt, Foxo1 and p38 MAPK were examined. Our data revealed significantly reduced pan protein expression of Akt together with absent Akt2 protein expression in Akt2^{-/-} mice, validating the murine model (Fig. 10A). In addition, Akt phosphorylation was significantly reduced in Akt2^{-/-} mice. Although there is little significant change with the pAkt-to-Akt ratio, the ratio tended to be elevated (although not reaching significance) in Akt2^{-/-} mice and trehalose treatment tended to reduce the pAkt-to-Akt ratio. Akt2 knockout significantly elevated phosphorylation of Foxo1 and p38 MAPK, the effect of which was significantly abrogated by trehalose treatment. Trehalose treatment itself significantly decreased the phosphorylated levels of both Foxo1 and p38 MAPK (but not that of Akt) in WT mice. Neither Akt2 knockout nor trehalose displayed any notable effect on pan expression of Foxo1 and p38 MAPK (Fig. 10).

3.9 Effect of p38 MAPK activator anisomycin, Foxo1 inhibitor AS1842856, and trehalose on Akt2 knockout induced cardiomyocyte contractile defects

To further elucidate a cause-effect relationship for trehalose, p38 MAPK and Foxo1 in Akt2 knockout-induced cardiac contractile response, freshly isolated cardiomyocytes from WT and Akt2^{-/-} mice were exposed to the p38 MAPK activator anisomycin and the Foxo1 inhibitor (through phosphorylation) AS1842856 in the presence or absence of trehalose for 2 hrs. Our data revealed that trehalose incubation significantly attenuated Akt2 knockout-induced cardiomyocyte contractile dysfunction including depressed peak shortening, and

maximal velocity of shortening /relgthening as well as prolonged TR₉₀. Treatment of anisomycin and AS1842856 significantly suppressed peak shortening, maximal velocity of shortening/relengthening without affecting resting cell length and duration of shortening/relengthening in cardiomyocytes from WT mice. Interestingly, the trehalose-elicited beneficial effects on Akt2^{-/-} mice were mitigated by the p38 MAPK activator anisomycin or the Foxo1 inhibitor AS1842856 (Fig. 11).

4. DISCUSSION

The salient findings from our current work suggested that Akt2 knockout elicits damages in myocardial geometry and function, and intracellular Ca²⁺ homeostasis (as evidenced by enlarged LVESD, normalized LV mass, enlarged cardiomyocyte cross-sectional area, upregulated levels of prohypertrophic genes and depressed fractional shortening, reduced peak shortening, maximum velocity of shortening/relengthening, prolonged duration of shortening/relengthening, decreased resting and electrically stimulated rise in intracellular Ca²⁺). Our data revealed that the compromised cardiac function and intracellular Ca²⁺ handling seen in Akt2^{-/-} mice were accompanied with activated apoptosis (elevated Caspase 3 activity, TUNEL staining and upregulated apoptotic markers), ER stress (upregulated ER stress (protein markers) as well as suppressed autophagy (downregulated autophagic protein markers). On one hand, our findings demonstrated that trehalose effectively attenuated apoptosis and ER stress, consistent with the earlier finding (43). On the other hand, the autophagy inducer trehalose effectively turned on autophagy in the autophagy-deficient Akt2^{-/-} insulin resistant mice. Our findings provided compelling evidence for the possible mechanism of trehalose on cardiac function, geometry and autophagy regulation in insulin resistance. In particular, phosphorylation of Foxo1 and p38 MAPK signaling molecules was elevated in Akt2 ablation-induced insulin resistant mouse hearts, the effects of which were significantly attenuated or negated by trehalose. Akt phosphorylation was dampened in Akt2^{-/-} murine hearts, the pattern of which was not affected by trehalose treatment. These data suggested that trehalose may activate autophagy possibly associated with dephosphorylation of Foxo1 and p38 MAPK.

Our findings validated Akt2 ablation as a murine model of insulin resistance as evidenced by increased area underneath IPGTT curve with unchanged baseline blood glucose levels (11). Akt2^{-/-} mice display a significant rise in serum insulin levels (11). Akt is known to play a pivotal role in glucose metabolism and myocardial function (12, 14, 58). Our echocardiographic data revealed that Akt2 knockout significantly increased LVESD and LV mass along with decreased fractional shortening, in concordance with our previous observations of myocardial dysfunction with Akt2 ablation (16, 17). Akt2 ablation-induced cardiac remodeling (elevated normalized LV mass) is further substantiated by enlarged cardiomyocyte cross-sectional area, upregulated levels of prohypertrophic genes. In addition, data from our study suggested cardiomyocyte contractile anomalies including reduced peak shortening and maximal velocity of shortening/relengthening, and prolonged duration of shortening/relengthening in mice with Akt2 ablation. Furthermore, intercellular Ca²⁺ homeostasis was disrupted in cardiomyocytes from Akt2^{-/-} mice, as evidenced by depressed resting and peak intracellular Ca²⁺ levels and prolonged intracellular Ca²⁺ clearance. These findings are in line with our previous reports on Akt2^{-/-} murine model (16, 17) and with

other types of insulin resistance murine models (such as sucrose- or high fat diet-induced insulin resistance) (4, 5, 59). Nonetheless, trehalose treatment significantly ameliorated or mitigated Akt2 knockout-induced myocardial contractile dysfunction and intracellular Ca^{2+} mishandling. These data indicate that Akt2 may participate in the regulation of glucose metabolism and myocardial function although the underlying mechanism remains unclear.

Findings from our current study revealed an essential role for apoptosis and ER stress in Akt2 ablation-induced cardiac anomalies. Emerging evidence has indicated a role for apoptosis and ER stress in insulin resistance-triggered myocardial contractile dysfunction (60–62). In this study, we observed that Akt2 knockout induced overt apoptosis and ER stress as evidenced by protein markers of apoptosis and ER stress, Caspase 3 activity and TUNEL staining, supporting a likely role for apoptosis and ER stress in cardiac contractile and geometric (hypertrophic remodeling) anomalies in Akt2 ablation-induced insulin resistance. This is consistent with the changes in pro- and anti-apoptotic signaling molecules including BAD, BAX, cleaved-Caspase 3/9 and Bcl-2 as well as the ER stress protein markers (GRP78, phosphorylation of IRE1 α and JNK) in hearts from Akt2^{-/-} mice, consistent with earlier findings from our group using the same model (17). Previous results have depicted an indispensable role for Akt2 in survival, apoptosis and ER protein synthesis (63–65). Most importantly, Cui and colleagues found that knockout of Akt2 in human glioblastoma cells promoted apoptosis (66).

Our results revealed, for the first time, that Akt2 knockout suppresses autophagy in the heart manifested as decreased expression level of Atg5, Beclin1, LC3II and LC3II-to-LC3I ratio (concurrent with functional defect), as well as the autophagosome cargo protein p62. Trehalose treatment effectively nullified Akt2 knockout-induced changes in autophagy makers. Inhibition of lysosomal enzymes using bafilomycin A1 cancelled off trehalose-induced changes in p62 and LC3B-II-to-LC3B-I ratio in WT and Akt2^{-/-} mice, suggesting a permissive role for lysosomal activity in trehalose-offered beneficial autophagy responses. Earlier evidence has demonstrated that inactivation of autophagy leads to cytoplasmic protein inclusions (composed of degenerated proteins and accumulation of deformed organelles), contributing to the onset and development of heart diseases, diabetes, liver injury and neurodegenerative diseases (67). Previous studies have indicated a unique role for defective cardiomyocyte autophagy in various cardiac anomalies such as cardiomyopathy and heart failure, as well as premature cardiac aging (68–71). Basal autophagy and autophagic flux are crucial for cardiomyocyte homeostasis (72). Cardiomyocyte-specific Atg5 knockout mice developed dilated cardiomyopathy in adulthoods (71). We observed suppressed basal autophagic levels in Akt2^{-/-} mice, manifested as the decreased autophagosome formation along with a low p62 accumulation (possibly due to the low available lysosomal degradation substrates). Bafilomycin A1 failed to alter LC3B-II-to-LC3B-I ratio in Akt2^{-/-} mice although the lysosomal inhibition approach enhanced p62 accumulation as expected.

Trehalose has been characterized as an effective autophagy inducer in various mammalian cells (42, 73, 74). Trehalose is shown to protect against hypoxic and anoxic injuries and suppress protein aggregation (44, 75). *In vivo* findings have revealed that trehalose-induced autophagy may facilitate the clearance of mutant forms of protein aggregates in

neurodegenerative disorders such as Huntington and Parkinson diseases in murine models (6, 44, 76, 77). As an autophagy inducer, trehalose may turn on autophagy in an mTOR-independent manner (42). Despite of the reported existence of mTOR-independent autophagy pathways, the precise mechanism behind trehalose-induced autophagy has not been well characterized. Casillo and colleagues found that trehalose enhanced nuclear translocation of Foxo1, an important transcription factor to turn on autophagy (45, 78). In our hands, a pronounced dephosphorylation of Foxo1 was identified following trehalose treatment, denoting a drop in inactive form of Foxo1, which would promote more active Foxo1 for nuclear translocation to promote autophagy. A novel mTOR-independent autophagy pathway was reported where p38 MAPK plays a crucial role in regulating autophagy through the p38 MAPK interacting protein (p38IP) (41). It is believed that p38IP possesses dual binding affinities for p38 MAPK and autophagy protein 9 (Atg9). Upon p38 MAPK activation, p38IP binds with Atg9 to downregulate autophagy. On the other hand, the p38IP-Atg9 interaction is disengaged with inhibition of p38 MAPK, leading to redistribution of Atg9 to the endosomes and autophagy induction. Our data seem to favor a possible role for p38 MAPK in the trehalose-facilitated autophagy (through blockade of p38 MAPK activation and interaction between p38 MAPK and p38IP, thus to release Atg9 to the endosomes).

Foxo family proteins play important roles in insulin signaling, glucose and lipid metabolism (79, 80). Elevated insulin levels inactivate Foxo1 via PI3K–Akt signaling, furthermore Foxo1 was able to control the upstream signaling elements to govern insulin sensitivity and glucose metabolism (79). It has been reported that both Akt1 and Akt2 isoforms may regulate Foxo1 activity (81). Among Akt family members, Akt2 has been extensively studied in the past decade (11, 63, 64, 66). Akt2 plays an important role in sensing insulin signaling, with impaired insulin sensitivity in Akt2^{-/-} mice despite intact Akt1 and Akt3 signaling. Our data revealed that Foxo1 was inactivated (elevated phosphorylation) in Akt2^{-/-} mice. However, phosphorylation of Akt was dampened only by a small portion in Akt2^{-/-} mice compared with their littermates, suggested that phosphorylation of Akt1 may compensate the loss function of Akt2.

Although it is beyond the scope of the present study, Akt2 kinase is known to negatively regulate p38 MAPK through ASK1, a member of the mitogen-activated protein kinase kinase kinase family which activated MKK3/MKK6–p38 MAPK signaling cascade (63). Akt2 kinase complex may phosphorylate ASK1 at Ser⁸³, resulting in inhibition of ASK1 activity and blockade of p38 MAPK activation, consistent with our current findings. In Akt2^{-/-} mice, activation of p38 MAPK may be due to the loss of function for Akt2. In addition, ASK1 and p38 MAPK may promote Bax conformational change (63) to regulate apoptosis and ER stress. Bax is upregulated to trigger apoptosis with Akt2 ablation. Therefore, loss of function in Akt2 and activation of p38 MAPK may both contribute to the induction of apoptosis and ER stress as well as the inhibition of autophagy. It is noteworthy that insulin resistance is associated with dampened cardiac energy utilization efficiency (glucose oxidation versus fatty acid oxidation) (1). Possible contribution of trehalose treatment to metabolic switch between glucose oxidation and fatty acid oxidation under insulin resistance cannot be excluded at this time.

Our *in vivo* data indicate that Akt2 knockout may inhibit autophagy and activate apoptosis in the heart accompanied with decreased Foxo1 and/or enhanced p38 MAPK activities, the effects of which were reversed by trehalose treatment, validating a likely role for Foxo1 and p38 MAPK in trehalose-offered cardiac protection. This is convincingly supported by our *in vitro* findings where trehalose-offered beneficial effect is nullified (masked) by either the Foxo1 inhibitor AS1842856 (via phosphorylation) or the p38 MAPK activator anisomycin. Anisomycin and AS1842856 both compromised cardiac contractile response in cardiomyocytes from WT mice, supporting the role for p38 MAPK and Foxo1 in the maintenance of cardiomyocyte homeostasis.

In summary, our study suggested that Akt2 knockout may induce cardiac functional and geometric anomalies via inhibition of autophagy as well as activation of apoptosis and ER stress. Our data revealed a pivotal and permissive role of Akt2 signaling molecule in the regulation of insulin signaling, autophagy, apoptosis and ER stress, possibly via Foxo1 and p38 MAPK. This is supported by our observation of decreased Foxo1 (higher phosphorylation) and increased p38 MAPK activity in hearts from Akt2^{-/-} mice. Our results suggested that trehalose treatment may promote key autophagic markers associated with Foxo1, an essential transcriptional factor in the regulation of autophagy, and the inhibition of p38 MAPK, a novel player for mTOR-independent autophagy signaling. The fact that lysosomal inhibition using bafilomycin A1 negated trehalose-induced autophagy response suggests a vital role for autophagy flux in trehalose-offered cardiac protection. Although it is still somewhat premature to discern the precise mechanism underlying Akt2 ablation- and trehalose-induced changes in cardiac function and geometry, our work should shed some lights towards a better understanding of the therapeutic potentials for autophagy, Foxo1 and p38 MAPK in Akt2 deficiency-induced insulin resistance.

Acknowledgments

This work was supported by NIGMS 5P20GM103432.

REFERENCE

1. Reaven GM. Pathophysiology of insulin resistance in human disease. *Physiological reviews*. 1995; 75:473–486. [PubMed: 7624391]
2. Wang CC, Goalstone ML, Draznin B. Molecular mechanisms of insulin resistance that impact cardiovascular biology. *Diabetes*. 2004; 53:2735–2740. [PubMed: 15504952]
3. Dong F, Fang CX, Yang X, Zhang X, Lopez FL, Ren J. Cardiac overexpression of catalase rescues cardiac contractile dysfunction induced by insulin resistance: Role of oxidative stress, protein carbonyl formation and insulin sensitivity. *Diabetologia*. 2006; 49:1421–1433. [PubMed: 16586065]
4. Fang CX, Dong F, Ren BH, Epstein PN, Ren J. Metallothionein alleviates cardiac contractile dysfunction induced by insulin resistance: role of Akt phosphorylation, PTB1B, PPARgamma and c-Jun. *Diabetologia*. 2005; 48:2412–2421. [PubMed: 16172869]
5. Roe ND, He EY, Wu Z, Ren J. Folic acid reverses nitric oxide synthase uncoupling and prevents cardiac dysfunction in insulin resistance: role of Ca²⁺/calmodulin-activated protein kinase II. *Free radical biology & medicine*. 2013; 65:234–243. [PubMed: 23820268]
6. Chaudhary RK, Kardani J, Singh K, Banerjee R, Roy I. Deciphering the roles of trehalose and Hsp104 in the inhibition of aggregation of mutant huntingtin in a yeast model of Huntington's disease. *Neuromolecular medicine*. 2014; 16:280–291. [PubMed: 24248470]

7. Sharma N, Arias EB, Sajan MP, MacKrell JG, Bhat AD, Farese RV, Cartee GD. Insulin resistance for glucose uptake and Akt2 phosphorylation in the soleus, but not epitrochlearis, muscles of old vs. adult rats. *Journal of applied physiology*. 2010; 108:1631–1640. [PubMed: 20339009]
8. Dummmlerand BA, Hemmings B. Physiological roles of PKB/Akt isoforms in development and disease. *Biochemical Society transactions*. 2007; 35:231–235. [PubMed: 17371246]
9. Leslie NR. The redox regulation of PI 3-kinase-dependent signaling. *Antioxidants & redox signaling*. 2006; 8:1765–1774. [PubMed: 16987030]
10. Hers I, Vincent EE, Tavaré JM. Akt signalling in health and disease. *Cellular signalling*. 2011; 23:1515–1527. [PubMed: 21620960]
11. Cho H, Mu J, Kim JK, Thorvaldsen JL, Chu Q, Crenshaw EB 3rd, Kaestner KH, Bartolomei MS, Shulman GI, Birnbaum MJ. Insulin resistance and a diabetes mellitus-like syndrome in mice lacking the protein kinase Akt2 (PKB beta). *Science*. 2001; 292:1728–1731. [PubMed: 11387480]
12. Semple D, Smith K, Bhandari S, Seymour AM. Uremic cardiomyopathy and insulin resistance: a critical role for akt? *Journal of the American Society of Nephrology : JASN*. 2011; 22:207–215. [PubMed: 20634295]
13. Fresno Vara JA, Casado E, de Castro J, Cejas P, Belda-Iniesta C, Gonzalez-Baron M. PI3K/Akt signalling pathway and cancer. *Cancer treatment reviews*. 2004; 30:193–204. [PubMed: 15023437]
14. Lu M, Wan M, Leavens KF, Chu Q, Monks BR, Fernandez S, Ahima RS, Ueki K, Kahn CR, Birnbaum MJ. Insulin regulates liver metabolism in vivo in the absence of hepatic Akt and Foxo1. *Nature medicine*. 2012; 18:388–395.
15. George S, Rochford JJ, Wolfrum C, Gray SL, Schinner S, Wilson JC, Soos MA, Murgatroyd PR, Williams RM, Acerini CL, Dunger DB, Barford D, Umpoleby AM, Wareham NJ, Davies HA, Schafer AJ, Stoffel M, O'Rahilly S, Barroso I. A family with severe insulin resistance and diabetes due to a mutation in AKT2. *Science*. 2004; 304:1325–1328. [PubMed: 15166380]
16. Li L, Hua Y, Ren J. Short-chain fatty acid propionate alleviates Akt2 knockout-induced myocardial contractile dysfunction. *Experimental diabetes research*. 2012; 2012:851717. [PubMed: 21960994]
17. Hu N, Dong M, Ren J. Hydrogen sulfide alleviates cardiac contractile dysfunction in an Akt2-knockout murine model of insulin resistance: role of mitochondrial injury and apoptosis. *American journal of physiology Regulatory, integrative and comparative physiology*. 2014; 306:R761–R771.
18. Klionskyand SD, Emr DJ. Autophagy as a regulated pathway of cellular degradation. *Science*. 2000; 290:1717–1721. [PubMed: 11099404]
19. Levineand DJ, Klionsky B. Development by self-digestion: molecular mechanisms and biological functions of autophagy. *Developmental cell*. 2004; 6:463–477. [PubMed: 15068787]
20. Mizushima N, Levine B, Cuervo AM, Klionsky DJ. Autophagy fights disease through cellular self-digestion. *Nature*. 2008; 451:1069–1075. [PubMed: 18305538]
21. Kassiotis C, Ballal K, Wellnitz K, Vela D, Gong M, Salazar R, Frazier OH, Taegtmeier H. Markers of autophagy are downregulated in failing human heart after mechanical unloading. *Circulation*. 2009; 120:S191–S197. [PubMed: 19752367]
22. Nakai A, Yamaguchi O, Takeda T, Higuchi Y, Hikoso S, Taniike M, Omiya S, Mizote I, Matsumura Y, Asahi M, Nishida K, Hori M, Mizushima N, Otsu K. The role of autophagy in cardiomyocytes in the basal state and in response to hemodynamic stress. *Nature medicine*. 2007; 13:619–624.
23. Nemchenko A, Chiong M, Turer A, Lavandero S, Hill JA. Autophagy as a therapeutic target in cardiovascular disease. *Journal of molecular and cellular cardiology*. 2011; 51:584–593. [PubMed: 21723289]
24. Barlowand DC, Thomas AD. Autophagy in diabetes: beta-cell dysfunction, insulin resistance, and complications. *DNA and cell biology*. 2015; 34:252–260. [PubMed: 25665094]
25. Kume S, Thomas MC, Koya D. Nutrient sensing, autophagy, and diabetic nephropathy. *Diabetes*. 2012; 61:23–29. [PubMed: 22187371]
26. Liu HY, Han J, Cao SY, Hong T, Zhuo D, Shi J, Liu Z, Cao W. Hepatic autophagy is suppressed in the presence of insulin resistance and hyperinsulinemia: inhibition of FoxO1-dependent expression of key autophagy genes by insulin. *The Journal of biological chemistry*. 2009; 284:31484–31492. [PubMed: 19758991]

27. Shi L, Zhang T, Liang X, Hu Q, Huang J, Zhou Y, Chen M, Zhang Q, Zhu J, Mi M. Dihydromyricetin improves skeletal muscle insulin resistance by inducing autophagy via the AMPK signaling pathway. *Molecular and cellular endocrinology*. 2015; 409:92–102. [PubMed: 25797177]
28. Sarkar S. Regulation of autophagy by mTOR-dependent and mTOR-independent pathways: autophagy dysfunction in neurodegenerative diseases and therapeutic application of autophagy enhancers. *Biochemical Society transactions*. 2013; 41:1103–1130. [PubMed: 24059496]
29. Inoki K, Li Y, Zhu T, Wu J, Guan KL. TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR signalling. *Nature cell biology*. 2002; 4:648–657. [PubMed: 12172553]
30. Sancak Y, Peterson TR, Shaul YD, Lindquist RA, Thoreen CC, Bar-Peled L, Sabatini DM. The Rag GTPases bind raptor and mediate amino acid signaling to mTORC1. *Science*. 2008; 320:1496–1501. [PubMed: 18497260]
31. Kim E, Goraksha-Hicks P, Li L, Neufeld TP, Guan KL. Regulation of TORC1 by Rag GTPases in nutrient response. *Nature cell biology*. 2008; 10:935–945. [PubMed: 18604198]
32. Petiot A, Ogier-Denis E, Blommaert EF, Meijer AJ, Codogno P. Distinct classes of phosphatidylinositol 3'-kinases are involved in signaling pathways that control macroautophagy in HT-29 cells. *The Journal of biological chemistry*. 2000; 275:992–998. [PubMed: 10625637]
33. Heras-Sandoval D, Perez-Rojas JM, Hernandez-Damian J, Pedraza-Chaverri J. The role of PI3K/AKT/mTOR pathway in the modulation of autophagy and the clearance of protein aggregates in neurodegeneration. *Cellular signalling*. 2014; 26:2694–2701. [PubMed: 25173700]
34. Meijerand P, Codogno AJ. AMP-activated protein kinase and autophagy. *Autophagy*. 2007; 3:238–240. [PubMed: 17224623]
35. Kim J, Kundu M, Viollet B, Guan KL. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nature cell biology*. 2011; 13:132–141. [PubMed: 21258367]
36. Gwinn DM, Shackelford DB, Egan DF, Mihaylova MM, Mery A, Vasquez DS, Turk BE, Shaw RJ. AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Molecular cell*. 2008; 30:214–226. [PubMed: 18439900]
37. Wang Q, Yang L, Hua Y, Nair S, Xu X, Ren J. AMP-activated protein kinase deficiency rescues paraquat-induced cardiac contractile dysfunction through an autophagy-dependent mechanism. *Toxicological sciences : an official journal of the Society of Toxicology*. 2014; 142:6–20. [PubMed: 25092649]
38. Sarkar S, Floto RA, Berger Z, Imarisio S, Cordenier A, Pasco M, Cook LJ, Rubinsztein DC. Lithium induces autophagy by inhibiting inositol monophosphatase. *The Journal of cell biology*. 2005; 170:1101–1111. [PubMed: 16186256]
39. Williams A, Sarkar S, Cuddon P, Ttofi EK, Saiki S, Siddiqi FH, Jahreiss L, Fleming A, Pask D, Goldsmith P, O'Kane CJ, Floto RA, Rubinsztein DC. Novel targets for Huntington's disease in an mTOR-independent autophagy pathway. *Nature chemical biology*. 2008; 4:295–305. [PubMed: 18391949]
40. Sarkar S, Korolchuk VI, Renna M, Imarisio S, Fleming A, Williams A, Garcia-Arencibia M, Rose C, Luo S, Underwood BR, Kroemer G, O'Kane CJ, Rubinsztein DC. Complex inhibitory effects of nitric oxide on autophagy. *Molecular cell*. 2011; 43:19–32. [PubMed: 21726807]
41. Henson SM, Lanna A, Riddell NE, Franzese O, Macaulay R, Griffiths SJ, Puleston DJ, Watson AS, Simon AK, Tooze SA, Akbar AN. p38 signaling inhibits mTORC1-independent autophagy in senescent human CD8(+) T cells. *The Journal of clinical investigation*. 2014; 124:4004–4016. [PubMed: 25083993]
42. Sarkar S, Davies JE, Huang Z, Tunnacliffe A, Rubinsztein DC. Trehalose, a novel mTOR-independent autophagy enhancer, accelerates the clearance of mutant huntingtin and alpha-synuclein. *The Journal of biological chemistry*. 2007; 282:5641–5652. [PubMed: 17182613]
43. Xu C, Li X, Wang F, Weng H, Yang P. Trehalose prevents neural tube defects by correcting maternal diabetes-suppressed autophagy and neurogenesis. *American journal of physiology Endocrinology and metabolism*. 2013; 305:E667–E678. [PubMed: 23880312]
44. Kara NZ, Toker L, Agam G, Anderson GW, Belmaker RH, Einat H. Trehalose induced antidepressant-like effects and autophagy enhancement in mice. *Psychopharmacology*. 2013; 229:367–375. [PubMed: 23644913]

45. Castillo K, Nassif M, Valenzuela V, Rojas F, Matus S, Mercado G, Court FA, van Zundert B, Hetz C. Trehalose delays the progression of amyotrophic lateral sclerosis by enhancing autophagy in motoneurons. *Autophagy*. 2013; 9:1308–1320. [PubMed: 23851366]
46. Davies JE, Sarkar S, Rubinstein DC. Trehalose reduces aggregate formation and delays pathology in a transgenic mouse model of oculopharyngeal muscular dystrophy. *Human molecular genetics*. 2006; 15:23–31. [PubMed: 16311254]
47. Hintz KK, Aberle NS, Ren J. Insulin resistance induces hyperleptinemia, cardiac contractile dysfunction but not cardiac leptin resistance in ventricular myocytes. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity*. 2003; 27:1196–1203.
48. Ren J, Privratsky JR, Yang X, Dong F, Carlson EC. Metallothionein alleviates glutathione depletion-induced oxidative cardiomyopathy in murine hearts. *Critical care medicine*. 2008; 36:2106–2116. [PubMed: 18552690]
49. Li Q, Yang X, Sreejayan N, Ren J. Insulin-like growth factor I deficiency prolongs survival and antagonizes paraquat-induced cardiomyocyte dysfunction: role of oxidative stress. *Rejuvenation research*. 2007; 10:501–512. [PubMed: 17979500]
50. Hazzalin CA, Le Panse R, Cano E, Mahadevan LC. Anisomycin selectively desensitizes signalling components involved in stress kinase activation and fos and jun induction. *Molecular and cellular biology*. 1998; 18:1844–1854. [PubMed: 9528756]
51. Chien PT, Lin CC, Hsiao LD, Yang CM. Induction of HO-1 by carbon monoxide releasing molecule-2 attenuates thrombin-induced COX-2 expression and hypertrophy in primary human cardiomyocytes. *Toxicology and applied pharmacology*. 2015; 289:349–359. [PubMed: 26385185]
52. Xu X, Hua Y, Nair S, Zhang Y, Ren J. Akt2 knockout preserves cardiac function in high-fat diet-induced obesity by rescuing cardiac autophagosome maturation. *Journal of molecular cell biology*. 2013; 5:61–63. [PubMed: 23258696]
53. Li SY, Gomelsky M, Duan J, Zhang Z, Gomelsky L, Zhang X, Epstein PN, Ren J. Overexpression of aldehyde dehydrogenase-2 (ALDH2) transgene prevents acetaldehyde-induced cell injury in human umbilical vein endothelial cells: role of ERK and p38 mitogen-activated protein kinase. *The Journal of biological chemistry*. 2004; 279:11244–11252. [PubMed: 14722101]
54. Zhang Y, Babcock SA, Hu N, Maris JR, Wang H, Ren J. Mitochondrial aldehyde dehydrogenase (ALDH2) protects against streptozotocin-induced diabetic cardiomyopathy: role of GSK3beta and mitochondrial function. *BMC medicine*. 2012; 10:40. [PubMed: 22524197]
55. Xu X, Bucala R, Ren J. Macrophage migration inhibitory factor deficiency augments doxorubicin-induced cardiomyopathy. *Journal of the American Heart Association*. 2013; 2:e000439. [PubMed: 24334905]
56. Roeand J, Ren ND. Oxidative activation of Ca(2+)/calmodulin-activated kinase II mediates ER stress-induced cardiac dysfunction and apoptosis. *American journal of physiology Heart and circulatory physiology*. 2013; 304:H828–H839. [PubMed: 23316062]
57. Li SY, Golden KL, Jiang Y, Wang GJ, Privratsky JR, Zhang X, Eason AR, Culver B, Ren J. Inhibition of sarco(endo)plasmic reticulum Ca2+-ATPase differentially regulates contractile function in cardiac myocytes from normotensive and spontaneously hypertensive rats: role of Ca2+ regulatory proteins. *Cell biochemistry and biophysics*. 2005; 42:1–12. [PubMed: 15673924]
58. Yu Q, Gao F, Ma XL. Insulin says NO to cardiovascular disease. *Cardiovascular research*. 2011; 89:516–524. [PubMed: 21051417]
59. Xuand J, Ren X. Macrophage migration inhibitory factor (MIF) knockout preserves cardiac homeostasis through alleviating Akt-mediated myocardial autophagy suppression in high-fat diet-induced obesity. *International journal of obesity*. 2015; 39:387–396. [PubMed: 25248618]
60. Velez M, Kohli S, Sabbah HN. Animal models of insulin resistance and heart failure. *Heart failure reviews*. 2014; 19:1–13. [PubMed: 23456447]
61. Abel ED, O'Shea KM, Ramasamy R. Insulin resistance: metabolic mechanisms and consequences in the heart. *Arteriosclerosis, thrombosis, and vascular biology*. 2012; 32:2068–2076.
62. Keku TO, Lund PK, Galanko J, Simmons JG, Woosley JT, Sandler RS. Insulin resistance, apoptosis, and colorectal adenoma risk. *Cancer epidemiology, biomarkers & prevention : a*

- publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology. 2005; 14:2076–2081.
63. Yuan ZQ, Feldman RI, Sussman GE, Coppola D, Nicosia SV, Cheng JQ. AKT2 inhibition of cisplatin-induced JNK/p38 and Bax activation by phosphorylation of ASK1: implication of AKT2 in chemoresistance. *The Journal of biological chemistry*. 2003; 278:23432–23440. [PubMed: 12697749]
 64. Fujio Y, Mitsuuchi Y, Testa JR, Walsh K. Activation of Akt2 Inhibits anoikis and apoptosis induced by myogenic differentiation. *Cell death and differentiation*. 2001; 8:1207–1212. [PubMed: 11753568]
 65. Jiang K, Coppola D, Crespo NC, Nicosia SV, Hamilton AD, Sebt SM, Cheng JQ. The phosphoinositide 3-OH kinase/AKT2 pathway as a critical target for farnesyltransferase inhibitor-induced apoptosis. *Molecular and cellular biology*. 2000; 20:139–148. [PubMed: 10594016]
 66. Cui Y, Lin J, Zuo J, Zhang L, Dong Y, Hu G, Luo C, Chen J, Lu Y. AKT2-knockdown suppressed viability with enhanced apoptosis, and attenuated chemoresistance to temozolomide of human glioblastoma cells in vitro and in vivo. *OncoTargets and therapy*. 2015; 8:1681–1690. [PubMed: 26185456]
 67. Komatsu M, Kurokawa H, Waguri S, Taguchi K, Kobayashi A, Ichimura Y, Sou YS, Ueno I, Sakamoto A, Tong KI, Kim M, Nishito Y, Iemura S, Natsume T, Ueno T, Kominami E, Motohashi H, Tanaka K, Yamamoto M. The selective autophagy substrate p62 activates the stress responsive transcription factor Nrf2 through inactivation of Keap1. *Nature cell biology*. 2010; 12:213–223. [PubMed: 20173742]
 68. Roe ND, Xu X, Kandadi MR, Hu N, Pang J, Weiser-Evans MC, Ren J. Targeted deletion of PTEN in cardiomyocytes renders cardiac contractile dysfunction through interruption of Pink1-AMPK signaling and autophagy. *Biochimica et biophysica acta*. 2015; 1852:290–298. [PubMed: 25229693]
 69. Xu X, Roe ND, Weiser-Evans MC, Ren J. Inhibition of mammalian target of rapamycin with rapamycin reverses hypertrophic cardiomyopathy in mice with cardiomyocyte-specific knockout of PTEN. *Hypertension*. 2014; 63:729–739. [PubMed: 24446058]
 70. Hua Y, Zhang Y, Ceylan-Isik AF, Wold LE, Nunn JM, Ren J. Chronic Akt activation accentuates aging-induced cardiac hypertrophy and myocardial contractile dysfunction: role of autophagy. *Basic research in cardiology*. 2011; 106:1173–1191. [PubMed: 21901288]
 71. Taneike M, Yamaguchi O, Nakai A, Hikoso S, Takeda T, Mizote I, Oka T, Tamai T, Oyabu J, Murakawa T, Nishida K, Shimizu T, Hori M, Komuro I, Takuji Shirasawa TS, Mizushima N, Otsu K. Inhibition of autophagy in the heart induces age-related cardiomyopathy. *Autophagy*. 2010; 6:600–606. [PubMed: 20431347]
 72. Liand JA, Hill DL. Cardiomyocyte autophagy and cancer chemotherapy. *Journal of molecular and cellular cardiology*. 2014; 71:54–61. [PubMed: 24239608]
 73. Wu Q, Jiang D, Huang C, van Dyk LF, Li L, Chu HW. Trehalose-mediated autophagy impairs the anti-viral function of human primary airway epithelial cells. *PloS one*. 2015; 10:e0124524. [PubMed: 25879848]
 74. Zhang X, Chen S, Song L, Tang Y, Shen Y, Jia L, Le W. MTOR-independent, autophagic enhancer trehalose prolongs motor neuron survival and ameliorates the autophagic flux defect in a mouse model of amyotrophic lateral sclerosis. *Autophagy*. 2014; 10:588–602. [PubMed: 24441414]
 75. Chenand GG, Haddad Q. Role of trehalose phosphate synthase and trehalose during hypoxia: from flies to mammals. *The Journal of experimental biology*. 2004; 207:3125–3129. [PubMed: 15299033]
 76. Fernandez-Estevez MA, Casarejos MJ, Lopez Sendon J, Garcia Caldentey J, Ruiz C, Gomez A, Perucho J, de Yebenes JG, Mena MA. Trehalose reverses cell malfunction in fibroblasts from normal and Huntington's disease patients caused by proteasome inhibition. *PloS one*. 2014; 9:e90202. [PubMed: 24587280]
 77. Schaefferand M, Goedert V. Stimulation of autophagy is neuroprotective in a mouse model of human tauopathy. *Autophagy*. 2012; 8:1686–1687. [PubMed: 22874558]
 78. Vidal RL, Figueroa A, Court FA, Thielen P, Molina C, Wirth C, Caballero B, Kiffin R, Segura-Aguilar J, Cuervo AM, Glimcher LH, Hetz C. Targeting the UPR transcription factor XBP1

protects against Huntington's disease through the regulation of FoxO1 and autophagy. Human molecular genetics. 2012; 21:2245–2262. [PubMed: 22337954]

79. Ni YG, Wang N, Cao DJ, Sachan N, Morris DJ, Gerard RD, Kuro OM, Rothermel BA, Hill JA. FoxO transcription factors activate Akt and attenuate insulin signaling in heart by inhibiting protein phosphatases. Proceedings of the National Academy of Sciences of the United States of America. 2007; 104:20517–20522. [PubMed: 18077353]
80. Barthel A, Schmoll D, Unterman TG. FoxO proteins in insulin action and metabolism. Trends in endocrinology and metabolism: TEM. 2005; 16:183–189. [PubMed: 15860415]
81. Gonzalez E, Flier E, Molle D, Accili D, McGraw TE. Hyperinsulinemia leads to uncoupled insulin regulation of the GLUT4 glucose transporter and the FoxO1 transcription factor. Proceedings of the National Academy of Sciences of the United States of America. 2011; 108:10162–10167. [PubMed: 21646544]

Fig. 1.

Assessment of cardiac hypertrophy and glucose tolerance in wild-type (WT) and Akt2^{-/-} mice with or without trehalose treatment [TLS, 1 mg/g/day, i.p. for two days followed by 2% (w/v) in drinking water for two months]. A: Representative images using Wheat Germ Agglutinin staining of left ventricular tissues; B: Heart weight-to-body weight ratio; C: Quantification data of cardiomyocyte cross-sectional area; D: ANP mRNA expression; E: BNP mRNA expression; F: α-MHC mRNA expression; G: β-MHC mRNA expression. H: Intraperitoneal glucose tolerance test (IPGTT, 2 g/kg) in WT and Akt2^{-/-} mice fasted for 12

hrs prior to the test; and I: Area underneath the IPGTT curve plotted in panel H. Mean $\pm$ SEM, n = 6–10 mice per group (or 400 randomly selected cells for panel C); *p < 0.05 vs. WT group, #p < 0.05 vs. Akt2^{-/-} group.

Fig. 2. Effect of Akt2 knockout and trehalose treatment on myocardial geometry and contractile function. WT and Akt2^{-/-} mice were treated with or without trehalose [TLS, 1 mg/g/day, i.p. for two days followed by 2% (w/v) in drinking water for two months] prior to the assessment of echocardiographic properties. A: Fractional shortening; B: Left ventricular (LV) wall thickness; C: LV end-systolic diameter (LVESD); D: LV end-diastolic diameter (LVESD); E: LV mass normalized to body weight (BWT); and F: Heart rate. Mean±SEM, n=8–11 mice per group; *p < 0.05 vs. WT group, #p < 0.05 vs. Akt2^{-/-} group.

Fig. 3.

Cardiomyocyte contractile properties of WT and Akt2^{-/-} mice treated with or without trehalose [TLS, 1 mg/g/day, i.p. for two days followed by 2% (w/v) in drinking water for two months]. A: Resting cell length; B: Peak shortening (normalized to cell length); C: Maximal velocity of shortening (+ dL/dt); D: Maximal velocity of relengthening (– dL/dt); E: Time-to-peak shortening (TPS); and F: Time-to-90% relengthening (TR₉₀). Mean ± SEM, n = 158–160 cells per group; *p < 0.05 vs. WT group, #p < 0.05 vs. Akt2^{-/-} group.

Fig. 4.

Intracellular Ca^{2+} transient properties in cardiomyocytes from WT and $\text{Akt2}^{-/-}$ mice treated with/without trehalose [TLS, 1 mg/g/day, i.p. for two days followed by 2% (w/v) in drinking water for two months]. A: Resting fura-2 fluorescence intensity (FFI); B: Peak FFI; C: Electrically-stimulated rise in FFI (ΔFFI); and D: Intracellular Ca^{2+} decay rate (single exponential). Mean $\pm$ SEM, $n=121$ cells per group; * $p < 0.05$ vs. WT group, # $p < 0.05$ vs. $\text{Akt2}^{-/-}$ group.

Fig. 5. Effect of Akt2 knockout and trehalose (TLS) treatment on cardiomyocyte apoptosis assessed using TUNEL staining and caspase 3 activity. A: Frozen myocardial sections were stained with desmin (red), TUNEL (green) and nucleus DAPI (blue); B: Quantification data of TUNEL staining; and C: Caspase 3 activity. Mean $\pm$ SEM, $n = 4-6$ mice per group; * $p < 0.05$ vs. WT group, # $p < 0.05$ vs. Akt2^{-/-} group.

Fig. 6.

Western blot of autophagic markers in WT and Akt2^{-/-} mice treated with or without trehalose [TLS, 1 mg/g/day, i.p. for two days followed by 2% (w/v) in drinking water for two months]. A: Representative gel blots depicting Atg5, Beclin1, LC3B-I, LC3B-II, p62 and GAPDH (loading control) using specific antibodies; B: Atg5 protein level; C: Beclin1 protein level; D: LC3B I protein level; E: LC3B II protein level; F: LC3B II-to-LC3B I ratio; and G: p62 protein level. Mean $\pm$ SEM, n = 6–10 mice per group, *p < 0.05 vs. WT group, #p < 0.05 vs. Akt2^{-/-} group.

Fig. 7.

Western blot of autophagic protein markers in isolated cardiomyocytes from WT and Akt2^{-/-} mice treated with or without the lysosomal inhibitor bafilomycin A1 (100 nM) in the presence or absence of trehalose (TLS, 100 mM) for 2 hrs. A: Representative gel blots depicting LC3B I, LC3B II, p62 and GAPDH (loading control) using specific antibodies; B: LC3B I protein level; C: LC3B II protein level; D: LC3B II-to-LC3B I ratio; and E: p62 protein level. Mean±SEM, n=4–5 mice per group, *p < 0.05 vs. WT group, #p < 0.05 vs. Akt2^{-/-} group, &p < 0.05 vs. corresponding TLS-treated group.

Fig. 8.

Western blot of apoptotic protein markers in WT and Akt2^{-/-} mice treated with or without trehalose [TLS, 1 mg/g/day, i.p. for two days followed by 2% (w/v) in drinking water for two months]. A: Representative gel blots depicting BAD, BAX, Bcl2, cleaved-Caspase 3, cleaved-Caspase 9 and GAPDH (loading control) using specific antibodies; B: BAD protein level; C: BAX protein level; D: Bcl2 protein level; E: Cleaved-Caspase 3 protein level; and F: Cleaved-Caspase 9 protein level. Mean $\pm$ SEM, n = 6–10 mice per group, *p < 0.05 vs. WT group, #p < 0.05 vs. Akt2^{-/-} group.

Fig. 9.

Western blot of the ER stress markers GRP78, IRE1α, JNK and GADD153 in WT and Akt2^{-/-} mice treated with or without trehalose [TLS, 1 mg/g/day, i.p. for two days followed by 2% (w/v) in drinking water for two months]. A: Representative gel blots depicting GRP78, pan and phosphorylated IRE1α and JNK, GADD153 and GAPDH (loading control) using specific antibodies; B: GRP78 protein level; C: pan IRE 1α protein level; D: phospho-IRE1α (pIRE1α Ser⁷²⁴) protein level; E: pIRE1α-to-IRE1α ratio; F: pan JNK protein level; G: phospho-JNK (pJNK Try¹⁸⁵) protein level; H: pJNK-to-JNK ratio; and I: GADD153

protein level. Mean $\pm$ SEM, n= 6–10 mice per group, *p < 0.05 vs. WT group, #p < 0.05 vs. Akt2^{-/-} group.

Fig. 10.

Western blot of Akt, Foxo1 and p38 MAPK in WT and Akt2^{-/-} mice treated with or without trehalose [TLS, 1 mg/g/day, i.p. for two days followed by 2% (w/v) in drinking water for two months]. A: Representative gel blots depicting pan and phosphorylated Akt, Akt2, Foxo1, p38 MAPK and GAPDH (loading control) using specific antibodies; B: pan Akt expression; C: phospho-Akt (pAkt Ser⁴⁷³) expression; D: pAkt-to-Akt ratio; E: pan Foxo1 expression; F: phospho-Foxo1 (pFoxo1) expression; G: pFoxo1-to-Foxo1 ratio; H: pan p38 MAPK expression; I: phospho-p38 MAPK (pp38 MAPK) expression; and J: pp38 MAPK-

to-p38 MAPK ratio. Mean $\pm$ SEM, n = 6–10 mice per group, *p < 0.05 vs. WT group, #p < 0.05 vs. Akt2^{-/-} group.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 11.

Effect of anisomycin and AS1842856 on trehalose (TLS)-induced cardiomyocyte contractile responses. Freshly isolated cardiomyocytes from WT and Akt2^{-/-} mice were incubated with trehalose (100 mM) for 2 hrs in the presence or absence of pretreatment (1 hr) with the p38 MAPK activator anisomycin (10 μg/ml) or the Foxo1 inhibitor AS1842856 (10 μM). A: Resting cell length; B: peak shortening; C: + dL/dt; D: - dL/dt; E: TPS; and F: TR₉₀. Mean ± SEM, n = 62–68 cells from two mice per group; *p < 0.05 vs. WT group, #p < 0.05 vs.

Akt2^{-/-} group, +p < 0.05 vs. Akt2^{-/-}-TLS group, &p < 0.05 vs. corresponding WT-anisomycin or WT-AS1842856 group.

Table 1

Primer sequence information for individual genes.

Primers	Forward sequence	Reverse Sequence
ANP	5'-ACAGCCAAGGAGGAAAAGC-3'	5'-CCACAGTGGCAATGTGACCA-3'
BNP	5'-TCCAGAGCAATTCAAGATGCA-3'	5'-CTTTTGTGAGGCCTTGGTCC-3'
α -MHC	5'-CCACCTGGGCAAGTCTAACAA-3'	5'-TGTAGTCCACGGTGCCAGC-3'
β -MHC	5'-GATGTTTTTGTGCCCCGATGA-3'	5'-ACCGTCTTGCCATTCTCCG-3'