Published in final edited form as:

Womens Health Issues. 2016 ; 26(5): 529–536. doi:10.1016/j.whi.2016.07.002.

DISCLOSURE OF DEPRESSION IN PRIMARY CARE: A QUALITATIVE STUDY OF WOMEN'S PERCEPTIONS

Abiola O. Keller, PA-C, MPH, PhD^a, Carmen R. Valdez, PhD^b, Rebecca J. Schwei, MPH^c, and Elizabeth A. Jacobs, MD, MPP^d

Carmen R. Valdez: cvaldez@wisc.edu; Rebecca J. Schwei: rschwei@medicine.wisc.edu; Elizabeth A. Jacobs: eajacobs@medicine.wisc.edu

^aAssistant Professor, Marquette University, Department of Physician Assistant Studies, 1700 W Wells St., Milwaukee, WI 53233, Phone: 414-288-8569, Fax: 414-288-7951

^bAssociate Professor, University of Wisconsin-Madison, Department of Counseling Psychology, 301 Education Building, 1000 Bascom Mall, Madison, WI 53706, (608) 263-4493

^cAssistant Researcher, Departments of Medicine, University of Wisconsin School of Medicine and Public Health, 800 University Bay Drive, Suite 210, Madison, WI 53705, Phone: (608) 262-8312, Fax: 888-263-2864

^dAssociate Vice Chair for Health Services Research, Departments of Medicine and Population Health Sciences, University of Wisconsin School of Medicine and Public Health, 800 University Bay Drive, Suite 210, Madison, WI 53705, Phone: 608-262-8302, Fax: 888-263-2864

Abstract

Background—Healthcare providers are better able to diagnose depression and initiate treatment when patients disclose symptoms. However, many women are reluctant to disclose depressive symptoms. Little is known about the experience of disclosing depression symptoms in primary care among racially and ethnically diverse women across the life course. We qualitatively explore experiences of disclosure of depressive symptoms to primary care providers among self-identified African American, Hispanic and non-Hispanic white women.

Methods—Twenty-four women with depression were recruited for open-ended interviews. We recorded, transcribed, and coded interviews using inductive content analysis.

Findings—Two distinct domains emerged: participant factors that hinder and facilitate disclosure and provider cues that encourage and dissuade discussing depression. Participants described perceptions about primary care not being the appropriate place, fear of not having a choice in

Correspondence concerning this article should be addressed to Abiola O. Keller, Department of Physician Assistant Studies, Marquette University, 1700 W. Wells St., Milwaukee, WI 53233, abiola.keller@marquette.edu, Phone: 414-288-8569.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Declaration of Conflicting Interests

The authors declare that there is no conflict of interest.

None of the authors have a conflict of interest with this research. The authors would like to thank all the women that participated in this study.

treatment decisions, and the emotional cost of retelling as impediments to disclosure; perceiving an increased likelihood of getting help was described as a facilitator. Women identified provider behaviors of asking about depression and showing concern as facilitators, and provider time constraints as a barrier to disclosure.

Conclusions—Women perceive that primary care is not the appropriate place to disclose depression symptoms. Increased public education about behavioral health management in primary care, as well as more robust integration of the two, is needed. Efforts to improve depression disclosure in primary care must also encompass systematic use of depression screening tools and implementation of targeted interventions to cultivate provider empathy.

Keywords

depression; disclosure; patient-provider interaction; primary care; disparities

Introduction

Depression disproportionately affects women (National Institute for Health Care Management (NIHCM) Foundation, 2010) and adversely affects their health and wellbeing (Pan, Sun, Okereke, Rexrode, & Hu, 2011; Song, Lee, Baek, & Miller, 2012; Zhang, McKeown, Hussey, Thompson, & Woods, 2005). Despite existing effective treatments, many do not receive adequate depression treatment, (Young, Klap, Sherbourne, & Wells, 2001) and disparities in the receipt of adequate treatment persist (González et al., 2010; Witt et al., 2011). Women may not receive treatment because they are not screened by healthcare providers or because they do not disclose symptoms (Cepoiu et al., 2008). Missed depression diagnoses are an important problem given that the cost of treating depression is far less than societal costs associated with lost productivity or suicide (Goldman, Nielsen, & Champion, 1999).

Research shows that healthcare providers are more likely to initiate treatment when patients disclose depressive symptoms or directly request help (Kravitz et al., 2005; Tylee, Freeling, Kerry, & Burns, 1995). However, disclosure of depression occurs infrequently during medical encounters (Robinson & Roter, 1999; Sleath & Rubin, 2002). While studies examining depression symptom disclosure among women have identified barriers including stigma, lack of time, and lack of trust in providers (Dennis & Chung-Lee, 2006; Goodman, 2009; Jesse, Dolbier, & Blanchard, 2008), these studies have focused on postpartum depression. Little is known about women's experiences when disclosing depressive symptoms outside of the perinatal period. Although a large number of adults seek depression care in primary care (Marcus & Olfson, 2010; Substance Abuse and Mental Health Services Administration, 2009), few studies have focused on depressive symptom disclosure in this setting. Those focused on this disclosure found that 43% of adults reported at least 1 reason for nondisclosure of depression symptoms (Bell et al., 2011) and that perceptions of primary care providers' mental health-related capabilities and interests (Kravitz et al., 2011) are barriers to disclosure among predominantly white samples.

Although these studies highlight important barriers to disclosure in the primary care setting, little is known about the experience of disclosing depression symptoms in this setting among

racially and ethnically diverse women across the life course. To address these gaps, this study employs qualitative methods to explore the experience of depression disclosure in primary care settings among self-identified African American, Hispanic and non-Hispanic white women.

Methods

Study Participants

We conducted semi-structured interviews with a convenience sample of English-speaking African American, Hispanic, and non-Hispanic white women. Women were age 18 and older and reported a current or previous diagnosis of depression or met diagnostic criteria for current depression (score ≥ 10) based on the eight-item Patient Health Questionnaire depression scale (PHQ-8) (Kroenke et al., 2009). The PHQ-8 includes eight of the nine DSM-5 diagnostic criteria for the diagnosis of depression. It is different than the PHQ-9, in that it omits a question which assesses suicidal or self-injurious thoughts. We used the PHQ-8 because the study interviews were conducted in public non-clinical settings thereby limiting our ability to provide adequate follow-up if the women expressed suicidal or other harmful ideation. The PHQ-8 is a valid measure of depression (Kroenke et al., 2009) and research has shown that identical scoring thresholds could be used for both the eight and nine-item scales (Kroenke & Spitzer, 2002). To focus on the experience of disclosure independent of access to care, we excluded women who had not seen a healthcare provider in the previous year.

The Institutional Review Board at the University of Wisconsin-Madison approved the study. Over a 6-month period, we recruited participants from a Midwest city in the U.S. through online postings on Craigslist, flyers in local clinics, and letters sent to African American and Hispanic patients from a local women's internal medicine clinic. Our goal was to recruit ten women within each group. Participants each received a \$50 stipend.

Data Collection

The principal investigator conducted all interviews using a semi-structured interview guide (Appendix). Interviews lasted an average of 56 minutes and were recorded and professionally transcribed. The coding team (AOK & RJS) reviewed transcripts for accuracy.

Data Analysis

We examined transcripts for patterns and key content-related categories using inductive content analysis. Inductive content analysis uses open and unstructured coding to allow for identification of previously unidentified or unexpected thematic categories (Elo & Kyngas, 2008). The coding team independently reviewed the first three transcripts line-by-line and generated initial codes. They then met to discuss similarities and differences in code definitions. After reaching consensus on the definition and meaning of each code, a revised code list was created by collapsing codes that were similar into broader higher order categories (Elo & Kyngas, 2008). The first three interviews were recoded using the agreed upon codebook. The coding team continued to modify codes and add codes and thematic categories throughout the analysis process. All transcripts were coded with the final code

list. Additionally, a co-investigator not involved in the initial phases of the analysis (EAJ) audited the final codes for consistency, clarity, and comprehensiveness.

As part of our analyses, we separated the transcripts by white, African American, and Hispanic patient groups and assessed similarities and differences in codes and thematic categories by race and ethnicity. This was based on previous research demonstrating that there are often meaningful differences in the substantive content of themes emerging from qualitative analyses about preferences and experiences of care across diverse racial and ethnic groups (Mulvaney-Day, Earl, Diaz-Linhart, & Alegria, 2011; Tucker et al., 2003). We used QSR International's NVivo 10 qualitative data analysis software ("NVivo qualitative data analysis software," 2012) for managing codes and themes.

Results and Discussion

Study Participants

Thirty-four women responded to recruitment efforts; 31 were eligible, 28 were scheduled for interviews, and 24 interviews were completed (10 white, 9 African American, and 5 Hispanic). Participants were between the ages of 18–58 (mean = 40), and their PHQ-8 scores ranged from 3–24 (mean 14.7; scores ≥ 10 identify current depression) (Table 1). Of the 24 women interviewed, 20 (83%) met the diagnostic criteria for current depression using the PHQ-8. The PHQ-8 scores for the four women who did not meet the criteria were 3, 9, 9, and 9; all four of these women reported a previous depression diagnosis.

When women spoke about disclosure, two distinct domains emerged: participant factors that hinder and facilitate disclosure and provider cues that encourage and dissuade discussing depression. We present the themes that emerged within each domain. Differences by race and ethnicity are discussed when present.

Domain 1: Participant Factors that Hinder and Facilitate Disclosure

Participants commonly brought up 4 factors that hindered disclosure of depressive symptoms: (1) the perception that primary care providers were not the right people to talk to about depression; (2) being afraid of not having a choice in treatment decisions; (3) the emotional cost of retelling their story; and (4) difficulty recognizing and accepting symptoms. They described one facilitator to disclosure: perceiving that they would likely get help.

Primary care providers are not the right people—Women emphasized the distinction between physical and mental health and talked about how primary care providers were not the appropriate people with whom to discuss their depressive symptoms. One white woman described this distinction by saying, "I feel like the [primary care doctor] should be working on stomach hurts, you know, stitches kinda thing, not really mental. I think they're more with the physical." Additionally, a white woman said: "You don't really want to necessarily talk about something like that to your [primary care doctor] because all your life you've associated them with just, Oh, I'm going in there for a broken arm, or I've got the flu."

Both of these quotes illustrate that although women made the assumption that primary care providers were only concerned with physical health, they were uncertain, as evidenced by their tentative language of “You don’t really want to necessarily talk about [mental health],” and “I think they’re more with the physical.” Rather than clarifying this assumption with providers, women chose not to engage in this type of clarification, suggesting that other factors may be involved in their intentions to disclose.

Being afraid of not having a choice in treatment decisions—Women described being afraid that they would not be included as active participants in decisions about treatment. A Hispanic woman spoke about how she did not trust that her provider would respect her choices if she fully disclosed her feelings.

If I tell them exactly that, the feelings that I’ve had lately where I’m feeling so overwhelmed, then they’ll wanna step in and either medicate me.... And I don’t wanna be at that point. I think what I want is the choice, the choice to go in and get the help without having it forced on me, which is why I don’t really tell her everything.

As this quote suggests, even when women disclose symptoms, they may not share the severity of their symptoms due to fear of not being included in decision-making processes.

Emotional cost of retelling—Even in cases where women perceived the initial disclosure of symptoms to be beneficial, having to change providers and go through the emotional process of retelling their story was a barrier. An African American woman explained,

...it was helpful. You know it made me feel a little bit better. But then going through the whole initial thing again and having to explain why you’re feeling this way—you know kinda feels like, mm, maybe I don’t want to go through this again because you don’t.

This suggests that women treat each opportunity to disclose symptoms as discrete and emotionally burdensome events. Even if they benefited from disclosing symptoms in the past, the need to repeatedly retell their story, often caused by discontinuity in care, can deter disclosure of recurrent or persistent symptoms.

Recognizing and Accepting—African American and white women, but not Hispanic women, spoke about having difficulties identifying the way they were feeling as symptoms of an illness. One white woman described the difficulty in this way, “[women] are not gonna recognize it a lot of times. I mean that’s the first thing is they’re not gonna know what’s happening. They’re not gonna know this isn’t normal.” An African American woman described an experience where she did not accept her depression diagnosis. “And I told [the doctor] how I was feeling, and he pinpointed it as being depressed. I didn’t see it as being that.” Women often described the increasing severity of symptoms as what ultimately led them to disclose symptoms to their primary care provider. One white woman described her reasons for disclosure by saying, “The only reason I finally told somebody about is, I mean I was just absolutely against the wall, could not do it anymore.”

Women appeared to be more willing to disclose when the severity of their symptoms exceeds their individual coping resources. However, if they did not believe in the biomedical model of the etiology of depression held by most clinicians, they may be less likely to believe that a clinician is the person to talk to and therefore not initiate or continue treatment.

Likelihood of getting help—Although women doubted the appropriateness of talking about depression with their primary care provider, women were more likely to do so if they perceived that disclosing symptoms would ensure receiving help. One African American woman said, “I would want them to offer the whatever it is—medication they offer... I would expect them to do that or recommend you to like a therapist or something.” A white woman described her expectations of her provider the following way:

It has always been hard for me to open up to people, and I think for that to happen with someone who’s kind of a complete stranger, I need to feel some kind of, I don’t know if connection is the right word, but to feel confident that telling them will actually do something.

This quote suggests that the perception of getting help may result from a sense of connection and trust with the provider, suggesting that provider cues are important in influencing patients’ beliefs about disclosure.

Domain 2: Provider Cues That Encouraged and Dissuaded Discussing Depression

Perceptions about the appropriateness of disclosing to primary care providers were likely reinforced by provider behaviors during disclosure attempts. Specifically, women identified the behaviors of asking about depression along with showing sincere concern as facilitators to disclosure and provider time constraints as a barrier to disclosure.

Asking Questions—One Hispanic woman described not telling her provider how she feels unless she is asked directly.

If you really wanna know what’s going on, I’ll tell you. But if I don’t get asked those questions then I kinda shut down because I- I’m like, ‘It’s fine,’ and then I don’t tell them exactly what’s really going.

Additionally, a white woman said, “I know if my medical doctor would have asked me something I would have—I would have told them something. Like it’s um—people are looking for that opportunity to get help.”

Both of these quotes illustrate the importance of providers initiating direct conversations about depression in women’s disclosure process. Along with asking about depression, women wanted providers to create a safe space where they felt that the information shared would be protected. Specifically, they described the need to keep that “very fragile piece of you in a protective stance” and that without assurances of such protection, they were less willing to disclose.

Sincere Concern—Although a critical component of women’s willingness to disclose, asking about depression in and of itself was not enough to encourage women to disclose

their feelings. Questions about depression needed to be combined with a demonstration of sincere concern. One Hispanic woman described a situation where she was asked questions about her mood, but the provider did nothing to address her response and therefore she determined the provider wasn't actually concerned about her well-being.

And I think they, they tried asking me, 'How are things at home?' And when I would say, 'Well, they really suck,' They'd be like, 'Oh, okay,' and then they would just move on. [*laughter*] So I'm like, 'Okay, you didn't really wanna know about that. Um, why did you ask me about that?'

Women perceived that their provider was sincerely concerned about them when providers listened to them *and* made them feel comfortable through a positive connection. One African American woman noted the importance of feeling comfortable in the following terms: "It has to be someone that's—makes you feel comfortable and have to go that little extra yard of asking you more questions about yourself and about them, and relating to you is real important too." A Hispanic woman described how her provider showed concern by listening: "This particular doctor, she's real open. So, like when you talk to her, you get the feeling that she's really concerned that she really wants to listen." Similarly, a white woman stated, "...you need somebody who listens to you and who understands what you're trying to say."

One mechanism that women used to evaluate their provider's level of sincerity was through the provider's listening behaviors. As evident by their desire for providers "to go that little extra yard of asking you more questions" and demonstrate that they "understand what you're trying to say," women valued providers who were actively engaged in both listening and probing so they could explore patient statements about symptoms in greater detail and reflect back what was said. When providers appeared to be genuinely concerned and engaged, women were more willing to share information.

Time—A main barrier to disclosure was lack of time with providers. Time was described as both length of visits and frequency of contact with providers. One African American woman described how she doesn't see her provider enough to feel comfortable talking about her depression symptoms.

I go to a doctor it's maybe, you know, it might be for the yearly, you know, pap smear or something like that, or breast exam, just little stuff or I might go to the doctor for if I get sick in the wintertime. You know, I feel like I don't see her enough to know—actually tell her what—how I'm feelin' or ask for help because I'm only there for, you know, maybe three or four times out of the year and I don't feel comfortable with talking to her or asking her could she help me with this.

Women also expressed concerns about visits not being long enough. One Hispanic woman said:

My experience has been that doctors don't really want to get into the nitty-gritty of it because of their time managed as it is... I used to work at a clinic, too, so I know how pressed for time you are.

Women with depression value having “face time” with primary care providers; therefore, lack of time with providers as a result of infrequent visits, short visit times and/or rushed providers prevents women from being able to fully present and discuss their feelings.

Our participants shared with us factors that hinder and facilitate disclosure of depression symptoms. Similar to previous work (Kravitz et al., 2011), this study found that women doubted the appropriateness of disclosing symptoms to primary care providers. Recent health policy efforts have focused on improving depression recognition in primary care; most notably, the US Preventive Services Task Force (USPSTF) now recommends screening for depressive disorders in the general adult population (U.S. Preventive Services Task Force). To increase compliance with USPSTF guidelines, investigators have focused on developing depression screening tools that are minimally burdensome to both patients and providers (Blozik, Scherer, Lacruz, Ladwig, & group, 2013; Corson, Gerrity, & Dobscha, 2004). Our study suggests that women are reluctant to disclose depressive symptoms to primary care providers, but that they are more open to discussing how they feel if providers ask directly about depression and show sincere concern for their responses. Empathy, “an understanding of the emotional states of others and expression of this understanding” (Kelm, Womer, Walter, & Feudtner, 2014), is not expressed in clinical encounters as often as patients would like (Levinson, Gorawara-Bhat, & Lamb, 2000). Efforts to improve depression disclosure in primary care will need to encompass increased use of depression screening tools and implementation of targeted interventions to cultivate provider empathy (Kelm et al., 2014).

We also found that women’s decision to disclose often depended on the severity of symptoms and the anticipated benefit of disclosure. Even when concerned about her mood, a woman may not disclose concerns until they become severe and a threat to her wellbeing. While the severity of symptoms and the desire to avoid consequences of depression might be powerful enough to encourage disclosure, we also heard that decisions about where and to whom they will disclose their distress are driven by perceptions of the type and amount of help they will receive. Yet, confidence in the provider’s ability to initiate a treatment plan was often tempered by the fear of not being included in the treatment decision-making process. Specifically, women shared their desire to have choices and not feel forced to accept a specific treatment. The desire to be an active participant in the treatment decision-making process is not unique to our study participants. Previous studies have found that the majority of individuals with mental illness desire greater participation in treatment decisions (Adams, Drake, & Wolford, 2007), particularly through shared decision-making interventions (Deegan, Rapp, Holter, & Riefer, 2008). Shared decision-making creates an environment that not only encourages symptom disclosure, but may also improve the likelihood of treatment adherence, as depression treatment plans that are consistent with patient preferences have been shown to have increased initiation and adherence rates (Raue, Schulberg, Heo, Klimstra, & Bruce, 2009).

Lack of time with providers was a barrier to symptom disclosure. One potential approach to overcoming this barrier is a team approach to providing patient care (Hopkins & Sinsky, 2014). Although team-based care does not necessarily increase the amount of time patients spend with providers, by including structured interactions with other care-team members

this model of care encourages establishing relational ties with team members aside from the provider. Additionally, purposefully involving other healthcare workers in the care delivery process increases the number of healthcare contacts patients experience which may result in more opportunities for disclosing symptoms. To ensure that individuals who disclose depressive symptoms are appropriately evaluated and treated, healthcare teams may benefit from evidence-based strategies for optimizing performance and communication, such as TeamSTEPPS (Agency for Healthcare Research and Quality, 2016).

While the two large thematic categories identified in the study were similar between the three diverse racial and ethnic groups, there were some differences for Hispanic women. These women did not experience difficulty with recognizing and accepting depressive symptoms as a barrier to disclosure; however, Hispanic women discussed interpersonal factors, such as showing concern and time with the provider, more frequently than their white and African American counterparts. These findings are consistent with the Hispanic cultural values of *simpatía* and *personalismo*, where emphasis is placed on politeness and pleasantness (Triandis, Lisansky, Marin, & Betancourt, 1984), and suggest that the need for a positive relationship and feeling connected with a primary care provider may be a particularly important facilitator for disclosure among Hispanic women. Future research using a larger sample of Hispanic participants is needed to know whether these differences hold.

Several potential limitations of this study should be noted. First, we excluded women who had not seen a provider in the year before enrollment in the study; therefore, our findings may not capture experiences of women with depression who did not previously seek care. This population should be a focus of future research given that women with depression who have not sought care in the previous year may be at greater risk for negative outcomes. Second, while the study found more similarities than differences among these groups of women, our ability to explore this was limited by the small number of Hispanic women in the sample. Moreover, we did not explicitly ask about how their race or ethnicity influenced their experience with disclosing depression symptoms. Race and ethnicity can be very sensitive topics, and participants may have been reluctant to raise them. Our study results show that women may not disclose depression in primary care unless the provider asks; perhaps the same dynamic may apply to issues around race and ethnicity in qualitative methods. It should also be noted that only 6 women reported being in poor health. Women in poor health may have more contact with their providers and therefore their experience of disclosing symptoms of depression might differ. Finally, non-English speaking women were not included in the study, which likely contributed to the difficulty in recruiting Hispanic women. Research has shown that limited English proficiency is a barrier to mental health service use (Sentell, Shumway, & Snowden, 2007) and receipt of adequate treatment (Keller, Gangnon, & Witt, 2013). Moreover, Spanish-speaking Hispanic and English-speaking Hispanic persons have been shown to differ in their beliefs about the etiology of depression and preferences for treatment (Fernandez, Franks, Jerant, Bell, & Kravitz, 2011). Future studies are needed to explore the process of disclosing mental health symptoms in primary care among this vulnerable population.

This study has important strengths. The sample included women from three racial and ethnic groups and allowed for the opportunity to compare content categories across groups. Although the history of depression diagnosis was self-reported, participants did complete the PHQ-8 at the time of enrollment, thereby providing valid information about their depression symptom status.

Implications for Practice and/or Policy

The perception that primary care is not the appropriate place to disclose depression symptoms persists. Increased public education about and more robust integration of behavioral health and primary care are likely necessary to foster women's willingness to disclose depression symptoms in primary care. Our study findings highlight how potentially modifiable aspects of the patient-provider relationship influenced women's willingness to disclose depression symptoms. Interventions that support providers asking about depression while augmenting their ability to demonstrate empathy and reflective listening may be key strategies for encouraging women to disclose depression symptoms in primary care settings.

Acknowledgments

Abiola Keller had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. We also thank Linda Park for her contributions to earlier versions of this manuscript and Sheri Johnson for her comments on drafts of this manuscript.

Funding Statement: This project was supported by the National Center for Advancing Translational Sciences (grant number UL1TR000427) and the National Institute on Minority Health and Health Disparities (grant number 5P60MD003428) of the National Institutes of Health (NIH), NIH/National Institute of Child Health and Human Development Kirschstein-National Research Service Award (grant number T32HD049302-06), and the NIH Loan Repayment Program for Health Disparities Research (grant number L60 MD008863). Additional support was provided by the Health Innovation Program. The funding agencies had no involvement in the study design, the collection, analysis and interpretation of data, the writing of the report, or in the decision to submit the article for publication. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Appendix Semi-Structured Interview Guide for Participant Interviews

Introduction

Thank you for agreeing to take time out of your busy schedule to speak with me today. As I mentioned over the phone, this project is about women's experiences with getting care for depression.

Before we begin the conversation we need to go over the consent form. This is a form that provides you with information about the conversation today. You will take a copy of it with you at the end of the day. We will take a couple of minutes for you to read it. Please let me know if you would like for me to read it.

[Complete the consent process]

I am going to ask you a few questions about your experience with depression or feelings of sadness and what makes it easier or harder for women to get the care that they need.

Ok, if you don't have any more questions, let's get started.

- Q1** What do women need from their health care providers when they are depressed?
- Q2** How well do health care providers meet those needs?
- Q3** What makes a good provider when it comes to treating depression?
- Q3a** Are there things about the provider that make it easier for women to talk about their mood? What are they?
- Q3b** Are there things that the provider can do to make it easier for women to talk about their mood? What are they?
- Q4** What kinds of things do you think health care providers should say to or ask a woman who is depressed?
- Q5** Some people think health care providers should involve them in making decisions about treating their depression. What do you think about that?
- Q6** Tell me about different types of health care providers or healers that you have encountered while seeking care for depression.

For each provider/healer mentioned:

- Q6a** When did you encounter this provider?
- Q6b** What prompted you to seek care?
- Q6c** How did you find out about this provider/healer?
- Q6d** How would you characterize the quality of your experience with this provider/healer?
- Q6e** How did this provider/healer help you?
- Q6f** What made their advice or suggestions helpful?

Biographies

Abiola O. Keller, MPH, PhD, is an Assistant Professor at the College of Health Sciences at Marquette University. Her research focuses on improving women's health through studying patient-provider interactions and associated outcomes.

Carmen R. Valdez, PhD, is an Associate Professor in the Department of Counseling Psychology at the University of Wisconsin-Madison. Her research examines the impact of family stress and sociocultural context on outcomes for Latino children.

Rebecca J Schwei, MPH, is an Assistant Researcher at the University of Wisconsin. Her research interests include health disparities, provision of linguistically appropriate health care, and the social determinants of health.

Elizabeth A. Jacobs MD, MPP, a Professor of Medicine and Population Health Sciences at the University of Wisconsin-Madison, research interests include access to and cultural

specificity of care, impact of interpreter service interventions on healthcare, and health literacy and numeracy.

References

- Adams JR, Drake RE, Wolford GL. Shared decision-making preferences of people with severe mental illness. *Psychiatr Serv.* 2007; 58(9):1219–1221. doi: 58/9/1219 [pii]10.1176/appi.ps.58.9.1219. [PubMed: 17766569]
- Agency for Healthcare Research and Quality. TeamSTEPPS®: Strategies and Tools to Enhance Performance and Patient Safety. 2016. Retrieved from <http://www.ahrq.gov/professionals/education/curriculum-tools/teamstepps/index.html>
- Bell RA, Franks P, Duberstein PR, Epstein RM, Feldman MD, Fernandez y Garcia E, Kravitz RL. Suffering in silence: reasons for not disclosing depression in primary care. *Ann Fam Med.* 2011; 9(5):439–446. DOI: 10.1370/afm.1277 [PubMed: 21911763]
- Blozik E, Scherer M, Lacruz ME, Ladwig KH, group, K. s. Diagnostic utility of a one-item question to screen for depressive disorders: results from the KORA F3 study. *BMC Fam Pract.* 2013; 14:198.doi: 10.1186/1471-2296-14-198 [PubMed: 24359193]
- Cepoiu M, McCusker J, Cole MG, Sewitch M, Belzile E, Ciampi A. Recognition of depression by non-psychiatric physicians—a systematic literature review and meta-analysis. *J Gen Intern Med.* 2008; 23(1):25–36. DOI: 10.1007/s11606-007-0428-5 [PubMed: 17968628]
- Corson K, Gerrity MS, Dobscha SK. Screening for depression and suicidality in a VA primary care setting: 2 items are better than 1 item. *Am J Manag Care.* 2004; 10(11 Pt 2):839–845. [PubMed: 15609737]
- Deegan PE, Rapp C, Holter M, Riefer M. Best practices: a program to support shared decision making in an outpatient psychiatric medication clinic. *Psychiatr Serv.* 2008; 59(6):603–605. [pii]. DOI: 10.1176/appi.ps.59.6.60359/6/603 [PubMed: 18511580]
- Dennis CL, Chung-Lee L. Postpartum depression help-seeking barriers and maternal treatment preferences: a qualitative systematic review. *Birth.* 2006; 33(4):323–331. DOI: 10.1111/j.1523-536X.2006.00130.x [PubMed: 17150072]
- Elo S, Kyngas H. The qualitative content analysis process. *J Adv Nurs.* 2008; 62(1):107–115. [pii]. DOI: 10.1111/j.1365-2648.2007.04569.xJAN4569 [PubMed: 18352969]
- Fernandez YGE, Franks P, Jerant A, Bell RA, Kravitz RL. Depression treatment preferences of Hispanic individuals: exploring the influence of ethnicity, language, and explanatory models. *J Am Board Fam Med.* 2011; 24(1):39–50. DOI: 10.3122/jabfm.2011.01.100118 [PubMed: 21209343]
- Goldman LS, Nielsen NH, Champion HC. Awareness, diagnosis, and treatment of depression. *J Gen Intern Med.* 1999; 14(9):569–580. [PubMed: 10491249]
- González HM, Vega WA, Williams DR, Tarraf W, West BT, Neighbors HW. Depression care in the United States: too little for too few. *Arch Gen Psychiatry.* 2010; 67(1):37–46. DOI: 10.1001/archgenpsychiatry.2009.168 [PubMed: 20048221]
- Goodman JH. Women's attitudes, preferences, and perceived barriers to treatment for perinatal depression. *Birth.* 2009; 36(1):60–69. DOI: 10.1111/j.1523-536X.2008.00296.x [PubMed: 19278385]
- Hopkins K, Sinsky CA. Team-based care: saving time and improving efficiency. *Fam Pract Manag.* 2014; 21(6):23–29. [PubMed: 25403048]
- Jesse DE, Dolbier CL, Blanchard A. Barriers to seeking help and treatment suggestions for prenatal depressive symptoms: focus groups with rural low-income women. *Issues Ment Health Nurs.* 2008; 29(1):3–19. DOI: 10.1080/01612840701748664 [PubMed: 18214775]
- Keller AO, Gangnon R, Witt WP. The Impact of Patient–Provider Communication and Language Spoken on Adequacy of Depression Treatment for U.S. Women. *Health Commun.* 2013; :1–10. DOI: 10.1080/10410236.2013.795885 [PubMed: 23330852]
- Kelm Z, Womer J, Walter JK, Feudtner C. Interventions to cultivate physician empathy: a systematic review. *BMC Med Educ.* 2014; 14:219.doi: 10.1186/1472-6920-14-219 [PubMed: 25315848]

- Kravitz RL, Epstein RM, Feldman MD, Franz CE, Azari R, Wilkes MS, Franks P. Influence of patients' requests for direct-to-consumer advertised antidepressants: a randomized controlled trial. *JAMA*. 2005; 293(16):1995–2002. doi: 293/16/1995 [pii]10.1001/jama.293.16.1995. [PubMed: 15855433]
- Kravitz RL, Paterniti DA, Epstein RM, Rochlen AB, Bell RA, Cipri C, Duberstein P. Relational barriers to depression help-seeking in primary care. *Patient Educ Couns*. 2011; 82(2):207–213. DOI: 10.1016/j.pec.2010.05.007 [PubMed: 20570462]
- Kroenke K, Spitzer RL. The PHQ-9: A new depression diagnostic and severity measure. *Psychiatric Annals*. 2002; 32(9):509–515.
- Kroenke K, Strine TW, Spitzer RL, Williams JB, Berry JT, Mokdad AH. The PHQ-8 as a measure of current depression in the general population. *J Affect Disord*. 2009; 114:1–3. 163–173. [pii]. DOI: 10.1016/j.jad.2008.06.026S0165-0327(08)00282-6 [PubMed: 18819715]
- Levinson W, Gorawara-Bhat R, Lamb J. A study of patient clues and physician responses in primary care and surgical settings. *JAMA*. 2000; 284(8):1021–1027. [PubMed: 10944650]
- Marcus SC, Olfson M. National trends in the treatment for depression from 1998 to 2007. *Arch Gen Psychiatry*. 2010; 67(12):1265–1273. [pii]. DOI: 10.1001/archgenpsychiatry.2010.15167/12/1265 [PubMed: 21135326]
- Mulvaney-Day NE, Earl TR, Diaz-Linhart Y, Alegria M. Preferences for Relational Style With Mental Health Clinicians: A Qualitative Comparison of African American, Latino and Non-Latino White Patients. *J Clin Psychol*. 2011; 67(1):31–44. DOI: 10.1002/Jclp.20739 [PubMed: 20939021]
- National Institute for Health Care Management (NIHCM) Foundation. Identifying and Treating Maternal Depression: Strategies & Considerations for Health Plans. Washington, DC: National Institute for Health Care Management (NIHCM) Foundation; 2010.
- NVivo qualitative data analysis software (Version 10). QSR International Pty Ltd; 2012.
- Pan A, Sun Q, Okereke OI, Rexrode KM, Hu FB. Depression and risk of stroke morbidity and mortality: a meta-analysis and systematic review. *JAMA*. 2011; 306(11):1241–1249. doi: 306/11/1241 [pii]10.1001/jama.2011.1282. [PubMed: 21934057]
- Raue PJ, Schulberg HC, Heo M, Klimstra S, Bruce ML. Patients' depression treatment preferences and initiation, adherence, and outcome: a randomized primary care study. *Psychiatr Serv*. 2009; 60(3): 337–343. [pii]. DOI: 10.1176/appi.ps.60.3.33760/3/337 [PubMed: 19252046]
- Robinson JW, Roter DL. Psychosocial problem disclosure by primary care patients. *Soc Sci Med*. 1999; 48(10):1353–1362. doi: S0277953698004390 [pii]. [PubMed: 10369436]
- Sentell T, Shumway M, Snowden L. Access to mental health treatment by English language proficiency and race/ethnicity. *J Gen Intern Med*. 2007; 22(Suppl 2):289–293. DOI: 10.1007/s11606-007-0345-7 [PubMed: 17957413]
- Sleath B, Rubin RH. Gender, ethnicity, and physician-patient communication about depression and anxiety in primary care. *Patient Educ Couns*. 2002; 48(3):243–252. doi: S0738399102001775 [pii]. [PubMed: 12477609]
- Song MR, Lee YS, Baek JD, Miller M. Physical activity status in adults with depression in the National Health and Nutrition Examination Survey, 2005–2006. *Public Health Nurs*. 2012; 29(3): 208–217. DOI: 10.1111/j.1525-1446.2011.00986.x [PubMed: 22512422]
- Substance Abuse and Mental Health Services Administration. The NSDUH Report: Major Depressive Episode and Treatment among Adults. Rockville, MD: 2009.
- Triandis HC, Lisansky J, Marin G, Betancourt H. Simpatia as a Cultural Script of Hispanics. *J Pers Soc Psychol*. 1984; 47(6):1363–1375. DOI: 10.1037//0022-3514.47.6.1363
- Tucker CM, Herman KC, Pedersen TR, Higley B, Montrichard M, Ivery P. Cultural sensitivity in physician-patient relationships: perspectives of an ethnically diverse sample of low-income primary care patients. *Med Care*. 2003; 41(7):859–870. DOI: 10.1097/01.MLR.0000068544.11426.4C [PubMed: 12835610]
- Tyler A, Freeling P, Kerry S, Burns T. How does the content of consultations affect the recognition by general practitioners of major depression in women? *Br J Gen Pract*. 1995; 45(400):575–578. [PubMed: 8554836]

- U.S. Preventive Services Task Force. Clinical Summary: Depression in Adults: Screening. Retrieved September 17, 2015, from <http://www.uspreventiveservicestaskforce.org/Page/Document/ClinicalSummaryFinal/depression-in-adults-screening>
- Witt WP, Keller A, Gottlieb C, Litzelman K, Hampton J, Maguire J, Hagen EW. Access to Adequate Outpatient Depression Care for Mothers in the USA: A Nationally Representative Population-Based Study. *J Behav Health Serv Res*. 2011; 38(2):191–204. DOI: 10.1007/s11414-009-9194-y [PubMed: 19838806]
- Young AS, Klap R, Sherbourne CD, Wells KB. The quality of care for depressive and anxiety disorders in the United States. *Arch Gen Psychiatry*. 2001; 58(1):55–61. [PubMed: 11146758]
- Zhang J, McKeown RE, Hussey JR, Thompson SJ, Woods JR. Gender differences in risk factors for attempted suicide among young adults: findings from the Third National Health and Nutrition Examination Survey. *Ann Epidemiol*. 2005; 15(2):167–174. DOI: 10.1016/j.annepidem.2004.07.095 [PubMed: 15652723]

Table 1

Demographic characteristics of participants overall and by race/ethnicity

	Overall (n=24)	African American (n=9)	Hispanic (n=5)	White (n=10)
Average Age, years (range)	40 (18,58)	42 (21,55)	43 (35,54)	38 (18,58)
Average PHQ-8 Score, (range)	15 (3,24)	16 (11,22)	14 (9,24)	14 (3,20)
Average # doctor's visits past year, (range)	5 (0,10+)	4 (0,10+)	5 (3,10+)	6 (1,10+)
Insurance Type (#)				
Private	11	2	3	6
Public	9	4	2	3
None	3	2	0	1
Education Level (#)				
< High School	3	3	0	0
High School/GED	11	5	3	3
Technical School	2	0	0	2
College Degree	8	1	2	5
Self Rated Health (#)				
Excellent	1	0	0	1
Very good/Good	17	7	4	6
Fair/Poor	6	2	1	3
Receiving Depression Treatment (#)				
Counseling	16	3	3	10
Medication	18	6	5	8