Discordant ability of the triglyceride to apolipoprotein B ratio to predict triglyceride-rich lipoprotein particle size in normal-weight and obese men

Miryam Amigo-Benavent¹, Laura Sinausia¹, Emilio Montero² and Javier S Perona¹

¹Instituto de la Grasa, Consejo Superior de Investigaciones Científicas (IG-CSIC), Campus Universidad Pablo de Olavide, Ctra. Utrera, km1. E-41013 Sevilla, Spain; ²HHUU Virgen del Rocío. Av. Manuel Siurot s/n. E-41013. Seville, Spain

Corresponding author: Javier S Perona. Email: perona@ig.csic.es

Abstract

The atherogenicity of triglyceride-rich lipoproteins (TRLs) is dependent of their particle size as it determines their metabolic fate. Since TRL possess a single apolipoprotein B (Apo B) molecule per particle, the triglyceride (TG)/Apo B ratio has been used as a convenient method to estimate TRL size. The aim of this study was to validate this approach by correlating the serum TG/Apo B ratio, and the TRL particle size measured by dynamic light scattering (DLS). Twenty-four male volunteers (12 normal-weight and 12 obese individuals) received a high-fat meal. Preprandial (0 h) and postprandial (2 and 4 h) serum samples were collected after meal ingestion, and TRLs were isolated. Serum TG and Apo B levels were quantified, and the TG/Apo B ratio was plotted against TRL particle size measured by DLS for correlation. A strong association between TRL particle size and serum TG/Apo B ratio for normal-weight subjects ($P \leq 0.001$) was observed but not for obese subjects ($P = 0.6116$). TG/Apo B ratio correlates with particle size in healthy normal-weight males but not in obese individuals. Whether this ratio is useful to estimate TRL size in females and in other dyslipidemic patients should be subject of future investigations.

Keywords: Apolipoprotein B, particle size, ratio, triglyceride, triglyceride-rich lipoprotein

Experimental Biology and Medicine 2016; 241: 1772–1775. DOI: 10.1177/1535370216639394

Introduction

It has become clear in the past years that postprandial lipemia, which is caused by raised levels of triglyceride-rich lipoproteins (TRLs) in the blood after a fatty meal, may be a risk factor for atherosclerosis.¹ TRL remnants can cross the endothelial barrier and enter into the vascular wall,² where they can interact with lipoprotein receptors on macrophages, enhancing lipid accumulation and leading to foam cell formation, one of the first steps of atherosclerosis.³ The magnitude and duration of the postprandial triglyceride (TG) response is influenced by a number of metabolic processes, including the rate of TG secretion from the intestine and the liver, the activity of enzymes involved in the TRL processing and the rate of clearance of TRL remnants by receptor-mediated pathways.¹

These processes are mediated, among others, by changes in particle size, which is an important determinant of TRL metabolic fate. We have previously reported that TRL particles with differences in size and composition can modulate TRL clearance postprandially, which may affect TG incorporation to the tissues.⁴ Moreover, Palmer et al.⁵

demonstrated that the highest TG uptake by macrophages takes place when cells were incubated with TRL obtained at the time point at which the particle size is intermediate (Sf 60–400), i.e. not as large as nascent particles (Sf > 400) and not as small as small – very low-density lipoprotein (VLDL) particles (Sf 20–60). Particle size is dependent upon TG hydrolysis by lipoprotein lipase (LPL), which is believed to be the predominant mechanism for TG uptake from TRL.⁶ TRL particles (chylomicrons and VLDL) share the LPL pathway, which is saturable. However, despite being present in a greater number in the postprandial state, VLDLs are much slowly cleared from plasma compared to chylomicrons, probably because large chylomicrons bind more avidly to LPL.⁷ LPL activity reduces TG but not apolipoprotein B (Apo B) content of TRL. This feature has been used to calculate the ratio between serum TG and Apo B concentrations as a rapid and convenient method to indirectly measure particle size.^{8,9} Small TRLs are cleared from the plasma at a slower rate than large TRL,¹⁰ increasing their opportunity to penetrate the arterial wall¹¹ and for the transfer of lipids from high-density lipoprotein (HDL) leading to a lowering of HDL cholesterol.¹² Therefore, the TRL particle

size can be of importance for processes involved in atherogenesis at different stages. However, it is presently unknown whether the serum TG/Apo B ratio correlates with other methods for particle size measurement and if it is useful for all kind of human subjects and physiological situations. Therefore, in the present study, we aimed to correlate the serum TG/Apo B ratio and the particle size measured by dynamic light scattering (DLS) in normal-weight and obese subjects at different postprandial times.

Materials and methods

Twenty-four healthy men, aged 31.8 ± 9.7 years participated in the study and were divided into two groups (normal weight and obese) according to their body mass index (BMI). Subjects were categorized as normal weight when BMI was lower than 25 kg/cm^2 and obese when BMI was higher than 30 kg/cm^2 . Sample size was calculated by power analysis, considering a type 1 error (α) = 0.05 and a power $(1-\beta)$ = 0.9 and resulting in a minimum number of 10 individuals per group. Subjects were excluded if they suffered from any digestive or metabolic disorder, were taking dietary supplements, or were under medication of any kind. Participants gave written, informed consent to a protocol approved by the Institutional Committee on Human Research (Hospital Universitario Virgen del Rocío, Seville, Spain). All procedures were in accordance with the Institutional and National ethical standards for human experimentation and the Helsinki declaration of 1975 (revised in 2000).

Participants were asked to have a low-fat dinner the evening prior to the postprandial study and to abstain from alcohol drinking and smoking for 24 h. On arrival, after an overnight fast (12 h), a cubital vein was catheterized and a baseline blood sample was taken immediately before consumption of the test meal that consisted of three slices of bread (71 g), 200 ml of skimmed-milk (fat content was 0.3 g/100 g), 55 g of butter, and 20 g of cocoa powder.

Blood samples were also collected 2 and 4 h after the intake. During the course of the experiment, subjects were allowed to drink water and to undertake only light activities. Serum was recovered by centrifugation ($1620 \times g$, 30 min, 4°C), and sodium azide, phenylmethylsulfonyl fluoride, and aprotinin (Sigma-Aldrich, Poole, UK) were added to a final concentration of 1 mmol/L, 10 $\mu\text{mol/L}$, and 0.5 mg/L, respectively.

Postprandial TRLs were isolated from 4.5 mL of serum that was layered under 8 mL of NaCl solution ($d = 1.006 \text{ kg/L}$), by a single ultracentrifugation spin (39,000 rpm, 16 h, 12°C), which ensures that all TRLs are collected. Ultracentrifugation was performed using a SW41Ti swinging bucket rotor in a Beckman L8-70 M preparative ultracentrifuge (Beckman Instruments, Palo Alto, USA). TRLs were rapidly frozen at -80°C and thawed just before isolation.

Serum Apo B concentrations were quantified by a turbidimetric immunoassay method (BioSystems, Spain, coefficient of variation [CV] ranged from 18.7% to 28.9%) following manual's instructions, and TG concentrations were measured by standard enzymatic methods (Roche

Diagnostics, Basilea, Switzerland, CV ranged from 10.6% to 21.0%). TRL size was determined by DLS using a Zetasizer Nano ZS instrument (Malvern Instruments, Malvern, UK; CV ranged from 5.0% to 7.4%). Samples were well homogenized and diluted with 0.9% (w/w) NaCl to an appropriate concentration (1:40 to 1:200 times). One milliliter of diluted TRL sample was loaded on a folded capillary cell and incubated at 37°C for 2 min before the reading. The conditions of the instrument were as follows: dispersant water, temperature 37°C , and refractive index 1.330. Results are expressed as the Z-average mean, which is the harmonic intensity of averaged particle diameter. Measurements were performed in triplicates. For the calculation of the TG/Apo B ratio, TG and Apo B values were expressed as mg/dL. Results were expressed as means $\pm$ standard error of the mean ($n = 12$). Data analyses were performed with the GraphPad Prism[®] 5 statistical package (GraphPad Software Inc., San Diego, CA). Correlations between variables were assessed using Pearson's correlation coefficients. Differences were considered statistically significant at $P < 0.05$.

Results and discussion

In the present study, we searched for correlation between serum TG/Apo B ratio and particle size in TRL from normal-weight and obese subjects in the postprandial state. Participants were males of a similar age, non-smokers, and moderate consumers of alcohol. Serum total cholesterol, LDL cholesterol, glucose, insulin, and TG concentrations as well as the homeostatic model assessment of insulin resistance (HOMA-IR) and systolic pressure were higher in obese individuals at baseline but no significant differences were observed for Apo B concentrations and TG/ApoB between the groups (Table 1).

Figure 1 shows the correlation of TRL particle size, measured as DLS and serum TG/Apo B ratio in all studied subjects and in the groups of normal-weight and obese subjects at the three time points (0, 2, and 4 h). Correlation was strongly significant in normal-weight subjects ($P < 0.001$) but not in obese individuals ($P = 0.6116$). As a consequence, in the whole group of participants significance was not achieved ($P = 0.0526$), and the correlation coefficient was very low ($r = 0.1745$). Al-Shayji et al.⁹ studied the effect of exercise on the TG/Apo B ratio of large VLDL in a group of overweight subjects, finding a non-significant increment that was associated with increased particle size. Our results show that the TG/Apo B ratio does not correlate in obese subjects, and these data might be different if Apo B was measured directly in VLDL or chylomicrons instead of serum. The absence of correlation in obese individuals might be related to the heterogeneity particle size in this group, showing coefficients of variation ranging from 17% at 0 h to 48% at 4 h, whereas in the normal-weight group, they ranged from 14% at 0 h to 22% at 4 h. Significant correlations were maintained when TG values were expressed as mmol/L, which is common in many laboratories. The ratio can be easily converted by applying the molecular weight (MW) of triolein (MW = 885).

TRL particle size and serum TG/Apo B ratio did not correlate in normal-weight subjects at 2 h after the intake of the experimental meal, when nascent, large chylomicrons are predominant. In contrast, correlation was significant in

Table 1 Anthropometric and fasting serum biochemical parameters of the participants in the study

	Normal weight	Obese
Age (y)	27.3 ± 10.0	35.8 ± 12.1
Weight (kg)	74.9 ± 8.0	112.9 ± 13.5***
Height (cm)	179.6 ± 6.9	180.8 ± 5.2
BMI (kg/m ²)	23.2 ± 1.0	34.7 ± 4.8***
Systolic pressure (mmHg)	110.9 ± 12.8	130.8 ± 7.6***
Diastolic pressure (mmHg)	68.9 ± 7.3	74.5 ± 10.0
Cholesterol (mg/dL)	172.5 ± 32.5	202.9 ± 34.6*
LDL-C (mg/dL)	94.7 ± 19.4	158.7 ± 16.6***
HDL-C (mg/dL)	57.5 ± 9.7	45.9 ± 8.7**
Glucose (mg/dL)	73.7 ± 4.8	83.6 ± 8.3**
Insulin (mU/mL)	4.9 ± 2.8	12.6 ± 4.2***
HOMA-IR	0.91 ± 0.51	2.41 ± 1.14***
Triglycerides (mg/dL)	78.7 ± 24.6	129.7 ± 60.2*
Apo B	96.4 ± 27.0	108.2 ± 43.4
TG/ApoB	0.9 ± 0.5	1.0 ± 0.4

Data are expressed as means ± SD.

BMI: body mass index; LDL-C: low-density lipoprotein cholesterol; HDL-C: high-density lipoprotein cholesterol; HOMA-IR: homeostatic model assessment of insulin resistance; Apo B: apolipoprotein B; TG/Apo B: triglyceride to apolipoprotein B ratio.

Statistical differences among all groups were assessed by Student's *t*-test.

P* < 0.05; *P* < 0.01; ****P* < 0.001 vs. normal-weight subjects.

fasting conditions and 4 h after the lipid challenge, time points at which the particle size is usually lower.⁸ Particles obtained from obese subjects were significantly larger at all time points studied. However, although TRL size was lower at 0 h in both groups of subjects, we did not find significant differences between 2 and 4 h, regardless the method used for its measurement (Supplementary Table).

DLS has been proposed as a reliable technique to measure TRL size and to study TG hydrolysis by lipases.^{13,14} The technique has been validated for LDL in comparison to electrophoresis and for larger particles,¹⁵ like TRL in comparison to gel chromatography,¹⁶ and thus it can be used to measure lipoprotein particle size.

The study has two limitations that need to be addressed: participants were all males, and the sample size is rather low. Therefore, future investigations should include females and larger samples to confirm these results. A possible limitation was the interference of TG in the Apo B assay. However, we consider this unlikely, since the CV (%) for the Apo B assay was similar in normal-weight and obese subjects and slightly higher in fasting conditions (0 h) than in the postprandial period (2 and 4 h).

Conclusion

In conclusion, our findings show that the TG/Apo B ratio correlates with particle size measured by DLS, and can be employed to estimate particle size, in healthy normal-weight males but not in obese individuals. Whether the ratio can be safely used in females and in subjects affected by dyslipidemia requires further investigation.

Figure 1 Correlation of serum triglyceride to apolipoprotein B ratio and particle size of triglyceride-rich lipoproteins from all, normal-weight and obese subjects in fasting conditions (0 h) and 2 and 4 h after a fat challenge (12 individuals per group), and composite of all time points. Correlations between variables were assessed using Pearson's correlation coefficients. Differences were considered statistically significant at *P* < 0.05

Author contributions: JSP designed the study. JSP and EM carried out the postprandial assay. MAB and LS performed the analyses. JSP and MAB drafted the manuscript. All authors approved the final article.

ACKNOWLEDGEMENTS

This work was supported by funds from Ministry of Economy and Competitiveness (Grant AGL2011-23810 and a FPI fellowship to Laura Sinausia). The funding body did not have any other role in the design, experimental work or writing of the manuscript.

DECLARATION OF CONFLICTING INTERESTS

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

REFERENCES

- Jackson KG, Poppitt SD, Minihane AM. Postprandial lipemia and cardiovascular disease risk: interrelationships between dietary, physiological and genetic determinants. *Atherosclerosis* 2012;**220**:22–33
- Mamo JC, Wheeler JR. Chylomicrons or their remnants penetrate rabbit thoracic aorta as efficiently as do smaller macromolecules, including low-density lipoprotein, high-density lipoprotein, and albumin. *Coron Artery Dis* 1994;**5**(8): 695–705
- Napolitano M, Avella M, Botham KM, Bravo E. Chylomicron remnant induction of lipid accumulation in J774 macrophages is associated with up-regulation of triacylglycerol synthesis which is not dependent on oxidation of the particles. *Biochim Biophys Acta* 2003;**1631**:255–64
- Cabello-Moruno R, Perona JS, Ruiz-Gutierrez V. Influence of minor components of olive oils on the composition and size of TRLs and on macrophage receptors involved in foam cell formation. *Biochem Soc Trans* 2008;**35**:470–1
- Palmer AM, Nova E, Anil E, Jackson K, Bateman P, Wolstencroft E, Williams CM, Yaqoob P. Differential uptake of subfractions of triglyceride-rich lipoproteins by THP-1 macrophages. *Atherosclerosis* 2005;**180**:233–44
- Milosavljevic D, Kontush A, Griglio S, Le Naour G, Thillet J, Chapman MJ. VLDL-induced triglyceride accumulation in human macrophages is mediated by modulation of LPL lipolytic activity in the absence of change in LPL mass. *Biochim Biophys Acta* 2003;**1631**(1): 51–60
- Xiang SQ, Cianflone K, Kalant D, Sniderman AD. Differential binding of triglyceride-rich lipoproteins to lipoprotein lipase. *J Lipid Res* 1999;**40**(9): 1655–63
- Cabello-Moruno R, Sinausia L, Botham KM, Montero E, Avella M, Perona JS. Postprandial phase time influences the uptake of TAG from postprandial TAG-rich lipoproteins by THP-1 macrophages. *Br J Nutr* 2014;**112**(9): 1469–77. DOI: 10.1017/S000711451400244X
- Al-Shayji IA, Caslake MJ, Gill JM. Effects of moderate exercise on VLDL₁ and intralipid kinetics in overweight/obese middle-aged men. *Am J Physiol Endocrinol Metab* 2012;**302**(3): E349–55. DOI: 10.1152/ajpendo.00498.2011
- Karpe F, Olivecrona T, Hamsten A, Hultin M. Chylomicron/chylomicron remnant turnover in humans: evidence for margination of chylomicrons and poor conversion of larger to smaller chylomicron remnants. *J Lipid Res* 1997;**38**(5): 949–61
- Deedwania P, Barter P, Carmena R, Fruchart JC, Grundy SM, Haffner S, Kastelein JJ, LaRosa JC, Schachner H, Shepherd J, Waters DD. Treating to new targets investigators. Reduction of low-density lipoprotein cholesterol in patients with coronary heart disease and metabolic syndrome: analysis of the treating to new targets study. *Lancet* 2006;**368**(9539): 919–28
- Hyson D, Rutledge HC, Berglund L. Postprandial lipemia and cardiovascular disease. *Curr Atheroscler Rep* 2003;**5**:437–44
- Ruf H, Gould BJ. Size distributions of chylomicrons from human lymph from dynamic light scattering measurements. *Eur Biophys J* 1999;**28**(1): 1–11
- Antelo A, Perona JS. Evaluation of a method of preparation of lipid emulsions as a model for chylomicron-like particles. *J Liposome Res* 2013;**23**(2): 126–33. DOI: 10.3109/08982104.2012.754464
- O'Neal D, Harrip P, Dragicevic G, Rae D, Best JD. A comparison of LDL size determination using gradient gel electrophoresis and light-scattering methods. *J Lipid Res* 1998;**39**:2086–90
- Sakurai T, Trirongitmoah S, Nishibata Y, Namita T, Tsuji M, Hui SP, Jin S, Shimizu K, Chiba H. Measurement of lipoprotein particle sizes using dynamic light scattering. *Ann Clin Biochem* 2010;**47**:476–81. DOI: 10.1258/acb.2010.010100

(Received July 28, 2015, Accepted February 22, 2016)