Published in final edited form as:

Behav Res Ther. 2016 October ; 85: 6–12. doi:10.1016/j.brat.2016.07.010.

Patient adherence and treatment outcome with exposure and response prevention for OCD: Which components of adherence matter and who becomes well?

Michael G. Wheaton^{1,2,3}, Hanga Galfalvy^{2,3}, Shari A. Steinman^{2,3}, Melanie M. Wall^{2,3}, Edna B. Foa⁴, and H. Blair Simpson^{2,3}

¹Ferkauf Graduate School of Psychology, Yeshiva University, Bronx, NY

²New York State Psychiatric Institute, New York, NY

³Columbia Psychiatry, Columbia University, New York, NY

⁴University of Pennsylvania, Philadelphia, PA

Abstract

Exposure and response prevention (EX/RP) is an evidence-based treatment for obsessive-compulsive disorder (OCD), yet not all patients achieve wellness with EX/RP. The degree to which patients adhere to EX/RP procedures outside of sessions has been found to predict therapy outcomes, including who achieves post-treatment wellness. We sought to investigate which components of treatment adherence most relate to outcome and to develop adherence benchmarks to identify who does and does not become well to provide clinicians with prognostic tools. Adherence data came from 37 adult patients with DSMIV OCD who received 17 sessions of EX/RP as part of a randomized controlled trial of augmentation strategies for incomplete response to serotonin reuptake inhibitors (SRIs). Therapists rated between-session patient adherence at each exposure session by quantifying: 1) the quantity of homework exposures attempted; 2) the quality of attempted exposures; and 3) the degree of success with response prevention. Each adherence item significantly correlated with post-treatment OCD severity. Success with response prevention proved particularly strongly linked to therapy outcome. Time course analysis of this item accurately identified relatively early in treatment who would achieve post-treatment wellness. These data provide an efficient method for differentiating between those patients who will and will not achieve wellness after EX/RP augmentation of SRIs. Limitations and clinical implications of the current findings are discussed.

Keywords

obsessive-compulsive disorder; Exposure and response prevention; patient adherence

Corresponding Author: H. Blair Simpson, MD/PhD, Director of the Anxiety Disorders Clinic and Center for Obsessive-Compulsive and Related Disorders, New York State Psychiatric Institute/Columbia University, 1051 Riverside Drive, Unit 69, New York, NY 10032, Phone: 646-774-8110, Fax: 646-774-8105, Simpson@nyspi.columbia.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Obsessive-compulsive disorder (OCD) affects up to 2% of the population and can be disabling when severe (Ruscio, Stein, Chu & Kessler, 2010). Cognitive behavioral therapy (CBT) consisting of exposure and response prevention (EX/RP) is an effective treatment for OCD, and is recommended in practice guidelines (Koran et al., 2007; Koran & Simpson, 2013; NICE, 2013). Yet, not all patients achieve minimal symptoms at the end of treatment (e.g., 75–80% of patients respond, yet only 40–52% achieve remission; Farris et al. 2013; Simpson et al., 2006; Simpson et al., 2008; Simpson, Foa et al., 2013). Identifying predictors of treatment response can improve patient care by providing markers to identify which individuals are likely to achieve wellness and who might require additional interventions or alternative treatments.

One factor known to affect EX/RP outcome is the degree to which patients adhere to treatment procedures (Abramowitz, Franklin, Zoellner, & DiBernardo, 2002; DeAraujo, Ito & Marks, 1996; Tolin, Maltby, Diefenbach, Hannan, & Worhunsky, 2004). Successful EX/RP requires patients to confront their fears (i.e., exposure component) as well as to voluntarily stop their rituals (i.e., response prevention component). During treatment sessions, therapists direct patients in these procedures. Patients are also asked to carry out these procedures between sessions as homework (Foa, Yadin, & Lichner, 2012). To quantify how well patients adhere to between session assignments (hereafter referred to as *patient adherence*), Simpson et al. (2010) developed the Patient EX/RP Adherence Scale (PEAS). With input from a panel of EX/RP experts, the authors devised the PEAS to tap the components of standard EX/RP homework practice (Kozak & Foa, 1997) thought to be necessary for good outcomes: confronting fears (exposures) and voluntarily stopping rituals (Foa et al., 1980). This measure includes three items that therapists rate at the beginning of an exposure session to quantify patient adherence to the previous session's EX/RP assignments: PEAS_A quantifies the number of exposures the patient attempted (as a percentage of those assigned), PEAS_B rates the quality of attempted exposures, and PEAS_C assesses the patient's degree of success with response prevention between sessions. As a global measure of patient adherence, the three PEAS items are averaged at each session and then across all sessions that include EX/RP homework assignments.

Preliminary evidence supports the use of the PEAS as a predictor of EX/RP response. In a small clinical trial of adults with OCD who were randomized to either EX/RP (N=15) or EX/RP augmented with motivational interviewing (EX/RP+MI; N=15), higher mean ratings on the PEAS predicted lower post-treatment OCD severity as assessed with the Yale-Brown Obsessive Compulsive Scale (YBOCS; Goodman et al., 1989a,b) across the full sample (Simpson et al. 2011). Moreover, mean adherence ratings predicted patient attainment of minimal OCD symptoms (YBOCS ≤ 12). Achieving post-treatment symptoms below this threshold has been associated with good quality of life and a high level of adaptive functioning (Farris et al., 2013). Thus, the PEAS may prove to be a useful tool for prognostic prediction of who is likely to become well after EX/RP and who may need additional treatment.

We capitalized on data from a large randomized controlled trial that included a group of patients undergoing manualized EX/RP to conduct three sets of analyses related to patient

EX/RP adherence. First, we sought to replicate that the PEAS predicts EX/RP outcome in an independent sample. Based on the data reviewed above, we hypothesized that the PEAS would predict degree of EX/RP response, as well as attainment of posttreatment wellness.

Second, we sought to extend previous findings by examining the predictive ability of individual PEAS items to determine which components of adherence most strongly relate to treatment outcome. Dismantling studies suggest that exposures and response prevention have independent effects and are each key to good EX/RP outcomes (Foa, Steketee, & Milby, 1980; Foa, Steketee, Grayson, Turner & Latimer, 1984), so we expected adherence to both components to predict outcome. However, a recent metaanalysis (Mausbach et al., 2010) of the relationship between homework compliance and CBT outcomes (across disorders and treatments) found relatively weaker effects for compliance ratings based on the percentage of homework completed (as opposed to Likert-based ratings of homework quality). Therefore we specifically hypothesized significant unique predictor effects for PEAS_B and PEAS_C (both of which incorporate ratings of quality), but not PEAS_A (which only involves rating quantity).

Third and finally, we evaluated the ability of the PEAS to forecast outcomes for individual patients through the use of clinically-relevant benchmarks. Specifically, we conducted time course analyses of adherence ratings across treatment to forecast who is likely to become well at the end of treatment and who is not. From this analysis, we evaluated PEAS benchmarks to determine how early in treatment they could make accurate predictions about post-treatment status (i.e., attainment of post-treatment wellness or failure to do so). In so doing, we aimed to provide tools to help treating clinicians identify patients unlikely to achieve post-treatment wellness as early in treatment as possible, so that additional or alternative interventions might be offered.

Method

Participants

Data came from 37 patients with DSM-IV OCD who completed 17 sessions of EX/RP as part of a randomized controlled trial comparing serotonin reuptake inhibitor (SRI) augmentation strategies (Simpson, Foa et al., 2013). Eligible patients were adults with a principal diagnosis of OCD determined by the Structured Clinical Interview for DSM-IV (SCID; First, Spitzer, Gibbon, & Williams, 2001), who remained symptomatic despite receiving an SRI at a maximally tolerated dose for 12 weeks or more. Exclusion criteria were: 1) diagnosis of bipolar or psychotic disorder; 2) substance abuse or dependence in the past 3 months; 3) clinically significant suicidal ideation; 4) severe depression (≥ 25 on the 17-item Hamilton Depression Rating Scale [HDRS; Hamilton, 1960]); 5) primary hoarding symptoms; or 6) previous trial of risperidone (≥ 0.5 mg/day for 8 weeks) or EX/RP (≥ 8 sessions over 2 months) while taking an SRI.

Procedures

Full description of study procedures appear elsewhere (Simpson, Foa, et al., 2013). Eligible participants were randomized to EX/RP, risperidone, or pill placebo; only participants who

completed EX/RP (n=37) are included in this report.¹ The study was conducted at two academic outpatient clinics in New York City, New York, and Philadelphia, Pennsylvania. Institutional Review Boards at both sites approved the study protocol, and patients provided written informed consent.

EX/RP sessions were 90 minutes long and comprised of two introductory sessions followed by 15 exposure sessions, daily homework assignments (self-directed exposures and response prevention), and phone check-ins between each session (Kozak & Foa, 1997). EX/RP was delivered by doctoral-level clinicians (PhD or PsyD), who participated in weekly group supervision phone calls in order to standardize treatment delivery across the two sites. Homework was assigned after the first exposure session; thus, adherence to homework was assessed (as described below) at sessions 4–17.

Independent evaluators blinded to treatment condition were assigned to individual patients across time points and assessed patients' OCD symptoms at baseline (week 0), mid-treatment (week 4) and post-treatment (week 8).

Measures

Yale Brown Obsessive Compulsive Scale (YBOCS; Goodman et al., 1989a, Goodman et al., 1989b)—The YBOCS is the “gold standard” OCD severity measure, a semi-structured clinician-administered interview used to assess symptom severity of obsessions and compulsions in the past week. Each item is rated on a 5-point Likert scale (0 = no symptoms, 4 = extreme). Total scores range from 0 to 40. The YBOCS has good internal consistency, excellent inter-rater reliability, and good test-retest reliability (Goodman et al., 1989a, Goodman et al., 1989b). Internal consistency in the present sample was acceptable ($\alpha = .73$).

Clinical Global Impression Scale (CGI; Guy 1976)—As a secondary measure, independent evaluators rated improvement in OCD severity using the CGI, a commonly used instrument for rating clinical impressions of overall illness severity and improvement. This measure uses a single item to assess illness improvement on a 7-point scale, from 1 (very much improved) to 7 (very much worse).

Patient EX/RP Adherence Scale (PEAS; Simpson et al., 2010)—The PEAS was rated by the treating clinician at each therapy session beginning after the first exposure session (i.e., sessions 4–17). The PEAS consists of three items: (a) the quantity of exposures (percentage of assigned HW exposures that were attempted), (b) quality of attempted exposures, and (c) degree of success with response prevention (percentage of urges to ritualize that the patient resisted since the previous therapy session). The therapist rates each item on a 7-point Likert-type scale. For item A, therapists ask patients how many of their assigned exposure practices they completed, which the therapist converts into a percentage, which is rated on a 7 point scale (e.g., 0% = 1; <10% = 2; ~25% = 3, etc). For PEAS_B,

¹The original trial comprised 40 OCD patients, but 3 dropped out of EX/RP. Two of these patients dropped out before PEAS data was collected while the third dropped out midway through treatment. The pattern of results in our regression analyses were identical carrying forward this patient's last observation. However, given that our analysis of PEAS over time required complete PEAS data, we report only the completer analyses in the present report.

therapists ask a semi-structured set of questions to rate how well the patient did the exposures they attempted (with prompts including “how did the _____ exposure go?” and “Were you able to complete the exposure without doing any compulsions during or afterwards?”). The therapist averages ratings across all attempted exposures since the last session and assigns a rating for PEAS_B on a 7-point scale (1-refused-did none of the assigned exposures to 7-excellent: all exposures performed as assigned by the therapist without rituals or safety aids and the patient looked for opportunities to extend the assignment to make it more challenging). For PEAS_C, therapists ask patients to rate the percent of urges to ritualize that the patient successfully resisted since the last visit (not solely during exposure practices). In rating PEAS_C, success with response prevention is also put onto a 7 point scale (e.g., 0% =1, <10%=2, ~25%=3, ~50%=4, ~75%=5, ~90%=6, ~100%=7). Mean PEAS total scores are calculated by averaging the scores on the three items (PEAS_A, PEAS_B and PEAS_C) for each session and then across sessions. The present sample mean PEAS ratings had good reliability ($\alpha = .89$).

Statistical Methods

To test whether EX/RP adherence predicts outcome, we employed subjects' mean PEAS scores (averaged across items and then across sessions) as the independent variable in two regression models. The first model used linear regression to predict post-treatment YBOCS *scores*. The second used logistic regression to predict post-treatment *wellness* (as indicated by YBOCS 12). Both models controlled for baseline YBOCS severity. Overall significance was set at $\alpha < .05$.

To investigate which PEAS items most relate to outcomes, these analyses were repeated by simultaneously entering PEAS_A, PEAS_B and PEAS_C as independent variables in lieu of mean PEAS. Because there is some overlap in item content between PEAS_C and the YBOCS items assessing compulsion severity (i.e., items assessing resistance and control over compulsive rituals), these results were followed up with sensitivity analyses predicting post-treatment YBOCS obsession and compulsion severity separately, as well as ordinal regression predicting our secondary outcome measure, post-treatment CGI-improvement.

Finally, we conducted a series of analyses to test the ability of PEAS benchmarks to forecast achievement of post-treatment wellness (i.e. YBOCS 12) throughout treatment. Specifically, we based our prediction on scoring below or above pre-specified cutoffs on cumulative average adherence scores. These were calculated for every session by averaging all the adherence scores for that subject up to that session. Based on our regression results (see below), we employed scores on PEAS_C (adherence to response prevention) for these analyses. Cumulative scores were used as a way to smooth the variability associated with individual sessions (e.g., patients may have a “good week”, followed by a “bad week”). In order to maximize interpretability and clinical utility we considered four cutoffs based on the item response anchors corresponding to approximately >90%, >75%, <75% and <50% success with response prevention. Our outcome variable was the post-treatment wellness indicator defined above (YBOCS 12). For each session, and for each cutoff (i.e., response prevention = ~>90%, >75%, <75% and <50%), we calculated the positive predictive value (i.e., the proportion of subjects who achieved post-treatment wellness, out of all those who

were above the predictor cutoff). Graphical displays of the positive predictive values by time and RP cutoff were used to demonstrate the how adherence cutoffs could be used to predict post-treatment wellness across treatment.

Results

Sample

Participant age ranged from 18 to 65 ($M=33.78$; $SD=12.54$). The sample was 51.4% female and 86.5% of participants were non-Hispanic White. Overall mean scores on the YBOCS decreased significantly from baseline ($M=27.03$, $SD=3.98$) to posttreatment ($M=13.0$, $SD=6.09$, $t(36)=12.0$, $p<.001$). The mean reduction in YBOCS score was $M=14.03$ ($SD=7.11$) points (minimum = 1; maximum=31). Seventeen participants (45.9%) attained post-treatment wellness (YBOCS ≤ 12). Descriptive statistics for PEAS items and mean scores are shown in Table 1. In addition, the three PEAS items were significantly intercorrelated (Table 1). Mean PEAS scores were not significantly related to baseline YBOCS severity, $r=-.16$, $p>.34$.

Patient adherence predicts EX/RP outcomes

In Step 1 of the regression, baseline YBOCS did not significantly account for variance in post-treatment YBOCS scores ($R^2=.002$, $F=.09$; $df=1, 35$; $p>.77$). In Step 2, mean PEAS scores explained 31% of the variance in post-treatment YBOCS scores, beyond baseline YBOCS ($R^2=.31$, $F=14.94$; $df=1, 34$; $p<.001$). As shown in Table 2, patients with higher mean PEAS (averaged across items and then across sessions) had lower post-treatment YBOCS ($\beta=-.56$, $p<.001$). Baseline YBOCS also did not predict whether a patient achieved wellness (YBOCS ≤ 12). However, mean PEAS significantly predicted wellness: a one unit increase in mean PEAS increased the odds of posttreatment wellness by a factor of 8.8 (95% CI [1.94, 39.9], $p=.005$) as shown in Table 2.

Which components of adherence predict EX/RP outcomes?

All three PEAS items were significantly negatively correlated with post-treatment YBOCS scores (with PEAS_A: $r=-.42$, $p=.010$, PEAS_B: $r=-.48$, $p=.003$, PEAS_C: $r=-.63$, $p<.001$). In a stepwise regression which entered baseline YBOCS in Step 1 and all three PEAS items simultaneously in Step 2 of the model, the three PEAS components accounted for 40% of post-treatment YBOCS variance ($R^2=.40$; $F=7.01$; $df=3, 32$; $p=.001$). As Table 3 shows, only PEAS_C emerged as a significant individual predictor, with higher values of PEAS_C associated with lower post-treatment severity.

Results were similar in sensitivity analyses (separately predicting YBOCS obsession and compulsions severity) as shown in Table 3. PEAS_C was also the only significant predictor in an ordinal regression predicting post-treatment CGI-Improvement scores from the three PEAS items (Table 4). Moreover, logistic regression revealed that higher overall mean PEAS_C across all sessions was associated with increased odds of achieving wellness (i.e., YBOCS ≤ 12) (Table 5).

Adherence benchmarks for predicting wellness throughout treatment

We analyzed PEAS_C data for our benchmark analyses because it predicted significant independent variation in outcomes and because it provides a parsimonious measure of adherence on its own. When averaged across the whole treatment, the mean PEAS_C was 5.3 (SD=.97), corresponding to somewhat greater than 75% response prevention. Those with PEAS_C scores above the sample mean were more likely to achieve post-treatment wellness than those below the mean (13 of 18 [72.2%] versus 4 of 19 [21.1%] respectively).

To aid prognostic decision making *during* treatment, we tested the predictive accuracy of *session-by-session* cumulative PEAS_C cutoffs as markers of post-treatment wellness. To do so we calculated the probability of achieving wellness at each session with PEAS data (i.e., beginning at session 4) for four different levels of ritual prevention that are captured by the PEAS_C anchors (specifically approximately >90%, >75%, <75% and <50% response prevention). Figure 1 plots the probability of achieving post-treatment wellness for each of these cutoffs. As can be seen, response prevention tended to increase across treatment and a mean response prevention of approximately 90% or higher was an excellent marker for achieving post-treatment wellness. Patients who achieved this were more than 80% likely to achieve post-treatment wellness, and this could be forecast as early as the ninth session of EX/RP (i.e., after only 6 sessions of EX/RP homework). On the other hand, mean response prevention of at least 75% was less predictive of post-treatment wellness (averaging just below 70% accuracy at predicting post-treatment wellness throughout treatment).

PEAS_C cutoffs also were able to identify individuals who were unlikely to achieve post-treatment wellness. For example, as shown in Figure 1, individuals whose cumulative PEAS_C scores indicated that they were able to resist less than half of their compulsions (i.e., worse than 50% response prevention) after five sessions of EX/RP (with EX/RP homework assigned three times) had less than a 20% probability of achieving post-treatment wellness. Later in treatment, individuals who did not achieve cumulative response prevention of 75% or better were also unlikely to achieve wellness: beginning after 10 sessions of EX/RP (8 of which included EX/RP homework), individuals with RP this low had less than an 18% likelihood of becoming well after treatment.

Discussion

We examined the relationship between patient adherence to EX/RP procedures and treatment outcome in a sample of OCD patients receiving manualized EX/RP as part of a randomized controlled trial. As hypothesized, we found a strong relationship between EX/RP outcome and mean patient adherence as measured by the PEAS, replicating a previous report (Simpson et al., 2011) in an independent sample. Expanding on that prior report, we found all three PEAS items correlated with outcome, but only success with response prevention (PEAS_C) predicted unique outcome variance. Time course analyses showed success with response prevention tended to increase as treatment progressed, and differentiated patients who ultimately achieved post-treatment wellness from those who did not. Analysis of cumulative success with response prevention yielded clinically relevant benchmarks capable of identifying individuals likely to achieve post-treatment wellness as well those unlikely to do so (e.g., 90% cumulative response prevention predicted likely

achievement of post-treatment wellness, and less than 50% cumulative response prevention predicted failure to achieve wellness).

Our replication that the PEAS predicts EX/RP outcome further validates this measure as a clinically useful tool and is noteworthy given that the extant OCD literature on predictors of response to CBT catalogues many inconsistent findings (Keeley et al., 2008; Knopp et al., 2013). As one example of a predictor with a mixed literature, some (Franklin et al., 2000; Keijsers, Hoogduin, & Schaap, 1994; Mataix-Cols, Marks, Greist, Kobak, & Baer, 2002) but not all (Foa et al., 1983; Maher et al., 2010; Rufer, Fricke, Moritz, Kloss, & Hand, 2006) findings suggest that greater baseline OCD severity predicts worse EX/RP outcome. As was previously reported on in this sample (Wheaton et al., 2015), baseline OCD severity did not relate to EX/RP outcome. This finding suggests that severe OCD does not in itself preclude an individual from benefitting from EX/RP, and that therapists should instead monitor patient adherence regardless of initial severity to determine who will likely respond well and who may need alternative treatment.

This study extends prior findings by analyzing the relationships between individual PEAS items and EX/RP outcome. Although all three PEAS items (assessing - quantity of homework exposures attempted [PEAS_A], quality of attempted exposure practices [PEAS_B], and success with response prevention between sessions [PEAS_C]) were significantly correlated with outcome (post-treatment YBOCS), only the response prevention item (PEAS_C) accounted for independent variance in our regression analyses. Importantly, item content overlaps somewhat between PEAS_C and the two YBOCS items assessing patients' ability to resist and control compulsions. However, sensitivity analyses suggested that item overlap did not account for the strong relationship between success with response prevention and treatment outcome. Specifically, PEAS_C independently predicted obsessional severity post-treatment, in addition to compulsion severity, as well as global improvement as rated on the CGI. These results suggest that adherence to response prevention robustly predicts EX/RP outcomes. Of course, in our study all patients engaged in both exposures and response prevention, making it impossible to disentangle the effects of each component in isolation. Thus, our results should not be interpreted to mean that response prevention is more important than exposure practice. Rather, our findings suggest that assessing success with response prevention serves as a strong marker of EX/RP outcome.

This finding has clinical implications, as this one item assessing success with response prevention (PEAS_C) could provide clinicians with a powerful prognostic tool for forecasting therapy outcome. For example, achieving 90% or better response prevention was linked to achievement of minimal symptoms (YBOCS = 12), a marker of overall wellness (Farris et al., 2013). Some patients were able to achieve this degree of control of their compulsions approximately midway through treatment, and these individuals were highly likely to achieve post-treatment wellness. Our data would suggest to therapists that good outcomes are likely for these patients.

Evaluating PEAS_C also proved capable of identifying individuals unlikely to achieve post-treatment wellness, relatively early in treatment. Since our data are cross-sectional and

cannot determine causality, future studies will need to determine whether adjunctive or alternative treatments can help these individuals. For example, motivational interviewing (MI) strategies have been used to enhance patient EX/RP adherence, albeit with mixed results to date (Simpson et al., 2008; Simson et al., 2010; Zuckoff, Balan & Simpson, 2014). Another approach to improve patient adherence could involve devising interventions to directly bolster the ability to resist compulsions. For example, preliminary data suggest that computerized response inhibition in conjunction with OCDstimuli training could be used to enhance response prevention to improve therapy outcomes (Kalanthoff, Steinman, & Simpson, 2016).

That PEAS_B (assessing quality of HW exposures) did not emerge as a significant individual predictor of outcome was unanticipated, given that dismantling studies have shown that both exposures and response prevention are important (Foa et al., 1980; Foa et al., 1984). One possible explanation is that the exact wording used in the PEAS to evaluate exposure practices may not precisely conform to the factors that make exposures effective. Specifically, the current PEAS_B item evaluates successful exposures based on patients' ability to complete the exercise for a sufficient amount of time without giving into compulsions or safety behaviors. However, evaluating other aspects of exposures, such as violating expectancies, learning something new, and practicing in a new context (as emphasized by Abramowitz & Arch, 2014; Craske et al., 2008; 2014) might reveal a stronger relationship between homework exposure quality and treatment outcome. Thus the PEAS might be refined to explicitly evaluate these aspects of exposure quality (i.e., formally evaluating the degree to which patients' expectancies are violated during an exposure).

Strengths of the present study include standardized delivery of EX/RP with highly trained therapists and independent evaluations of therapy outcomes. In addition, we assessed adherence using a validated instrument that was rated by therapists on a session-by-session basis, which improves upon alternative methods of assessing adherence (i.e., self-report measures, retrospective measurement at post-treatment).

However, our study also had limitations. First, this is a secondary analysis of data drawn from only one arm of a larger treatment study, and thus is subject to replication in an independent sample, in a study designed and powered for the specific aim of testing the association between adherence and outcome. Second, our assessment of adherence involved therapists rating the PEAS based on patients' self-reports, which may be subject to reporting biases (i.e., demand characteristics and social desirability) on the part of both the patient and therapist. In particular, PEAS_C involves having patients reflect back on their ability to resist compulsions during the entire period of elapsed time since the last session, which could be challenging. Guidelines for administering the PEAS (Simpson et al., 2010) suggest that therapists review self-monitoring of rituals forms (completed on a daily basis by patients) to help form accurate estimates for this item.

Third, all of the patients in our study were taking SRI medications. Thus additional study is needed to determine if our results generalize to non-medicated patients, and to investigate whether medication may influence adherence. Similarly, most of our participants were non-Hispanic White and so future work in more diverse samples is warranted. Fourth, because

treatment response was so common in our sample, we were unable to differentiate responders from non-responders (as opposed to differentiating patients who got well from those who did not get well). Similarly, the low dropout rate in the present study precluded us from investigating whether the PEAS could be used to predict dropout.

It is important to note that we cannot make causal determinations about the adherence-outcome relationship given that there was no experimental manipulation of adherence. While it could be the case that higher levels of adherence lead to better outcomes, it could also be that patients who notice improvements in their symptoms might find the treatment to be more credible and thus begin to adhere to EX/RP procedures to a greater degree. Thus adherence and outcome might influence one another in a reciprocal fashion, perhaps also involving patients' treatment expectations and how credible they find EX/RP to be. Future studies might employ session-by-session ratings of OCD severity (e.g., using the adapted version of the Florida Obsessive-Compulsive Inventory [FOCI] as recommended in the DSM-5; APA, 2013) in order to investigate whether changes in symptoms precede or follow changes in adherence. Time lag analysis with such data might elucidate the longitudinal and reciprocal influences by which adherence and outcome influence each other.

In summary, patient EX/RP adherence measured by the PEAS strongly predicted treatment outcome. The PEAS measure and administration and scoring instructions are publicly available for clinical use (published as appendix in Simpson et al., 2010). The PEAS item assessing success with response prevention proved particularly strongly linked to therapy outcome. This single item could provide clinicians with an efficient tool for identifying who is and is not likely to become well with EX/RP and it would be relatively easy for clinicians to calculate cumulative PEAS_C throughout treatment. Future study is needed to examine the relationship between adherence and outcome in other practice settings (e.g., private practice and intensive outpatient programs) with modalities of EX/RP delivery other than the twice weekly format used in the present study (e.g., once weekly, intensive outpatient, inpatient). As previously hypothesized (Simpson et al., 2010), between-session adherence might prove an even stronger predictor of outcome when therapy is less frequent. In addition, whether increasing adherence leads causally to better therapy outcomes remains an important topic for future work.

Acknowledgments

Funding/Support: This study was funded by National Institute of Mental Health grants R01 MH045436 (Dr Simpson) and R01 MH45404 (Dr Foa) and by K24 MH091555 (Dr. Simpson) and the New York State Office of Mental Hygiene.

Role of the Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Conflict of Interest Disclosures: During the parent study, Dr Simpson received research funds from Transcept Pharmaceuticals (2011–2013) and Neuropharm Ltd (2009), served on a scientific advisory board for Pfizer (for Lyrica, 2009–2010) and Jazz Pharmaceuticals (for Luvox CR [controlled release], 2007–2008), and received royalties from Cambridge University Press and UpToDate Inc. Dr Foa was a consultant to Jazz Pharmaceuticals (for Acetelion), and she receives royalties from Bantam and Oxford University Press for book sales, including a manual of cognitive-behavioral therapy for OCD.

References

- Abramowitz JS, Franklin ME, Zoellner LA, Dibernardo CL. Treatment compliance and outcome in obsessive-compulsive disorder. *Behavior Modification*. 2002; 26(4):447–463. [PubMed: 12205821]
- Abramowitz JS, Arch JJ. Strategies for improving long-term outcomes in cognitive behavioral therapy for obsessive-compulsive disorder: insights from learning theory. *Cognitive and Behavioral Practice*. 2014; 21(1):20–31.
- Craske MG, Kircanski K, Zelikowsky M, Mystkowski J, Chowdhury N, Baker A. Optimizing inhibitory learning during exposure therapy. *Behaviour research and therapy*. 2008; 46(1):5–27. [PubMed: 18005936]
- Craske MG, Treanor M, Conway CC, Zbozinek T, Vervliet B. Maximizing exposure therapy: an inhibitory learning approach. *Behaviour research and therapy*. 2014; 58:10–23. [PubMed: 24864005]
- De Araujo LA, Ito LM, Marks IM. Early compliance and other factors predicting outcome of exposure for obsessive-compulsive disorder. *The British Journal of Psychiatry*. 1996; 169(6):747–752. [PubMed: 8968633]
- Farris SG, McLean CP, Van Meter PE, Simpson HB, Foa EB. Treatment response, symptom remission and wellness in obsessive-compulsive disorder. *The Journal of clinical psychiatry*. 2013; 74(7):685. [PubMed: 23945445]
- First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JB. Structured Clinical Interview for DSM-IV Axis I Disorders, Patient Edition (SCID-P), version 2. New York, NY: Biometrics Research Department, New York State Psychiatric Institute; 1996.
- Foa EB, Grayson JB, Steketee GS, Doppelt HG, Turner RM, Latimer PR. Success and failure in the behavioral treatment of obsessive-compulsives. *Journal of consulting and clinical psychology*. 1983; 51(2):287. [PubMed: 6841773]
- Foa EB, Steketee G, Grayson JB, Turner RM, Latimer PR. Deliberate exposure and blocking of obsessive-compulsive rituals: Immediate and long-term effects. *Behavior Therapy*. 1984; 15(5):450–472.
- Foa EB, Steketee G, Milby JB. Differential effects of exposure and response prevention in obsessive-compulsive washers. *Journal of Consulting and Clinical Psychology*. 1980; 48(1):71. [PubMed: 7365045]
- Foa, EB.; Yadin, E.; Lichner, TK. Exposure and response (ritual) prevention for obsessive-compulsive disorder. 2nd. New York, NY: Oxford University Press; 2012.
- Franklin ME, Abramowitz JS, Kozak MJ, Levitt JT, Foa EB. Effectiveness of exposure and ritual prevention for obsessive-compulsive disorder: randomized compared with nonrandomized samples. *Journal of consulting and clinical psychology*. 2000; 68(4):594. [PubMed: 10965635]
- Goodman WK, Price LH, Rasmussen SA, Mazure C, Fleischmann RL, Hill CL, Charney DS. The Yale-Brown obsessive compulsive scale: I. Development, use, and reliability. *Archives of general psychiatry*. 1989a; 46(11):1006. [PubMed: 2684084]
- Goodman WK, Price LH, Rasmussen SA, Mazure C, Delgado P, Heninger GR, Charney DS. The Yale-Brown obsessive compulsive scale: II. Validity. *Archives of general psychiatry*. 1989b; 46(11):1012. [PubMed: 2510699]
- Guy W. Clinical global impression scale. *The ECDEU Assessment Manual for Psychopharmacology-Revised*. Volume DHEW Publ No ADM 76. 1976; 338:218–222.
- Hamilton M. A rating scale for depression. *Journal of neurology, neurosurgery, and psychiatry*. 1960; 23(1):56.
- Kalanthroff E, Steinman SA, Simpson HB. Personalized computerized inhibitory training for treatment refractory obsessive compulsive patients: A tool to augment exposure and response prevention. Manuscript in preparation. 2016
- Keeley ML, Storch EA, Merlo LJ, Geffken GR. Clinical predictors of response to cognitive-behavioral therapy for obsessive-compulsive disorder. *Clinical psychology review*. 2008; 28(1):118–130. [PubMed: 17531365]

- Keijsers GP, Hoogduin CA, Schaap CP. Predictors of treatment outcome in the behavioural treatment of obsessive-compulsive disorder. *The British Journal of Psychiatry*. 1994; 165(6):781–786. [PubMed: 7661936]
- Knopp J, Knowles S, Bee P, Lovell K, Bower P. A systematic review of predictors and moderators of response to psychological therapies in OCD: Do we have enough empirical evidence to target treatment? *Clinical psychology review*. 2013; 33(8):1067–1081. [PubMed: 24077387]
- Koran LM, Hanna GL, Hollander E, Nestadt G, Simpson HB. American Psychiatric Association. Practice guideline for the treatment of patients with obsessive-compulsive disorder. *American Journal of Psychiatry*. 2007; 164(suppl):5–53. [PubMed: 17849776]
- Koran, L.; Simpson, H. APA Practice Guidelines. American Psychiatric Publishing, Inc; 2013. Guideline Watch (March 2013): Practice Guideline for the Treatment of Patients with Obsessive-Compulsive Disorder.
- Kozak, MJ.; Foa, EB. Mastery of obsessive-compulsive disorder: A cognitive-behavioral approach. Graywind Publications; 1997.
- Maher MJ, Huppert JD, Chen H, Duan N, Foa EB, Liebowitz MR, Simpson HB. Moderators and predictors of response to cognitive-behavioral therapy augmentation of pharmacotherapy in obsessive-compulsive disorder. *Psychological medicine*. 2010; 40(12):2013–2023. [PubMed: 20416137]
- Mataix-Cols D, Marks IM, Greist JH, Kobak KA, Baer L. Obsessive-compulsive symptom dimensions as predictors of compliance with and response to behaviour therapy: results from a controlled trial. *Psychotherapy and psychosomatics*. 2002; 71(5):255–262. [PubMed: 12207105]
- Mausbach BT, Moore R, Roesch S, Cardenas V, Patterson TL. The relationship between homework compliance and therapy outcomes: an updated meta-analysis. *Cognitive Therapy and Research*. 2010; 34(5):429–438. [PubMed: 20930925]
- NICE. Obsessive-compulsive disorder: Evidence update September 2013. 2013 NICE clinical guideline 31. Available at <https://www.nice.org.uk/guidance/cg31/evidence/evidence-update-194847085>.
- Ruscio AM, Stein DJ, Chiu WT, Kessler RC. The epidemiology of obsessive-compulsive disorder in the National Comorbidity Survey Replication. *Molecular psychiatry*. 2010; 15(1):53–63. [PubMed: 18725912]
- Rufer M, Fricke S, Moritz S, Kloss M, Hand I. Symptom dimensions in obsessive-compulsive disorder: prediction of cognitive-behavior therapy outcome. *Acta Psychiatrica Scandinavica*. 2006; 113(5):440–446. [PubMed: 16603035]
- Simpson HB, Foa EB, Liebowitz MR, Huppert JD, Cahill S, Maher MJ, Campeas R. Cognitive-behavioral therapy vs risperidone for augmenting serotonin reuptake inhibitors in obsessive-compulsive disorder: a randomized clinical trial. *JAMA Psychiatry*. 2013; 70(11):1190–1199. [PubMed: 24026523]
- Simpson HB, Huppert JD, Petkova E, Foa EB, Liebowitz MR. Response versus remission in obsessive-compulsive disorder. *The Journal of clinical psychiatry*. 2006; 67(2):1–478.
- Simpson HB, Maher M, Page JR, Gibbons CJ, Franklin ME, Foa EB. Development of a patient adherence scale for exposure and response prevention therapy. *Behavior therapy*. 2010; 41(1):30–37. [PubMed: 20171325]
- Simpson HB, Maher MJ, Wang Y, Bao Y, Foa EB, Franklin M. Patient adherence predicts outcome from cognitive behavioral therapy in obsessive-compulsive disorder. *Journal of consulting and clinical psychology*. 2011; 79(2):247. [PubMed: 21355639]
- Simpson HB, Zuckoff AM, Maher MJ, Page JR, Franklin ME, Foa EB, Wang Y. Challenges using motivational interviewing as an adjunct to exposure therapy for obsessive-compulsive disorder. *Behaviour research and therapy*. 2010; 48(10):941–948. [PubMed: 20609435]
- Simpson HB, Zuckoff A, Page JR, Franklin ME, Foa EB. Adding motivational interviewing to exposure and ritual prevention for obsessive-compulsive disorder: An open pilot trial. *Cognitive Behaviour Therapy*. 2008; 37(1):38–49. [PubMed: 18365797]
- Tolin DF, Maltby N, Diefenbach GJ, Hannan SE, Worhunsky P. Cognitive-behavioral therapy for medication nonresponders with obsessive-compulsive disorder: a wait-list-controlled open trial. *Journal of Clinical Psychiatry*. 2004; 65(7):922–931. [PubMed: 15291681]

- Wheaton MG, Rosenfield D, Foa EB, Simpson HB. Augmenting Serotonin Reuptake Inhibitors in Obsessive–Compulsive Disorder: What Moderates Improvement? *Journal of Consulting and Clinical Psychology*. 2015 Epub ahead of print.
- Zuckoff, A.; Balan, IC.; Simpson, HB. Enhancing the Effectiveness of Exposure and Response Prevention in the Treatment of Obsessive-Compulsive Disorder: Exploring a Role for Motivational Interviewing. In: Arkowitz, H.; Westra, HA.; Miller, WR.; Rollnick, S., editors. *Motivational Interviewing in the Treatment of Psychological Disorders*. Guilford Press; 2014.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- This study examined patient adherence to between-session EX/RP assignments
- Adherence predicted treatment outcome and attainment of post-treatment wellness
- Adherence to response prevention was a strong individual predictor
- Benchmarks based on response prevention predicted outcomes for individual patients
- These results could help clinicians make prognostic decisions

Figure 1.

Note. RP=response prevention. The first exposure was conducted at session 3 (solid line), after which between sessions EX/RP homework began. Thus the first data point collected on the PEAS was at session 4 (dashed line).

Table 1

Descriptive statistics for Patient EX/RP Adherence Scale (PEAS)

Measure	M (SD)	Range	Correlations (r)		
			PEAS_A	PEAS_B	PEAS_C
PEAS_total	5.33 (.89)	3.05 – 6.73	.93	.92	.89
PEAS_A	5.33 (1.14)	2.71 – 7.0	--	.80	.70
PEAS_B	5.34 (.82)	3.5 – 6.58		--	.75
PEAS_C	5.30 (.97)	2.93 – 7.0			--

Note. PEAS total scores calculated by averaging the scores on the three items (PEAS_A, PEAS_B and PEAS_C) for each session and then across treatment. All correlations significant, $p<.001$.

Table 2

Patient Adherence and EX/RP Outcomes

Linear regression predicting post-treatment YBOCS score					
Predictor	β	<i>B</i>	95 % CI	sr^2	<i>p</i>
Baseline YBOCS	-.04	-.06	[-.51, .39]	<.01	.79
Mean PEAS	-.56	-3.83	[-5.84, -1.82]	.31	<.001
Logistic Regression Predicting Post-treatment Wellness (YBOCS 12)					
Predictor	<i>B (SE)</i>	OR	95 % CI	Wald	<i>p</i>
Baseline YBOCS	.14 (.11)	1.15	[.92, 1.44]	1.51	.219
Mean PEAS	2.18 (.77)	8.80	[1.94, 39.9]	7.95	.005

Note. YBOCS = Yale-Brown Obsessive Compulsive Scale; PEAS = Patient EX/RP Adherence Scale; CI = confidence interval; sr^2 = squared semipartial correlation, a measure of the unique variance explained by each predictor that is equivalent to the R^2 change in a hierarchical model when each predictor is entered previously.

Table 3**Components of Patient Adherence and EX/RP Outcomes**

Predictor	β	B	95 % CI	sr^2	p
Predicting post-treatment YBOCS total score					
Baseline YBOCS	-.05	-.08	[-.52, .36]	<.01	.724
PEAS_A	.08	.45	[-2.14, 3.03]	<.01	.727
PEAS_B	-.08	-.61	[-4.52, 3.29]	<.01	.751
PEAS_C	-.63	-3.95	[-6.66, -1.24]	.166	.006
Predicting post-treatment YBOCS Obsessional Severity					
Baseline YBOCS	-.08	-.06	[-.30, .18]	<.01	.625
PEAS_A	.18	.49	[-.92, 1.89]	=.01	.486
PEAS_B	-.13	-.47	[-2.59, 1.65]	<.01	.653
PEAS_C	-.55	-1.69	[-3.16, -.23]	.123	.025
Predicting post-treatment YBOCS Compulsion Severity					
Baseline YBOCS	-.02	-.02	[-.25, .22]	<.01	.870
PEAS_A	-.01	-.04	[-1.42, 1.35]	<.01	.956
PEAS_B	-.03	-.14	[-2.23, 1.95]	<.01	.892
PEAS_C	-.64	-2.26	[-3.71, -.81]	.239	.003

Note. YBOCS = Yale-Brown Obsessive Compulsive Scale; PEAS = Patient EX/RP Adherence Scale; PEAS_A = percentage of assigned HW exposures that were attempted, PEAS_B = quality of attempted exposures; PEAS_C = degree of success with response prevention. CI = confidence interval; sr^2 = squared semipartial correlation, a measure of the unique variance explained by each predictor that is equivalent to the R^2 change in a hierarchical model when each predictor is entered previously.

Table 4

Components of Patient Adherence and CGI-Improvement

predictor	<i>B (SE)</i>	95 % CI	Wald	<i>p</i>
Baseline YBOCS	-.07 (.09)	[-.25, .11]	.59	.444
PEAS_A	.18 (.54)	[-.89, 1.25]	.11	.739
PEAS_B	-.05 (.84)	[-1.69, 1.6]	.01	.957
PEAS_C	-1.69 (.67)	[-2.99, -.37]	6.33	.012

Note. YBOCS = Yale-Brown Obsessive Compulsive Scale; CGI=Clinical Global Impressions Scale; CI = confidence interval.

Table 5

Components of Patient Adherence and Post-treatment Wellness

predictor	B (SE)	OR	95 % CI	Wald	p
Baseline YBOCS	.19 (.13)	1.21	[.94, 1.55]	2.13	.145
PEAS_A	.07 (.59)	1.07	[.34, 3.37]	.01	.906
PEAS_B	.50 (1.06)	1.65	[.21, 13.08]	.23	.633
PEAS_C	2.12 (.87)	8.32	[1.52, 45.59]	5.96	.015

Note. YBOCS = Yale-Brown Obsessive Compulsive Scale; Wellness = Post-treatment YBOCS 12; CI = confidence interval.