Parathyroid Neoplasms: The Army Hospital (Research & Referral) Experience

Surg Capt B Fanthome*, Lt Col R Bharadwaj⁺, Air Cmde KM Suryanarayana[#]

Abstract

Background: Neoplasms of the parathyroid are common but parathyroid carcinoma is exceptionally rare. In contrast to most other malignant endocrine tumours that are usually less hormonally active, malignant parathyroid tumours are hyper functional. Malignant parathyroid tumours pose a diagnostic dilemma for the pathologist.

Objective: To study the clinicopathological profile of a case series of parathyroid neoplasms and determine features which facilitate a malignant diagnosis.

Methods: A retrospective analysis of seven cases of surgically treated parathyroid tumours over a three-year period at a single centre was done. Clinical, haematological, biochemical, and radiological data was accrued from medical records. The histopathology slides were reviewed along with the clinicopathological profile in an attempt to delineate markers of malignancy.

Results: Patients ranged from 30 to 58 years of age. Males and females were approximately equal. Weakness and bone pain were the commonest presenting symptoms. Over 50% had significant hypercalcaemia and all had elevated serum parathormone. Clinically apparent mass was seen in only one. All tumours were successfully localised using CT scan and MRI. Thick fibrous capsule and broad septal fibrosis was seen in both the carcinomas; these were thin in the adenomas. Mitotic counts of 1-3 per high power field (HPF), capsular invasion and nodal metastasis were noted in the malignant tumours.

Conclusion: Elevated serum calcium and parathormone values point to a parathyroid neoplasm. Current imaging modalities are successful in localising the tumour preoperatively. Markedly elevated serum calcium, broad fibrous bands, mitotic counts and capsular invasion are indicators of malignancy.

MJAFI 2006; 62 : 312-315

Key Words: Parathyroid; Adenoma; Carcinoma

Introduction

Neoplasms of the parathyroid are common, but parathyroid carcinoma is exceptionally rare [1]. In contrast to other malignant endocrine tumours that are hormonally less active, malignant parathyroid tumours are hyper functional and are characterised by high levels of serum parathormone. The hyperparathyroidism is severe with high serum calcium levels, bone disease and renal stones. The clinical picture varies with the stage at presentation and is characterised by the consequences of prolonged elevation of serum parathormone, serum calcium and local recurrences in the neck. Metastases to lungs, bone and liver are a late feature. With the increasing use of automated screening techniques along with sophisticated imaging modalities, the detection of parathyroid disease is increasing. The endocrinologist is faced with complex presentations and cryptic manifestations of a rare disease. In the absence of metastasis, parathyroid carcinoma may be suspected, but cannot be confirmed prior to operation, complicating the treatment algorithm. In this case series apart from

documenting the clinical profile, an attempt has been made to evaluate criteria useful in differentiating adenomas from carcinomas by clinicopathological correlation.

Materials and Method

Consecutive patients with parathyroid tumours presenting at the Army Hospital (Research & Referral) from Dec 1999 to Dec 2002 were evaluated. Clinical, haematological, biochemical, and radiological data was accrued from medical records. The clinical presentation, serum calcium, phosphate and parathormone levels, radiological findings, operative findings, histopathology findings and course of the disease were noted. The per-operative tumour size, adherence to surrounding structures, lymph node involvement and extension to thyroid were also assessed. The histopathology slides were reviewed to identify morphologic criteria like fibrous capsule, broad fibrous trabeculae, cellular architecture, cell type, presence of mitosis and invasion. Based on these criteria, they were categorised into adenomas or carcinomas. An attempt was made to identify the most reliable biochemical, surgical or histological indicator of aggressive disease.

*Senior Advisor (Surgery & Oncosurgery), INHS Asvini, Mumbai. ⁺Reader (Pathology & Oncopathology), Armed Forces Medical College, Pune. [#]Consultant (Medicine and Endocrinology), Command Hospital (Air Force), Bangalore.

Results

A total of seven cases of parathyroid disease were diagnosed and treated from Dec 1999 to Dec 2002. Our patients ranged from 30-58 years of age. Females and males were four and three respectively. Out of seven, four had significant hypercalcaemia ranging from 13.8 to 19.6 mg/dl. Serum parathormone was raised in all seven cases. The predominant clinical features were generalised weakness and joint pains, resulting in inability to walk. A palpable neck mass was present only in one case.

Imaging located the tumour in all cases. Renal calculi were noted in four, with associated medullary sponge kidney in one and a simple cyst of the liver in another. Generalised osteopenia associated with fracture was seen in two cases (Table 1). Osteitis fibrosa cystica was confirmed on bone marrow biopsy in one. All seven cases underwent excision of tumour mass with or without nodal dissection. The gross size of the lesion varied from 0.8 to 2.5 cms. There were five adenomas and two carcinomas. The carcinomas showed a thick fibrous capsule with broad septal fibrosis in both cases. The fibrous bands in adenomas were thin and wavy (Fig 1). The pattern was trabecular in three adenomas and in two carcinomas, making it the commonest pattern encountered in our study, but excluding it as a reliable feature to differentiate between adenomas and carcinomas. We encountered predominantly clear cell adenomas but the carcinomas were both clear cell (one case) and mixed chief and oxyphil cell type (one case). Mitotic count was 1-3 per high power field in the carcinomas while it was negligible in the adenomas. Cellular pleomorphism was seen in both adenomas and carcinomas. Capsular invasion was noted in one but was not so distinct in the second case (Fig 2). This case however, showed tumour deposits in the neck nodes along with caseating granulomas suggestive of concurrent tubercular infection.

Discussion

The rarity of parathyroid carcinoma limits reports to small institutional series. Even Massachusetts General Hospital or the Mayo Clinic have reported one to two dozen cases in clinical reviews spanning four to five decades [1,2,3]. Age distribution in most series ranged

between 14-88 years with a mean of 54 years. Our patients in the age group 30 – 58 years, reflected the demographic profile of our service population. The equal male: female ratio is comparable to most series. These tumours are difficult to diagnose clinically unless a high index of suspicion is maintained [3,4]. In most situations, symptoms are vague and often ignored both by the patient and the doctor. Patients in our series had been symptomatic from six months to five years. Bone pains, pathological fracture, or other evidence of bone diseases are seen in approximately 90%, followed by renal stones (50-80 % cases) as reported in most series. Symptoms of hypercalcaemia are vague and include fatigue, weakness or constipation. We encountered a neck mass in two patients and both had carcinoma. All the malignant cases in our series had extreme elevations of serum parathormone and serum calcium, were symptomatic with renal and bone symptoms with one having mental obtundation and chronic renal failure (CRF).

Non-functioning parathyroid carcinomas have been reported in literature, and a more aggressive course is attributed to them. All the patients in our series had hyper

Fig. 1 : Microphotograph showing the border between adenoma and normal parathyroid gland

Fig.2 : Malignant cells infiltrating surrounding fat

Table 1

Symptoms	Number
Weakness	5/7
Bone Pains	5/7
Pathological fractures	2/7
Palpable neck mass	2/7
Nephrocalcinosis	3/7
CRF	1/7
Serum Calcium	10.8 - 14.1 mg/dl (N=9-11 mg/dl)
Parathormone	980-1962 pg/ml (N=10-55pg/ml)
CT localisation	6/6
MRI localisation	1/1

Duration of symptoms ranged from six months to five years

functioning tumours. Fine needle aspiration cytology (FNAC) is not very helpful because they often present as a solitary thyroid nodule and mimic follicular thyroid neoplasms and was not used in our series [5]. No preoperative test reliably distinguishes parathyroid cancer from benign primary hyperparathyroidism. Diagnosis is based on the histological appearance of the excised parathyroid gland. The operating surgeon usually encounters a large lesion that is more firm than adenomas and is frequently grey brown versus the red brown of benign lesions, due to increased fibrous stroma within these tumours [4,6,7]. Tumours are usually encapsulated and often have fibrous septa extending into the gland. The parenchyma has a predominance of one type of cell. The carcinomas have a bland cytological appearance. The most reliable histological indicators of malignancy include broad fibrous bands, trabeculae and mitotic counts. Capsular invasion should ideally be present, but may not be seen in third of the cases. Capsular invasion and vascular invasion are not diagnostic of malignancy in parathyroid neoplasm since spontaneous haemorrhage in adenomas may result in a similar histological picture. Trabecular pattern can be encountered in both adenomas and carcinomas as in our series. Mitotic counts and capsular invasion were the most reliable indicators of malignancy in our cases [7]. Aneuploidy has been suggested as a marker of malignancy and carries poor prognosis [8]. The number of cases in our study are few for any definitive conclusions but are useful for the documentation of a significant cluster of a rare disease.

Medical management of hypercalcaemia is often difficult, only transient responses are noted and is usually the cause of death in patients with metastatic disease. In such situations, surgical intervention is necessary to control hypercalcaemia [9]. The first patient of our series was shifted in a moribund condition from the ICU for excision of a parathyroid carcinoma and made a dramatic recovery after surgery.

The goal of surgery is to remove the tumour en bloc with adherent tissue, the ipsilateral thyroid lobe and any enlarged lymph nodes. Parathyroid carcinomas are often adherent to the surrounding structures, while adenomas are easily mobilised. Recommendations on the precise extent of the dissection cannot be made because of the rarity of this condition but en bloc excision with negative margins and removal of the involved lymph nodes is probably adequate [2,9,10,11]. Parathyroid cancer is rare and no staging systems have been established. Postoperatively, bone hunger may be profound and requires substantial doses of postoperative calcium. The postoperative serum calcium levels show a satisfying decline in most cases as noted in our study.

The delayed recurrence of hypercalcaemia after successful surgery is suggestive of parathyroid carcinoma. After initial surgery, 65 % patients with parathyroid carcinomas have cervical recurrence [1,4]. One recurrent lesion was noted in our series. Whatever the primary operative procedure, all investigators agree that cervical recurrence and hypercalcaemia are best treated operatively and provide substantial palliation for many years. In our experience, two recurrences were noted in one case manifesting with hypercalcaemia. In large series from Mayo Clinic and the Lahey Clinic, no patient who underwent re-resection were cured [2,10]. Authors have found situations in which hypercalcaemia is the only evidence of recurrence and neck exploration often negative. Local recurrence after surgery does not mean parathyroid carcinoma and may be due to regrowth of incompletely resected benign lesion or due to diffuse seeding of cervical tissue called parathyromatosis, which occurs if the adenoma ruptures during excision. In neoplasms of the parathyroid 35% develop metastases, of which 30 % are in cervical lymph nodes, 40 % in lungs and 10 % in liver. Flow cytometry to analyse DNA content and molecular techniques to differentiate the genetic abnormalities identifying carcinoma from adenoma may someday elucidate diagnosis [8,12-15]. Radio-guided surgery is going to make future surgical intervention easier [16].

Morbidity and mortality are usually related to hypercalcaemia. Overall 5-year survival rate ranges from 50-75 %, with the median survival ranging between three and five years and isolated case reports of patients surviving several years with intermittent surgical debulking. A significant number of patients die of recurrent disease within 5 years or longer after the initial treatment [1,2,3,10,13].

To conclude, with the ready availability of hormone assays and better imaging modalities, more lesions of the parathyroid are being diagnosed. Patients presenting with vague symptoms are diagnosed quickly, if serum calcium or parathormone levels are ordered based on a high index of suspicion. Supporting evidence of malignancy in primary hyperparathyroidism are markedly elevated calcium levels and large gland size. The mitotic counts and broad fibrous bands along with capsular invasion are reliable morphological indicators of malignancy. Morbidity and mortality is usually related to the hypercalcaemia.

Conflicts of Interest

None identified

References

1. Fraker DL. Parathyroid Tumours. In: DeVita VT, Hellman S, Rosenberg SA, editors. *Cancer: Principles and Practice of*

- Oncology. 6th ed. Philadelphia : Lippincott Williams & Wilkins, 2001; 1763-70.
2. Van Heerden JA, Weiland LH, Remine WH, et al. Cancer of the parathyroid glands. Arch Surg 1979; 114: 475-80.
 3. Wang C, Gaz RD. Natural history of parathyroid carcinoma: diagnosis, treatment and results. Am J Surg 1985; 149:522-7.
 4. Fujimoto Y, Obara T. How to recognise and treat parathyroid carcinoma. Surg Clin North Am 1987; 67:343-57.
 5. Weymouth MD, Serpell JW, Chambers D. Palpable parathyroid adenomas presenting clinically as solitary thyroid nodule and cytologically as follicular thyroid neoplasm. ANZ J Surg 2003; 73: 36-9.
 6. Levin KE, Galante M, Clark OH. Parathyroid carcinoma versus parathyroid adenoma in patients with profound hypercalcemia. Surgery 1987; 101:649-60.
 7. Smith JF, Coombs RRH. Histologic diagnosis of carcinoma of the parathyroid gland. J Clin Pathol 1984; 37:1370-8.
 8. August DA, Flynn SD, Jones MA, et al. Parathyroid carcinoma: the relationship of nuclear DNA content to clinical outcome. Surgery 1993; 113:290-6.
 9. Obara T, Okamoto T, Ito Y, et al. Surgical and medical management of patients with pulmonary metastasis from parathyroid carcinoma. Surgery 1993; 114:1040-9.
 10. Cohn K, Silverman M, Corrado J, et al. Parathyroid Carcinoma: the Lahey Clinic experience. Surgery 1985; 98:1095-100.
 11. Vestergaard P, Mosekilde I. Cohort study on effects of parathyroid surgery on multiple outcomes in primary hyperparathyroidism. Br Med J 2003; 327:530-4.
 12. Vetto JT, Brennan MF, Woodruff J. Parathyroid carcinoma: diagnosis and clinical history. Surgery 1993; 114: 882-92.
 13. Hansen CP, Pedersen ML, Christensen L. Diagnosis, treatment and outcome of parathyroid cancer: a report of eight patients. Eu J Surg 1991; 157:517-20.
 14. Obara T, Fujimoto Y, Hirayama A, et al. Flow cytometric DNA analysis of parathyroid tumours with special reference to its diagnostic and prognostic value in parathyroid carcinoma. Cancer 1990; 65:1789-93.
 15. Sandelin K, Auer G, Bondeson L, et al. Prognostic factors in parathyroid cancer: a review of 95 cases. World J Surg 1992; 16:724-31.
 16. Takeyama H, Shioya H, Mori Y, et al. Usefulness of radio guided surgery using technetium-99m methoxyisobutylsonitrite for primary and secondary hyperparathyroidism. World J Surg 2004; 28: 576-82.

ATTENTION ADVERTISERS

Rates of advertisements in Medical Journal Armed Forces India :

	Black & White	Colour	Size
Full page (per issue)	Rs. 5000/-	Rs. 10000/-	25 x 18 cms
Half page (per issue)	Rs. 3000/-	Rs. 6000/-	18 x 12 cms

(Positive for advertisement to be supplied by client)

Inserts (two pages) Rs. 5000/- per insert (Black and White / Colour)

(Printed inserts to be supplied by client)

Address for Correspondence

EDITOR-IN-CHIEF

MEDICAL JOURNAL ARMED FORCES INDIA

C/O ARMED FORCES MEDICAL COLLEGE, PUNE - 411 040.