Published in final edited form as:

Heart Rhythm. 2016 October ; 13(10): e295–e324. doi:10.1016/j.hrthm.2016.05.024.

J Wave Syndromes Consensus Conference: Emerging Concepts & Gaps in Knowledge

Charles Antzelevitch, PhD¹, Gan-Xin Yan, MD, PhD², Michael J. Ackerman, MD, PhD³, Martin Borggrefe, MD⁴, Domenico Corrado, MD, PhD⁵, Jihong Guo, MD⁶, Ihor Gussak, MD, PhD⁷, Can Hasdemir, MD⁸, Minoru Horie, MD⁹, Heikki Huikuri, MD¹⁰, Changsheng Ma, MD¹¹, Hiroshi Morita, MD, PhD¹², Gi-Byoung Nam, MD, PhD¹³, Frederic Sacher, MD, PhD¹⁴, Wataru Shimizu, MD, PhD¹⁵, Sami Viskin, MD¹⁶, and Arthur A. M. Wilde, MD, PhD¹⁷

¹Lankenau Institute for Medical Research, Wynnewood, PA, USA ²Lankenau Medical Center, Wynnewood, PA, USA ³Departments of Cardiovascular Diseases, Pediatrics, and Molecular Pharmacology & Experimental Therapeutics; Divisions of Heart Rhythm Services and Pediatric Cardiology; Windland Smith Rice Sudden Death Genomics Laboratory, Mayo Clinic, Rochester, MN, USA ⁴1st Department of Medicine-Cardiology, University Medical Centre Mannheim, Mannheim, Germany and DZHK (German Centre for Cardiovascular Research), partner site Heidelberg/Mannheim, Mannheim, Germany ⁵Dept. of Cardiac, Thoracic and Vascular Sciences University of Padua Medical School, Padua, Italy ⁶Division of Cardiology, Peking University of People's Hospital, Beijing, China ⁷Rutgers University, New Brunswick, NJ USA ⁸Department of Cardiology, Ege University School of Medicine, Izmir, Turkey ⁹Shiga University of Medical Sciences, Ohtsu, Shiga, Japan ¹⁰Research Unit of Internal Medicine, Medical Research Center, Oulu University Hospital, and University of Oulu, Finland ¹¹Department of Cardiology, Beijing Anzhen Hospital, Capital Medical University, National Clinical Research Center for Cardiovascular Diseases, Beijing, China ¹²Department of Cardiovascular Therapeutics, Okayama University Graduate School of Medicine. Okayama, Japan ¹³Heart Institute, Asan Medical Center and Department of Internal Medicine, University of Ulsan College of Medicine Seoul, Korea ¹⁴Bordeaux University Hospital, LIRYC Institute/INSERM 1045, Bordeaux, France ¹⁵Department of Cardiovascular Medicine, Nippon Medical School, Tokyo, Japan ¹⁶Tel-Aviv Medical Center and Sackler School of Medicine, Tel Aviv University, Israel ¹⁷Heart Center, Department of Clinical and Experimental Cardiology, Academic Medical Center, University of Amsterdam, the Netherlands and Princess Al-Jawhara Al-Brahim Centre of Excellence in Research of Hereditary Disorders, Jeddah, Kingdom of Saudi Arabia

Address for correspondence: Charles Antzelevitch, PhD, FACC, FHRS, FAHA, Professor and Executive Director, Cardiovascular Research, Lankenau Institute for Medical Research, Director of Research, Lankenau Heart Institute, 100 East Lancaster Avenue, Wynnewood, PA 19096, cantzelevitch@gmail.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Sudden cardiac death; J wave; Brugada Syndrome; Early Repolarization Syndrome; cardiac arrhythmias; ventricular fibrillation; inherited cardiac arrhythmias syndromes

Preamble

The J wave syndromes, consisting of the Brugada (BrS) and Early Repolarization Syndromes (ERS), have captured the interest of the cardiology community over the past two decades, following the identification of BrS as a new clinical entity by Pedro and Josep Brugada in 1992.¹ The clinical impact of ERS was not fully appreciated until 2008.^{2,3,4} Consensus conferences dedicated to BrS were held in 2000 and 2004.^{5, 6} A consensus conference specifically focused on ERS has not previously been convened other than that dealing with terminology, and Guidelines for both syndromes were last considered in 2013.⁷ A great deal of new information has emerged since. The present forum was organized to evaluate new information and highlight emerging concepts with respect to differential diagnosis, prognosis, cellular and ionic mechanisms, and approaches to therapy of the J wave syndromes. Leading experts, including members of the Heart Rhythm Society (HRS), the European Heart Rhythm Association (EHRA) and the Asian-Pacific Heart Rhythm Society (APHRS) met in Shanghai, China, in April 2015. The task force was charged with a review of emerging concepts and assessment of new evidence for or against particular diagnostic procedures and treatments. Every effort was made to avoid any actual, potential, or perceived conflict of interest that might arise as a result of outside relationships or personal interest. This consensus report is intended to assist healthcare providers in clinical decision making. The ultimate judgment regarding care of a particular patient, however, must be made by the healthcare provider based on all of the facts and circumstances presented by the patient.

Members of this Task Force were selected to represent professionals involved with the medical care of patients with the J wave syndromes, as well as those involved in research into the mechanisms underlying these syndromes. These selected experts in the field undertook a comprehensive review of the literature. Critical evaluation of methods of diagnosis, risk stratification, approaches to therapy and mechanistic insights was performed, including assessment of the risk–benefit ratio. The level of evidence and the strength of the recommendation of particular management options were weighed and graded. Recommendations with Class designations are taken from HRS, EHRA, APHRS, and/or ESC consensus statements or guidelines.^{8, 9} Recommendations without Class designations are derived from unanimous consensus of the authors. The consensus recommendations in this document use the commonly used Class I, IIa, IIb and III classifications and the corresponding language: “is recommended” for Class I consensus recommendation; “can be useful” or “is reasonable” for a Class IIa; “may be considered” to signify a Class IIb; and “is not recommended” for a Class III consensus recommendation.

Introduction

The appearance of prominent J waves in the electrocardiogram (ECG) have long been reported in cases of hypothermia¹⁰⁻¹² and hypercalcemia.^{13, 14} More recently, accentuation of the J wave has been associated with life-threatening ventricular arrhythmias.¹⁵ Under these circumstances, the accentuated J wave typically may be so broad and tall as to appear as an ST segment elevation, as in cases of Brugada syndrome (BrS). In humans, the normal J wave often appears as a J point elevation, with part of the J wave buried inside the QRS. An early repolarization (ER) pattern in the ECG, consisting of a distinct J wave or J point elevation, a notch or slur of the terminal part of the QRS with and without an ST segment elevation, has traditionally been viewed as benign.^{16, 17} The benign nature of an ER pattern was challenged in 2000¹⁸ on the basis of experimental data showing that this ECG manifestation predisposes to the development of polymorphic ventricular tachycardia and fibrillation (VT/VF) in coronary-perfused wedge preparations.^{15, 18-20} Validation of this hypothesis was provided eight years later by Haïssaguerre et al.,² Nam et al.³ and Rosso et al.⁴ These seminal studies together with numerous additional case control and population-based studies have provided clinical evidence for an increased risk for development of life-threatening arrhythmic events and sudden cardiac death (SCD) among patients presenting with an ER pattern, particularly in inferior and infero-lateral leads. The lack of agreement regarding the terminology relative to ER has led to a great deal of confusion and inconsistency in reporting.²¹⁻²³ A recent expert consensus report focused on the terminology of early repolarization recommends that peak of an end QRS notch and/or the onset of an end QRS slur be designated as J_p and that J_p should exceed 0.1mV in 2 contiguous inferior and/or lateral leads of a standard 12-lead ECG for early repolarization to be present.²⁴ It was further recommended that the start of the end QRS notch or J wave be designated as J_o and the termination as J_t.

Early repolarization syndrome (ERS) and BrS are thought to represent two manifestations of the J wave syndromes. Both syndromes are associated with vulnerability to development of polymorphic ventricular tachycardia (VT) and ventricular fibrillation (VF) leading to SCD^{1-3, 15} in young adults with no apparent structural heart disease and occasionally to sudden infant death syndrome.²⁵⁻²⁷ The region generally most affected in BrS is the anterior right ventricular outflow tract; in ERS, it is the inferior region of the left ventricle.^{2, 4, 28-32} As a consequence, BrS is characterized by accentuated J waves appearing as a coved-type ST segment elevation in the right precordial leads V1-V3, whereas ERS is characterized by J waves, J_o elevation, notch or slur of the terminal part of the QRS and ST-segment or J_t elevation in the lateral (type I), infero-lateral (type II) or in infero-lateral + anterior or right ventricular leads (type III).¹⁵ Early repolarization pattern (ERP) is often encountered in ostensibly healthy individuals, particularly in young males, black individuals and athletes. ERP is also observed in acquired conditions, including hypothermia and ischemia.^{15, 33, 34} When associated with VT/VF in the absence of organic heart disease, ERP is referred to as early repolarization syndrome (ERS).

The prevalence of BrS with a type 1 ECG in adults is higher in Asian countries, such as Japan (0.15 to 0.27%),^{35, 36} the Philippines (0.18%),³⁷ and among Japanese-Americans in North America (0.15%)³⁸ than in Western countries, including Europe (0 to 0.017%)³⁹⁻⁴¹ or

North America (0.005 to 0.1%).^{42, 43} In contrast, the prevalence of an ER pattern in the inferior and/or lateral leads with a J point elevation of 0.1 mV ranges between 1% and 24%, and for J point elevation of > 0.2 mV, it ranges between 0.6% to 6.4%.⁴⁴⁻⁴⁶ No significant regional differences have been reported in the prevalence of an ER pattern.⁴⁷ However, ER pattern is significantly more common in blacks than in Caucasians. Little in the way of regional differences in the manifestation of ERS has been reported. Early repolarization pattern appears to be more common in Aboriginal Australians than in Caucasian Australians.⁴⁸

Updates on the Diagnosis of BrS

According to the 2013 Consensus statement on inherited cardiac arrhythmias⁸ and 2015 Guidelines for the management of patients with ventricular arrhythmias and prevention of sudden cardiac death⁹: “BrS is diagnosed in patients with ST-segment elevation with type 1 morphology 2 mm in 1 lead among the right precordial leads V1, V2, positioned in the 2nd, 3rd or 4th intercostal space occurring either spontaneously or after provocative drug test with intravenous administration of Class I antiarrhythmic drugs. BrS is diagnosed in patients with type 2 or type 3 ST-segment elevation in 1 lead among the right precordial leads V1, V2 positioned in the 2nd, 3rd or 4th intercostal space when a provocative drug test with intravenous administration of Class I antiarrhythmic drugs induces a type I ECG morphology.”

The present task force is concerned that this could result in an over-diagnosis of BrS, particularly in patients displaying a Type 1 ECG only after a drug challenge. Data suggest the latter population is at very low risk, and that the presumed false positive rate of pharmacologic challenge is not trivial⁴⁹. Though a rigorous process was undertaken to establish the preceding guidelines, there remains no gold standard for establishing a diagnosis, particularly in patients with weak evidence of disease. Accordingly, we recommend adoption of the following diagnostic criteria and score system for BrS. Consistent with the recommendation of the 2013 and 2015 guidelines, only a Type 1 (“coved type”) ST segment elevation is considered diagnostic of BrS (**Figure 1**), and BrS is characterized by ST segment elevation 2 mm (0.2 mV) in 1 right precordial leads (V1-V3) positioned in the 4th, 3rd or 2nd intercostal space. However, as a departure from the guidelines, this consensus report recommends that when a Type 1 ST segment elevation is unmasked using sodium channel blockers (**Table 1**), diagnosis of BrS should require that the patient also present with one of the following: documented VF or polymorphic ventricular tachycardia, syncope of probable arrhythmic cause, a family history of sudden cardiac death at <45 years old with negative autopsy, coved-type ECGs in family members, or nocturnal agonal respiration. Inducibility of VT/VF with one or two premature supports the diagnosis of BrS under these circumstances.⁵⁰

A Type 2 (“saddle-back type”) or Type 3 ST segment elevation cannot substitute for a Type 1, unless converted to Type 1 with fever or sodium drug challenge. A drug-challenge-induced Type 1 can be used to diagnose BrS only if accompanied by one of the criteria specified above. Type 2 is characterized by ST segment elevation of 0.5 mm (generally 2 mm in V2) in 1 right precordial lead (V1-V3), followed by a convex ST. The ST segment is

followed by a positive T wave in V2 and variable morphology V1. Type 3 is characterized by either a saddleback or coved appearance with an ST-segment elevation of <1 mm. The placement of the right precordial leads in more cranial positions (in the 3rd or 2nd intercostal space) in a 12-lead resting ECG or 12-lead Holter ECG increases the sensitivity of ECG.⁵¹⁻⁵³ It is recommended that ECG recordings be obtained in standard and superior positions for the V1 and V2 leads. Veltman et al. showed that RV outflow tract (RVOT) localization using MRI correlates with Type 1 ST segment elevation in BrS and that lead positioning according to RVOT location improves the diagnosis of BrS. Interestingly, a Type I pattern was in most cases found in the 3rd intercostal space in sternal and left-parasternal positions.⁵⁴ In reviewing ECGs of a large cohort of BrS patients, Richter et al. concluded that lead V3 does not yield diagnostic information in BrS.⁵⁵

A proposed Diagnostic Score System for BrS, referred to as the Proposed Shanghai BrS Score, is presented in **Table 2**. These recommendations are based on the available literature and on the clinical experience of members of the task force.^{8, 56-60} Weighting of variables is based on expert opinion informed by cohort studies that typically do not include all variables presented. Thus, rigorous, objectively weighted coefficients were not derived from large-scale risk factor and outcome-informed datasets. Nonetheless, the authors felt that some inferential weighting would be of benefit when applied to patients. As with all such recommendations they will need to undergo initial and ongoing validation by future studies.

Pharmacological tests and other diagnostic tools

When there is clinical suspicion of BrS in the absence of spontaneous Type 1 ST segment elevation, a pharmacological challenge using a sodium channel blocker is recommended. A list of agents used for this purpose is presented in **Table 1** (also see www.brugadadrugs.org). The test is considered positive only if a Type 1 ECG pattern is obtained, and it should be discontinued in case of frequent ventricular extrasystoles or other arrhythmias, or widening of the QRS >130% over the baseline value.⁶ As an alternative, the “full stomach test” has been proposed for diagnosing BrS.⁶¹ In this case, ECGs are performed before and after a large meal. The use of “high-electrodes” increases the sensitivity for recognizing spontaneous Type I ST segment elevation at night or after heavy meals.⁶² A Type 1 ST segment elevation recorded using a Holter is a spontaneous Type 1, and it is reasonable to assume that a spontaneous Type 1 recorded by Holter at night or after a large meal has more value — both diagnostic and prognostic — than a drug-induced Type 1.

Drug challenge is not indicated in asymptomatic patients displaying the Type 1 ECG under baseline conditions because of the lack of the additional diagnostic value. These provocative drug tests are also not recommended in cases in which fever has been documented to induce a Type I ECG, other than for research purposes. Much debate has centered around the definition of a false positive sodium channel block challenge.⁶³ The consensus is that it is difficult to define a false positive because of the lack of a gold standard. The development of a Type 1 ST segment elevation in response to sodium block challenge should be considered as probabilistic, rather than binary, in nature. As will be discussed below, a similar approach is recommended in evaluating the ability of genetic variants to promote the BrS phenotype.

Asymptomatic patients with a family history of BrS or SCD should be informed of the availability of a sodium channel blocker challenge test to provide a more definitive diagnosis of BrS. However, patients should be advised that no therapy may be recommended regardless of the outcome, because the long-term risk of patients with BrS diagnosed by this test is significantly lower than the risk of patients with spontaneous Type 1. Patients should also be informed about the risk of the test and about the emotional consequences of having a positive test not followed by definitive therapy. The decision as to whether to undergo the drug challenge ultimately should be left up to the well-informed patient.

Performing an ajmaline test in children is problematic for two reasons: First, the test is apparently less sensitive in children than in adults. In fact, in one study, a repeat ajmaline challenge performed after puberty unmasked BrS in 23% of relatives with a previously negative drug test performed during childhood.⁶⁴ Second, the test is associated with greater risk than in adults. In one series, 10% of children undergoing the ajmaline test, including 3% of the asymptomatic subgroup, developed sustained VT.^{64, 65} Caution should also be exercised when performing a sodium blocker challenge in adults with a known pathogenic sodium channel mutation or in patients with prolonged PR intervals, pointing to a carrier of such a mutation.⁶⁶

Differential diagnosis

Other causes of ST segment elevation should be excluded before establishing the diagnosis of BrS (**Table 3**). Artifacts secondary to low-pass filtering should be ruled out.⁶⁷

Circumstances that produce a Type 1 Brugada-like ECG include right bundle branch block (RBBB), pectus excavatum, arrhythmogenic right ventricular cardiomyopathy (ARVC), as well as occlusion of the left anterior descending artery or the conus branch of the right coronary artery, which supplies the RVOT (**Table 3A**).

Discrimination between BrS and ARVC is particularly challenging. Although debate continues as to the extent to which structural abnormalities are present in BrS, most investigators consider BrS to be a channelopathy. Concealed structural abnormalities, such as histologic myocardial fibrosis of the RVOT, which may not become evident using conventional imaging techniques, have been proposed to account for or contribute to delayed conduction and ventricular arrhythmias in BrS. MRI and electron beam computed tomography studies of BrS patients consistently show subtle abnormalities, including wall motion abnormalities and reduced contractile function of the right ventricle (RV) and to a lesser extent of the left ventricle (LV), and dilatation of the RVOT.⁶⁸⁻⁷¹ In the only study that discriminated between patients with and without *SCN5A* mutations, no difference was observed in RVOT dimensions or RV ejection fraction between such patients. Slightly greater depression of LV dimensions and ejection fraction were observed in patients with *SCN5A* mutations. Significant differences were observed in RV and LV dimensions and ejection fraction when compared to healthy controls.⁷² Cardiac dilatation and reduced contractility in all of these studies were attributed to structural changes (fibrosis, fatty degeneration). However, as noted by van Hoorn and co-workers, virtually no signs of

fibrosis or fatty degeneration could be detected, perhaps due to the fact that the spatial resolution of imaging used was too low to detect such subtle changes.⁷²

Antzelevitch and colleagues have long suggested an alternative explanation.^{31, 73, 74} Loss of the action potential (AP), which has been shown in experimental models to create the arrhythmogenic substrate in BrS, leads to contractile changes that could explain the wall motion abnormalities observed. The all-or-none repolarization at the end of phase 1 of the epicardial AP responsible for loss of the dome causes the calcium channel to inactivate very soon after it activates. As a consequence, calcium channel current is dramatically reduced, the cell becomes depleted of calcium, and contractile function ceases in those cells. This is expected to lead to wall motion abnormalities, particularly in the RVOT, dilatation of the RVOT region and reduced ejection fraction observed in patients with BrS. It has also been proposed that the loss of the AP dome, because it creates a hibernation-like state, may, over long periods of time, lead to mild structural changes, including intracellular lipid accumulation, vacuolization and connexin 43 redistribution. These structural changes may, in turn, contribute to the arrhythmogenic substrate of BrS, although they are very different from those encountered in ARVC/D.^{31, 75} This hypothesis would predict that some of the changes observed by recent studies⁷⁶ may be the result of, rather than the cause of, the BrS phenotype.

A recent study by Nademanee et al. reports additional evidence pointing to pathological changes in the RVOT of patients with BrS that have proved undetectable by echocardiography or MRI.⁷⁶

In contrast, imaging techniques in ARVC clearly display morphological and functional changes (e.g., dilatation, bulging/aneurysms, wall motion abnormalities). ARVC is an inherited cardiac disease resulting from genetically defective desmosomal (DS) proteins,^{77, 78} characterized by fibro-fatty myocardial replacement predisposing to scar-related ventricular arrhythmias that may lead to SCD, mostly in young people and athletes.⁷⁹ Life-threatening ventricular arrhythmias may occur early, during the “concealed phase” of the disease, prior to overt structural changes.^{77, 78, 80} Recent experimental studies demonstrated that loss of expression of DS proteins may induce electrical ventricular instability by causing sodium channel dysfunction and current reduction as a consequence of the cross-talk between these molecules at the intercalated discs, which predisposes to sodium current-dependent lethal arrhythmias, similar to those leading to SCD in patients with J-wave/Brugada syndromes.⁸⁰⁻⁸² Further evidence of the overlap between phenotypic manifestation of ARVC and J-wave/Brugada syndromes comes from: 1) clinic-pathologic studies showing that a subset of ARVC patients may share ECG changes and patterns of ventricular arrhythmias with BrS;⁸³ and 2) genotype-phenotype correlation studies demonstrating that PKP2 mutation may cause a Brugada phenotype in the human heart by reducing sodium current.⁸⁴ These findings support the concept that specific DS-gene mutations involved in the pathogenesis of ARVC can lead to a decreased depolarization reserve that manifests as J-wave/Brugada syndromes. Thus, ARVC and J-wave/Brugada syndromes are not completely different conditions, but the ends of a spectrum of structural myocardial abnormalities and sodium current deficiency that share a common origin as diseases of the connexome.⁸⁴ The ECG abnormalities in ARVC are not dynamic and display

a constant T-wave inversion, epsilon waves and, in the progressive stage, reduction of the R amplitude. End-stage ARVC is usually associated with monomorphic VT with left bundle branch morphology and is precipitated by catecholamines,⁸⁵ whereas BrS is associated with polymorphic VT predominantly during sleep or rest.⁸⁶ A positive ajmaline challenge has been reported in 16% of patients with ARVC.^{87, 88}

Modulating factors

Sympatho-vagal balance, hormones, metabolic factors, and pharmacological agents are thought to modulate not only the ECG morphology but also explain the development of ventricular arrhythmias under certain conditions.⁸⁹ If any of these modulating factors is present, it should be promptly corrected (**Table 3B**).

Acquired Brugada pattern and phenocopies

The Brugada ECG is often concealed and can be unmasked with a wide variety of drugs and conditions, including a febrile state, vagotonic agents and maneuvers, α adrenergic agonists, β adrenergic blockers, Class IC antiarrhythmic drugs, tricyclic or tetracyclic antidepressants, hyperkalemia, hypokalemia, hypercalcemia, as well as by alcohol and cocaine toxicity.⁹⁰⁻¹⁰⁰ Pre-excitation of RV can unmask the BrS phenotype in cases of RBBB.¹⁰¹ An up-to-date list of agents known to unmask the Brugada ECG and that should be avoided by patients with BrS can be found at www.brugadadrugs.org.⁸⁹

Environmental factors leading to the appearance of an ECG similar or identical to a Type 1 BrS pattern in the absence of any apparent genetic dysfunction has been suggested to represent a Brugada ECG *phenocopy*.¹⁰² Features of the Brugada phenocopies include: 1) Brugada-like ECG pattern; 2) presence of an identifiable underlying condition; 3) disappearance of the ECG pattern after resolution of the condition; 4) absence of family history of SD in relatively young first-degree relatives (< 45 years) or of Type 1 BrS pattern; 5) absence of symptoms such as syncope, seizures or nocturnal agonal respiration; and 6) a negative sodium channel-blocker challenge test. Debate continues as to the appropriateness of this terminology given that it is very difficult to rule out a genetic predisposition, which is a prerequisite for designating the ECG manifestation as a phenocopy. Designation of these conditions as acquired forms of Brugada ECG pattern or BrS may be more appropriate and better aligned with the terminology used in the long QT syndrome.

Update on the diagnosis of ERS

ERS is generally diagnosed in patients who display ER in the inferior and/or lateral leads presenting with aborted cardiac arrest, documented VF or polymorphic ventricular tachycardia. Consistent with the recent Consensus Report on ER pattern,²⁴ early repolarization is recognized if: 1) there is an end QRS notch (J wave) or slur on the downslope of a prominent R wave with and without an ST segment elevation; 2) the peak of the notch or J wave (Jp) ≥ 0.1 mV in two or more contiguous leads of the 12 lead ECG, excluding leads V1-V3; and 3) QRS duration (measured in leads in which a notch or slur is absent) < 120 ms. **Table 4** lists the exclusion criteria in the differential diagnosis of ERS.

A proposed Diagnostic Score System for ERS, referred to as the Proposed Shanghai ERS Score, is presented in Table 5. The scoring system is based on evidence available in the literature to date. As in BrS, weighting of variables is based on expert opinion informed by cohort studies that do not include all variables presented. Thus, rigorous, objectively weighted coefficients were not derived from large-scale risk factor- and outcome-informed datasets. Nonetheless, the authors felt that some inferential weighting would be of benefit when applied to patients. As with all such recommendations, they will need to undergo initial and ongoing validation by future studies.

Similarities and difference between BrS and ERS

BrS and ERS display several clinical similarities, suggesting similar pathophysiology (Table 6).^{19, 21, 103-105} Males predominate in both syndromes: in BrS, presenting in 71%-80% among Caucasians and 94%-96% among Japanese.^{106, 107} In the setting of ER pattern, VF occurred mainly in males (72%) when studied in an international cohort,² but in a much higher percentage in a report by Japanese investigators.¹⁰⁸ BrS and ERS patients may be totally asymptomatic until presenting with cardiac arrest. In both syndromes, the highest incidence of VF or SCD occurs in the third decade of life, perhaps tied to testosterone levels in males.¹⁰⁹ In both syndromes, the appearance of accentuated J waves and ST segment elevation are generally associated with bradycardia or pauses.^{110, 111} This can explain why VF in both syndromes often occurs during sleep or at a low level of physical activities.^{108, 112} The QT interval is relatively short in patients with ERS^{2, 113} and BrS who carry mutations in calcium channel genes.¹¹⁴

As will be discussed in more detail below, ERS and BrS also share similarities with respect to the response to pharmacological therapy. In both, electrical storms and associated J wave manifestations can be suppressed using β -adrenergic agonists.¹¹⁵⁻¹¹⁸ Chronic oral pharmacological therapy using quinidine,^{119, 120} bepridil,¹¹⁷ denopamine,^{115, 121} and cilostazol^{115, 117, 121-125} is reported to suppress the development of VT/VF in both ERS and BrS secondary to inhibition of I_{to} , augmentation of I_{Ca} , or both.^{3, 122, 126}

Differences between the two syndromes include: 1) the region of the heart most affected (RVOT vs. inferior LV); 2) the presence of (discrete) structural abnormalities in BrS and not in ERS; 3) the incidence of late potentials in signal-averaged ECGs (SAECG: BrS [60%] > ERS [7%])¹⁰⁸; and 4) greater elevation of J_O , J_P or J_t (ST segment elevation) in response to sodium channel blockers in BrS vs. ERS and higher prevalence of atrial fibrillation in BrS vs. ERS.¹²⁷ Early studies suggested a different pathophysiological basis for ERS and BrS based on the observation that sodium channel blockers unmask or accentuate J wave manifestation in BrS, but reduces the amplitude in ERS.¹⁰⁸ The recent study by Nakagawa et al., however, showed that J waves recorded using unipolar LV epicardial leads introduced into the left lateral coronary vein in ERS patients are indeed augmented, even though J waves recorded in the lateral precordial leads are diminished, due principally to engulfment of the surface J wave by the widened QRS.^{29, 108} The case report of Nakagawa et al. has recently been supplemented with additional cases in which this technique was used; two of these three cases showed pilsicainide-induced accentuation of the J waves in electrograms recorded from the epicardial surface of the LV (Morita, unpublished observation). Also in

support of the thesis that these ECG patterns and syndromes are closely related are reports of cases in which ERS transitions into ERS plus BrS.^{105, 128}

The principal difference between BrS and ERS has to do with the region of the ventricle most affected. Epicardial mapping studies in BrS patients report accentuated J waves, fragmented and/or late potentials in the epicardial region of the RVOT,¹²⁹⁻¹³¹ whereas in ERS, only accentuated J waves, particularly in the inferior wall of LV, are observed.²⁹ Fractionated electrogram activity and late potentials have been observed in experimental models of ERS,³⁰ but as yet have not been reported clinically. Non-invasive mapping electro-anatomical studies have reported very steep localized repolarization gradients across the inferior/lateral regions of LV of ERS patients, preceded by normal ventricular activation,¹³² whereas in BrS both slow, discontinuous conduction and steep dispersion of repolarization are present in the RVOT.¹³³ Another presumed difference is the presence of structural abnormalities in BrS, which as yet have not been described in ERS.⁷⁶

Although J waves are accentuated or induced by both hypothermia and fever,^{33, 34, 134-139} the development of arrhythmias in ERS is much more sensitive to hypothermia, and arrhythmogenesis in BrS appears to be promoted only by fever.^{33, 34, 138, 139} Hypothermia has been reported to increase the risk of VF in ERS,^{33, 34, 134, 135, 140} and fever is well recognized as a major risk factor in BrS.^{138, 139} It is noteworthy that hypothermia can diminish the manifestation of a BrS ECG when already present.^{141, 142}

An ER pattern is associated with an increased risk for VF in patients with acute myocardial infarction¹⁴³ and hypothermia.^{33, 144} A concomitant ER pattern in the infero-lateral leads has also been reported to be associated with an increased risk of arrhythmic events in patients with BrS. Kawata et al. reported that the prevalence of ER in infero-lateral leads was high (63%) in BrS patients with documented VF.¹⁴⁵

Genetics

BrS has been associated with variants in 18 genes (**Table 7**). To date, more than 300 BrS-related variants in *SCN5A* have been described^{21, 146-148} **Figure 2** shows the overlap syndromes attributable to genetic defects in *SCN5A*. Loss-of-function mutations in *SCN5A* contribute to the development of both BrS and ERS, as well as to a variety of conduction diseases, Lenegre disease and Sick Sinus Syndrome. The available evidence suggests that the presence of a prominent I_{to} determines whether loss-of-function mutations resulting in a reduction in I_{Na} will manifest as BrS/ERS or as conduction disease.^{59, 149-151}

Variants in *CACNA1C* (Cav1.2), *CACNB2b* (Cavβ2b) and *CACNA2D1* (Cavα2δ) have been reported in up to 13% of probands.¹⁵²⁻¹⁵⁵ Mutations in glycerol-3-phosphate dehydrogenase 1-like enzyme gene (*GPD1L*), *SCN1B* (β₁-subunit of Na channel), *KCNE3* (MiRP2), *SCN3B* (β3-subunit of Na channel), *KCNJ8* (Kir 6.1), *KCND3* (Kv4.3), *RANGRF* (MOG1), *SLMAP*, *ABCC9* (SUR2A), (Navβ2), *PKP2* (Plakophilin-2), *FGF12* (FHAF1), *HEY2*, and *SEMA3A* (Semaphorin) are relatively rare.¹⁵⁶⁻¹⁷⁶ An association of BrS with *SCN10A*, a neuronal sodium channel, was recently reported.^{167, 177, 178} A wide range of yields of variants were reported by the two studies that examined the prevalence of

pathogenic *SCN10A* mutations and rare variants (5%-16.7%).¹⁷⁷⁻¹⁷⁹ Mutations in these genes lead to loss of function in sodium (I_{Na}) and calcium (I_{Ca}) channel currents, as well as to a gain of function in transient outward potassium current (I_{to}) or ATP-sensitive potassium current (I_{K-ATP}).¹⁷⁸

New susceptibility genes recently proposed and awaiting confirmation include the Transient Receptor Potential Melastatin Protein 4 gene (*TRPM4*)¹⁸⁰ and the *KCND2* gene. The mutation uncovered in *KCND2* in a single patient was shown to cause a gain of function in I_{to} when heterologously expressed.¹⁸¹

Variants in *KCNH2*, *KCNE5*, *SEMA3A*, although not causative, have been identified as capable of modulating the substrate for the development of BrS.¹⁸²⁻¹⁸⁵ Loss-of-function mutations in *HCN4* causing a reduction in pacemaker current, I_f , can unmask BrS by reducing heart rate.¹⁸⁶

An ER pattern in the ECG has been shown to be familial.¹⁸⁷⁻¹⁸⁹ The ER pattern and ERS have been associated with variants in 7 genes. Consistent with the findings that I_{K-ATP} activation can generate an ER pattern in canine ventricular wedge preparations, variants in *KCNJ8* and *ABCC9*, responsible for the pore forming and ATP-sensing subunits of the I_{K-ATP} channel, have been reported in patients with ERS.^{156, 158, 190} Loss-of-function variations in the $\alpha 1$, $\beta 2$ and $\alpha 2\delta$ subunits of the cardiac L-type calcium channel (*CACNA1C*, *CACNB2*, and *CACNA2D1*) and the 1 subunit of Nav1.5 and Nav1.8 (*SCN5A* and *SCN10A*) have been reported in patients with ERS.^{113, 152, 177}

It is important to point out that only a small fraction of identified genetic variants in the genes associated with BrS and ERS have been examined using functional expression studies to ascertain causality and establish a plausible contribution to pathogenesis. Only a handful have been studied in genetically engineered animal models, and very few have been studied in native cardiac cells or in induced pluripotent stem cell-derived cardiac myocytes isolated from ERS and BrS patients. Computational strategies developed to predict the functional consequences of mutations are helpful, but these methods have not been rigorously tested. The lack of functional or biological validation of mutation effects remains the most severe limitation of genetic test interpretation, as recently highlighted by Schwartz et al.¹⁹¹

Recent technological advances have resulted in expansion of disease-specific panels.¹⁹² Large public databases of genetic variation from next-generation sequencing programs such as the 1000 Genomes Project, the National Heart Lung and Blood Institute Grand Opportunity Exome Sequencing Project (GO-ESP), and the Exome Aggregation Consortium (ExAC), have challenged drastically our understanding of the “normal” burden and extent of background genetic variation within cardiac channelopathy-susceptibility genes.¹⁹³⁻¹⁹⁵

Although *SCN5A* variants account for 18%-28% of BrS,¹⁹⁶ *SCN5A* genetic testing is complicated by a ~3%-5% “benign” variant frequency in the general population.¹⁹⁴ Therefore, even in the most common genetic cause of BrS, 1 in 10 “positive” tests could be a “false positive” even if found in an individual with a robust BrS phenotype. To date, there are over 20 J wave syndrome (JWS)-susceptibility genes.^{146, 195, 197} However, these additional genes have only magnified the issues of interpretation by adding to the overall

“genetic noise,” without significantly increasing the true mutation yield.^{178, 198-200} In fact, one study revealed that 1:23 individuals in the GO-ESP population possess a previously published BrS-associated variant that would prompt a “positive” genetic test had it been identified in a patient.²⁰¹

These issues reinforce the necessity to interpret JWS genetic test results as strictly probabilistic, rather than binary/deterministic, in nature. Additional lines of evidence²⁰² can be amassed to aid in the probabilistic interpretation of variants in JWS-susceptibility genes such as case phenotype,²⁰³ segregation, functional studies,²⁰⁴ *in silico* predictions,²⁰⁵⁻²⁰⁸ variant type and location,¹⁹⁴ and variant frequency in cases and control databases.¹⁹³ Despite these aids, a large number of variants remain in “genetic purgatory,” and this number will only increase as the use of exome/genome sequencing becomes more utilized. This then demands the development and utilization of a uniform variant repository that would include clinical assertions and evidence for variant classification. Even with these issues, the emergence of exome/genome sequencing holds promise for the opportunity to study genetic variation like never before, holding the promise of improvements in diagnostic, prognostic, and therapeutics for the JWSs and the other heritable cardiac channelopathies. Kapplinger et al. recently reported the synergistic use of up to 7 *in silico* tools to help promote or demote a variant’s pathogenic status and alter its relegation to genetic purgatory.²⁰⁹

It is noteworthy that in a recent study Le Scouarnec and colleagues estimated the burden of rare coding variation in arrhythmia-susceptibility genes among 167 BrS index patients and compared that with 167 individuals 65 years old and older with no history of cardiac arrhythmia.¹⁹⁹ The authors concluded that, except for *SCN5A*, rare coding variations in all previously reported BrS-susceptibility genes do not contribute significantly to the occurrence of BrS in a population with European ancestry, emphasizing that caution should be taken when interpreting genetic variations in these other genes because rare coding variants are observed to a similar extent in both cases and controls.¹⁹⁹ Similar data were obtained and a similar conclusion was reached by Kapplinger et al., by analyzing the prevalence of rare variants in the BrS-susceptibility genes in the publicly available ExAC exomes.²⁰⁹

Collectively, these data suggest the possibility that, in the individual patient, BrS and the susceptibility to VF and SCD may not be due to a single mutation (classical Mendelian view), but rather to inheritance of multiple BrS-susceptibility variants (oligogenic) acting in concert through one or more mechanistic pathways.¹⁶⁷ This also fits with the findings of Probst et al. that in 5 out of 13 large families with a putative *SCN5A* mutation, the genotype did not co-segregate with the phenotype.²¹⁰ In addition to the multifactorial nature of the genetics, expressivity of the syndrome may be multifactorial in that the genetic predisposition can be modulated by hormonal (testosterone,^{211, 212} thyroxine²¹³) and other environmental factors, as well as morphological changes (fibrosis).⁷⁶

Update on the ionic and cellular mechanisms underlying BrS and ERS

The J wave syndromes are so named because they involve accentuation of the electrocardiographic J wave. Experimental evidence indicates that the J wave is inscribed as a consequence of a transmural voltage gradient caused by the manifestation of an AP notch in epicardium but not endocardium due to a heterogeneous transmural distribution of I_{to} .¹⁰⁴ An end of QRS notch, resembling a J wave, has been proposed to be due to intra-ventricular conduction delays. The two ECG manifestations can be distinguished on the basis of their response to rate, with the latter showing accentuation at faster rates.^{24, 59}

The cellular mechanisms underlying JWS have long been a matter of debate.^{214, 215} In the case of BrS, two principal hypotheses have been advanced: 1) The **repolarization hypothesis** asserts that an outward shift in the balance of currents in right ventricular epicardium can lead to repolarization abnormalities resulting in the development of phase 2 reentry, which generates closely coupled premature beats capable of precipitating VT/VF; and 2) The **depolarization hypothesis** suggests that slow conduction in the RVOT, secondary to fibrosis, reduced Cx43 leading to discontinuities in conduction, plays a primary role in the development of the electrocardiographic and arrhythmic manifestations of the syndrome. Conduction slowing is not necessarily limited to the RVOT area. Some investigators have postulated that changes in ion channel current responsible for BrS (i.e., loss of function I_{Na} and I_{Ca} and gain of function of I_{to}) can alter AP morphology so as to reduce the safety of conduction at high-resistance junctions, such as regions of extensive fibrosis.^{216, 217} Others have argued that this is highly unlikely because conduction at critical junctions of current-to-load mismatch is exquisitely sensitive to changes in rate. The typical behavior of patients with BrS to acceleration of rate is diminution of ST segment elevation, opposite to that expected at a site of discontinuous conduction. The diminution of ST segment elevation is consistent with the reduced availability of I_{to} at the faster rate due to slow recovery of the current from inactivation.^{59, 214} The repolarization and depolarization theories are not necessarily mutually exclusive and may indeed be synergistic.

The most compelling apparent evidence in support of the depolarization hypothesis derives from the seminal studies of Nademanee et al.¹²⁹ showing that radiofrequency ablation (RFA) of epicardial sites displaying late potentials and fractionated bipolar electrograms (EGs) in the RVOT of patients with BrS significantly reduced the arrhythmia-vulnerability as well as the ECG-manifestation of the syndrome. Similar results were reported by Brugada et al.,¹³¹ and in an isolated case by Sacher and co-workers,¹³⁰ who also observed that accentuation of the Brugada ECG by ajmaline was associated with increasing area of low-voltage and fragmented electrogram activity. A wider area of low-voltage activity was associated with a more prominent ST-segment elevation.¹³¹ These authors concluded that the late potential and fractionated electrogram activity are due to conduction delays within the RVOT/RV anterior wall and that ablation of the sites of slow conduction is the basis for the ameliorative effect of ablation therapy.¹²⁹⁻¹³¹ In a direct test of this hypothesis, Szel and Antzelevitch¹⁵⁰ provided evidence for an alternative mechanism using an experimental model of BrS. The low-voltage fractionated electrogram activity was shown to develop due to regional desynchronization in the appearance of the second AP upstroke, secondary to accentuation of the epicardial AP notch, and high-frequency late potentials develop in the RV epicardium

secondary to concealed phase 2 reentry. Delayed conduction of the primary beat was never observed in a wide variety of BrS models created by exposing canine right ventricular wedge preparations to drugs mimicking the different genetic defects known to give rise to BrS.¹⁵⁰ In more recent studies, ablation of the RV epicardium was shown to diminish the manifestation of J waves and ST segment elevation and to abolish all arrhythmic activity by destroying the cells with the most prominent AP notch, thus eliminating the cells responsible for the repolarization abnormalities that give rise to phase 2 reentry and VT/VF.^{15, 218} Confirmation of all of the above results in *in vivo* animal models is desirable. In an attempt to create such a model, Park et al. recently genetically engineered Yucatan minipigs to heterozygously express a nonsense mutation in SCN5A (E558X) originally identified in a child with BrS.¹⁴⁹ Patch clamp analysis of atrial myocytes isolated from the SCN5A^{E558X/+} pigs showed a loss of function of I_{Na} . Conduction abnormalities consisting of prolongation of P wave, QRS complex and PR interval were observed, but a BrS phenotype was not observed, not even after the administration of flecainide. These observations are expected owing to the lack of I_{to} in the pig, which is a prerequisite for the development of the repolarization abnormalities associated with BrS. Some have argued that the absence of a BrS phenotype is due to the young age of the minipigs (22 months).²¹⁹ It is however difficult to reconcile why the minipigs manifest major conduction delays at this age but not a BrS phenotype, if indeed the latter depends on the former. Finally, it is noteworthy that monophasic APs recorded from the epicardial and endocardial surfaces of the RVOT of a patient with BrS are nearly identical to transmembrane APs recorded from the epicardial and endocardial surfaces of the wedge model of BrS.^{220, 221} These differences were not observed in an isolated heart explanted from a BrS patient after transplantation of a new heart. However, the epicardium of this heart was very depressed, perhaps as a result of the 129 shocks delivered by the ICD in an attempt to control the multiple electrical storms.³²

Zhang et al. recently performed noninvasive ECG imaging (ECGI) on 25 BrS and 6 RBBB patients.¹³³ The authors concluded that both slow, discontinuous conduction and steep dispersion of repolarization are present in the right ventricular outflow tract of patients with BrS. ECGI was able to differentiate between BrS and RBBB. Unlike BrS, RBBB showed delayed activation in the entire right ventricle, without ST-segment elevation, fractionation, or repolarization abnormalities showing on the electrograms. Importantly, in 6 BrS patients the response to an increase in rate was studied. Increasing rate increased fractionation of the electrogram but reduced ST-segment elevation, indicating that the conduction impairment was not the principal cause of the BrS ECG.

The congruence between BrS and ERS with respect to clinical manifestations and response to therapy lend further support for the repolarization hypothesis. Using an experimental model of ERS, Koncz et al.³⁰ recently provided evidence in support of the hypothesis that, similar to the mechanism operative in BrS, an accentuation of transmural gradients in the LV wall are responsible for the repolarization abnormalities underlying ERS, giving rise to J point elevation, distinct J waves, or slurring of the terminal part of the QRS. The repolarization defect is accentuated by cholinergic agonists and reduced by quinidine, isoproterenol, cilostazol and milrinone, accounting for the ability of these agents to reverse the repolarization abnormalities responsible for ERS.^{30, 222} Higher intrinsic levels of I_{to} in the inferior LV were also shown to underlie the greater vulnerability of the inferior LV wall

to VT/VF.³⁰ The advent and implementation of electrocardiographic imaging (ECGI) by Rudy and colleagues provided additional evidence for repolarization abnormalities by identifying abnormally short activation-recovery intervals in the inferior and lateral regions of LV and a marked dispersion of repolarization.¹³² More recent studies involving ECGI mapping in an ERS patient during VF have demonstrated VF rotors anchored in the inferior-lateral left ventricular wall.²²

Conduction delays are known to give rise to notching of the QRS complex. When it occurs on the rising phase of the R wave, it is due to a conduction defect within the ventricle. When it occurs at the terminal portion of the QRS, thus masquerading as a J wave, it may be due either to a conduction or repolarization defect.^{21, 223} The response to prematurity or to an increase in rate can differentiate between the two.⁵⁹ Delayed conduction invariably becomes more exaggerated at faster rates or during premature beats, thus leading to an accentuation of the QRS notch, whereas repolarization defects are usually mitigated resulting in a diminution of the J wave at faster rates. Although typical J waves are usually accentuated with bradycardia or long pauses, the opposite has also been described.^{224, 225} J waves are often seen in young males with no apparent structural heart diseases, whereas intra-ventricular conduction delay is often observed in older individuals or those with a history of myocardial infarction or cardiomyopathy.^{223, 224} The prognostic value of a fragmented QRS has been demonstrated in BrS,^{49, 226} although fragmentation of the QRS is not associated with increased risk in the absence of cardiac disease.²²⁷ Factors that may aid in the differential diagnosis of J wave vs. IVCD-mediated syndromes are summarized in **Table 8**.

Risk stratification of the J wave syndromes

A great deal of attention has been devoted to risk assessment for the development of life-threatening arrhythmias in BrS and ERS.^{1, 228} The incidental discovery of a J wave on routine screening should not be interpreted as a marker of “high risk” for SCD since the odds for this fatal disease are approximately 1:10,000.²²⁹ Rosso et al. indicated that the presence of a J wave on the ECG increases the probability of VF from 3.4:100,000 to 11:100,000.^{4, 230} However, careful attention needs to be paid to subjects with “high risk” ER or J waves. **Figure 4** presents a graphic representation of the prevalence and arrhythmic risk associated with the appearance of ECG J waves and clinical manifestations of BrS and ERS. **Tables 9-11** present the available data from studies designed to identify patients at high risk for BrS and ERS. Among these risk stratifiers, some are highly predictive, including: 1) history of cardiac events or syncope likely due to VT/VF, and 2) prominent J waves in global leads including Type 1 ST segment elevation in the right precordial leads (**Figure 5**).

Risk stratification of patients with ERS

The majority of the studies employing the criteria of Haissaguerre et al. for diagnosing the ER pattern² have shown that ER, especially in the inferior ECG leads, predicts cardiac and arrhythmic death. Negative studies are few and may be attributable to exclusion criteria employed (e.g., atrial fibrillation, flutter, acute coronary syndrome), a relatively short follow-up period,^{231, 232} or different definitions of ER pattern.²³³ The recent consensus paper by Macfarlane et al. dealing with terminology of J wave-related phenomena in the

setting of ER should enable us to avoid such confusion in the future.²⁴ The inclusion of Africans or African-Americans, in whom ER is prevalent but apparently not associated with high risk, may alter outcomes as well.²³⁴

Huikuri and colleagues reported in a series of seminal papers the results of a population-based study in Finland involving long-term prognosis of subjects with an ER pattern in the ECG.⁴⁴ Tikkanen et al. showed that J point elevation ≥ 0.1 mV was present in 5.8% of the population and that only 0.3% of the population had significant J point elevation ≥ 0.2 mV. J point elevation ≥ 0.1 mV in the inferior leads was associated with cardiovascular death (relative risk [RR]: 1.28) and arrhythmic death (RR: 1.43), and J point elevation ≥ 0.2 mV had a markedly elevated risk of death from cardiac causes (RR: 2.92) and from arrhythmia (RR: 2.92). Subsequent studies confirmed the association of J wave or ER with death from all causes, death from cardiovascular disease, sudden/unexpected death and death from arrhythmias.^{44-46, 127, 235-238} A horizontal or descending ST segment is associated with a worse prognosis than an ascending ST segment (**Figure 3**).^{237, 239} Individuals with a high amplitude J wave ≥ 0.2 mV followed by a horizontal or descending ST segment in the inferior/infero-lateral leads have a higher risk of lethal arrhythmias than those with a lower amplitude J-wave, especially those with a rapidly ascending ST segment following the J wave.

The appearance of J wave or ER is now recognized to predispose to the development of arrhythmogenesis when associated with other cardiac disorders, such as ischemia, heart failure and hypothermia. The J wave might predict prognosis of cardiac events in various heart diseases, and the appearance of a new J wave during acute ischemia seems to be a messenger of VF.^{240, 241}

Family history of sudden death in subjects with ER pattern has been identified as a risk factor.^{189, 242} The presence of co-existing Brugada ECG pattern (J waves in V1-V3) or short QT intervals in subjects with ER also suggests a more malignant nature.^{243, 244}

ER is commonly observed in the young, especially in fit and highly trained athletes ranging to a prevalence of up to 40%. In the majority of cases, the ensuing ST segment is rapidly ascending, suggesting that this is a benign ECG manifestation.^{237, 245}

Mahida and co-workers recently reported that electrophysiologic study using current programmed stimulation protocols do not enhance risk stratification in ER syndrome.²⁴⁶

Risk stratification of patients with BrS

Numerous studies consistently show that clinical presentation is the strongest predictor of risk in BrS, overshadowing all other risk factors. The risk of recurrent VF among patients presenting with cardiac arrest is considerable: $\approx 35\%$ at 4 years,^{247, 248} 44% at 7 years²⁴⁹ and 48% at 10 years.²⁵⁰ Fortunately, only a minority of BrS (6% in Europe²⁴⁷ but 18% in Japan²⁴⁸) diagnosed nowadays have a history of cardiac arrest.

Approximately one-third of contemporary BrS cohorts present with syncope.²⁴⁷ Their risk of arrhythmic events during followup is intermediate: approximately four times higher than

the risk of asymptomatic patients^{49, 247, 251} but four times lower than that of patients diagnosed post-cardiac arrest.²⁴⁷ One explanation for this observation is that the syncope population consists of two different groups, one with arrhythmic syncope and bad prognosis, and a second group with vagal syncope and good prognosis. Although a detailed clinical history may be of great value in differentiating between these 2 groups, it is not infallible.²⁵² In reviewing the records of 342 BrS patients, Olde Nordkamp et al.⁶⁰ concluded that arrhythmic and nonarrhythmic syncope can be distinguished by clinical characteristics, including the absence of prodromes and specific triggers. Compared to suspected non-arrhythmic syncope patients, patients presenting with presumed arrhythmic syncope were more likely to be male (relative risk 2.1), to have urinary incontinence (relative risk 4.6) and were less likely to report prodromes. They were also older at first event (45 vs. 20 years), and their syncope was never triggered by hot/crowded surroundings, pain or other emotional stress, sight of blood, or prolonged standing as in the case of non-arrhythmic syncope. During follow-up, all the spontaneous arrhythmic events occurred in patients originally presenting with presumed arrhythmic syncope; patients with benign syncope had an excellent long-term prognosis.

Asymptomatic patients represent a majority (~63%) of newly diagnosed Brugada patients today.^{247, 248} Their risk of developing symptoms is relatively low (0.5% per year).^{247, 248} Unfortunately, for most, the first symptom is cardiac arrest or sudden cardiac death. Therefore, risk stratification of asymptomatic patients is of utmost importance, and strategies for doing so are discussed below. In cardiac arrest patients or patients with presumed arrhythmic syncope, these strategies are of little use since they are recognized to be at high risk.

Age and gender

The mean age at the time of cardiac arrest in Brugada patients is 39-48 years, and the vast majority develop symptoms between 20 and 65 years of age.²⁴⁷⁻²⁴⁹ Asymptomatic elderly patients with Brugada syndrome are thought to be at a relatively low risk for future cardiac events.²⁵³ BrS in children is very rare, but sudden death in this population has been described.^{65, 254, 255} As in adults, cardiac arrest survivors are at high risk of recurrence, but data regarding risk stratification of asymptomatic children are limited. In large series, 64%-94% of patients with BrS presenting with cardiac arrest were male.^{247, 248, 250} Males are also at increased risk of displaying a spontaneous Type I Brugada ECG and of having inducible VF during electrophysiologic (EP) studies.¹⁰⁶ Nevertheless, because the majority of asymptomatic patients are also male, gender is not an independent predictor of arrhythmic events.^{106, 248}

Familial and genetic background

Neither family history of SCD nor the presence of a mutation (of any type) in the SCN5A gene have consistently been demonstrated to be of value in risk stratification.²⁴⁷⁻²⁴⁹ One study, however, has shown that SCN5A mutations resulting in protein truncation do confer greater risk.²⁵⁶ Certain rare variants and polymorphisms in SCN5A and in other genes have

also been associated with prognosis.^{167, 257-261} Nevertheless, the data are limited, and genetic testing is not generally used for risk stratification at this time.

Spontaneous vs. drug-induced Type I Brugada pattern

A consistent finding in nearly all BrS series is that patients with spontaneous Type I ECG at time of diagnosis have a greater risk of arrhythmic events than patients who develop such an ECG pattern *only* when challenged with a sodium channel blocker.^{247, 262} This observation is true for asymptomatic patients²⁴⁷ as well as patients presenting with syncope^{49, 247, 251, 263} and remains an independent predictor of arrhythmic events in multivariate analysis. The problem is that only a minority of patients have a consistent spontaneous Type 1 pattern when repeated ECGs are analyzed.^{264, 265} Therefore, caution should be employed when using a single ECG for risk stratification.

Ventricular arrhythmias are rarely induced during a sodium-channel-block challenge^{65, 262, 266, 267} and are likely to be dose-dependent (i.e., will not occur when the infusion is stopped for safety reasons prior to reaching the full dose). Their long-term prognostic significance remains unclear. Extra caution should be exercised when administering sodium channel blockers to patients with significant conduction disease.

Placing the right precordial leads over the 2nd and 3rd intercostal space, in addition to the standard 4th intercostal space, increases the sensitivity for detecting the coved ST-segment elevation of the Type I Brugada pattern. Available data suggest that this increased detection of spontaneous Type I pattern does not affect the value of this parameter for predicting VF.^{52, 268, 269}

Electrophysiologic (EP) studies with programmed ventricular stimulation

The role of programmed ventricular stimulation during EP studies continues to be passionately debated.²⁷⁰⁻²⁷² The question is not whether VF inducibility correlates with arrhythmic risk; in all series, the VF inducibility rate is highest for cardiac-arrest survivors, intermediate for those with syncope and lowest for those asymptomatic at presentation.²⁷³ The central question is whether the prognostic information provided by VF inducibility is robust enough for clinical decision making. Some studies suggest that this is the case,^{262, 274, 275} but others do not.^{248, 249} One central issue suggested as an explanation for the discrepancy between studies is the extrastimulation protocol used. Specifically, the prognostic impact of site (RV apex vs. RV apex + RVOT)^{49, 270, 276} and number of extrastimuli (2 vs. 3)^{49, 276} has been analyzed but with inconsistent results. The only prospective study specifically designed to examine the prognostic yield of EP studies in BrS was the PRELUDE registry.⁴⁹ It did not show that sustained VF induction identifies high-risk patients but did demonstrate that a short, effective ventricular refractory period (< 200 ms) is a risk marker.

Other risk markers

Several electrocardiographic markers have been associated with risk in BrS. These include: a) fragmentation of the QRS,^{226, 277, 278} b) the concomitant finding of a Type 1 Brugada

pattern and an early repolarization pattern in infero-lateral leads^{145, 248, 279-281} and c) dynamic changes in the manifestation of prominent J waves or ST segment elevation.²⁸²

Other markers associated with increased risk but with limited or inconsistent data include: a) late potentials recorded using signal-averaged electrocardiography;²⁸³⁻²⁸⁵ b) microscopic T-wave alternans (TWA);²⁸⁶ c) macroscopic TWA during a sodium-blocker challenge test;^{285, 287, 288} d) increased QRS width;^{49, 248, 277, 289, 290} e) prominent R wave in aVR;²⁸⁹⁻²⁹¹ and f) augmented ST-segment elevation of a Type 1 Brugada pattern during the recovery phase of an exercise test.²⁹² Prolonged Tpeak-Tend²⁹³⁻²⁹⁶ and relatively steep QT/RR slope have been associated with higher risk in cases of BrS.^{297, 298} Combining several risk factors (e.g., fragmented QRS + ER pattern²⁷⁷) appears to confer an additive risk, however, data supporting this are limited.

Update on approaches to therapy of BrS and ERS

Figures 6 and 7 graphically present recommendations for the management of BrS and ERS as modified from the 2013 HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes and the 2015 ESC Guidelines for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death.^{8, 9} Those recommendations are based on the available literature and on the clinical experience of members of the task force. As with all such recommendations they will need to undergo continuous validation by future studies.

Education and lifestyle changes for the prevention of arrhythmias are critical in BrS. Patients should be informed of the various modulators and precipitating factors that could cause malignant arrhythmias. Fever should be treated aggressively with antipyretics, and contraindicated substances should be avoided (www.brugadadrugs.org).⁸⁹ Referral for ECG is recommended during high fever. Family members may be referred for CPR training and advised to consider purchase of an AED for home use. Because malignant ventricular arrhythmias are infrequent in asymptomatic patients with BrS²⁴⁷ or ER pattern⁴⁴ and usually unrelated to physical activity, the presence of these patterns does not contraindicate participation in sports.

It is noteworthy, however, that the Brugada pattern is accentuated immediately after exercise, due presumably to an increase in vagal tone.^{292, 299, 300} In reviewing 98 case of BrS studies dealing with exercise, Masrur et al. concluded that there are insufficient data on the risks of exercise in BrS to make recommendations for exercise.²⁹⁹

Implantable Cardioverter Defibrillator

The only proven effective therapeutic strategy for the prevention of SCD in high risk BrS and ERS patients is an implantable cardioverter defibrillator (ICD).^{301, 302} It is important to recognize that ICDs are associated with complications, especially in young active individuals.^{249, 303} At 10 years post-implant, the rate of inappropriate shock and lead failure are 37% and 29%, respectively. Remote monitoring can identify lead failure and avoid inappropriate shocks.³⁰⁴ Sub-cutaneous ICDs are thought to represent the future for this

indication in that they are expected to be associated with fewer complications over a lifetime.³⁰⁵

Implantation of an ICD is first-line therapy for JWS patients presenting with aborted SCD or documented VT/VF with or without syncope (Class I recommendation).^{301, 306} ICDs can be useful (Class IIa) in symptomatic BrS patients with Type 1 pattern, in whom syncope was likely caused by VT/VF. The HRS/EHRA/APHRS Expert Consensus states that ICD may be considered (Class IIb) in asymptomatic patients with inducible VF during programmed electrical stimulation (PES).⁸ Some studies suggest that the predictive value of EP studies may be improved by limiting the PES protocol to 2 extrastimuli,^{50, 276} but that observation is not supported by other studies.^{49, 307} Similarly, some studies advocate that PES should be limited to the RVA and credit this limited PES strategy for a very high positive predictive value found in some series.²⁷⁵ Again, that observation is not confirmed by other studies.²⁷⁶

The current Task Force proposes that ICDs are reasonable (Class IIa) in symptomatic BrS patients with Type 1 pattern, but that implantation be considered on a case-by-case basis by an electrophysiologist experienced in BrS, taking into consideration age, gender, clinical presentation, ECG characteristics (QRS fragmentation, Jp amplitude) and patient's preference. The current Task Force also proposes that electrophysiologic study may be considered in asymptomatic individuals with spontaneous Type 1 Brugada pattern. If VT/VF is inducible, an ICD should be considered.⁷ More recent studies argue in favor of using two or less extrastimuli to induce VT/VF.^{50, 276} ICDs are not indicated in asymptomatic patients without any of the above characteristics. At present, there is no clear role for PES in patients with ERS.

Pacemaker therapy

Arrhythmic events and sudden cardiac death in both BrS and ERS generally occur during sleep or at rest and are associated with slow heart rates. These observations notwithstanding, a potential therapeutic role for cardiac pacing remains largely unexplored.³⁰⁸ A few case reports are available.^{309, 310}

Radiofrequency Ablation Therapy

Nademanee et al.¹²⁹ showed that RFA of **epicardial** sites displaying late potentials and fractionated bipolar electrograms (EGs) in the RVOT of BrS patients can significantly reduce arrhythmia vulnerability and the ECG manifestation of the disease. Ablation at these sites was reported to render VT/VF non-inducible and to normalize the Brugada ECG pattern in the vast majority of patients over a period of weeks or months. Long-term follow-up (20±6months) showed no recurrent VT/VF, with only 1 patient on medical therapy with amiodarone. Case reports have been published in support of these effects.³¹¹ Additional evidence in support of the effectiveness of epicardial substrate ablation was provided by Sacher et al. and Shah et al.^{130, 312}

More recently, Brugada and co-workers used flecainide to identify the full extent of low voltage electrogram activity in the anterior RV and RVOT and targeted this region for RFA. In all, 14 BrS patients, RFA eliminated abnormal bipolar electrograms, normalized ST

segment elevation on right precordial leads of ECG, and VT/VF was no longer inducible.¹³¹ Ablation therapy can be life-saving in otherwise uncontrollable cases. RF ablation may be considered (**Class IIb** recommendation) in BrS patients with frequent appropriate ICD-shocks due to recurrent electrical storms.⁷ There are no clinical reports of ablation of the LV substrate in patients with ERS. In patients in whom BrS combines with ERS, ablation of the anterior RV epicardium (including the RVOT) is not ameliorative.

Pharmacologic approach to therapy

Pharmacologic Approach to therapy of BrS

ICD implantation may be problematic in infants or young children due to the high complication rate. ICDs are also economically out of reach for patients in some regions of the world. A pharmacologic approach to therapy, based on a rebalancing of currents active during the early phases of the epicardial AP in the right ventricle so as to reduce the magnitude of the AP notch and/or restore the AP dome, has been a focus of basic and clinical research in recent years. Antiarrhythmic agents such as amiodarone and β blockers have been shown to be ineffective.³¹³ Class IC antiarrhythmic drugs (such as flecainide and propafenone) and class IA agents, such as procainamide, are contraindicated because of their effects to unmask BrS and induce arrhythmogenesis. Disopyramide is a class IA antiarrhythmic that has been demonstrated to normalize ST segment elevation in some Brugada patients but to unmask the syndrome in others.³¹⁴

Because the presence of a prominent I_{to} is a prerequisite for the development of both BrS and ERS, partial inhibition of this current is thought to be effective regardless of the ionic or genetic basis for the disease. Unfortunately, cardio-selective and I_{to} specific blockers are not available.

The only agent available in the United States and around the world with significant I_{to} blocking properties is quinidine.^{19, 73} Experimental studies have shown that quinidine is effective in restoring the epicardial AP dome, thus normalizing the ST segment and preventing phase 2 reentry and polymorphic VT in a variety of different experimental models of BrS.^{19, 150, 315-317} A recent experimental study suggests that quinidine, owing to its effect to block I_{to} , can also exert a protective effect against hypothermia-induced VT/VF in a J wave syndrome model.¹⁴⁴ It is noteworthy that, historically, quinidine was used to prevent ventricular fibrillation in patients who required hypothermia for surgical procedures.³¹⁷

Clinical evidence for the effectiveness of quinidine in normalizing ST segment elevation and/or preventing arrhythmic events in patients with BrS has been reported in numerous studies and case reports.^{117, 119, 120, 124, 318-331} Hermida et al. reported 76% efficacy in prevention of VF induced by PES.¹¹⁹ Belhassen and colleagues³³² recently reported a 90% efficacy in prevention of VF induction following treatment with quinidine, despite the use of very aggressive protocols of extrastimulation. Furthermore, there were no arrhythmic events among BrS patients treated with quinidine during a mean follow-up period of 10 years.

In a recent trial conducted at two French centers, 44 asymptomatic BrS patients with inducible VT/VF were enrolled (47 ± 10 years, 95% male).³³³ Of these, 34 (77%) were no longer inducible while treated with 600 mg/day hydroquinidine (HQ) for 6.2 ± 3 years. Among the 10 other patients (22%), who remained inducible and received ICD (Group PVS+), none received appropriate therapy during a mean followup of 7.7 ± 2 years.

A prospective registry of empiric quinidine for asymptomatic BrS has been established. The study appears at the National Institutes of Health website (ClinicalTrials.gov) and can be accessed at http://clinicaltrials.gov/ct2/show/NCT00789165?term_brugada&rank_2. Doses between 600 and 900 mg were recommended, if tolerated.³²²

Quinidine may be considered (Class IIb indication) in BrS patients presenting with electrical storms and in patients implanted with an ICD and experiencing repeated appropriate shocks. Quinidine can also be useful in asymptomatic BrS patients displaying a spontaneous Type I ECG, if they qualify for an ICD and the device is refused or is contraindicated (Class IIa recommendation).

Agents that augment the L-type calcium channel current, such as β adrenergic agents like isoproterenol, denopamine or orciprenaline, are useful as well.^{19, 117, 121, 327, 334, 335}

Isoproterenol, at times in combination with quinidine, has been utilized successfully to control VF storms and in normalizing ST elevation, particularly in children.^{116, 319, 320, 331, 336, 337, 93, 115, 117, 118, 279, 325, 338-343} Spontaneous VF in patients with BrS is often related to increases in vagal tone and is amenable to treatment by an increase of sympathetic tone via isoproterenol administration. Administration of isoproterenol is a Class IIa recommendation for BrS patients presenting with electrical storms.⁷

Another promising pharmacologic approach for BrS is the administration of phosphodiesterase III inhibitor cilostazol,^{117, 121, 123} which normalizes the ST segment, most likely by augmenting calcium current (I_{Ca}) as well as by reducing I_{to} secondary to an increase in cAMP and heart rate.³⁴⁴ Other effects of cilostazol may contribute to its actions (e.g., adenosine, NO, mitochondrial I_{KATP}).³⁴⁵ Its efficacy in combination with bepridil in preventing VF episodes was recently reported by Shinohara et al.¹²⁵ The failure of cilostazol in the treatment of BrS has been described in a single case report.³⁴⁶

Milrinone is another phosphodiesterase III inhibitor recently identified as a more potent alternative to cilostazol in suppressing ST elevation and arrhythmogenesis in an experimental model of BrS.^{150, 347} No clinical reports have appeared as yet.

Wenxin Keli, a traditional Chinese medicine has recently been shown to inhibit I_{to} and thus to suppress polymorphic VT in experimental models of BrS when combined with low concentrations of quinidine (5 μ M).³¹⁶

Agents that augment peak and late I_{Na} , including bepridil and dimethyl lithospermate B (dmLSB), are suggested to be of value in BrS. Bepridil has been reported to suppress VT/VF in several studies of patients with BrS.^{117, 297, 298, 348} The drug's action are thought to be mediated by: 1) inhibition of I_{to} ; 2) augmentation of I_{Na} via up-regulation of the sodium channels³⁴⁹; and 3) prolongation of QT interval at slow rates thus increasing the QT/RR

slope.^{297, 298} Dimethyl lithospermate B, an extract of Danshen, a traditional Chinese herbal remedy, has been reported to slow inactivation of I_{Na} , thus increasing I_{Na} during the early phases of the AP and suppressing arrhythmogenesis in experimental models of BrS.³⁵⁰

Because malignant ventricular arrhythmias are infrequent in asymptomatic patients with BrS²⁴⁷ or ER pattern⁴⁴ and usually unrelated to physical activity, the presence of these patterns does not contraindicate participation in sports, although as previously discussed, insufficient data are currently available to make definitive recommendations for participation in sports.

Approach to therapy of ERS

It is not surprising that the approach to therapy of ERS is similar to that of BrS, since the mechanisms underlying the two syndromes are potentially similar. Quinidine, phosphodiesterase III inhibitors and isoproterenol have all been shown to exert an ameliorative effect in preventing or quieting arrhythmias associated with ERS. Isoproterenol has been shown to be effective in quieting electrical storms developing in patients with either BrS^{117, 338} or ERS.¹⁹⁰ Isoproterenol has been shown to act by reversing the repolarization abnormalities responsible for the disease phenotype secondary to restoration of the epicardial AP dome in experimental models of both BrS^{19, 315} and ERS.³⁰ This action of the β adrenergic agonist is expected due to its actions to potently increase I_{Ca} .

The phosphodiesterase (PDE) III inhibitor cilostazol has been reported to reduce the ECG and arrhythmic manifestations of ERS.¹²² PDE inhibitors are known to activate I_{Ca} secondary to an increase in cAMP.^{121, 344, 351-355} The augmentation of I_{Ca} is thought to prevent arrhythmias associated with JWS by reversing the repolarization defects and restoring electrical homogeneity across the ventricular wall secondary to restoration of the epicardial AP dome in both BrS³⁴⁷ and ERS.¹⁴⁴ Cilostazol has been hypothesized to also block I_{to} . Augmentation of I_{Ca} together with inhibition of I_{to} are expected to produce an inward shift in the balance of currents active during the early phases of the epicardial AP that should be especially effective in suppressing J wave activity. The effectiveness of bepridil in ERS has been reported in a single patient thus far.³⁵⁶

No clinical data are available regarding the effectiveness of radiofrequency (RF) ablation in the setting of ERS, despite the fact that low-voltage fractionated electrogram activity and high-frequency late potentials are observed in the LV in patients with ERS³⁵⁷ and in experimental models of ERS (Yoon and Antzelevitch, unpublished observation). Nakagawa et al.³⁵⁷ reported the results of a study in which they recorded epicardial electrograms directly from the left ventricle of patients diagnosed with ERS by introducing a multipolar catheter into the left lateral (marginal) coronary vein, anterior interventricular vein, and middle cardiac vein (MCV) via the coronary sinus. The authors reported late potentials in the bipolar electrograms recorded from the LV epicardium of the ERS patients.³⁵⁷

Acknowledgments

We are grateful to Buchang Pharmaceutical for providing the funds to host the J wave Consensus Conference. We are also grateful to Dr. Mina Chung for helpful discussion and advice and to Ms. Donna Loyle for proofing the manuscript.

Funding: The consensus conference was organized with the assistance of the Chinese Heart Rhythm Society and funded by an unrestricted educational grant from Buchang Pharmaceuticals. We acknowledge support by grants from NHLBI (HL47678, CA), Sharpe-Strumia Research Foundation (GXY), National Natural Science Foundation of China (NSFC-81370289, GXY), the Mayo Clinic Windland Smith Rice Comprehensive Sudden Cardiac Death Program (MJA), Ministry of Health, Labor and Welfare of Japan for Clinical Research on Intractable Diseases; H24-033, H26-040, H27-032 (MH, HM, WS); Finnish Academy of Science and Sigrid Juselius Foundation, Helsinki, Finland (HH), JSPS KAKENHI (24591051 and 15K09082) (HM), Ministry of Science and Technology of China (2013BAI09B02, 2013DFB30310, CM), University of Padua, Italy (TRANSAC, DC); Netherlands CardioVascular Research Initiative: the Dutch Heart Foundation, Dutch Federation of University Medical Centres, the Netherlands Organization for Health Research and Development and the Royal Netherlands Academy of Sciences (AW). Investment of the Future grant, ANR-10-IAHU-04, Agence National de la Recherche, French Government (FS).

Disclosures: Dr. Antzelevitch was a consultant for Gilead Sciences and has received grant funding and speaker fees from Gilead Sciences and Buchang Pharma. Dr. Yan is a consultant for GlaxoSmithKline and has received funding from GlaxoSmithKline, Johnson&Johnson, Novartis, Astellas and Pfizer. Dr Sacher is consultant to St. Jude Medical, Boston Scientific, and Bayer Healthcare, and has received speaking honorarium from Medtronic, Biosense Webster, Sorin Group, Boehringer Ingelheim, and Buchang Pharma. Dr. Viskin is member of the European Scientific Advisory Board of Boston Scientific. Dr. Huikuri received funding from Medtronic Bakken Research Center. Dr Morita is affiliated with a department endowed by Japan Medtronic Inc.; Dr. Wilde is a member of the scientific advisory board of Sorin. Dr. Ackerman is a consultant for Boston Scientific, Gilead Sciences, Medtronic, and St. Jude Medical and has received royalties from Transgenomic Inc.

References

1. Brugada P, Brugada J. Right bundle branch block, persistent ST segment elevation and sudden cardiac death: a distinct clinical and electrocardiographic syndrome: a multicenter report. *JAmCollCardiol*. 1992; 20:1391–1396.
2. Haissaguerre M, Derval N, Sacher F, et al. Sudden cardiac arrest associated with early repolarization. *NEnglJ Med*. 2008; 358:2016–2023.
3. Nam GB, Kim YH, Antzelevitch C. Augmentation of J waves and electrical storms in patients with early repolarization. *NEnglJ Med*. 2008; 358:2078–2079.
4. Rosso R, Kogan E, Belhassen B, Rozovski U, Scheinman MM, Zeltser D, Halkin A, Steinvil A, Heller K, Glikson M, Katz A, Viskin S. J-point elevation in survivors of primary ventricular fibrillation and matched control subjects: incidence and clinical significance. *J Am Coll Cardiol*. 2008; 52:1231–1238. [PubMed: 18926326]
5. Wilde AA, Antzelevitch C, Borggrefe M, Brugada J, Brugada R, Brugada P, Corrado D, Hauer RN, Kass RS, Nademanee K, Priori SG, Towbin JA. Proposed diagnostic criteria for the Brugada syndrome: consensus report. *Circulation*. 2002; 106:2514–2519. [PubMed: 12417552]
6. Antzelevitch C, Brugada P, Borggrefe M, et al. Brugada syndrome: report of the second consensus conference: endorsed by the Heart Rhythm Society and the European Heart Rhythm Association. *Circulation*. 2005; 111:659–670. [PubMed: 15655131]
7. Priori SG, Wilde AA, Horie M, et al. Executive Summary: HRS/EHRA/APHRS Expert Consensus Statement on the Diagnosis and Management of Patients with Inherited Primary Arrhythmia Syndromes. *Heart Rhythm*. 2013; 15:1389–1406.
8. Priori SG, Wilde AA, Horie M, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes: document endorsed by HRS, EHRA, and APHRS in May 2013 and by ACCF, AHA, PACES, and AEPC in June 2013. *Heart Rhythm*. 2013; 10:1932–1963. [PubMed: 24011539]
9. Priori SG, Blomstrom-Lundqvist C, Mazzanti A, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: The Task Force for the Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of Cardiology (ESC) Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC). *Eur Heart J*. 2015
10. Clements SD, Hurst JW. Diagnostic value of ECG abnormalities observed in subjects accidentally exposed to cold. *AmJCardiol*. 1972; 29:729–734.
11. Thompson R, Rich J, Chmelik F, Nelson WL. Evolutionary changes in the electrocardiogram of severe progressive hypothermia. *JElectrocardiol*. 1977; 10:67–70. [PubMed: 833527]

12. Eagle K. Images in clinical medicine. Osborn waves of hypothermia. *NEnglJMed*. 1994; 10:680.
13. Kraus F. Ueber die wirkung des kalziums auf den kreislauf 1). *DtschMedWochenschr*. 1920; 46:201–203.
14. Sridharan MR, Horan LG. Electrocardiographic J wave of hypercalcemia. *AmJCardiol*. 1984; 54:672–673.
15. Antzelevitch C, Yan GX. J wave syndromes. *Heart Rhythm*. 2010; 7:549–558. [PubMed: 20153265]
16. Wasserburger RH, Alt WJ. The normal RS-T segment elevation variant. *Am J Cardiol*. 1961; 8:184–192. [PubMed: 13783301]
17. Mehta MC, Jain AC. Early repolarization on scalar electrocardiogram. *Am J MedSci*. 1995; 309:305–311.
18. Gussak I, Antzelevitch C. Early repolarization syndrome: clinical characteristics and possible cellular and ionic mechanisms. *JElectrocardiol*. 2000; 33:299–309. [PubMed: 11099355]
19. Yan GX, Antzelevitch C. Cellular basis for the Brugada syndrome and other mechanisms of arrhythmogenesis associated with ST segment elevation. *Circulation*. 1999; 100:1660–1666. [PubMed: 10517739]
20. Shu J, Zhu T, Yang L, Cui C, Yan GX. ST-segment elevation in the early repolarization syndrome, idiopathic ventricular fibrillation, and the Brugada syndrome: cellular and clinical linkage. *J Electrocardiol*. 2005; 38:26–32. [PubMed: 16226071]
21. Antzelevitch C. J wave syndromes: molecular and cellular mechanisms. *J Electrocardiol*. 2013; 46:510–518. [PubMed: 24011992]
22. Mahida S, Derval N, Sacher F, et al. History and clinical significance of early repolarization syndrome. *Heart Rhythm*. 2015; 12:242–249. [PubMed: 25257090]
23. Wellens HJ, Schwartz PJ, Lindemans FW, et al. Risk stratification for sudden cardiac death: current status and challenges for the future. *EurHeart J*. 2014; 35:1642–1651.
24. Macfarlane P, Antzelevitch C, Haissaguerre M, Sacher F, Tikkanen J, Wellens H, Yan G-X. Consensus Paper-Early Repolarization Pattern. *J Amer Coll Cardiol*. 2015 Huikuri HPMRR. In Press.
25. Kanter RJ, Pfeiffer R, Hu D, Barajas-Martinez H, Carboni MP, Antzelevitch C. Brugada-like syndrome in infancy presenting with rapid ventricular tachycardia and intraventricular conduction delay. *Circulation*. 2012; 125:14–22. [PubMed: 22090166]
26. Antzelevitch C. Molecular biology and cellular mechanisms of brugada and long QT syndromes in infants and young children. *JElectrocardiol*. 2001; 34:177–181. [PubMed: 11781953]
27. Wedekind H, Smits JP, Schulze-Bahr E, et al. De novo mutation in the *SCN5A* gene associated with early onset of sudden infant death. *Circulation*. 2001; 104:1158–1164. [PubMed: 11535573]
28. Nagase S, Kusano KF, Morita H, Fujimoto Y, Kakishita M, Nakamura K, Emori T, Matsubara H, Ohe T. Epicardial electrogram of the right ventricular outflow tract in patients with the Brugada syndrome: using the epicardial lead. *J Am Coll Cardiol*. 2002; 39:1992–1995. [PubMed: 12084598]
29. Nakagawa K, Nagase S, Morita H, Ito H. Left ventricular epicardial electrogram recordings in idiopathic ventricular fibrillation with inferior and lateral early repolarization. *Heart Rhythm*. 2014; 11:314–317. [PubMed: 24184784]
30. Koncz I, Gurabi Z, Patocskaï B, Panama BK, Szel T, Hu D, Barajas-Martinez H, Antzelevitch C. Mechanisms underlying the development of the electrocardiographic and arrhythmic manifestations of early repolarization syndrome. *J Mol Cell Cardiol*. 2014; 68C:20–28. [PubMed: 24378566]
31. Antzelevitch C. Brugada syndrome. *Pacing ClinElectrophysiol*. 2006; 29:1130–1159.
32. Coronel R, Casini S, Koopmann TT, et al. Right ventricular fibrosis and conduction delay in a patient with clinical signs of Brugada syndrome: a combined electrophysiological, genetic, histopathologic, and computational study. *Circulation*. 2005; 112:2769–2777. [PubMed: 16267250]
33. Bastiaenen R, Hedley PL, Christiansen M, Behr ER. Therapeutic hypothermia and ventricular fibrillation storm in early repolarization syndrome. *Heart Rhythm*. 2010; 7:832–834. [PubMed: 20206297]

34. Federman NJ, Mechulan A, Klein GJ, Krahn AD. Ventricular fibrillation induced by spontaneous hypothermia in a patient with early repolarization syndrome. *J Cardiovasc Electrophysiol.* 2013; 24:586–588. [PubMed: 23140469]
35. Sakabe M, Fujiki A, Tani M, Nishida K, Mizumaki K, Inoue H. Proportion and prognosis of healthy people with coved or saddle-back type ST segment elevation in the right precordial leads during 10 years follow-up. *EurHeart J.* 2003; 24:1488–1493.
36. Tsuji H, Sato T, Morisaki K, Iwasaka T. Prognosis of subjects with Brugada-type electrocardiogram in a population of middle-aged Japanese diagnosed during a health examination. *Am J Cardiol.* 2008; 102:584–587. [PubMed: 18721516]
37. Gervacio-Domingo G, Isidro J, Tirona J, Gabriel E, David G, Amarillo ML, Morales D, Dans A. The Brugada type 1 electrocardiographic pattern is common among Filipinos. *J Clin Epidemiol.* 2008; 61:1067–1072. [PubMed: 18562166]
38. Ito H, Chen R, He Q, Curb JD. The prevalence and prognosis of a Brugada-type electrocardiogram in a population of middle-aged Japanese-American men with follow-up of three decades. *Am J Med Sci.* 2006; 331:4. YK. [PubMed: 16415656]
39. Letsas KP, Weber R, Astheimer K, Kalusche D, Arentz T. Tpeak-Tend interval and Tpeak-Tend/QT ratio as markers of ventricular tachycardia inducibility in subjects with Brugada ECG phenotype. *Europace.* 2010; 12:271–274. [PubMed: 19897501]
40. Gallagher MM, Forleo GB, Behr ER, Magliano G, De LL, Morgia V, De LF, Romeo F. Prevalence and significance of Brugada-type ECG in 12,012 apparently healthy European subjects. *IntJ Cardiol.* 2008; 130:44–48. [PubMed: 18054807]
41. Pecini R, Cedergreen P, Theilade S, Haunso S, Theilade J, Jensen GB. The prevalence and relevance of the Brugada-type electrocardiogram in the Danish general population: data from the Copenhagen City Heart Study. *Europace.* 2010; 12:982–986. [PubMed: 20356912]
42. Patel SS, Anees SS, Ferrick KJ. Prevalence of a Brugada pattern electrocardiogram in an urban population in the United States. *Pacing Clin Electrophysiol.* 2009; 32:704–708. [PubMed: 19545330]
43. Lee C, Soni A, Tate RB, Cuddy TE. The incidence and prognosis of Brugada electrocardiographic pattern in the Manitoba Follow-Up Study. *Can J Cardiol.* 2005; 21:1286–1290. [PubMed: 16341298]
44. Tikkanen JT, Anttonen O, Junttila MJ, Aro AL, Kerola T, Rissanen HA, Reunanen A, Huikuri HV. Long-term outcome associated with early repolarization on electrocardiography. *NEnglJ Med.* 2009; 361:2529–2537.
45. Sinner MF, Reinhard W, Muller M, et al. Association of early repolarization pattern on ecg with risk of cardiac and all-cause mortality: a population-based prospective cohort study (MONICA/KORA). *PLoS Med.* 2010; 7:e1000314. [PubMed: 20668657]
46. Haruta D, Matsuo K, Tsuneto A, Ichimaru S, Hida A, Sera N, Imaizumi M, Nakashima E, Maemura K, Akahoshi M. Incidence and prognostic value of early repolarization pattern in the 12-lead electrocardiogram. *Circulation.* 2011; 123:2931–2937. [PubMed: 21646495]
47. Hayashi M, Shimizu W, Albert CM. The spectrum of epidemiology underlying sudden cardiac death. *Circ Res.* 2015; 116:1887–1906. [PubMed: 26044246]
48. Brosnan MJ, Kumar S, LaGerche A, Brown A, Stewart S, Kalman JM, Prior DL. Early repolarization patterns associated with increased arrhythmic risk are common in young non-Caucasian Australian males and not influenced by athletic status. *Heart Rhythm.* 2015; 12:1576–1583. [PubMed: 25839111]
49. Priori SG, Gasparini M, Napolitano C, Della Bella P, Ottonelli AG, Sassone B, Giordano U, Pappone C, Mascioli G, Rossetti G, De Nardis R, Colombo M. Risk stratification in Brugada syndrome: results of the PRELUDE (PRogrammed ELectrical stimulation preDICTive value) registry. *Journal of the American College of Cardiology.* 2012; 59:37–45. [PubMed: 22192666]
50. Sroubek J, Probst V, Mazzanti A, et al. Programmed Ventricular Stimulation for Risk Stratification in the Brugada Syndrome: A Pooled Analysis. *Circulation.* 2016
51. Sangwatanaroj S, Prechawat S, Sunsaneewitayakul B, Sitthisook S, Tosukhowong P, Tungsanga K. Right ventricular electrocardiographic leads for detection of Brugada syndrome in sudden unexplained death syndrome survivors and their relatives. *ClinCardiol.* 2001; 24:776–781.

52. Miyamoto K, Yokokawa M, Tanaka K, Nagai T, Okamura H, Noda T, Satomi K, Suyama K, Kurita T, Aihara N, Kamakura S, Shimizu W. Diagnostic and prognostic value of a type 1 Brugada electrocardiogram at higher (third or second) V1 to V2 recording in men with Brugada syndrome. *Am J Cardiol.* 2007; 99:53–57. [PubMed: 17196462]
53. Nagase S, Hiramatsu S, Morita H, Nishii N, Murakami M, Nakamura K, Kusano KF, Ito H, Ohe T. Electroanatomical correlation of repolarization abnormalities in Brugada syndrome: detection of type 1 electrocardiogram in the right ventricular outflow tract. *J Am Coll Cardiol.* 2010; 56:2143–2145. [PubMed: 21144977]
54. Veltmann C, Papavassiliu T, Konrad T, Doesch C, Kuschyk J, Streitner F, Haghi D, Michaeli HJ, Schoenberg SO, Borggrefe M, Wolpert C, Schimpf R. Insights into the location of type I ECG in patients with Brugada syndrome: correlation of ECG and cardiovascular magnetic resonance imaging. *Heart Rhythm.* 2012; 9:414–421. [PubMed: 22119454]
55. Richter S, Sarkozy A, Paparella G, Henkens S, Boussy T, Chierchia GB, Brugada R, Brugada J, Brugada P. Number of electrocardiogram leads displaying the diagnostic coved-type pattern in Brugada syndrome: a diagnostic consensus criterion to be revised. *Eur Heart J.* 2010; 31:1357–1364.
56. Veerakul G, Nademanee K. Brugada syndrome: two decades of progress. *Circ J.* 2012; 76:2713–2722. [PubMed: 23149437]
57. Brugada P, Brugada J, Roy D. Brugada syndrome 1992-2012: 20 years of scientific excitement, and more. *Eur Heart J.* 2013; 34:3610–3615. [PubMed: 24186898]
58. Huikuri HV, Juhani Junttila M. Clinical aspects of inherited J-wave syndromes. *Trends Cardiovasc Med.* 2015; 25:24–30. [PubMed: 25442737]
59. Antzelevitch C, Yan GX. J-wave syndromes: Brugada and early repolarization syndromes. *Heart Rhythm.* 2015; 12:1852–1866. [PubMed: 25869754]
60. Olde Nordkamp LR, Vink AS, Wilde AA, de Lange FJ, de Jong JS, Wieling W, van Dijk N, Tan HL. Syncope in Brugada syndrome: prevalence, clinical significance, and clues from history taking to distinguish arrhythmic from nonarrhythmic causes. *Heart Rhythm.* 2015; 12:367–375. [PubMed: 25311410]
61. Ikeda T, Abe A, Yusa S, Nakamura K, Ishiguro H, Mera H, Yotsukura M, Yoshino H. The full stomach test as a novel diagnostic technique for identifying patients at risk for Brugada Syndrome. *J Cardiovasc Electrophysiol.* 2006; 17:602–607. [PubMed: 16836706]
62. Shimeno K, Takagi M, Maeda K, Tatsumi H, Doi A, Yoshiyama M. Usefulness of multichannel Holter ECG recording in the third intercostal space for detecting type 1 Brugada ECG: comparison with repeated 12-lead ECGs. *J Cardiovasc Electrophysiol.* 2009; 20:1026–1031. [PubMed: 19470036]
63. Viskin S, Rosso R, Friedensohn L, Havakuk O, Wilde AA. Everybody has Brugada syndrome until proven otherwise? *Heart Rhythm.* 2015; 12:1595–1598. [PubMed: 25998201]
64. Conte G, de Asmundis C, Cicconte G, Julia J, Sieira J, Chierchia GB, Brugada P. Follow-up from childhood to adulthood of individuals with family history of Brugada syndrome and normal electrocardiograms. *JAMA.* 2014; 312:2039–2041. [PubMed: 25399282]
65. Conte G, Dewals W, Sieira J, et al. Drug-induced Brugada syndrome in children: clinical features, device-based management and long-term follow-up. *J Am Coll Cardiol.* 2014; 63:2272–2279. [PubMed: 24681144]
66. Gandjbakhch E, Fressart V, Duthoit G, Marquie C, Deharo JC, Pousset F, Hebert JL, Simon F, Himbert C, Klug D, Charron P, Hidden-Lucet F. Malignant response to ajmaline challenge in SCN5A mutation carriers: experience from a large familial study. *Int J Cardiol.* 2014; 172:256–258. [PubMed: 24476701]
67. Kligfield P, Gettes LS, Bailey JJ, et al. Recommendations for the standardization and interpretation of the electrocardiogram: part I: the electrocardiogram and its technology a scientific statement from the American Heart Association Electrocardiography and Arrhythmias Committee, Council on Clinical Cardiology; the American College of Cardiology Foundation; and the Heart Rhythm Society endorsed by the International Society for Computerized Electrocardiology. *Journal of the American College of Cardiology.* 2007; 49:1109–1127. [PubMed: 17349896]

68. Papavassiliu T, Wolpert C, Fluchter S, Schimpf R, Neff W, Haase KK, Duber C, Borggrefe M. Magnetic resonance imaging findings in patients with Brugada syndrome. *J CardiovascElectrophysiol*. 2004; 15:1133–1138.
69. Papavassiliu T, Veltmann C, Doesch C, Haghi D, Germans T, Schoenberg SO, van Rossum AC, Schimpf R, Brade J, Wolpert C, Borggrefe M. Spontaneous type 1 electrocardiographic pattern is associated with cardiovascular magnetic resonance imaging changes in Brugada syndrome. *Heart Rhythm*. 2010; 7:1790–1796. [PubMed: 20828633]
70. Takagi M, Aihara N, Kuribayashi S, Taguchi A, Kurita T, Suyama K, Kamakura S, Takamiya M. Abnormal response to sodium channel blockers in patients with Brugada syndrome: augmented localised wall motion abnormalities in the right ventricular outflow tract region detected by electron beam computed tomography. *Heart*. 2003; 89:169–174. [PubMed: 12527670]
71. Catalano O, Antonaci S, Moro G, et al. Magnetic resonance investigations in Brugada syndrome reveal unexpectedly high rate of structural abnormalities. *EurHeart J*. 2009; 30:2241–2248.
72. van Hoorn F, Campian ME, Spijkerboer A, Blom MT, Planken RN, van Rossum AC, de Bakker JM, Wilde AA, Groenink M, Tan HL. SCN5A mutations in Brugada syndrome are associated with increased cardiac dimensions and reduced contractility. *PLoS One*. 2012; 7:e42037. [PubMed: 22876298]
73. Antzelevitch, C.; Brugada, P.; Brugada, J.; Brugada, R.; Nademanee, K.; Towbin, JA. The Brugada syndrome. Vol. 10. Futura Publishing Company, Inc.; Armonk, NY: 1999. Clinical approaches to tachyarrhythmias.
74. Antzelevitch C. Brugada syndrome: historical perspectives and observations. *EurHeart J*. 2002; 23:676–678.
75. Slezak J, Tribulova N, Okruhlicova L, Dhingra R, Bajaj A, Freed D, Singal P. Hibernating myocardium: pathophysiology, diagnosis, and treatment. *Can J Physiol Pharmacol*. 2009; 87:252–265. [PubMed: 19370079]
76. Nademanee K, Raju H, de Noronha SV, et al. Fibrosis, Connexin-43, and Conduction Abnormalities in the Brugada Syndrome. *J Am Coll Cardiol*. 2015; 66:1976–1986. [PubMed: 26516000]
77. Corrado D, Thiene G. Arrhythmogenic right ventricular cardiomyopathy/dysplasia: clinical impact of molecular genetic studies. *Circulation*. 2006; 113:1634–1637. [PubMed: 16585401]
78. Corrado D, Basso C, Pilichou K, Thiene G. Molecular biology and clinical management of arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Heart*. 2011; 97:530–539. [PubMed: 20930047]
79. Thiene G, Nava A, Corrado D, Rossi L, Pennelli N. Right ventricular cardiomyopathy and sudden death in young people. *The New England journal of medicine*. 1988; 318:129–133. [PubMed: 3336399]
80. Delmar M, McKenna WJ. The cardiac desmosome and arrhythmogenic cardiomyopathies: from gene to disease. *Circ Res*. 2010; 107:700–714. [PubMed: 20847325]
81. Delmar M, Makita N. Cardiac connexins, mutations and arrhythmias. *Curr OpinCardiol*. 2012; 27:236–241.
82. Cerrone M, Noorman M, Lin X, Chkourko H, Liang FX, van der Nagel R, Hund T, Birchmeier W, Mohler P, van Veen TA, van Rijen HV, Delmar M. Sodium current deficit and arrhythmogenesis in a murine model of plakophilin-2 haploinsufficiency. *Cardiovasc Res*. 2012; 95:460–468. [PubMed: 22764151]
83. Corrado D, Basso C, Buja G, Nava A, Rossi L, Thiene G. Right bundle branch block, right precordial ST-segment elevation, and sudden death in young people. *Circulation*. 2001; 103:710–717. [PubMed: 11156883]
84. Cerrone M, Lin X, Zhang M, et al. Missense mutations in plakophilin-2 cause sodium current deficit and associate with a Brugada syndrome phenotype. *Circulation*. 2014; 129:1092–1103. [PubMed: 24352520]
85. Corrado D, Leoni L, Link MS, et al. Implantable cardioverter-defibrillator therapy for prevention of sudden death in patients with arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Circulation*. 2003; 108:3084–3091. [PubMed: 14638546]

86. Matsuo K, Kurita T, Inagaki M, Kakishita M, Aihara N, Shimizu W, Taguchi A, Suyama K, Kamakura S, Shimomura K. The circadian pattern of the development of ventricular fibrillation in patients with Brugada syndrome. *EurHeart J*. 1999; 20:465–470.
87. Peters S, Trummel M, Denecke S, Koehler B. Results of ajmaline testing in patients with arrhythmogenic right ventricular dysplasia-cardiomyopathy. *IntJ Cardiol*. 2004; 95:207–210. [PubMed: 15193821]
88. Peters S, Trummel M, Koehler B. QRS fragmentation in standard ECG as a diagnostic marker of arrhythmogenic right ventricular dysplasia-cardiomyopathy. *Heart Rhythm*. 2008; 5:1417–1421. [PubMed: 18783995]
89. Postema PG, Wolpert C, Amin AS, Probst V, Borggrefe M, Roden DM, Priori SG, Tan HL, Hiraoka M, Brugada J, Wilde AA. Drugs and Brugada syndrome patients: review of the literature, recommendations, and an up-to-date website (www.brugadadrugs.org). *Heart Rhythm*. 2009; 6:1335–1341. www.brugadadrugs.org [PubMed: 19716089]
90. Shimizu W. Acquired forms of the Brugada syndrome. *J Electrocardiol*. 2005; 38(Suppl):22–25. [PubMed: 16226070]
91. Brugada P, Brugada J, Brugada R. Arrhythmia induction by antiarrhythmic drugs. *Pacing ClinElectrophysiol*. 2000; 23:291–292.
92. Brugada R, Brugada J, Antzelevitch C, Kirsch GE, Potenza D, Towbin JA, Brugada P. Sodium channel blockers identify risk for sudden death in patients with ST-segment elevation and right bundle branch block but structurally normal hearts. *Circulation*. 2000; 101:510–515. [PubMed: 10662748]
93. Miyazaki T, Mitamura H, Miyoshi S, Soejima K, Aizawa Y, Ogawa S. Autonomic and antiarrhythmic drug modulation of ST segment elevation in patients with Brugada syndrome. *JAmCollCardiol*. 1996; 27:1061–1070.
94. Babalarios VC, Hurst JW. Tricyclic antidepressants and the Brugada syndrome: an example of Brugada waves appearing after the administration of desipramine. *ClinCardiol*. 2002; 25:395–398.
95. Goldgran-Toledano D, Sideris G, Kevorkian JP. Overdose of cyclic antidepressants and the Brugada syndrome. *NEnglJMed*. 2002; 346:1591–1592.
96. Tada H, Sticherling C, Oral H, Morady F. Brugada syndrome mimicked by tricyclic antidepressant overdose. *JCardiovasElectrophysiol*. 2001; 12:275–275.
97. Ortega-Carnicer J, Bertos-Polo J, Gutierrez-Tirado C. Aborted sudden death, transient Brugada pattern, and wide QRS dysrhythmias after massive cocaine ingestion. *JElectrocardiol*. 2001; 34:345–349. [PubMed: 11590577]
98. Nogami A, Nakao M, Kubota S, et al. Enhancement of J-ST-segment elevation by the glucose and insulin test in Brugada syndrome. *Pacing ClinElectrophysiol*. 2003; 26:332–337.
99. Araki T, Konno T, Itoh H, Ino H, Shimizu M. Brugada syndrome with ventricular tachycardia and fibrillation related to hypokalemia. *CircJ*. 2003; 67:93–95. [PubMed: 12520160]
100. Pastor A, Nunez A, Cantale C, Cosio FG. Asymptomatic Brugada syndrome case unmasked during dimenhydrinate infusion. *JCardiovasc Electrophysiol*. 2001; 12:1192–1194. [PubMed: 11699532]
101. Chiale PA, Garro HA, Fernandez PA, Elizari MV. High-degree right bundle branch block obscuring the diagnosis of Brugada electrocardiographic pattern. *Heart Rhythm*. 2012; 9:974–976. [PubMed: 22306100]
102. Baranchuk A, Nguyen T, Ryu MH, Femenia F, Zareba W, Wilde AA, Shimizu W, Brugada P, Perez-Riera AR. Brugada phenocopy: new terminology and proposed classification. *Ann NoninvasiveElectrocardiol*. 2012; 17:299–314.
103. Nam GB. Idiopathic ventricular fibrillation, early repolarization and other J wave-related ventricular fibrillation syndromes. *Circ J*. 2012; 76:2723–2731. [PubMed: 23131759]
104. Yan GX, Antzelevitch C. Cellular basis for the electrocardiographic J wave. *Circulation*. 1996; 93:372–379. [PubMed: 8548912]
105. McIntyre WF, Perez-Riera AR, Femenia F, Baranchuk A. Coexisting early repolarization pattern and Brugada syndrome: recognition of potentially overlapping entities. *J Electrocardiol*. 2012; 45:195–198. [PubMed: 22178622]

106. Benito B, Sarkozy A, Mont L, Henkens S, Berruezo A, Tamborero D, Arzamendi D, Berne P, Brugada R, Brugada P, Brugada J. Gender differences in clinical manifestations of Brugada syndrome. *J Am Coll Cardiol*. 2008; 52:1567–1573. [PubMed: 19007594]
107. Kamakura T, Kawata H, Nakajima I, et al. Significance of non-type 1 anterior early repolarization in patients with inferolateral early repolarization syndrome. *J Am Coll Cardiol*. 2013; 62:1610–1618. [PubMed: 23850930]
108. Kawata H, Noda T, Yamada Y, Okamura H, Satomi K, Aiba T, Takaki H, Aihara N, Isobe M, Kamakura S, Shimizu W. Effect of sodium-channel blockade on early repolarization in inferior/lateral leads in patients with idiopathic ventricular fibrillation and Brugada syndrome. *Heart Rhythm*. 2012; 9:77–83. [PubMed: 21855521]
109. Matsumoto AM. Fundamental aspects of hypogonadism in the aging male. *Reviews in urology*. 2003; 5(Suppl 1):S3–s10. [PubMed: 16985941]
110. Kalla H, Yan GX, Marinchak R. Ventricular fibrillation in a patient with prominent J (Osborn) waves and ST segment elevation in the inferior electrocardiographic leads: a Brugada syndrome variant? *J Cardiovasc Electrophysiol*. 2000; 11:95–98. [PubMed: 10695469]
111. Aizawa Y, Sato A, Watanabe H, et al. Dynamicity of the J-wave in idiopathic ventricular fibrillation with a special reference to pause-dependent augmentation of the J-wave. *J Am Coll Cardiol*. 2012; 59:1948–1953. [PubMed: 22624834]
112. Nademanee K. Sudden unexplained death syndrome in southeast Asia. *American Journal of Cardiology*. 1997; 79(6A):10–11. [PubMed: 9080856]
113. Watanabe H, Nogami A, Ohkubo K, et al. Electrocardiographic characteristics and *SCN5A* mutations in idiopathic ventricular fibrillation associated with early repolarization. *Circ ArrhythmElectrophysiol*. 2011; 4:874–881.
114. Antzelevitch C, Pollevick GD, Cordeiro JM, et al. Loss-of-function mutations in the cardiac calcium channel underlie a new clinical entity characterized by ST-segment elevation, short QT intervals, and sudden cardiac death. *Circulation*. 2007; 115:442–449. [PubMed: 17224476]
115. Shimizu W, Kamakura S. Catecholamines in children with congenital long QT syndrome and Brugada syndrome. *J Electrocardiol*. 2001; 34(Suppl):173–175. [PubMed: 11781952]
116. Suzuki H, Torigoe K, Numata O, Yazaki S. Infant case with a malignant form of Brugada syndrome. *J Cardiovasc Electrophysiol*. 2000; 11:1277–1280. [PubMed: 11083249]
117. Ohgo T, Okamura H, Noda T, Satomi K, Suyama K, Kurita T, Aihara N, Kamakura S, Ohe T, Shimizu W. Acute and chronic management in patients with Brugada syndrome associated with electrical storm of ventricular fibrillation. *Heart Rhythm*. 2007; 4:695–700. [PubMed: 17556186]
118. Watanabe A, Fukushima KK, Morita H, Miura D, Sumida W, Hiramatsu S, Banba K, Nishii N, Nagase S, Nakamura K, Sakuragi S, Ohe T. Low-dose isoproterenol for repetitive ventricular arrhythmia in patients with Brugada syndrome. *EurHeart J*. 2006; 27:1579–1583.
119. Hermida JS, Denjoy I, Clerc J, Extramiana F, Jarry G, Milliez P, Guicheney P, Di Fusco S, Rey JL, Cauchemez B, Leenhardt A. Hydroquinidine therapy in Brugada syndrome. *J Am Coll Cardiol*. 2004; 43:1853–1860. [PubMed: 15145111]
120. Belhassen B, Glick A, Viskin S. Efficacy of quinidine in high-risk patients with Brugada syndrome. *Circulation*. 2004; 110:1731–1737. [PubMed: 15381640]
121. Tsuchiya T, Ashikaga K, Honda T, Arita M. Prevention of ventricular fibrillation by cilostazol, an oral phosphodiesterase inhibitor, in a patient with Brugada syndrome. *JCardiovascElectrophysiol*. 2002; 13:698–701.
122. Iguchi K, Noda T, Kamakura S, Shimizu W. Beneficial effects of cilostazol in a patient with recurrent ventricular fibrillation associated with early repolarization syndrome. *Heart Rhythm*. 2013; 10:604–606. [PubMed: 23142636]
123. Agac MT, Erkan H, Korkmaz L. Conversion of Brugada type I to type III and successful control of recurrent ventricular arrhythmia with cilostazol. *ArchCardiovasc Dis*. 2013
124. Hasegawa K, Ashihara T, Kimura H, Jo H, Itoh H, Yamamoto T, Aizawa Y, Horie M. Long-term pharmacological therapy of Brugada syndrome: is J-wave attenuation a marker of drug efficacy? *InternMed*. 2014; 53:1523–1526.

125. Shinohara T, Ebata Y, Ayabe R, Fukui A, Okada N, Yufu K, Nakagawa M, Takahashi N. Combination therapy of cilostazol and bepridil suppresses recurrent ventricular fibrillation related to J-wave syndromes. *Heart Rhythm*. 2014; 11:1441–1445. [PubMed: 24813378]
126. Haissaguerre M, Sacher F, Nogami A, et al. Characteristics of recurrent ventricular fibrillation associated with inferolateral early repolarization role of drug therapy. *J Am Coll Cardiol*. 2009; 53:612–619. [PubMed: 19215837]
127. Junttila MJ, Tikkanen JT, Kentta T, Anttonen O, Aro AL, Porthan K, Kerola T, Rissanen HA, Knekt P, Huikuri HV. Early repolarization as a predictor of arrhythmic and nonarrhythmic cardiac events in middle-aged subjects. *Heart Rhythm*. 2014; 11:1701–1706. [PubMed: 24858812]
128. Nam GB, Ko KH, Kim J, Park KM, Rhee KS, Choi KJ, Kim YH, Antzelevitch C. Mode of onset of ventricular fibrillation in patients with early repolarization pattern vs. Brugada syndrome. *Eur Heart J*. 2010; 31:330–339. [PubMed: 19880418]
129. Nademanee K, Veerakul G, Chandanamattha P, Chaothawee L, Ariyachaipanich A, Jirasirojanakorn K, Likittanasombat K, Bhuripanyo K, Ngarmukos T. Prevention of ventricular fibrillation episodes in Brugada syndrome by catheter ablation over the anterior right ventricular outflow tract epicardium. *Circulation*. 2011; 123:1270–1279. [PubMed: 21403098]
130. Sacher F, Jesel L, Jais P, Haissaguerre M. Insight into the mechanism of Brugada syndrome: epicardial substrate and modification during ajmaline testing. *Heart Rhythm*. 2014; 11:732–734. [PubMed: 23732228]
131. Brugada J, Pappone C, Berruezo A, Vicedomini G, Manguso F, Ciconte G, Giannelli L, Santinelli V. Brugada Syndrome Phenotype Elimination by Epicardial Substrate Ablation. *Circ Arrhythm Electrophysiol*. 2015
132. Ghosh S, Cooper DH, Vijayakumar R, Zhang J, Pollak S, Haissaguerre M, Rudy Y. Early repolarization associated with sudden death: insights from noninvasive electrocardiographic imaging. *Heart Rhythm*. 2010; 7:534–537. [PubMed: 20153422]
133. Zhang J, Sacher F, Hoffmayer K, et al. Cardiac electrophysiological substrate underlying the ECG phenotype and electrogram abnormalities in Brugada syndrome patients. *Circulation*. 2015; 131:1950–1959. [PubMed: 25810336]
134. Kowalczyk E, Kasprzak JD, Lipiec P. Giant J-wave and Brugada-like pattern in a patient with severe hypothermia. *Acta Cardiol*. 2014; 69:66–67. [PubMed: 24640526]
135. RuDusky BM. The electrocardiogram in hypothermia-the J wave and the Brugada syndrome. *Am J Cardiol*. 2004; 93:671–672. [PubMed: 14996612]
136. Adler A, Topaz G, Heller K, Zeltser D, Tzioni-Ohayon T, Rozovski U, Halkin A, Rosso R, Ben-Shachar S, Antzelevitch C, Viskin S. Fever-induced Brugada pattern: How common is it and what does it mean? *Heart Rhythm*. 2013; 10:1375–1382. [PubMed: 23872691]
137. Ansari E, Cook JR. Profound hypothermia mimicking a Brugada type ECG. *J Electrocardiol*. 2003; 36:257–260. [PubMed: 12942489]
138. Amin AS, Meregalli PG, Bardai A, Wilde AA, Tan HL. Fever increases the risk for cardiac arrest in the Brugada syndrome. *AnnInternMed*. 2008; 149:216–218.
139. Rattanawong P, Vutthikraivit W, Charoensri A, Jongraksak T, Prombandankul A, Kanjanahattakij N, Rungaramsin S, Wisaratapong T, Ngarmukos T. Fever-Induced Brugada Syndrome Is More Common Than Previously Suspected: A Cross-Sectional Study from an Endemic Area. *Annals of noninvasive electrocardiology : the official journal of the International Society for Holter and Noninvasive Electrocardiology, Inc*. 2015
140. Noda T, Shimizu W, Tanaka K, Chayama K. Prominent J wave and ST segment elevation: serial electrocardiographic changes in accidental hypothermia. *J CardiovascElectrophysiol*. 2003; 14:223.
141. Tan HL, Meregalli PG. Lethal ECG changes hidden by therapeutic hypothermia. *Lancet*. 2007; 369:78. [PubMed: 17208645]
142. Kurisu S, Inoue I, Kawagoe T, Ishihara M, Shimatani Y, Nakama Y, Maruhashi T, Kagawa E, Dai K, Aokage T, Matsushita J, Ikenaga H. Therapeutic hypothermia after out-of-hospital cardiac arrest due to Brugada syndrome. *Resuscitation*. 2008; 79:332–335. [PubMed: 18620795]
143. Patel RB, Ng J, Reddy V, Chokshi M, Parikh K, Subacius H, sheikh-Ali AA, Nguyen T, Link MS, Goldberger JJ, Ilkhanoff L, Kadish AH. Early repolarization associated with ventricular

- arrhythmias in patients with chronic coronary artery disease. *Circ ArrhythmElectrophysiol.* 2010; 3:489–495.
144. Gurabi Z, Koncz I, Patocskaï B, Nesterenko VV, Antzelevitch C. Cellular mechanism underlying hypothermia-induced VT/VF in the setting of early repolarization and the protective effect of quinidine, cilostazol and milrinone. *Circ Arrhythm Electrophysiol.* 2014; 7:134–142. [PubMed: 24429494]
 145. Kawata H, Morita H, Yamada Y, et al. Prognostic significance of early repolarization in inferolateral leads in Brugada patients with documented ventricular fibrillation: a novel risk factor for Brugada syndrome with ventricular fibrillation. *Heart Rhythm.* 2013; 10:1161–1168. [PubMed: 23587501]
 146. Antzelevitch C. Genetic, molecular and cellular mechanisms underlying the J wave syndromes. *Circ J.* 2012; 76:1054–1065. [PubMed: 22498570]
 147. Kapplinger JD, Tester DJ, Alders M, et al. An international compendium of mutations in the SCN5A encoded cardiac sodium channel in patients referred for Brugada syndrome genetic testing. *Heart Rhythm.* 2010; 7:33–46. [PubMed: 20129283]
 148. Ackerman MJ, Priori SG, Willems S, et al. HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies this document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Heart Rhythm.* 2011; 8:1308–1339. [PubMed: 21787999]
 149. Park DS, Cerrone M, Morley G, et al. Genetically engineered SCN5A mutant pig hearts exhibit conduction defects and arrhythmias. *J Clin Invest.* 2015; 125:403–412. [PubMed: 25500882]
 150. Szel T, Antzelevitch C. Abnormal repolarization as the basis for late potentials and fractionated electrograms recorded from epicardium in experimental models of brugada syndrome. *J Am Coll Cardiol.* 2014; 63:2037–2045. [PubMed: 24657694]
 151. Patocskaï, B.; Szel, T.; Yoon, N.; Antzelevitch, C. Cellular mechanisms underlying the fractionated and late potentials on epicardial electrograms and the ameliorative effect of epicardial radiofrequency ablation in an experimental model of Brugada syndrome; Program and abstracts of the 24th Annual Upstate New York Cardiac Electrophysiology Society Meeting; Buffalo, NY. Nov 3. 2014 2014Abstract 006
 152. Burashnikov E, Pfeiffer R, Barajas-Martinez H, et al. Mutations in the cardiac L-type calcium channel associated J wave syndrome and sudden cardiac death. *Heart Rhythm.* 2010; 7:1872–1882. [PubMed: 20817017]
 153. Cordeiro JM, Marieb M, Pfeiffer R, Calloe K, Burashnikov E, Antzelevitch C. Accelerated inactivation of the L-type calcium due to a mutation in *CACNB2b* due to a mutation in *CACNB2b* underlies Brugada syndrome. *J MolCell Cardiol.* 2009; 46:695–703.
 154. Antzelevitch C, Pollevick GD, Cordeiro JM, et al. Loss-of-function mutations in the cardiac calcium channel underline a new clinical entity characterized by ST segment elevation, short QT intervals, and sudden cardiac death. *CircRes.* 2006; 99:1279.
 155. Gurnett CA, De WM, Campbell KP. Dual function of the voltage-dependent Ca²⁺ channel alpha 2 delta subunit in current stimulation and subunit interaction. *Neuron.* 1996; 16:431–440. [PubMed: 8789958]
 156. Barajas-Martinez H, Hu D, Ferrer T, et al. Molecular genetic and functional association of Bugada and early repolarization syndromes with S422L missense mutation in *KCNJ8*. *Heart Rhythm.* 2012; 9:548–555. [PubMed: 22056721]
 157. Delaney JT, Muhammad R, Blair MA, Kor K, Fish FA, Roden DM, Darbar D. A KCNJ8 mutation associated with early repolarization and atrial fibrillation. *Europace.* 2012; 14:1428–1432. [PubMed: 22562657]
 158. Medeiros-Domingo A, Tan BH, Crotti L, et al. Gain-of-function mutation S422L in the KCNJ8-encoded cardiac K(ATP) channel Kir6.1 as a pathogenic substrate for J-wave syndromes. *Heart Rhythm.* 2010; 7:1466–1471. [PubMed: 20558321]
 159. Hu D, Barajas-Martinez H, Medeiros-Domingo A, et al. Novel mutations in the sodium channel 2 subunit gene (SCN2B) associated with Brugada syndrome and atrial fibrillation. *Circulation.* 2012; 126:A16521.

160. Riuro H, Beltran-Alvarez P, Tarradas A, et al. A missense mutation in the sodium channel $\beta 2$ subunit reveals *SCN2B* as a new candidate gene for Brugada syndrome. *Hum Mutat.* 2013; 34:961–966. [PubMed: 23559163]
161. Giudicessi JR, Ye D, Tester DJ, Crotti L, Mugione A, Nesterenko VV, Albertson RM, Antzelevitch C, Schwartz PJ, Ackerman MJ. Transient outward current (Ito) gain-of-function mutations in the KCND3-encoded Kv4.3 potassium channel and Brugada syndrome. *Heart Rhythm.* 2011; 8:1024–1032. [PubMed: 21349352]
162. Delpón E, Cordeiro JM, Núñez L, Thomsen PEB, Guerchicoff A, Pollevick GD, Wu Y, Kanter JK, Larsen CT, Burashnikov A, Christiansen M, Antzelevitch C. Functional effects of *KCNE3* mutation and its role in the development of Brugada syndrome. *Circ Arrhythm Electrophysiol.* 2008; 1:209–218. [PubMed: 19122847]
163. Olesen MS, Jensen NF, Holst AG, Nielsen JB, Tfelt-Hansen J, Jespersen T, Sajadieh A, Haunso S, Lund JT, Calloe K, Schmitt N, Svendsen JH. A novel nonsense variant in Nav1.5 cofactor MOG1 eliminates its sodium current increasing effect and may increase the risk of arrhythmias. *Can J Cardiol.* 2011; 27:523–523. [PubMed: 21621375]
164. Kattynarath D, Maugendre S, Neyroud N, et al. MOG1: a new susceptibility gene for Brugada syndrome. *Circ Cardiovasc Genet.* 2011; 4:261–268. [PubMed: 21447824]
165. Cerrone M, Lin X, Zhang M, et al. Missense mutations in plakophilin-2 cause sodium current deficit and associate with a Brugada syndrome phenotype. *Circulation.* 2013; 129:1092–1103. [PubMed: 24352520]
166. Hennessey JA, Marcou CA, Wang C, Wei EQ, Wang C, Tester DJ, Torchio M, Dagradi F, Crotti L, Schwartz PJ, Ackerman MJ, Pitt GS. FGF12 is a candidate Brugada syndrome locus. *Heart Rhythm.* 2013; 10:1886–1894. [PubMed: 24096171]
167. Bezzina CR, Barc J, Mizusawa Y, et al. Common variants at *SCN5A-SCN10A* and *HEY2* are associated with Brugada syndrome, a rare disease with high risk of sudden cardiac death. *Nat Genet.* 2013; 45:1044–1049. [PubMed: 23872634]
168. Boukens BJ, Sylva M, de Gier-de VC, Remme CA, Bezzina C, Christoffels VM, Coronel R. Reduced sodium channel function unmasks residual embryonic slow conduction in the adult right ventricular outflow tract. *Circ Res.* 2013; 113:137–141. [PubMed: 23661717]
169. Hartman ME, Liu Y, Zhu WZ, Chien WM, Weldy CS, Fishman GI, Laflamme MA, Chin MT. Myocardial deletion of transcription factor CHF1/Hey2 results in altered myocyte action potential and mild conduction system expansion but does not alter conduction system function or promote spontaneous arrhythmias. *FASEB J.* 2014; 28:3007–3015. [PubMed: 24687990]
170. Ishikawa T, Takahashi N, Ohno S, Sakurada H, Nakamura K, On YK, Park JE, Makiyama T, Horie M, Arimura T, Makita N, Kimura A. Novel *SCN3B* mutation associated with brugada syndrome affects intracellular trafficking and function of Nav1.5. *Circ J.* 2013; 77:959–967. [PubMed: 23257389]
171. Hu D, Barajas-Martinez H, Burashnikov E, Springer M, Wu Y, Varro A, Pfeiffer R, Koopmann TT, Cordeiro JM, Guerchicoff A, Pollevick GD, Antzelevitch C. A mutation in the beta 3 subunit of the cardiac sodium channel associated with Brugada ECG phenotype. *Circ Cardiovasc Genet.* 2009; 2:270–278. [PubMed: 20031595]
172. Watanabe H, Koopmann TT, et al. Sodium channel $\beta 1$ subunit mutations associated with Brugada syndrome and cardiac conduction disease in humans. *J ClinInvest.* 2008; 118:2260–2268. S. LS.
173. Valdivia CR, Ueda K, Ackerman MJ, Makielski JC. GPD1L links redox state to cardiac excitability by PKC-dependent phosphorylation of the sodium channel SCN5A. *Am J Physiol Heart Circ Physiol.* 2009; 297:H1446–H1452. [PubMed: 19666841]
174. Shy D, Gillet L, Abriel H. Cardiac sodium channel Nav1.5 distribution in myocytes via interacting proteins: the multiple pool model. *BiochimBiophys Acta.* 2013; 1833:886–894.
175. Weiss R, Barmada MM, Nguyen T, Seibel JS, Cavlovich D, Kornblit CA, Angelilli A, Villanueva F, McNamara DM, London B. Clinical and molecular heterogeneity in the Brugada syndrome: a novel gene locus on chromosome 3. *Circulation.* 2002; 105:707–713. [PubMed: 11839626]
176. Hu D, Barajas-Martinez H, Medeiros-Domingo A, et al. A novel rare variant in *SCN1Bb* linked to Brugada syndrome and SIDS by combined modulation of Na(v)1.5 and K(v)4.3 channel currents. *Heart Rhythm.* 2012; 9:760–769. [PubMed: 22155597]

177. Hu D, Barajas-Martinez H, Pfeiffer R, et al. Mutations in *SCN10A* are responsible for a large fraction of cases of Brugada syndrome. *J Am Coll Cardiol*. 2014; 64:66–79. [PubMed: 24998131]
178. Behr ER, Savio-Galimberti E, Barc J, et al. Role of common and rare variants in *SCN10A*: results from the Brugada syndrome QRS locus gene discovery collaborative study. *Cardiovasc Res*. 2015
179. Antzelevitch C. Cardiac repolarization. The long and short of it. *Europace*. 2005; 7(Suppl 2):3–9. [PubMed: 16102498]
180. Liu H, Chatel S, Simard C, et al. Molecular genetics and functional anomalies in a series of 248 Brugada cases with 11 mutations in the TRPM4 channel. *PLoS ONE*. 2013; 8:e54131. [PubMed: 23382873]
181. Perrin MJ, Adler A, Green S, et al. Evaluation of genes encoding for the transient outward current (I_{to}) identifies the *KCND2* gene as a cause of J wave syndrome associated with sudden cardiac death. *Circ Cardiovasc Genet*. 2014
182. Verkerk AO, Wilders R, Schulze-Bahr E, et al. Role of sequence variations in the human *ether-a-go-go*-related gene (*HERG*, *KCNH2*) in the Brugada syndrome 1. *Cardiovasc Res*. 2005; 68:441–453.
183. Wilders R, Verkerk AO. Role of the R1135H *KCNH2* mutation in Brugada syndrome. *Int J Cardiol*. 2010; 144:149–151. [PubMed: 19174314]
184. Ohno S, Zankov DP, Ding WG, et al. *KCNE5* (*KCNE1L*) variants are novel modulators of Brugada syndrome and idiopathic ventricular fibrillation. *Circ Arrhythm Electrophysiol*. 2011; 4:352–361.
185. Boczek NJ, Ye D, Johnson EK, et al. Characterization of *SEMA3A*-encoded semaphorin as a naturally occurring Kv4.3 protein inhibitor and its contribution to Brugada syndrome. *Circ Res*. 2014; 115:460–469. [PubMed: 24963029]
186. Ueda K, Nakamura K, Hayashi T, et al. Functional characterization of a trafficking-defective *HCFN4* mutation, D553N, associated with cardiac arrhythmia. *J Biol Chem*. 2004; 279:27194–27198. [PubMed: 15123648]
187. Noseworthy PA, Tikkanen JT, Porthan K, et al. The early repolarization pattern in the general population clinical correlates and heritability. *J Am Coll Cardiol*. 2011; 57:2284–2289. [PubMed: 21600720]
188. Reinhard W, Kaess BM, Debiec R, Nelson CP, Stark K, Tobin MD, Macfarlane PW, Tomaszewski M, Samani NJ, Hengstenberg C. Heritability of early repolarization: a population-based study. *Circ Cardiovasc Genet*. 2011; 4:134–138. [PubMed: 21282333]
189. Nunn LM, Bhar-Amato J, Lowe MD, Macfarlane PW, Rogers P, McKenna WJ, Elliott PM, Lambiase PD. Prevalence of J-point elevation in sudden arrhythmic death syndrome families. *J Am Coll Cardiol*. 2011; 58:286–290. [PubMed: 21737021]
190. Haissaguerre M, Chatel S, Sacher F, et al. Ventricular fibrillation with prominent early repolarization associated with a rare variant of *KCNJ8/K_{ATP}* channel. *J Cardiovasc Electrophysiol*. 2009; 20:93–98.
191. Schwartz PJ, Ackerman MJ, George AL Jr, Wilde AA. Impact of genetics on the clinical management of channelopathies. *J Am Coll Cardiol*. 2013; 62:169–180. [PubMed: 23684683]
192. Raffan E, Semple RK. Next generation sequencing--implications for clinical practice. *Br Med Bull*. 2011; 99:53–71. [PubMed: 21705347]
193. Refsgaard L, Holst AG, Sadjadieh G, Haunso S, Nielsen JB, Olesen MS. High prevalence of genetic variants previously associated with LQT syndrome in new exome data. *European journal of human genetics : EJHG*. 2012; 20:905–908. [PubMed: 22378279]
194. Kapa S, Tester DJ, Salisbury BA, Harris-Kerr C, Pungliya MS, Alders M, Wilde AA, Ackerman MJ. Genetic testing for long-QT syndrome: distinguishing pathogenic mutations from benign variants. *Circulation*. 2009; 120:1752–1760. [PubMed: 19841300]
195. Andreassen C, Refsgaard L, Nielsen JB, Sadjadieh A, Winkel BG, Tfelt-Hansen J, Haunso S, Holst AG, Svendsen JH, Olesen MS. Mutations in genes encoding cardiac ion channels previously associated with sudden infant death syndrome (SIDS) are present with high frequency in new exome data. *Can J Cardiol*. 2013; 29:1104–1109. [PubMed: 23465283]

196. Kapplinger J, Tester D, Alders M, et al. An International Compendium of Mutations in the SCN5A-Encoded Cardiac Sodium Channel in Patients Referred for Brugada Syndrome Genetic Testing. *Heart Rhythm*. 2010; 7:33–46. [PubMed: 20129283]
197. Hu D, Barajas-Martinez H, Terzic A, et al. *ABCC9* is a novel Brugada and early repolarization syndrome susceptibility gene. *Int J Cardiol*. 2014; 171:431–442. [PubMed: 24439875]
198. Alfares AA, Kelly MA, McDermott G, et al. Results of clinical genetic testing of 2,912 probands with hypertrophic cardiomyopathy: expanded panels offer limited additional sensitivity. *Genet Med*. 2015
199. Le Scouarnec S, Karakachoff M, Gourraud JB, et al. Testing the burden of rare variation in arrhythmia-susceptibility genes provides new insights into molecular diagnosis for Brugada syndrome. *Hum Mol Genet*. 2015
200. Veeramah KR, Karafet TM, Wolf D, Samson RA, Hammer MF. The KCNJ8-S422L variant previously associated with J-wave syndromes is found at an increased frequency in Ashkenazi Jews. *European journal of human genetics*. *EJHG*. 2014; 22:94–98. [PubMed: 23632791]
201. Risgaard B, Jabbari R, Refsgaard L, Holst AG, Haunso S, Sadjadieh A, Winkel BG, Olesen M, Tfelt-Hansen J. High prevalence of genetic variants previously associated with Brugada Syndrome in new exome data. *Clin Genet*. 2013; 84:489–495. [PubMed: 23414114]
202. Campuzano O, Allegue C, Fernandez A, Iglesias A, Brugada R. Determining the pathogenicity of genetic variants associated with cardiac channelopathies. *Sci Rep*. 2015; 5:7953. [PubMed: 25608792]
203. Crotti L, Marcou CA, Tester DJ, Castelletti S, Giudicessi JR, Torchio M, Medeiros-Domingo A, Simone S, Will ML, Dagradi F, Schwartz PJ, Ackerman MJ. Spectrum and prevalence of mutations involving BrS1-through BrS12-susceptibility genes in a cohort of unrelated patients referred for Brugada syndrome genetic testing: implications for genetic testing. *J Am Coll Cardiol*. 2012; 60:1410–1418. [PubMed: 22840528]
204. Song W, Shou W. Cardiac sodium channel Nav1.5 mutations and cardiac arrhythmia. *Pediatric cardiology*. 2012; 33:943–949. [PubMed: 22460359]
205. Ruklisa D, Ware JS, Walsh R, Balding DJ, Cook SA. Bayesian models for syndrome- and gene-specific probabilities of novel variant pathogenicity. *Genome medicine*. 2015; 7:5. [PubMed: 25649125]
206. Juang JM, Lu TP, Lai LC, et al. Utilizing multiple in silico analyses to identify putative causal SCN5A variants in Brugada syndrome. *Sci Rep*. 2014; 4:3850. [PubMed: 24463578]
207. Walsh R, Peters NS, Cook SA, Ware JS. Paralog annotation identifies novel pathogenic variants in patients with Brugada syndrome and catecholaminergic polymorphic ventricular tachycardia. *J Med Genet*. 2014; 51:35–44. [PubMed: 24136861]
208. Kapplinger JD, Giudicessi JR, Tester DJ, Callis TE, Ackerman MJ. Enhanced Classification of Non-synonymous Single Nucleotide Variants in the SCN5A-Encoded Nav1.5 Cardiac Sodium Channel. *Heart Rhythm*. 2012; 9:1912.
209. Kapplinger JD, Giudicessi JR, Ye D, Tester DJ, Callis TE, Valdivia CR, Makielski JC, Wilde AA, Ackerman MJ. Enhanced Classification of Brugada Syndrome-Associated and Long-QT Syndrome-Associated Genetic Variants in the SCN5A-Encoded Nav1.5 Cardiac Sodium Channel. *Circ Cardiovasc Genet*. 2015; 8:582–595. [PubMed: 25904541]
210. Probst V, Wilde AA, Barc J, et al. SCN5A mutations and the role of genetic background in the pathophysiology of Brugada syndrome. *Circ Cardiovasc Genet*. 2009; 2:552–557. [PubMed: 20031634]
211. Matsuo K, Akahoshi M, Seto S, Yano K. Disappearance of the Brugada-type electrocardiogram after surgical castration: a role for testosterone and an explanation for the male preponderance? *Pacing ClinElectrophysiol*. 2003; 1151–1153. In press.
212. Antzelevitch C. Androgens and male predominance of the Brugada syndrome phenotype. *Pacing ClinElectrophysiol*. 2003; 26:1429–1431.
213. Korte AK, Derde L, van Wijk J, Tjan DH. Sudden cardiac arrest as a presentation of Brugada syndrome unmasked by thyroid storm. *BMJ case reports*. 2015; 2015

214. Wilde AA, Postema PG, Di Diego JM, Viskin S, Morita H, Fish JM, Antzelevitch C. The pathophysiological mechanism underlying Brugada syndrome: depolarization versus repolarization. *J Mol Cell Cardiol.* 2010; 49:543–553. [PubMed: 20659475]
215. Morita H, Zipes DP, Wu J. Brugada syndrome: insights of ST elevation, arrhythmogenicity, and risk stratification from experimental observations. *Heart Rhythm.* 2009; 6:S34–S43. [PubMed: 19880072]
216. Hoogendijk MG, Opthof T, Postema PG, Wilde AA, de Bakker JM, Coronel R. The Brugada ECG pattern: a marker of channelopathy, structural heart disease, or neither? Toward a unifying mechanism of the Brugada syndrome. *Circ ArrhythmElectrophysiol.* 2010; 3:283–290.
217. Hoogendijk MG, Potse M, Linnenbank AC, et al. Mechanism of right precordial ST-segment elevation in structural heart disease: Excitation failure by current-to-load mismatch. *Heart Rhythm.* 2010; 7:238–248. [PubMed: 20022821]
218. Patocsikai B, Antzelevitch C. Expert Opinion on Orphan Drugs. Expert opinion on orphan drugs. 2015
219. Wilde AA, Postema PG. Bringing home the bacon? The next step in cardiac sodium channelopathies. *J Clin Invest.* 2015; 125:99–101. [PubMed: 25500878]
220. Antzelevitch C, Brugada P, Brugada J, Brugada R, Shimizu W, Gussak I, Perez Riera AR. Brugada syndrome: a decade of progress. *CircRes.* 2002; 91:1114–1119.
221. Kurita T, Shimizu W, Inagaki M, Suyama K, Taguchi A, Satomi K, Aihara N, Kamakura S, Kobayashi J, Kosakai Y. The electrophysiologic mechanism of ST-segment elevation in Brugada syndrome. *JAmCollCardiol.* 2002; 40:330–334.
222. Gurabi Z, Koncz I, Patocsikai B, Nesterenko VV, Antzelevitch C. Cellular mechanism underlying hypothermia-induced ventricular tachycardia/ventricular fibrillation in the setting of early repolarization and the protective effect of quinidine, cilostazol, and milrinone. *Circ Arrhythm Electrophysiol.* 2014; 7:134–142. [PubMed: 24429494]
223. Huikuri HV. Separation of Benign from Malignant J waves. *Heart Rhythm.* 2015; 12:384–385. [PubMed: 25460858]
224. Aizawa Y, Sato M, Kitazawa H, Aizawa Y, Takatsuki S, Oda E, Okabe M, Fukuda K. Tachycardia-dependent augmentation of "notched J waves" in a general patient population without ventricular fibrillation or cardiac arrest: Not a repolarization but a depolarization abnormality? *Heart Rhythm.* 2015; 12:376–383. [PubMed: 25460863]
225. Badri M, Patel A, Yan G. Cellular and ionic basis of J-wave syndromes. *Trends Cardiovasc Med.* 2015; 25:12–21. [PubMed: 25446046]
226. Morita H, Kusano KF, Miura D, Nagase S, Nakamura K, Morita ST, Ohe T, Zipes DP, Wu J. Fragmented QRS as a marker of conduction abnormality and a predictor of prognosis of Brugada syndrome. *Circulation.* 2008; 118:1697–1704. [PubMed: 18838563]
227. Terho HK, Tikkanen JT, Junttila JM, Anttonen O, Kentta TV, Aro AL, Kerola T, Rissanen HA, Reunanen A, Huikuri HV. Prevalence and prognostic significance of fragmented QRS complex in middle-aged subjects with and without clinical or electrocardiographic evidence of cardiac disease. *Am J Cardiol.* 2014; 114:141–147. [PubMed: 24819902]
228. Osher HL, Wolff L. Electrocardiographic pattern simulating acute myocardial injury. *AmJMedSci.* 1953; 226:541–545.
229. Viskin S, Adler A, Halkin A, Rosso R. Reply: is the J wave or the ST slope malignant...or neither? *J Am Coll Cardiol.* 2014; 63:1812–1813. [PubMed: 24315912]
230. Rosso R, Adler A, Halkin A, Viskin S. Risk of sudden death among young individuals with J waves and early repolarization: putting the evidence into perspective. *Heart Rhythm.* 2011; 8:923–929. [PubMed: 21295159]
231. Aagaard P, Shulman E, Di Biase L, Fisher JD, Gross JN, Kargoli F, Kim SG, Palma EC, Ferrick KJ, Krumerman A. Prognostic value of automatically detected early repolarization. *Am J Cardiol.* 2014; 114:1431–1436. [PubMed: 25306428]
232. Uberoi A, Jain NA, Perez M, Weinkopff A, Ashley E, Hadley D, Turakhia MP, Froelicher V. Early repolarization in an ambulatory clinical population. *Circulation.* 2011; 124:2208–2214. [PubMed: 21986288]

233. Muramoto D, Yong CM, Singh N, Aggarwal S, Perez M, Ashley E, Hadley D, Froelicher V. Patterns and prognosis of all components of the J-wave pattern in multiethnic athletes and ambulatory patients. *Am Heart J*. 2014; 167:259–266. [PubMed: 24439988]
234. Perez MV, Uberoi A, Jain NA, Ashley E, Turakhia MP, Froelicher V. The prognostic value of early repolarization with ST-segment elevation in African Americans. *Heart Rhythm*. 2012; 9:558–565. [PubMed: 22094072]
235. Klatsky AL, Oehm R, Cooper RA, Udaltsova N, Armstrong MA. The early repolarization normal variant electrocardiogram: correlates and consequences. *Am J Med*. 2003; 115:171–177. [PubMed: 12935822]
236. Rollin A, Maury P, Bongard V, Sacher F, Delay M, Duparc A, Mondoly P, Carrie D, Ferrieres J, Ruidavets JB. Prevalence, prognosis, and identification of the malignant form of early repolarization pattern in a population-based study. *Am J Cardiol*. 2012; 110:1302–1308. [PubMed: 22819431]
237. Tikkanen JT, Junttila MJ, Anttonen O, Aro AL, Luttinen S, Kerola T, Sager SJ, Rissanen HA, Myerburg RJ, Reunanen A, Huikuri HV. Early repolarization: electrocardiographic phenotypes associated with favorable long-term outcome. *Circulation*. 2011; 123:2666–2673. [PubMed: 21632493]
238. Olson KA, Viera AJ, Soliman EZ, Crow RS, Rosamond WD. Long-term prognosis associated with J-point elevation in a large middle-aged biracial cohort: the ARIC study. *Eur Heart J*. 2011; 32:3098–3106. [PubMed: 21785106]
239. Rosso R, Halkin A, Viskin S. J waves and early repolarization: do not confuse me with the facts! *Heart Rhythm*. 2012; 9:1603–1604. [PubMed: 22842117]
240. Demidova MM, Martin-Yebra A, van der Pals J, Koul S, Erlinge D, Laguna P, Martinez JP, Platonov PG. Transient and rapid QRS-widening associated with a J-wave pattern predicts impending ventricular fibrillation in experimental myocardial infarction. *Heart Rhythm*. 2014; 11:1195–1201. [PubMed: 24691451]
241. Shinde R, Shinde S, Makhale C, Grant P, Sathe S, Durairaj M, Lokhandwala Y, Di Diego JM, Antzelevitch C. Occurrence of "J waves" in 12-lead ECG as a marker of acute ischemia and their cellular basis. *Pacing Clin Electrophysiol*. 2007; 30:817–819.
242. Gourraud JB, Le Scouarnec S, Sacher F, et al. Identification of large families in early repolarization syndrome. *J Am Coll Cardiol*. 2013; 61:164–172. [PubMed: 23273290]
243. Watanabe H, Makiyama T, Koyama T, et al. High prevalence of early repolarization in short QT syndrome. *Heart Rhythm*. 2010; 7:647–652. [PubMed: 20206319]
244. Letsas KP, Charalampous C, Korantzopoulos P, Tsikrikas S, Bramos D, Kollias G, Efremidis M, Sideris A. Novel indexes of heterogeneity of ventricular repolarization in subjects with early repolarization pattern. *Europace*. 2012; 14:877–878. [PubMed: 22186777]
245. Rosso R, Glikson E, Belhassen B, Katz A, Halkin A, Steinvil A, Viskin S. Distinguishing "benign" from "malignant early repolarization": The value of the ST-segment morphology. *Heart Rhythm*. 2012; 9:225–229. [PubMed: 21914497]
246. Mahida S, Derval N, Sacher F, et al. Role of electrophysiological studies in predicting risk of ventricular arrhythmia in early repolarization syndrome. *J Am Coll Cardiol*. 2015; 65:151–159. [PubMed: 25593056]
247. Probst V, Veltmann C, Eckardt L, et al. Long-term prognosis of patients diagnosed with Brugada syndrome: Results from the FINGER Brugada Syndrome Registry. *Circulation*. 2010; 121:635–643. [PubMed: 20100972]
248. Kamakura S, Ohe T, Nakazawa K, et al. Long-term prognosis of probands with Brugada-pattern ST-elevation in leads V1-V3. *Circ Arrhythm Electrophysiol*. 2009; 2:495–503. [PubMed: 19843917]
249. Conte G, Sieira J, Ciconte G, et al. Implantable cardioverter-defibrillator therapy in brugada syndrome: a 20-year single-center experience. *Journal of the American College of Cardiology*. 2015; 65:879–888. [PubMed: 25744005]
250. Sacher F, Probst V, Maury P, et al. Outcome after implantation of a cardioverter-defibrillator in patients with Brugada syndrome: a multicenter study-part 2. *Circulation*. 2013; 128:1739–1747. [PubMed: 23995538]

251. Priori SG, Gasparini M, Napolitano C, Della BP, Ottonelli AG, Sassone B, Giordano U, Pappone C, Mascioli G, Rossetti G, De NR, Colombo M. Risk stratification in Brugada syndrome: results of the PRELUDE (PROgrammed ELEctrical stimUlation preDictive valuE) registry. *J Am Coll Cardiol.* 2012; 59:37–45. [PubMed: 22192666]
252. Take Y, Morita H, Toh N, Nishii N, Nagase S, Nakamura K, Kusano KF, Ohe T, Ito H. Identification of high-risk syncope related to ventricular fibrillation in patients with Brugada syndrome. *Heart Rhythm.* 2012; 9:752–759. [PubMed: 22123317]
253. Conte G, C DEA, Sieira J, Levinstein M, Chierchia GB, G DIG, Baltogiannis G, Ciconte G, Saitoh Y, Casado-Arroyo R, Pappaert G, Brugada P. Clinical characteristics, management, and prognosis of elderly patients with Brugada syndrome. *J Cardiovasc Electrophysiol.* 2014; 25:514–519. [PubMed: 24400668]
254. Priori SG, Napolitano C, Glordano U, Collisani G, Memmi M. Brugada syndrome and sudden cardiac death in children. *Lancet.* 2000; 355:808–809. [PubMed: 10711933]
255. Probst V, Denjoy I, Meregalli PG, et al. Clinical aspects and prognosis of Brugada syndrome in children. *Circulation.* 2007; 115:2042–2048. [PubMed: 17404158]
256. Meregalli PG, Tan HL, Probst V, et al. Type of *SCN5A* mutation determines clinical severity and degree of conduction slowing in loss-of-function sodium channelopathies. *Heart Rhythm.* 2009; 6:341–348. [PubMed: 19251209]
257. Cordeiro JM, Barajas-Martinez H, Hong K, et al. Compound heterozygous mutations P336L and I1660V in the human cardiac sodium channel associated with the Brugada syndrome. *Circulation.* 2006; 114:2026–2033. [PubMed: 17075016]
258. Nunez L, Barana A, Amoros I, et al. p.D1690N Nav1.5 rescues p.G1748D mutation gating defects in a compound heterozygous Brugada syndrome patient. *Heart Rhythm.* 2013; 10:264–272. [PubMed: 23085483]
259. Poelzing S, Forleo C, Samodell M, et al. *SCN5A* polymorphism restores trafficking of a Brugada syndrome mutation on a separate gene. *Circulation.* 2006; 114:368–376. [PubMed: 16864729]
260. Viswanathan PC, Benson DW, Balser JR. A common *SCN5A* polymorphism modulates the biophysical effects of an *SCN5A* mutation. *J ClinInvest.* 2003; 111:341–346.
261. Sommariva E, Pappone C, Martinelli BF, Di RC, Rosaria CM, Salvi E, Vergara P, Sala S, Cusi D, Ferrari M, Benedetti S. Genetics can contribute to the prognosis of Brugada syndrome: a pilot model for risk stratification. *Eur J Hum Genet.* 2013; 21:911–917. [PubMed: 23321620]
262. Delise P, Allocca G, Marras E, et al. Risk stratification in individuals with the Brugada type 1 ECG pattern without previous cardiac arrest: usefulness of a combined clinical and electrophysiologic approach. *EurHeart J.* 2011; 32:169–176.
263. Yokokawa M, Okamura H, Noda T, Satomi K, Suyama K, Kurita T, Aihara N, Kamakura S, Shimizu W. Neurally mediated syncope as a cause of syncope in patients with Brugada electrocardiogram. *J Cardiovasc Electrophysiol.* 2010; 21:186–192. [PubMed: 19793146]
264. Richter S, Sarkozy A, Veltmann C, Chierchia GB, Boussy T, Wolpert C, Schimpf R, Brugada J, Brugada R, Borggreffe M, Brugada P. Variability of the diagnostic ECG pattern in an ICD patient population with Brugada syndrome. *J CardiovascElectrophysiol.* 2009; 20:69–75.
265. Veltmann C, Schimpf R, Echternach C, Eckardt L, Kuschyk J, Streitner F, Spehl S, Borggreffe M, Wolpert C. A prospective study on spontaneous fluctuations between diagnostic and non-diagnostic ECGs in Brugada syndrome: implications for correct phenotyping and risk stratification. *EurHeart J.* 2006; 27:2544–2556.
266. Rolf S. The ajmaline challenge in Brugada syndrome: Diagnostic impact, safety, and recommended protocol. *European Heart Journal.* 2003; 24:1104–1112. [PubMed: 12804924]
267. Veltmann C, Wolpert C, Sacher F, Mabo P, Schimpf R, Streitner F, Brade J, Kyndt F, Kuschyk J, Le MH, Borggreffe M, Probst V. Response to intravenous ajmaline: a retrospective analysis of 677 ajmaline challenges. *Europace.* 2009; 11:1345–1352. [PubMed: 19589796]
268. Okamura H, Kamakura T, Morita H, et al. Risk stratification in patients with Brugada syndrome without previous cardiac arrest - prognostic value of combined risk factors. *Circ J.* 2015; 79:310–317. [PubMed: 25428522]

269. Govindan M, Batchvarov VN, Raju H, Shanmugam N, Bizrah M, Bastiaenen R, Kiotsekoglou A, Camm J, Behr ER. Utility of high and standard right precordial leads during ajmaline testing for the diagnosis of Brugada syndrome. *Heart*. 2010; 96:1904–1908. [PubMed: 20962343]
270. Brugada J, Brugada R, Brugada P. Electrophysiologic testing predicts events in Brugada syndrome patients. *Heart Rhythm*. 2011; 8:1595–1597. [PubMed: 21762670]
271. Wilde AA, Viskin S. EP testing does not predict cardiac events in Brugada syndrome. *Heart Rhythm*. 2011; 8:1598–1600. [PubMed: 21767517]
272. Viskin S, Rogowski O. Asymptomatic Brugada syndrome: a cardiac ticking time-bomb? *Europace*. 2007; 9:707–710. [PubMed: 17715171]
273. Paul M, Gerss J, Schulze-Bahr E, Wichter T, Vahlhaus C, Wilde AA, Breithardt G, Eckardt L. Role of programmed ventricular stimulation in patients with Brugada syndrome: a meta-analysis of worldwide published data. *EurHeart J*. 2007; 28:2126–2133.
274. Brugada P, Brugada R, Brugada J. Patients with an asymptomatic Brugada electrocardiogram should undergo pharmacological and electrophysical testing. *Circulation*. 2005; 112:279–285. [PubMed: 16009809]
275. Sieira J, Conte G, Ciconte G, et al. Prognostic Value of Programmed Electrical Stimulation in Brugada Syndrome: 20 Years Experience. *Circ Arrhythm Electrophysiol*. 2015
276. Makimoto H, Kamakura S, Aihara N, et al. Clinical impact of the number of extrastimuli in programmed electrical stimulation in patients with Brugada type 1 electrocardiogram. *Heart Rhythm*. 2012; 9:242–248. [PubMed: 21939629]
277. Tokioka K, Kusano KF, Morita H, Miura D, Nishii N, Nagase S, Nakamura K, Kohno K, Ito H, Ohe T. Electrocardiographic parameters and fatal Arrhythmic events in patients with Brugada syndrome: combination of depolarization and repolarization abnormalities. *J Am Coll Cardiol*. 2014; 63:2131–2138. [PubMed: 24703917]
278. Take Y, Morita H. Fragmented QRS: What Is The Meaning? *Indian Pacing Electrophysiol J*. 2012; 12:213–225. [PubMed: 23071383]
279. Kaneko Y, Horie M, Niwano S, et al. Electrical storm in patients with Brugada syndrome is associated with early repolarization. *Circ Arrhythm Electrophysiol*. 2014
280. Takagi M, Aonuma K, Sekiguchi Y, Yokoyama Y, Aihara N, Hiraoka M, Japan Idiopathic Ventricular Fibrillation Study I. The prognostic value of early repolarization (J wave) and ST-segment morphology after J wave in Brugada syndrome: multicenter study in Japan. *Heart Rhythm*. 2013; 10:533–539. [PubMed: 23274366]
281. Rollin A, Sacher F, Gourraud JB, et al. Prevalence, characteristics, and prognosis role of type 1 ST elevation in the peripheral ECG leads in patients with Brugada syndrome. *Heart Rhythm*. 2013; 10:1012–1018. [PubMed: 23499630]
282. Marquez MF, Bisteni A, Medrano G, De Micheli A, Guevara M, Iturralde P, Colin L, Hermosillo G, Cardenas M. Dynamic electrocardiographic changes after aborted sudden death in a patient with Brugada syndrome and rate-dependent right bundle branch block. *J Electrocardiol*. 2005; 38:256–259. [PubMed: 16003712]
283. Kasanuki H, Ohnishi S, Ohtuka M, Matsuda N, Nirei T, Isogai R, Shoda M, Toyoshima Y, Hosoda S. Idiopathic ventricular fibrillation induced with vagal activity in patients without obvious heart disease. *Circulation*. 1997; 95:2277–2285. [PubMed: 9142005]
284. Nademanee K, Veerakul G, Nimmannit S, Chaowakul V, Bhuripanyo K, Likittanasombat K, Tunsanga K, Kuasirikul S, Malasit P, Tansupasawadikul S, Tatsanavivat P. Arrhythmogenic marker for the sudden unexplained death syndrome in Thai men. *Circulation*. 1997; 96:2595–2600. [PubMed: 9355899]
285. Ikeda T, Takami M, Sugi K, Mizusawa Y, Sakurada H, Yoshino H. Noninvasive risk stratification of subjects with a brugada-type electrocardiogram and no history of cardiac arrest. *AnnNoninvasiveElectrocardiol*. 2005; 10:396–403.
286. Yoshioka K, Amino M, Zareba W, Shima M, Matsuzaki A, Fujii T, Kanda S, Deguchi Y, Kobayashi Y, Ikari Y, Kodama I, Tanabe T. Identification of high-risk brugada syndrome patients by combined analysis of late potential and T-wave amplitude variability on ambulatory electrocardiograms. *Circ J*. 2013; 77:610–618. [PubMed: 23439592]

287. Tada T, Kusano KF, Nagase S, Banba K, Miura D, Nishii N, Watanabe A, Nakamura K, Morita H, Ohe T. Clinical significance of macroscopic T-wave alternans after sodium channel blocker administration in patients with Brugada syndrome. *J CardiovascElectrophysiol*. 2008; 19:56–61.
288. Uchimura-Makita Y, Nakano Y, Tokuyama T, et al. Time-domain T-wave alternans is strongly associated with a history of ventricular fibrillation in patients with Brugada syndrome. *J Cardiovasc Electrophysiol*. 2014; 25:1021–1027. [PubMed: 24761970]
289. Junttila MJ, Brugada P, Hong K, Lizotte E, de Zutter M, Sarkozy A, Brugada J, Benito B, Perkiomaki JS, Makikallio TH, Huikuri HV, Brugada R. Differences in 12-lead electrocardiogram between symptomatic and asymptomatic Brugada syndrome patients. *J CardiovascElectrophysiol*. 2008; 19:380–383.
290. Maury P, Rollin A, Sacher F, et al. Prevalence and Prognostic Role of Various Conduction Disturbances in Patients With the Brugada Syndrome. *Am J Cardiol*. 2013
291. Bigi MA, Aslani A, Shahrzad S. Clinical predictors of atrial fibrillation in Brugada syndrome. *Europace*. 2007; 9:947–950. [PubMed: 17540664]
292. Makimoto H, Nakagawa E, Takaki H, Yamada Y, Okamura H, Noda T, Satomi K, Suyama K, Aihara N, Kurita T, Kamakura S, Shimizu W. Augmented ST-segment elevation during recovery from exercise predicts cardiac events in patients with Brugada syndrome. *J Am Coll Cardiol*. 2010; 56:1576–1584. [PubMed: 21029874]
293. Castro Hevia J, Antzelevitch C, Tornes Barzaga F, Dorantes Sanchez M, Dorticos Balea F, Zayas Molina R, Quinones Perez MA, Fayad Rodriguez Y. Tpeak-Tend and Tpeak-Tend dispersion as risk factors for ventricular tachycardia/ventricular fibrillation in patients with the Brugada syndrome. *J Am Coll Cardiol*. 2006; 47:1828–1834. [PubMed: 16682308]
294. Karim Talib A, Sato N, Sakamoto N, Tanabe Y, Takeuchi T, Saijo Y, Kawamura Y, Hasebe N. Enhanced transmural dispersion of repolarization in patients with J wave syndromes. *J Cardiovasc Electrophysiol*. 2012; 23:1109–1114. [PubMed: 22612896]
295. Lambiase PD. Tpeak-Tend interval and Tpeak-Tend/QT ratio as markers of ventricular tachycardia inducibility in subjects with Brugada ECG phenotype. *Europace*. 2010; 12:158–159. [PubMed: 20045864]
296. Maury P, Sacher F, Gourraud JB, et al. Increased Tpeak-Tend interval is highly and independently related to arrhythmic events in Brugada syndrome. *Heart Rhythm*. 2015
297. Sugao M, Fujiki A, Nishida K, Sakabe M, Tsuneda T, Iwamoto J, Mizumaki K, Inoue H. Repolarization dynamics in patients with idiopathic ventricular fibrillation: pharmacological therapy with bepridil and disopyramide. *Journal of Cardiovascular Pharmacology*. 2005; 45:545–549. [PubMed: 15897781]
298. Aizawa Y, Yamakawa H, Takatsuki S, et al. Efficacy and safety of bepridil for prevention of ICD shocks in patients with Brugada syndrome and idiopathic ventricular fibrillation. *IntJ Cardiol*. 2013; 168:5083–5085. [PubMed: 23954005]
299. Masrur S, Memon S, Thompson PD. Brugada syndrome, exercise, and exercise testing. *Clin Cardiol*. 2015; 38:323–326. [PubMed: 25955277]
300. Amin AS, de Groot EA, Ruijter JM, Wilde AA, Tan HL. Exercise-induced ECG changes in Brugada syndrome. *Circ Arrhythm Electrophysiol*. 2009; 2:531–539. [PubMed: 19843921]
301. Brugada J, Brugada R, Brugada P. Pharmacological and device approach to therapy of inherited cardiac diseases associated with cardiac arrhythmias and sudden death. *JElectrocardiol*. 2000; 33(Suppl):41–47. [PubMed: 11265735]
302. Nademanee K, Veerakul G, Mower M, Likittanasombat K, Krittayapong R, Bhuripanyo K, Sitthisook S, Chaothawee L, Lai MY, Azen SP. Defibrillator Versus beta-Blockers for Unexplained Death in Thailand (DEBUT): a randomized clinical trial. *Circulation*. 2003; 107:2221–2226. [PubMed: 12695290]
303. Sacher F, Probst V, Maury P, et al. Outcome after Implantation of Cardioverter-Defibrillator in Patients with Brugada Syndrome: A Multicenter Study - Part 2. *Circulation*. 2013; 128:1739–1747. [PubMed: 23995538]
304. Sacher F, Probst V, Bessouet M, et al. Remote implantable cardioverter defibrillator monitoring in a Brugada syndrome population. *Europace*. 2009; 11:489–494. [PubMed: 19218576]

305. De Maria E, Olaru A, Cappelli S. The entirely subcutaneous defibrillator (s-icd): state of the art and selection of the ideal candidate. *Current cardiology reviews*. 2015; 11:180–186. [PubMed: 25158682]
306. Brugada P, Brugada R, Brugada J, Geelen P. Use of the prophylactic implantable cardioverter defibrillator for patients with normal hearts. *AmJCardiol*. 1999; 83:98D–100D.
307. Takagi M, Tatsumi H, Yoshiyama M. Approach to the asymptomatic patients with Brugada syndrome. *Indian Pacing ElectrophysiolJ*. 2007; 7:73–76. [PubMed: 17538698]
308. van Den Berg MP, Wilde AA, Viersma TJW, Brouwer J, Haaksma J, van der Hout AH, Stolte-Dijkstra I, Bezzina TCR, Van Langen IM, Beaufort-Krol GC, Cornel JH, Crijns HJ. Possible bradycardic mode of death and successful pacemaker treatment in a large family with features of long QT syndrome type 3 and Brugada syndrome. *JCardiovascElectrophysiol*. 2001; 12:630–636.
309. Bertomeu-Gonzalez V, Ruiz-Granell R, Garcia-Civera R, Morell-Cabedo S, Ferrero A. Syncopal monomorphic ventricular tachycardia with pleomorphism, sensitive to antitachycardia pacing in a patient with Brugada syndrome. *Europace*. 2006; 8:1048–1050. [PubMed: 17098780]
310. Lee KL, Lau C, Tse H, Wan S, Fan K. Prevention of ventricular fibrillation by pacing in a man with Brugada syndrome. *J Cardiovasc Electrophysiol*. 2000; 11:935–937. [PubMed: 10969759]
311. Cortez-Dias N, Placido R, Marta L, Bernardes A, Sobral S, Carpinteiro L, de SJ. Epicardial ablation for prevention of ventricular fibrillation in a patient with Brugada syndrome. *RevPortCardiol*. 2014; 33:305–305.
312. Shah AJ, Hocini M, Lamaison D, Sacher F, Derval N, Haissaguerre M. Regional substrate ablation abolishes Brugada syndrome. *J Cardiovasc Electrophysiol*. 2011; 22:1290–1291. [PubMed: 21457386]
313. Brugada J, Brugada R, Brugada P. Right bundle-branch block and ST-segment elevation in leads V₁ through V₃. A marker for sudden death in patients without demonstrable structural heart disease. *Circulation*. 1998; 97:457–460. [PubMed: 9490240]
314. Chinushi M, Aizawa Y, Ogawa Y, Shiba M, Takahashi K. Discrepant drug action of disopyramide on ECG abnormalities and induction of ventricular arrhythmias in a patient with Brugada syndrome. *JElectrocardiol*. 1997; 30:133–136. [PubMed: 9141608]
315. Minoura Y, Di Diego JM, Barajas-Martinez H, Zygmunt AC, Hu D, Sicouri S, Antzelevitch C. Ionic and cellular mechanisms underlying the development of acquired Brugada syndrome in patients treated with antidepressants. *J Cardiovasc Electrophysiol*. 2012; 23:423–432. [PubMed: 22034916]
316. Minoura Y, Panama BK, Nesterenko VV, Betzenhauser M, Barajas-Martinez H, Hu D, Di Diego JM, Antzelevitch C. Effect of Wenxin Keli and quinidine to suppress arrhythmogenesis in an experimental model of Brugada syndrome. *Heart Rhythm*. 2013; 10:1054–1062. [PubMed: 23499631]
317. Johnson P, Lesage A, Floyd WL, Young WG Jr, Sealy WC. Prevention of ventricular fibrillation during profound hypothermia by quinidine. *Ann Surg*. 1960; 151:490–495. [PubMed: 14407499]
318. Belhassen B, Shapira I, Shoshani D, Paredes A, Miller H, Laniado S. Idiopathic ventricular fibrillation: inducibility and beneficial effects of class I antiarrhythmic agents. *Circulation*. 1987; 75:809–816. [PubMed: 3829343]
319. Belhassen B, Viskin S, Antzelevitch C. The Brugada syndrome: is an implantable cardioverter defibrillator the only therapeutic option? *Pacing Clin Electrophysiol*. 2002; 25:1634–1640. [PubMed: 12494624]
320. Alings M, Dekker L, Sadee A, Wilde A. Quinidine induced electrocardiographic normalization in two patients with Brugada syndrome. *Pacing ClinElectrophysiol*. 2001; 24:1420–1422.
321. Belhassen B, Viskin S. Pharmacologic approach to therapy of Brugada syndrome: quinidine as an alternative to ICD therapy?. In: Antzelevitch, C.; Brugada, P.; Brugada, J.; Brugada, R., editors. *The Brugada Syndrome: From Bench to Bedside*. Blackwell Futura; Oxford: 2004. p. 202–211.
322. Viskin S, Wilde AA, Tan HL, Antzelevitch C, Shimizu W, Belhassen B. Empiric quinidine therapy for asymptomatic Brugada syndrome: time for a prospective registry. *Heart Rhythm*. 2009; 6:401–404. [PubMed: 19251219]

323. Belhassen B, Glick A, Viskin S. Excellent long-term reproducibility of the electrophysiologic efficacy of quinidine in patients with idiopathic ventricular fibrillation and Brugada syndrome. *Pacing Clin Electrophysiol.* 2009; 32:294–301. [PubMed: 19272057]
324. Marquez MF, Bonny A, Hernandez-Castillo E, De SA, Gomez-Flores J, Nava S, Hidden-Lucet F, Iturralde P, Cardenas M, Tonet J. Long-term efficacy of low doses of quinidine on malignant arrhythmias in Brugada syndrome with an implantable cardioverter-defibrillator: A case series and literature review. *Heart Rhythm.* 2013; 9:1995–2000. [PubMed: 23059185]
325. Pellegrino PL, Di BM, Brunetti ND. Quinidine for the management of electrical storm in an old patient with Brugada syndrome and syncope. *Acta Cardiol.* 2013; 68:201–203. [PubMed: 23705565]
326. Probst V, Allouis M, Sacher F, et al. Progressive cardiac conduction defect is the prevailing phenotype in carriers of a Brugada syndrome SCN5A mutation. *J Cardiovasc Electrophysiol.* 2006; 17:270–275. [PubMed: 16643399]
327. Schweizer PA, Becker R, Katus HA, Thomas D. Successful acute and long-term management of electrical storm in Brugada syndrome using orciprenaline and quinine/quinidine. *Clin Res Cardiol.* 2010; 99:467–470. [PubMed: 20221832]
328. Marquez MF, Rivera J, Hermosillo AG, Iturralde P, Colin L, Moragrega JL, Cardenas M. Arrhythmic storm responsive to quinidine in a patient with Brugada syndrome and vasovagal syncope. *Pacing ClinElectrophysiol.* 2005; 28:870–873.
329. Viskin S, Antzelevitch C, Marquez MF, Belhassen B. Quinidine: a valuable medication joins the list of 'endangered species'. *Europace.* 2007; 12:1105–1106. [PubMed: 17761793]
330. Rosso R, Glick A, Glikson M, et al. Outcome after implantation of cardioverter defibrillator in patients with Brugada syndrome: a multicenter Israeli study (ISRABRU). *IsrMed AssocJ.* 2008; 10:435–439.
331. Mok NS, Chan NY, Chi-Suen CA. Successful use of quinidine in treatment of electrical storm in Brugada syndrome. *Pacing ClinElectrophysiol.* 2004; 27:821–823.
332. Belhassen B, Rahkovich M, Michowitz Y, Glick A, Viskin S. Management of Brugada Syndrome: A 33-Year Experience Using Electrophysiologically-Guided Therapy with Class 1A Antiarrhythmic Drugs. *Circ Arrhythm Electrophysiol.* 2015
333. Bouzeman A, Traulle S, Messali A, Extramiana F, Denjoy I, Narayanan K, Marijon E, Hermida JS, Leenhardt A. Long-term follow-up of asymptomatic Brugada patients with inducible ventricular fibrillation under hydroquinidine. *Europace.* 2014; 16:572–577. [PubMed: 24068450]
334. Antzelevitch C. The Brugada syndrome: ionic basis and arrhythmia mechanisms. *JCardiovasc Electrophysiol.* 2001; 12:268–272. [PubMed: 11232628]
335. Kyriazis K, Bahlmann E, van der SH, Kuck KH. Electrical storm in Brugada syndrome successfully treated with orciprenaline; effect of low-dose quinidine on the electrocardiogram. *Europace.* 2009; 11:665–666. [PubMed: 19346290]
336. Shimizu W, Matsuo K, Takagi M, Tanabe Y, Aiba T, Taguchi A, Suyama K, Kurita T, Aihara N, Kamakura S. Body surface distribution and response to drugs of ST segment elevation in Brugada syndrome: clinical implication of eighty-seven-lead body surface potential mapping and its application to twelve-lead electrocardiograms. *J Cardiovasc Electrophysiol.* 2000; 11:396–404. [PubMed: 10809492]
337. Tanaka H, Kinoshita O, Uchikawa S, et al. Successful prevention of recurrent ventricular fibrillation by intravenous isoproterenol in a patient with Brugada syndrome. *Pacing ClinElectrophysiol.* 2001; 24:1293–1294.
338. Maury P, Couderc P, Delay M, Boveda S, Brugada J. Electrical storm in Brugada syndrome successfully treated using isoprenaline. *Europace.* 2004; 6:130–133. [PubMed: 15018871]
339. Maury P, Hocini M, Haissaguerre M. Electrical storms in Brugada syndrome: review of pharmacologic and ablative therapeutic options. *Indian Pacing Electrophysiol J.* 2005; 5:25–34. [PubMed: 16943940]
340. Jongman JK, Jekes-Bruin N, Ramdat Misier AR, Beukema WP, Delnoy PP, Oude LH, Dambrink JH, Hoorntje JC, Elvan A. Electrical storms in Brugada syndrome successfully treated with isoproterenol infusion and quinidine orally. *NethHeart J.* 2007; 15:151–155.

341. Sharif-Kazemi MB, Emkanjoo Z, Tavoosi A, Kafi M, Kheirkhah J, Alizadeh A, Sadr-Ameli MA. Electrical storm in Brugada syndrome during pregnancy. *Pacing Clin Electrophysiol.* 2011; 34:e18–e21. [PubMed: 20353417]
342. Furniss G. Isoprenaline and quinidine to calm Brugada VF storm. *BMJ CaseRep.* 2012
343. Roten L, Derval N, Sacher F, et al. Heterogeneous response of J wave syndromes to beta-adrenergic stimulation. *Heart Rhythm.* 2012; 9:1970–1976. [PubMed: 22864265]
344. Kanlop N, Chattipakorn S, Chattipakorn N. Effects of cilostazol in the heart. *J Cardiovasc Med (Hagerstown).* 2011; 12:88–95. [PubMed: 21200326]
345. Bai Y, Muqier, Murakami H, Iwasa M, Sumi S, Yamada Y, Ushikoshi H, Aoyama T, Nishigaki K, Takemura G, Uno B, Minatoguchi S. Cilostazol protects the heart against ischaemia reperfusion injury in a rabbit model of myocardial infarction: focus on adenosine, nitric oxide and mitochondrial ATP-sensitive potassium channels. *Clin Exp Pharmacol Physiol.* 2011; 38:658–665.
346. Abud A, Bagattin D, Goyeneche R, Becker C. Failure of cilostazol in the prevention of ventricular fibrillation in a patient with Brugada syndrome. *J CardiovascElectrophysiol.* 2006; 17:210–212.
347. Szel T, Koncz I, Antzelevitch C. Cellular mechanisms underlying the effects of milrinone and cilostazol to suppress arrhythmogenesis associated with Brugada syndrome. *Heart Rhythm.* 2013; 10:1720–1727. [PubMed: 23911896]
348. Murakami M, Nakamura K, Kusano KF, et al. Efficacy of low-dose bepridil for prevention of ventricular fibrillation in patients with Brugada syndrome with and without SCN5A mutation. *J Cardiovasc Pharmacol.* 2010; 56:389–395. [PubMed: 20625312]
349. Kang L, Zheng MQ, Morishima M, Wang Y, Kaku T, Ono K. Bepridil up-regulates cardiac Na⁺ channels as a long-term effect by blunting proteasome signals through inhibition of calmodulin activity. *Br J Pharmacol.* 2009; 157:404–414. [PubMed: 19371335]
350. Fish JM, Welchons DR, Kim YS, Lee SH, Ho WK, Antzelevitch C. Dimethyl lithospermate B, an extract of danshen, suppresses arrhythmogenesis associated with the Brugada syndrome. *Circulation.* 2006; 113:1393–1400. [PubMed: 16534004]
351. Kanlop N, Shinlapawittayatorn K, Sungnoon R, Weerateerangkul P, Chattipakorn S, Chattipakorn N. Cilostazol attenuates ventricular arrhythmia induction and improves defibrillation efficacy in swine. *Can J Physiol Pharmacol.* 2010; 88:422–428. [PubMed: 20555410]
352. Endoh M, Yanagisawa T, Taira N, Blinks JR. Effects of new inotropic agents on cyclic nucleotide metabolism and calcium transients in canine ventricular muscle. *Circulation.* 1986; 73:III117–III133. [PubMed: 2417745]
353. Rapundalo ST, Grupp I, Grupp G, Abdul MM, Solaro RJ, Schwartz A. Myocardial actions of milrinone: characterization of its mechanism of action. *Circulation.* 1986; 73:III134–III144. [PubMed: 2417746]
354. Atarashi H, Endoh Y, Saitoh H, Kishida H, Hayakawa H. Chronotropic effects of cilostazol, a new antithrombotic agent, in patients with bradyarrhythmias. *J Cardiovasc Pharmacol.* 1998; 31:534–539. [PubMed: 9554801]
355. Matsui K, Kiyosue T, Wang JC, Dohi K, Arita M. Effects of pimobendan on the L-type Ca²⁺ current and developed tension in guinea-pig ventricular myocytes and papillary muscle: comparison with IBMX, milrinone, and cilostazol. *Cardiovasc Drugs Ther.* 1999; 13:105–113. [PubMed: 10372225]
356. Aizawa Y, Chinushi M, Hasegawa K, et al. Electrical storm in idiopathic ventricular fibrillation is associated with early repolarization. *J Am Coll Cardiol.* 2013; 62:1015–1019. [PubMed: 23747791]
357. Nakagawa K, Nagase S, Morita H, Ito H. Left ventricular epicardial electrogram recordings in idiopathic ventricular fibrillation with inferior and lateral early repolarization. *Heart Rhythm.* 2013; 11:314–317. [PubMed: 24184784]
358. Morita H, Zipes DP, Morita ST, Wu J. Temperature modulation of ventricular arrhythmogenicity in a canine tissue model of Brugada syndrome. *Heart Rhythm.* 2007; 4:188–197. [PubMed: 17275755]
359. Probst V, Veltmann C, Eckardt L, et al. Long-term prognosis of patients diagnosed with Brugada syndrome: results from the FINGER Brugada Syndrome Registry. *Circulation.* 2010; 121:635–643. [PubMed: 20100972]

360. Takagi M, Aonuma K, Sekiguchi Y, Yokoyama Y, Aihara N, Hiraoka M. The prognostic value of early repolarization (J wave) and ST-segment morphology after J wave in Brugada syndrome: Multicenter study in Japan. *Heart Rhythm*. 2013; 10:533–539. [PubMed: 23274366]
361. Brugada J, Brugada R, Brugada P. Determinants of sudden cardiac death in individuals with the electrocardiographic pattern of Brugada syndrome. *Pacing ClinElectrophysiol*. 2003; 26:957.
362. Huang Z, Patel C, Li W, Xie Q, Wu R, Zhang L, Tang R, Wan X, Ma Y, Zhen W, Gao L, Yan GX. Role of signal-averaged electrocardiograms in arrhythmic risk stratification of patients with Brugada syndrome: a prospective study. *Heart Rhythm*. 2009; 6:1156–1162. [PubMed: 19632627]
363. Priori SG. Natural History of Brugada Syndrome: Insights for Risk Stratification and Management. *Circulation*. 2002; 105:1342–1347. [PubMed: 11901046]
364. Kamakura S, Ohe T, Nakazawa K, et al. Long-term prognosis of probands with Brugada-pattern ST-elevation in leads V1-V3. *Circ ArrhythmElectrophysiol*. 2009; 2:495–503.
365. Brugada J, Brugada R, Brugada P. Determinants of sudden cardiac death in individuals with the electrocardiographic pattern of Brugada syndrome and no previous cardiac arrest. *Circulation*. 2003; 108:3092–3096. [PubMed: 14623800]
366. Cappato R, Furlanello F, Giovinazzo V, et al. J wave, QRS slurring and ST elevation in athletes with cardiac arrest in the absence of heart disease: marker of risk, or innocent bystander? *Circ Arrhythm Electrophysiol*. 2010; 3:305–311. [PubMed: 20511538]
367. Perez-Riera AR, Abreu LC, Yanowitz F, Barros RB, Femenia F, McIntyre WF, Baranchuk A. "Benign" early repolarization versus malignant early abnormalities: Clinical-electrocardiographic distinction and genetic basis. *Cardiol J*. 2012; 19:337–346. [PubMed: 22825893]

Figure 1.

Three types of ST segment elevation associated with Brugada syndrome. Only Type 1 is diagnostic of BrS.

Figure 2.

Schematic showing the three categories of overlap syndromes resulting from genetic defects resulting in loss of function of sodium (I_{Na}) and/or calcium (I_{Ca}) channel current. In the absence of prominent I_{to} or I_{K-ATP} , loss-of-function mutations in the inward currents result in various manifestations of conduction disease. In the presence of prominent I_{to} or I_{K-ATP} , loss-of-function mutations in inward currents cause conduction disease as well as the J wave syndromes (Brugada and Early Repolarization Syndromes). Early Repolarization Syndrome is believed to be caused by loss-of-function mutations of inward current in the presence of prominent I_{to} in certain regions of the left ventricle (LV), particularly the inferior wall of the LV. The genetic defects that contribute to BrS and ERS can also contribute to the development of long QT and conduction system disease, in some cases causing multiple expressions of these overlap syndromes. In some cases, structural defects contribute to the phenotype.

Figure 3.

Different manifestations of Early Repolarization. **A.** The J wave may be distinct or appear as a slur. In the latter case, part of the J wave is buried inside the QRS, resulting in an elevation of J_0 . Patients with a distinct J wave have a worse prognosis than patients with a slurred J wave. **B.** The ST segment may be upsloping, horizontal or descending. Horizontal and descending ST segments are associated with a worse prognosis.

Figure 4.

Prevalence and arrhythmic risk associated with the appearance of ECG J waves and clinical manifestations of Brugada and Early Repolarization syndromes. The yellow highlighted region estimates the prevalence of the J wave syndromes. J waves in the lateral ECG leads have a high prevalence but are associated with a very low arrhythmic risk in a relatively small fraction of the cohort of individuals displaying J waves. On the other extreme, J waves appearing globally in the ECG have a very low prevalence, but are associated with a very high level of arrhythmic risk in a large fraction of the cohort presenting with J waves. Likewise, individuals displaying rapidly ascending ST segment elevation have a high prevalence but low risk, whereas subjects resuscitated from cardiac arrest have a very low prevalence by the highest level of arrhythmic risk.

Figure 5. Global Early Repolarization (Type 3 ER). J waves are apparent in the inferior, lateral and anterior (right precordial) leads.

Figure 6.

Indications for therapy of patients with Brugada syndrome. Recommendations with Class designations are taken from Priori SG, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes,⁸ and Priori SG, Blomstrom-Lundqvist C, Mazzanti A, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death.⁹ Recommendations without Class designations are derived from unanimous consensus of the authors.

ES= extrastimuli at RV apex, ICD=implantable cardioverter defibrillator. ILR=implantable loop recorder, NAR= nocturnal agonal respiration, VT=ventricular tachycardia

Figure 7.

Indications for therapy of patients with Early Repolarization Syndrome. Recommendations with Class designations are taken from Priori SG, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes. *Heart Rhythm* Dec 2013;10:1932-1963⁸ and Priori SG, Blomstrom-Lundqvist C, Mazzanti A, et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death. *Eur. Heart J* Aug 29 2015⁹ Recommendations without Class designations are derived from unanimous consensus of the authors.

ER=early repolarization, ICD=implantable cardioverter defibrillator, ILR=implantable loop recorder, NAR=nocturnal agonal respiration, VT=ventricular tachycardia.

Table 1

Drugs used to unmask the Brugada electrocardiogram.

Drug	Dose	Administration
Ajmaline	1 mg/kg over 10 minutes	Intravenous
Flecainide	2 mg/kg over 10 minutes 200-300 mg	Intravenous Oral (>1 hour)
Procainamide	10 mg/kg over 10 minutes	Intravenous
Pilsicainide	1 mg/kg over 10 minutes	Intravenous

Table 2**Proposed Shanghai Score System for Diagnosis of Brugada Syndrome**

I. ECG (12-Lead/Ambulatory)	Points
A. Spontaneous Type 1 Brugada ECG pattern at nominal or high leads	3.5
B. Fever-induced Type 1 Brugada ECG pattern at nominal or high leads	3
C. Type 2 or 3 Brugada ECG pattern that converts w/ provocative drug challenge	2
<i>* Only award points once for highest score within this category. One item from this category must apply.</i>	
II. Clinical History*	
A. Unexplained cardiac arrest or documented VF/polymorphic VT	3
B. Nocturnal agonal respirations	2
C. Suspected arrhythmic syncope	2
D. Syncope of unclear mechanism/unclear etiology	1
E. Atrial flutter/fibrillation in patients < 30 years without alternative etiology	0.5
<i>* Only award points once for highest score within this category.</i>	
III. Family History	
A. 1 st or 2 nd degree relative with definite BrS	2
B. Suspicious SCD (fever, nocturnal, Brugada aggravating drugs) in a 1 st or 2 nd degree relative	1
C. Unexplained SCD < 45 years in first/second degree relative with negative autopsy	0.5
<i>* Only award points once for highest score within this category.</i>	
IV. Genetic Test Result	
A. Probable pathogenic mutation in BrS-susceptibility gene	0.5
SCORE (Requires at least 1 ECG finding)	
> 3.5 points – Probable/Definite BrS	
2 – 3 points – Possible BrS	
< 2 points – Non-Diagnostic	

Table 3

Differential diagnosis and modulating factors in Brugada Syndrome.

A. Differential diagnosis	B. Modulating factors
<ul style="list-style-type: none"> • Atypical right bundle branch block • Ventricular hypertrophy • Early repolarization (especially in athletes) • Acute pericarditis/myocarditis • Acute myocardial ischemia or infarction (especially of the right ventricle) • Pulmonary thromboembolism • Prinzmetal angina • Dissecting aortic aneurism • Central and autonomic nervous systems abnormalities • Duchenne's muscular dystrophy • Friedreich's ataxia • Spinobulbar Muscular Atrophy • Myotonic Dystrophy • Arrhythmogenic right ventricular dysplasia • Mechanical compression of the right ventricular outflow tract (e.g., <i>pectus excavatum</i>, mediastinal tumor, hemopericardium, • Hypothermia • Post-defibrillation ECG 	<ul style="list-style-type: none"> • Electrolyte abnormalities: <ul style="list-style-type: none"> ○ Hyperkalemia ○ Hypokalemia ○ Hypercalcemia ○ Hyponatremia • Temperature: hyperthermia (fever), hypothermia • Hypertestosteronemia • Treatment with: <ul style="list-style-type: none"> ○ Antiarrhythmic drugs: sodium channel blockers (class IC, class IA), calcium antagonists, beta-blockers ○ Antianginal drugs: calcium antagonists, nitrates, potassium channel openers ○ Psychotropic drugs: tricyclic/tetracyclic antidepressants, phenothiazines, selective serotonin reuptake inhibitor, lithium, benzodiazepines ○ Anesthetics/analgesics: propofol, bupivacaine, procaine ○ Others: histamine H1 antagonist, alcohol intoxication, cocaine, cannabis, ergonovine

Table 4

Differential diagnosis of ER Pattern

Other causes of ER pattern include:	
•	Juvenile ST pattern
•	Pericardial disease (pericarditis, pericardial cyst or pericardial tumor)
•	Hypothermia
•	Hyperthermia
•	Myocardial tumor (Lipoma)
•	Hypertensive heart disease
•	Athlete's heartMyocardial ischemia
•	STEMI (i.e., anteroseptal MI)
•	Fragmented QRS (terminal notching)
•	Hypocalcemia
•	Hyperpotassemia
•	Thymoma
•	Aortic Dissection
•	ARVC
•	TakotsubO cardiomyopathy
•	Neurological causes (intracerebral bleeding, acute brain injury)
•	Myocarditis
•	Chagas disease
•	Cocaine use

Table 5

Proposed Shanghai score system for diagnosis of Early Repolarization Syndrome

I. Clinical History	
A. Unexplained cardiac arrest, documented VF or polymorphic VT	3
B. Suspected arrhythmic syncope	2
C. Syncope of unclear mechanism/unclear etiology	1
<i>* Only award points once for highest score within this category</i>	
II. 12-Lead ECG	
A. ER 0.2 mV in > 2 inferior and/or lateral ECG leads with horizontal/descending ST segment	2
B. Dynamic changes in J-point elevation (> 0.1 mV) in > 2 inferior and/or lateral ECG leads	1.5
C. 0.1 mV J-point elevation in at least 2 inferior and/or lateral ECG leads	1
<i>* Only award points once for highest score within this category</i>	
III. Ambulatory ECG monitoring	
A. Short coupled PVCs with R on ascending limb or peak of T wave	2
IV. Family History	
A. Relative with definite ERS	2
B. > 2 first-degree relatives with a II.A. ECG pattern	2
C. A first-degree relative with a II.A. ECG pattern	1
D. Unexplained sudden cardiac death < 45 years in a first/second-degree relative	0.5
<i>* Only award points once for highest score within this category</i>	
V. Genetic Test Result	
A. Probable pathogenic ERS-susceptibility mutation	0.5
SCORE (Requires at least one ECG finding)	
> 5 points – Probable/Definite ERS	
3 – 4.5 points – Possible ERS	
< 3 points – Non-Diagnostic	

Table 6

Similarities and differences between Brugada and Early Repolarization Syndromes and possible underlying mechanisms

	BrS	ERS	Possible Mechanism(s)
Similarities between BrS and ERS			
Male Predominance	Yes (>75%)	Yes (>80%)	Testosterone modulation of ion currents underlying the epicardial AP notch
Average age of first event	30-50	30-50	
Associated with mutations or rare variants in <i>KCNJ8</i>, <i>CACNA1C</i>, <i>CACNB2</i>, <i>CACNA2D</i>, <i>SCN5A</i>, <i>ABCC9</i>, <i>SCN10A</i>	Yes	Yes	Gain of function in outward currents (I_{K-ATP}) or loss of function in inward currents (I_{Ca} or I_{Na})
Relatively short QT intervals in subjects with Ca channel mutations	Yes	Yes	Loss of function of I_{Ca}
Dynamicity of ECG	High	High	Autonomic modulation of ion channel currents underlying early phases of the epicardial AP
VF often occurs during sleep or at a low level of physical activity	Yes	Yes	Higher level of vagal tone and higher levels of I_{to} at the slower heart rates
VT/VF trigger	Short-coupled PVC	Short-coupled PVC	Phase 2 reentry
Ameliorative response to quinidine and bepridil	Yes	Yes	Inhibition of I_{to} and possible vagolytic effect
Ameliorative response to isoproterenol denopamine and milrinone	Yes	Yes	Increased I_{Ca} and faster heart rate
Ameliorative response to cilostazol	Yes	Yes	Increased I_{Ca} , reduced I_{to} and faster heart rate
Ameliorative response to pacing	Yes	Yes	Reduced availability of I_{to} due to slow recovery from inactivation
Vagally mediated accentuation of ECG pattern	Yes	Yes	Direct effect to inhibit I_{Ca} and indirect effect to increase I_{to} (due to slowing of heart rate)
Effect of sodium channel blockers on unipolar epicardial electrogram	Augmented J waves	Augmented J wave	Outward shift of balance of current in the early phases of the epicardial action potential
Fever	Augmented J waves	Augmented J waves (rare)	Accelerated inactivation of I_{Na} and accelerated recovery of I_{to} from inactivation.
Hypothermia	Augmented J waves mimicking BrS	Augmented J waves	Slowed activation of I_{Ca} , leaving I_{to} unopposed. Increased phase 2 reentry, but reduced pVT due to prolongation of APD ^{R358}
Differences between BrS and ERS			
Region most involved	RVOT	Inferior LV wall	Higher levels of I_{to} and/or differences in conduction
Leads affected	V1-V3	II, II a, VF, V4, V5, V6; I, aVL, Both:	

	BrS	ERS	Possible Mechanism(s)
		infero-lateral	
Regional difference in prevalence			Europe: BrS = ERS Asia: BrS > ERS
Incidence of late potential in SAECD	Higher	Lower	
Prevalence of atrial fibrillation	Higher	Lower	
Effect of sodium channel blockers on the surface ECD	Increased J wave manifestation	Reduced J wave Manifestation	Reduction of J wave in the setting of ER is thought to be due largely to prolongation of QRS. Accentuation of repolarization defects predominates in BrS, whereas accentuation of depolarization defects predominates in ERS.
Structural changes—including mild fibrosis and reduced expression of Cx43 in RVOT or fibro-fatty infiltration in cases of ARVC. Imaging studies have also revealed wall motion abnormalities and mild dilatation in the region of the RVOT.	Higher in some forms of the syndrome	Unknown	Some investigators have hypothesized that some of these changes may be the result of, rather than the cause of, the BrS substrate, which may create a hibernation-like state due to loss of contractility in the RVOT secondary to loss of the AP dome.
Structural changes—including mild fibrosis and reduced expression of Cx43 in RVOT or fibro-fatty infiltration in cases of ARVC. Imaging studies have also revealed wall motion abnormalities and mild dilatation in the region of the RVOT.	Higher in some forms of the syndrome	Unknown	Some investigators have hypothesized that some of these changes may be the result of, rather than the cause of, the BrS substrate, which may create a hibernation-like state due to loss of contractility in the RVOT secondary to loss of the AP dome.

RVOT=right ventricular outflow tract, AP=action potential; PVC=premature ventricular contraction

Table 7

Gene defects associated with the Early Repolarization (ERS) and Brugada (BrS) Syndromes

Genetic Defects Associated with ERS				
	Locus	Gene/Protein	Ion Channel	% of Probands
ERS1	12p11.23	<i>KCNJ8, Kir6.1</i>	↑ I _{K-ATP}	Rare
ERS2	12p13.3	<i>CACNA1C, Ca_v1.2</i>	↓ I _{Ca}	4.1%
ERS3	10p12.33	<i>CACNB2b, Ca_vβ2b</i>	↓ I _{Ca}	8.3%
ERS4	7q21.11	<i>CACNA2D1, Ca_vα2σ1</i>	↓ I _{Ca}	4.1%
ERS5	12p12.1	<i>ABCC9, SUR2A</i>	↑ I _{K-ATP}	Rare
ERS6	3p21	<i>SCN5A, Na_v1.5</i>	↓ I _{Na}	Rare
ERS7	3p22.2	<i>SCN10A, Na_v1.8</i>	↓ I _{Na}	Rare
Genetic Defects Associated with BrS				
	Locus	Gene/Protein	Ion Channel	% of Probands
BrS1	3p21	<i>SCN5A, Na_v1.5</i>	↓ I _{Na}	11-28%
BrS2	3p24	<i>GPD1L</i>	↓ I _{Na}	Rare
BrS3	12p13.3	<i>CACNA1C, Ca_v1.2</i>	↓ I _{Ca}	6.6%
BrS4	10p12.33	<i>CACNB2b, Ca_vβ2b</i>	↓ I _{Ca}	4.8%
BrS5	19q13.1	<i>SCN1B, Na_vβ1</i>	↓ I _{Na}	1.1%
BrS6	11q13-14	<i>KCNE3, MiRP2</i>	↑ I _{to}	Rare
BrS7	11q23.3	<i>SCN3B, Na_vβ3</i>	↓ I _{Na}	Rare
BrS8	12p11.23	<i>KCNJ8, Kir6.1</i>	↑ I _{K-ATP}	2%
BrS9	7q21.11	<i>CACNA2D1, Ca_vα2σ1</i>	↓ I _{Ca}	1.8%
BrS10	1p13.2	<i>KCND3, K_v4.3</i>	↑ I _{to}	Rare
BrS11	17p13.1	<i>RANGRF, MOG1</i>	↓ I _{Na}	Rare
BrS12	3p21.2-p14.3	<i>SLMAP</i>	↓ I _{Na}	Rare
BrS13	12p12.1	<i>ABCC9, SUR2A</i>	↑ I _{K-ATP}	Rare
BrS14	11q23	<i>SCN2B, Na_vβ2</i>	↓ I _{Na}	Rare
BrS15	12p11	<i>PKP2, Plakophilin-2</i>	↓ I _{Na}	Rare
BrS16	3q28	<i>FGF12, FHAF1</i>	↓ I _{Na}	Rare
BrS17	3p22.2	<i>SCN10A, Na_v1.8</i>	↓ I _{Na}	5-16.7%
BrS18	6q	<i>HEY2 (transcriptional factor)</i>	↑ I _{Na}	Rare

Listed in chronological order of their discovery

Table 8

Differential diagnosis of J wave vs. Intra-ventricular Conduction Defect-mediated Notch Syndromes (IVCD)

	J wave	IVCD-induced end QRS notch
Male Predominance	Yes	No
Average Age at Initial Presentation	Young adults	Older adults
Most common Morphology	Dome-like smooth appearance	Relatively sharp appearance
Response to Change in Heart Rate	Bradycardia- and pause-dependent augmentation of J wave, which may be accompanied by T wave inversion	Tachycardia and prematurity-dependent augmentation of the notch
Structural Heart Diseases	Rare	Common History of myocardial infarction and/or cardiomyopathy

TABLE 9

Clinical variables associated with an increased risk of major arrhythmic events in Brugada Syndrome *

Variable	Pts N	Prevention	Study end-point	Multivariable analysis HR (95% CI), p	Reference
History					
Previous VF	93	P/S	SD, cardiac arrest or sustained VT/VF (N=25)	N/A (N/A), 0.005	Makimoto ^{R292}
Cardiac arrest	1029	P/S	SD (N=7), appropriate ICD shocks (N=44) or sustained VT/VF (N=0)	11 (4.8-24.3), 0.001	Probst ^{R359}
Syncope or cardiac arrest	460	P/S	VF or SD (N=38)	12.7 (4.5-53.4), <0.0001	Takagi ^{R360}
Syncope of unknown origin	547	P	SD (N=16), VF (N=29)	2.5 (1.2-5.3), 0.017	Brugada ^{R361}
Syncope	44	P/S	SCD (N=5), polymorphic VT or VF (recorded by ECG, Holter, or ICD) (N=7) or syncope of unknown etiology (11)	3.6 (1.09-11.7), 0.035	Huang ^{R362}
Syncope + spontaneous type 1 ECG	200	P/S	VF or SD from birth (N=22)	6.4 (1.9-21), <0.002	Priori ^{R363}
Syncope of probable arrhythmic origin	1029	P/S	SD (N=7), appropriate ICD shocks (N=44) or sustained VT/VF (N=0)	3.4 (1.6-7.4), 0.002	Kamakura ^{R364}
Syncope	320	P	SD (N=3), appropriate ICD shocks (N=14) or sustained VT/VF (N=0)	2.8 (1.1-8.1), 0.03	Delise ^{R262}
Syncope + spontaneous type 1 ECG	308	P	VF (N=1) or appropriate ICD intervention (N=13)	4.2 (1.4-12.8), 0.012	Priori ^{R49}
Ventricular refractoriness					
Ventricular refractory period <200ms	308	P	VF (N=1) or appropriate ICD intervention (N=13)	3.9 (1.03-12.8), 0.045	Priori ^{R49}
ECG characteristics					
Spontaneous type 1 ECG	1029	P/S	SD (N=7), appropriate ICD shocks (N=44) or sustained VT/VF (N=0)	1.8 (1.03-3.3), 0.04	Probst ^{R247}
Spontaneous type 1 ECG	320	P	SD (N=3), appropriate ICD shocks (N=14) or sustained VT/VF (N=0)	6.2 (1.8-40), 0.002	Delise ^{R262}
QRS fragmentation (2 or more spikes within the QRS complex in leads V1 to V3)	308	P	VF (N=1) or appropriate ICD intervention (N=13)	4.9 (1.5-1.8), 0.007	Priori ^{R49}
Family history of sudden cardiac death at age 45 years	330	P	VF (N=56), syncope (N=67), or asymptomatic (N=207)	3.28 (1.4-7.6), 0.005	Kamakura ^{R364}
J wave in inferior and lateral leads	330	P	VF (N=56), syncope (N=67), or asymptomatic (N=207)	2.66 (1.1-6.7), 0.005	Kamakura ^{R364}
QRS duration > 90 ms in lead V2	460	P/S	VF or SD (N=38)	3.6 (1.4-12.2), 0.007	Takagi ^{R360}
Horizontal ST segment after J wave (ST-segment elevation 0.1 mV within 100 ms after J point and continued as a flat ST	460	P/S	VF or SD (N=38)	>10 (1.9-20.2), 0.02	Takagi ^{R360}

Variable	Pts N	Prevention	Study end-point	Multivariable analysis HR (95% CI), p	Reference
<i>segment until the onset of the T-wave in 1 lead with J-wave)</i>					
Late potentials (root mean square voltage of the terminal 40 ms of the filtered QRS complex <20 μ V + duration of low amplitude signals <40 μ V of QRS in the terminal filtered QRS complex >38 ms)	44	P/S	SCD (N=5), polymorphic VT or VF (recorded by ECG, Holter, or ICD) (N=7) or syncope of unknown etiology (11)	10.9 (1.1–104), 0.038	Huang ^{R362}
ST-segment augmentation at early recovery of exercise test (ST-segment amplitude increase 0.05 mV in at least 1 of V1 to V3 leads at 1 to 4 min at recovery compared with the ST-segment amplitude at pre-exercise)	93	P/S	SD, cardiac arrest or sustained VT/VF (N=25).	N/A (N/A), 0.007	Makimoto ^{R292}

P=primary prevention patients only, P/S=primary and secondary prevention patients, SD=sudden death, VF=ventricular fibrillation, VT=ventricular tachycardia

* The list includes predictor variables that have been associated with an increased risk of major arrhythmic events (i.e, SCD, appropriate ICD interventions, or ICD therapy on fast VT/VF) in at least one published multivariable analysis in prospective studies.

TABLE 10

Prognostic value of programmed ventricular stimulation resulting from multivariate analysis in large multicenter studies on Brugada syndrome

N.	Prevention	Inducibility %	Multivariable HR (95% CI), p	Reference
408	P	40%	5.88 (2.0-16.7), <0.001	Brugada ^{R365}
308	P	41%	0.89 (0.3-2.6), 0.84	Priori ^{R49}
638	P/S	62%	N/A, 0.48	Probst ^{R247}
245	P	39%	Not tested on multivariable analysis, but an analysis based on C-statistics demonstrated that EPS results in combination with other risk factors provided additional value for risk stratification	Delise ^{R262}
334	P/S	67%	0.63 (0.3-1.3), 0.20 [*]	Takagi ^{R360}
1312 [*]	P/S	42%	2.66 (1.44-4.92), 0.002	Sroubek ^{R50}

P=primary prevention patients only, P/S=primary and secondary prevention patients

^{*} univariate analysis; not included in multivariable analysis

^{*} Pooled analysis from 8 international databases

TABLE 11

Early repolarization patterns associated to idiopathic ventricular fibrillation, cardiac death or all causes mortality

Study design	Patient population	Early repolarization patterns	End-point	OR/HR*	Reference
Case-control	206 idiopathic VF 412 matched controls	J-point elevation 0.1 mV	Idiopathic VF	10.9 (6.3-18.9)	Haissaguerre ^{R2}
Case-control	45 idiopathic VF 124 matched controls 121 noncompetitive athletes	J-point elevation in inferior leads	Idiopathic VF	3.2 (1.4-7.5), p=0.006	Rosso ^{R4}
		J-point elevation in I/aVL	Idiopathic VF	16.9 (2.0-140), p=0.009	
		J-point elevation in V4-V6	Idiopathic VF	NS	
Case-control	45 idiopathic VF 124 matched controls 121 noncompetitive athletes	J-point elevation	Idiopathic VF	4.0 (2.0-7.9)	Rosso ^{R245}
		J-point elevation + horizontal ST-segment	Idiopathic VF	13.8 (5.1-37.2)	
Case-control	21 athletes with idiopathic VF 365 controls athletes	J-point elevation 0.1 mV in infero-lateral leads	Idiopathic VF	4.63 (1.67-12.9), p=0.007	Cappato ^{R366}
		QRS slurring in any lead	Idiopathic VF	4.81 (1.73-13.4), p=0.007	
Prospective	10,864 middle-aged people enrolled in the Finnish Social Insurance Institution's Coronary Heart Disease Study (CHD study) between 1966 and 1972	J-point elevation 0.1 mV in inferior leads	Death from cardiac causes	1.28 (1.04-1.59) p=0.03	Tikkanen ^{R44}
			Death from arrhythmias	1.43 (1.06-1.94) p=0.03	
		J-point elevation 0.2 mV in inferior leads	Death from any cause	1.54 (1.06-2.24) p=0.03	
			Death from cardiac causes	2.98 (1.85-4.92) p<0.001	
			Death from arrhythmias	2.92 (1.45-5.89) p=0.01	
Prospective	10,864 middle-aged people enrolled in the Finnish Social Insurance Institution's Coronary Heart Disease Study (CHD study) between 1966 and 1972	J-point elevation 0.1 mV and horizontal/descending ST-segment	Sudden death	1.43 (1.05-1.94)	Tikkanen ^{R237}
		J-point elevation 0.1 mV and upsloping ST-segment	Sudden death	NS	
Prospective	1,161 middle-aged people enrolled in the 3 rd French Monitoring Trends and Determinants in Cardiovascular Disease (MONICA) Project between 1994 and 1997	J-point elevation 0.1 mV	Total mortality	2.45 (1.44-4.15), p=0.001	Rollin ^{R236}
			Cardiovascular mortality	5.6 (2.27-11.8), p=0.001	
		J-point elevation 0.2 mV	Total mortality	NS	
			Cardiovascular mortality	5.14 (1.72-15.4), p=0.004	
		J-point elevation 0.1 mV in inferior leads	Total mortality	2.85 (1.62-5.02), p=0.001	
			Cardiovascular mortality	5.28 (1.96-14.2), p=0.001	
		J-point elevation 0.1 mV in lateral leads	Total mortality	NS	

Study design	Patient population	Early repolarization patterns	End-point	OR/HR*	Reference
		J-point elevation 0.1 mV and horizontal ST-segment elevation	Cardiovascular mortality	6.27 (1.85-21.3), p=0.003	
			Total mortality	3.04 (1.71-5.41), p=0.001	
		J-point elevation 0.1 mV and ascending ST-segment elevation	Cardiovascular mortality	6.93 (2.75-17.4), p=0.001	
			Total mortality	NS	
		J-point elevation 0.1 mV with notching pattern	Cardiovascular mortality	NS	
			Total mortality	3.11 (1.72-5.6), p=0.001	
		J-point elevation 0.1 mV with slurring pattern	Cardiovascular mortality	8.32 (3.32-20.8), p=0.001	
			Total mortality	NS	
		J-point elevation 0.1 mV in white men	Cardiovascular mortality	NS	
			Total mortality	NS	
Prospective	15,792 middle-aged biracial people enrolled in the U.S. Atherosclerosis Risk in Communities (ARIC) between 1987 and 1989	J-point elevation 0.1 mV in white men	Sudden death	NS	Olson ^{R238}
			Coronary events	NS	
			All cause mortality	NS	
		J-point elevation 0.1 mV in white women	Sudden death	8.77 (3.19-24.13)	
			Coronary events	NS	
			All cause mortality	NS	
		J-point elevation 0.1 mV in black men	Sudden death	NS	
			Coronary events	NS	
			All cause mortality	NS	
		J-point elevation 0.1 mV in black women	Sudden death	NS	
			Coronary events	1.47 (1.03-2.09)	
			All cause mortality	NS	
Prospective	29,281 subjects evaluated at the Palo Alto Veterans Affairs Hospital	J-point elevation >0.1 mV in black individuals	Cardiovascular death	NS	Perez-Riera ^{R367}
		J-point elevation >0.1 mV in non-black individuals	Cardiovascular death	1.6, p=0.02	

* adjusted OR/HR are reported when available