Published in final edited form as:

Am J Med Genet A. 2015 October ; 167A(10): 2470–2473. doi:10.1002/ajmg.a.37173.

A Second Locus for Schneckenbecken Dysplasia Identified by a Mutation in the Gene Encoding *Inositol Polyphosphate Phosphatase-Like 1 (INPPL1)*

Hane Lee¹, Lisette Nevarez², Ralph S. Lachman³, William R. Wilcox⁴, Deborah Krakow^{3,6,7,8}, Daniel H. Cohn^{2,3,6,*}, and University of Washington Center for Mendelian Genomics⁵

¹Department of Pathology and Laboratory Medicine, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California

²Department of Molecular, , Cell and Developmental Biology, University of California Los Angeles, Los Angeles, California

³International Skeletal Dysplasia Registry, University of California Los Angeles, Los Angeles, California

⁴Division of Medical Genetics, Department of Human Genetics, Emory University, Decatur, Georgia

⁵University of Washington Center for Mendelian Genomics, University of Washington, Seattle, Washington

⁶Departments of Orthopaedic Surgery, Los Angeles, California

⁷Human Genetics, Los Angeles, California

⁸Obstetrics and Gynecology, University of California Los Angeles, Los Angeles, California

TO THE EDITOR

Schneckenbecken dysplasia [OMIM 269250] is an autosomal recessive, perinatal lethal skeletal dysplasia that was first characterized by [Borochowitz et al., 1986], and is phenotypically similar to affected individuals in a family described by Knowles et al. [1986]. Clinical findings include relative macrocephaly, a very flat midface, cleft palate, a short neck, a narrow thorax, brachydactyly, and a high incidence of polyhydramnios. The most characteristic radiographic abnormality, as viewed on an anterior-posterior radiograph, is a medial projection from the ilia that resembles a snail (thus snail pelvis or Schneckenbecken in German). Additional findings include handlebar clavicles, a hypoplastic scapula, generalized platyspondyly with small, rounded vertebral bodies, very short long bones with relatively widened metaphyses and precociously ossified carpal and tarsal bones [Borochowitz et al., 1986; Nikkels et al., 2001].

*Correspondence to: Daniel H. Cohn, Ph.D., Department of Molecular, Cell and Developmental Biology, 615 Charles E. Young Dr. South, Room 550, University of California, Los Angeles, Los Angeles, CA 90095. dcohn@mcdb.ucla.edu.

Conflict of interest: The authors have no conflicts of interest.

Schneckenbecken dysplasia has been classified among the severe spondylodysplastic disorders that also include the autosomal recessive disorders achondrogenesis type 1A [OMIM 200600], spondylometaphyseal dysplasia, Sedaghatian type [OMIM 250220], and opsismodysplasia [OMIM 258480] [Warman et al., 2011]. Mutations in *SLC35D1* [OMIM 610804], the gene encoding the Solute Carrier Family 35 UDP-Glucuronic Acid/ UDP-N-Acetylgalactosamine Dual Transporter, Member D1, were found in 2 of 6 cases of Schneckenbecken dysplasia [Hiraoka et al., 2007]. In a subsequent report *SLC35D1* mutations were identified in another three of five Schneckenbecken dysplasia cases but were absent among 15 cases of other severe spondylodysplastic skeletal dysplasias [Furuichi et al., 2009]. Most of the *SLC35D1* mutations were predicted to lead to absence of the protein. *SLC35D1* has been localized to the endoplasmic reticulum (ER) where it translocates two nucleotide sugars, UDP-glucuronic acid and UDP-N-acetylgalactosamine, into microsomal membrane vesicles to be used in the synthesis of chondroitin sulfate [Muraoka et al., 2001]. A mouse deficient for *Slc35d1* showed a phenotype similar to Schneckenbecken dysplasia and had defective chondroitin sulfate synthesis on the core protein of cartilage proteoglycans [Hiraoka et al., 2007].

Under an approved human subjects protocol, exome sequence analysis using a previously described pipeline [Lee et al., 2012] was carried out in a case of Schneckenbecken dysplasia (International Skeletal Dysplasia Registry reference number R02–170) in which *SLC35D1* was previously excluded (subject 4 in [Hiraoka et al., 2007]). The case presented as a 33 week gestation hydropic stillborn. The radiographic findings were characteristic for Schneckenbecken dysplasia and included midface hypoplasia, handlebar clavicles, short ribs, hypoplastic vertebrae with rounded anterior ends, short long bones with widened metaphyseal ends and a snail-like medial projection in the ileum (Fig. 1).

Analysis of the exome identified homozygosity for a variant, c.2415+1G>A in intron 21 of *INPPL1* [OMIM 600829], which encodes Inositol Polyphosphate Phosphatase-like 1. The mutation predicts skipping of exon 21, which would lead to a frameshift and a premature termination codon. Recessively inherited mutations in *INPPL1* also produce opsismodysplasia [OMIM 258480], another severe autosomal recessive spondylodysplastic disorder [Below et al., 2013; Huber et al., 2013]. Opsismodysplasia is generally non-lethal and is characterized by dysmorphic craniofacial features, a large anterior fontanelle, short stature and small hands and feet. Radiographic findings include very delayed bone maturation particularly at the knee, marked shortness of the metacarpals/ metatarsals and phalanges with metaphyseal cupping and markedly flat vertebral bodies [Cormier-Daire et al., 2003; Maroteaux et al., 1984]. Some affected individuals have severe renal phosphate wasting and there are variable degrees of respiratory insufficiency [Below et al., 2013] which may be secondary to the skeletal manifestations. A variety of other clinical findings have been described but an insufficient number of cases have been studied to know whether these are consistently associated with *INPPL1* mutations. The c.2415+1G>A mutation found in the homozygous state in this case of Schneckenbecken dysplasia was also identified in a surviving opsismodysplasia case in compound heterozygosity with a frameshift mutation. Thus, apparent loss of *INPPL1* function can produce either the perinatal lethal Schneckenbecken dysplasia phenotype or non-lethal opsismodysplasia.

Histomorphology on the cartilage growth plate was performed by staining the tissue with toluidine blue and Goldner's reagent. The findings included poor hypertrophic column formation (Fig. 2A) with fibrous septae between the columns, areas of hypercellularity with decreased extracellular matrix and hypervascularity (Fig. 2B), chondrocytes with central round nuclei (Fig. 2C), as well as thickened primary trabeculae and a fibrous band entering the growth plate from the bone collar (Fig. 2D). With the exception of the dense fibrous band in the growth plate, these findings have been previously described in Schneckenbecken dysplasia [Varkey and Jones, 2004], but the gene(s) involved were unknown. Similar findings were described in the *Slc35d1* knockout mice as well as a Schneckenbecken dysplasia case due to defects in *SLC35D1* [Hiraoka et al., 2007]. The histopathology in opsismodysplasia [Cormier-Daire et al., 2003] is similar, showing poorly organized growth plates, numerous vascular canals, increased chondrocyte density, and many round, ballooned chondrocytes and thickened trabeculae in the mineralized matrix. Thus the growth plate findings, while not necessarily pathognomonic for either Schneckenbecken dysplasia or opsismodysplasia, do show significant overlap. The only finding in this case that is distinct is the fibrous band that extends from the bone collar.

The similarities in the growth plates between this case of Schneckenbecken dysplasia due to homozygosity for an *INPPL1* mutation and those due to *SLC35D1* mutations could suggest possible mechanistic overlap. Defects in both genes could lead to reduced extracellular matrix synthesis and a hypercellular matrix containing chondrocytes with central nuclei. In cases due to *SLC35D1* mutations, reduced post-translational modification of matrix proteoglycans appears to be the underlying abnormality. How defects in *INPPL1* affect extracellular matrix production is unknown, but signaling cascades mediated by phosphatidylinositols including PI(3,4,5)P3 might contribute to matrix production and regulation.

The functional role of INPPL1 in skeletal biology is unknown. The protein is an inositol-1,4,5-trisphosphate 5-phosphatase and through its phosphatase activity converts phosphatidylinositol 3,4,5-triphosphate (PtdIns(3,4,5)P3) to PtdIns(3,4)P2. PtdIns-(3,4,5)P3 is widely expressed and acts as second messenger that participates in numerous cellular functions [Riehle et al., 2013]. Mouse models show that *Inpp1l* is a negative regulator of insulin sensitivity and signaling and mice deficient for *Inpp1l* are small and have craniofacial abnormalities [Sleeman et al., 2005; Dubois et al., 2012], implying a yet uncharacterized effect on skeletogenesis.

Our findings identify a second locus for Schneckenbecken dysplasia and demonstrate that the phenotype is part of an *INPPL1* allelic spectrum of disease. It remains to be determined whether the remaining *SLC35D1* negative cases harbor mutations in *INPPL1* or whether there is further locus heterogeneity in this disorder.

ACKNOWLEDGMENTS

We thank the family for their participation in this study. This work was supported by NIH grants RO1 DE019567 and RO1 AR062651 to DHC, RO1 AR066124 to DK and 1U54 HG006493 to Drs. Debbie Nickerson, Jay Shendure, and Michael Bamshad. We also thank the March of Dimes Birth Defects Foundation, the Joseph Drown Foundation and the Orthopaedic Institute for Children/Orthopaedic Hospital Research Center for their support.

Support was also provided by NIH/National Center for Advancing Translational Science (NCATS) UCLA CTSI Grant Number UL1TR000124.

Grant sponsor: NIH; Grant numbers: RO1 DE019567, R01 AR062651;

Grant sponsor: NIH/National Center for Advancing Translational Science; Grant number: UL1TR000124.

REFERENCES

- Below JE, Earl DL, Shively KM, McMillin MJ, Smith JD, Turner EH, Stephan MJ, Al-Gazali LI, Hertecant JL, Chitayat D. Whole-genome analysis reveals that mutations in inositol polyphosphate phosphatase-like 1 cause opsismodysplasia. *Am J Hum Genet.* 2013; 92:137–143. [PubMed: 23273567]
- Borochowitz Z, Jones K, Silbey R, Adomian G, Lachman R, Rimoin D, Opitz JM, Reynolds JF. A distinct lethal neonatal chondrodysplasia with snail-like pelvis: Schneckenbecken dysplasia. *Am J med Genet.* 1986; 25:47–59. [PubMed: 3799723]
- Cormier-Daire V, Delezoide A, Philip N, Marcorelles P, Casas K, Hillion Y, Faivre L, Rimoin D, Munnich A, Maroteaux P. Clinical, radiological, and chondro-osseous findings in opsismodysplasia: Survey of a series of 12 unreported cases. *J Med Genet.* 2003; 40:195–200. [PubMed: 12624139]
- Dubois E, Jacoby M, Blockmans M, Pernot E, Schiffmann SN, Foukas LC, Henquin J-C, Vanhaesebroeck B, Erneux C, Schurmans S. Developmental defects and rescue from glucose intolerance of a catalytically-inactive novel Ship2 mutant mouse. *Cell Signalling.* 2012; 24:1971–1980. [PubMed: 22750293]
- Furuichi T, Kayserili H, Hiraoka S, Nishimura G, Ohashi H, Alanay Y, Lerena J, Aslanger AD, Koseki H, Cohn DH. Identification of loss-of-function mutations of SLC35D1 in patients with Schneckenbecken dysplasia, but not with other severe spondylodysplastic dysplasias group diseases. *J Med Genet.* 2009; 46:562–568. [PubMed: 19508970]
- Hiraoka S, Furuichi T, Nishimura G, Shibata S, Yanagishita M, Rimoin DL, Superti-Furga A, Nikkels PG, Ogawa M, Katsuyama K. Nucleotide-sugar transporter SLC35D1 is critical to chondroitin sulfate synthesis in cartilage and skeletal development in mouse and human. *Nat Med.* 2007; 13:1363–1367. [PubMed: 17952091]
- Huber C, Faqeih EA, Bartholdi D, Bole-Feysot C, Borochowitz Z, Cavalcanti DP, Frigo A, Nitschke P, Roume J, Santos HG. Exome Sequencing Identifies *INPPL1* Mutations as a Cause of Opsismodysplasia. *Am J Hum Genet.* 2013; 92:144–149. [PubMed: 23273569]
- Knowles S, Winter R, Rimoin D, Opitz JM, Reynolds JF. A new category of lethal short-limbed dwarfism. *Am J Med Genet.* 1986; 25:41–46. [PubMed: 3799722]
- Lee H, Graham JM Jr, Rimoin DL, Lachman RS, Krejci P, Thompson SW, Nelson SF, Krakow D, Cohn DH. Exome Sequencing Identifies *PDE4D* Mutations in Acrodysostosis. *Am J Hum Genet.* 2012; 90:746–751. [PubMed: 22464252]
- Maroteaux P, Stanescu V, Stanescu R, Le Marec B, Moraine C, Lejarraga H. Opsismodysplasia: A new type of chondrodysplasia with predominant involvement of the bones of the hand and the vertebrae. *Am J Med Genet.* 1984; 19:171–182. [PubMed: 6496568]
- Muraoka M, Kawakita M, Ishida N. Molecular characterization of human UDP-glucuronic acid/UDP-N-acetylgalactosamine transporter, a novel nucleotide sugar transporter with dual substrate specificity. *FEBS Lett.* 2001; 495:87–93. [PubMed: 11322953]
- Nikkels P, Stigter RH, Knol IE, van der Harten HJ. Schneckenbecken dysplasia, radiology, and histology. *Pediatr Radiol.* 2001; 31:27–30. [PubMed: 11200994]
- Riehle RD, Cornea S, Degterev A. Role of Phosphatidylinositol 3, 4, 5-Trisphosphate in Cell Signaling. *Adv Exp Med Biol.* 2013; 991:105–139. [PubMed: 23775693]
- Sleeman MW, Wortley KE, Lai K-MV, Gowen LC, Kintner J, Kline WO, Garcia K, Stitt TN, Yancopoulos GD, Wiegand SJ. Absence of the lipid phosphatase SHIP2 confers resistance to dietary obesity. *Nat Med.* 2005; 11:199–205. [PubMed: 15654325]
- Varkey J, Jones R. Perinatally lethal, short-limbed dwarfism with distinct features—Schneckenbecken dysplasia. *Ultrasound Obstetrics Gynecol.* 2004; 24:575–577.

Warman ML, Cormier-Daire V, Hall C, Krakow D, Lachman R, LeMerrer M, Mortier G, Mundlos S, Nishimura G, Rimoin DL, Robertson S, Savarirayan R, Sillence D, Spranger J, Unger S, Zabel B, Superti-Furga A. Nosology and classification of genetic skeletal disorders: 2010 revision. *Am J Med Genet Part A*. 2011; 155A:943–968. [PubMed: 21438135]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

FIG. 1.

ISDR case R02-170, a 33 week stillborn with Schneckenbecken dysplasia. (A) AP full body radiograph showing a small thorax, micromelia, metaphyseal widening of the long bones, and characteristic snail-shaped iliac wings. (B) Lateral radiograph showing characteristic ovoid anterior vertebral body ossification with non-ossified posterior portions, as well as midface hypoplasia.

FIG. 2.

ISDR case R02-170 cartilage growth plate histology. (A) Cartilage growth plate (10X) showing poor hypertrophic column formation (arrow) with fibrous septae between the columns; B, bone, H, hypertrophic zone, P, proliferative zone. (B) Proliferative zone (10X) showing increased vascularity (arrows), decreased extracellular matrix and hypercellularity. (C) Reserve chondrocytes (40X) with arrows identifying chondrocytes with central round nuclei. (D) Cartilage stained with Goldner's reagent showing a fibrous band at the distal end of the growth plate (arrow) and thickened trabeculae in the primary zone of ossification.