Published in final edited form as:

Fetal Matern Med Rev. 2014 November ; 25(3-4): 332–338. doi:10.1017/S0965539515000066.

Intracellular Organisms as Placental Invaders

Marguerite B. Vigliani, MD¹ and Anna I. Bakardjiev, MD²

¹Department of Obstetrics and Gynecology, Warren Alpert Medical School of Brown University, Providence, Rhode Island ²Department of Pediatrics, Microbial Pathogenesis and Host Defense Program, University of California, San Francisco, California

Abstract

In this article we present a novel model for how the human placenta might get infected via the hematogenous route. We present a list of diverse placental pathogens, like *Listeria monocytogenes* or Cytomegalovirus, which are familiar to most obstetricians, but others, like *Salmonella typhi*, have only been reported in case studies or small case series. Remarkably, all of these organisms on this list are either obligate or facultative intracellular organisms. These pathogens are able to enter and survive inside host immune cells for at least a portion of their life cycle. We suggest that many blood-borne pathogens might arrive at the placenta via transportation inside of maternal leukocytes that enter the decidua in early pregnancy. We discuss mechanisms by which extravillous trophoblasts could get infected in the decidua and spread infection to other layers in the placenta. We hope to raise awareness among OB/GYN clinicians that organisms not typically associated with the TORCH list might cause placental infections and pregnancy complications.

Keywords

Intracellular pathogens; TORCH; placenta

A Surprise Pathogen

In an era of evidence-based medicine, physicians sometimes forget the value of anecdotes in stimulating thought about clinical problems. Our recent report of typhoid fever in a pregnant woman at 12 weeks of gestation is a good example.¹ Despite culture-proven diagnosis and appropriate treatment of the mother with antibiotics, fetal loss occurred at 16 weeks of gestation. *Salmonella typhi* was found in the fetal lung on autopsy, consistent with vertical transmission of the organism. None of the clinicians caring for the patient had imagined that Gram-negative bacteria could cross the placenta and kill the fetus despite early diagnosis and treatment with appropriate antibiotics.

In marked contrast, large animal veterinarians are trained to quarantine aborting animals until contaminated areas are disinfected, prophylactic antibiotics are given, and specimens of

Corresponding author: Marguerite B. Vigliani MD, 450 Veterans Memorial Parkway, Suite 101, East Providence, Rhode Island 02914. Phone (401) 438-1448. Cell (401) 261-4109. Fax (401) 434-6734. mbvmd@fullchannel.net. Reprints are not available.

Disclosure: None of the authors have a conflict of interest

placenta, fetus, blood, vaginal swabs, feces, and fetal stomach contents are systematically collected and sent to veterinary diagnostic labs for necropsy, histopathology, culture and sensitivity, and polymerase chain reaction (PCR) analysis for various organisms. This practice is normal for veterinarians in order to prevent huge economic damages that might occur should one animal ingest an infected placenta and pass the infection on to the whole herd. In humans *Mycoplasma hominis* or *Brucella* spp. might be in the differential diagnosis of recurrent miscarriage, but culture techniques are not generally used for evaluation of miscarriage. We don't think about infection very often, and we don't culture routinely. Indeed culturing every spontaneous human miscarriage might add significantly to the cost of medical care without yielding much useful information. Thus in our case, even the admitting physician who had knowledge of this patient's typhoid fever in the first trimester did not order Gram stain or bacterial cultures of the placenta at 16 weeks of gestation because infection is not in the differential diagnosis for second trimester fetal loss.

Beyond TORCH

Perhaps we clinicians depend too much on the TORCH mnemonic we learned in medical school. Although it is well known that the TORCH list addresses only a few of the blood-borne pathogens known to enter the fetal compartment, it is important to remember that the "O" in TORCH suggests that "other" microbes can infect the placenta as well. Table 1² lists the diverse placental pathogens that have been associated with pregnancy complications and fetal damage in humans.

This list contains viruses, bacteria, and protozoan microbes. Many of these are the same pathogens that cause embryonic and fetal demise in cattle, sheep, goats, and camelids.³ All these zoonotic infections are known in the veterinary literature to be caused by intracellular organisms. Likewise every organism listed in Table 1 is able to survive in host cells. Some, like the viruses, have obligate intracellular lifecycles; but others are facultative intracellular. Whether obligate or facultative, intracellular microbes have evolved many ways to elude host recognition. Many have learned to take up residence inside of circulating blood cells that facilitate systemic dissemination. For example, the bacterial pathogens *L. monocytogenes* and *Salmonella* can hitch a ride in monocytes to reach the central nervous system⁴ or the liver.⁵ Thus, it is plausible that infected monocytes can provide transportation for these organisms on their way to the maternal-fetal interface, giving new meaning to the concept of hematogenous spread.

The Decidua is the Initial site of Infection

In vivo studies with murine and guinea pig models have all pointed to the decidua as the preferred site of initial placental colonization for the hematogenously spread organisms *Toxoplasma gondii*, *Chlamydia psittaci*, *Coxiella burnetii*, *Fusobacterium nucleatum*, *Salmonella typhimurium*, *Brucella abortus*, *L. monocytogenes*, and Cytomegalovirus.^{2,6} The decidua actively recruits leukocytes to the endometrium during the secretory phase of the menstrual cycle and during the implantation phase of early pregnancy⁷. Once blood monocytes are localized in the decidua, they differentiate into decidual macrophages and microbes hiding inside of these have the opportunity to infect the trophoblasts. We know that

trophoblast infection does happen based on histological examination of human placentas infected by *L. monocytogenes*^{8–10} and by the recent finding of intracellular bacteria of diverse morphologies inside of trophoblasts at the basal plates of 27% of all placentas¹¹ including those without clinical or pathological evidence of chorioamnionitis.

Entering the Trophoblast

Trophoblast infection occurs most likely via multiple mechanisms depending on the armamentarium of the specific pathogen such as virulence factors that facilitate adherence or invasion of host cells (e.g. adhesins, invasins, and Type III secretion systems). Another possibility is bacterial cell-to-cell spread from infected maternal cells to trophoblasts without exposure to the extracellular environment. Inoculation of pregnant guinea pigs with wild-type and mutant *L. monocytogenes* has shown that the predominant mechanism for placental invasion by this organism is cell-to-cell spread.¹²

Figure 1 shows the maternal-fetal interface at the placenta. The largest surface area of fetal contact with maternal blood is the syncytiotrophoblast (SYN). The SYN is a continuous layer of fused multinucleated trophoblasts undergirded by cytotrophoblasts and a basement membrane. SYN has a distinct polarity and there are no intercellular junctions. Microbes have difficulty adhering to SYN and in human placental organ cultures SYN is highly resistant to infection by several organisms. Those that have been studied include *L. monocytogenes*,¹³ *T. gondii*,¹⁴ Herpes simplex virus,¹⁵ and Cytomegalovirus.¹⁶

Also shown is the extravillous trophoblast (EVT) that forms the anchoring villi to attach the placenta to the decidua and the uterine wall. EVT is intimately juxtaposed to a variety of leukocytes at the decidua. *In vitro* placental culture techniques have directly shown the ability of *L. monocytogenes*,¹³ *T. gondii*,¹⁴ Herpes simplex virus,¹⁵ and Cytomegalovirus¹⁶ to infect extravillous trophoblast (EVT).

Immune Tolerance

Maternal leukocytes are plentiful in the decidua, but their functions are modulated to help with angiogenesis or fetal immune tolerance.⁷ Most of the leukocytes in the decidua are specialized natural killer cells (dNK) and macrophages. T-cells and B-cells are scarce, due to epigenetic silencing of chemoattractants in the decidua.¹⁷ Cytolytic functions of dNKs, macrophages, and T-cells in the decidua are all suppressed perhaps due to cross talk with EVT, which expresses the immunosuppressant human leukocyte antigen G (HLA-G),⁷ but not the Class I histocompatibility molecules HLA-A or HLA-B.⁷ The intimate relationship of EVT to maternal macrophages in the decidua might be permissive to infection transmitted from these cells directly to trophoblasts. EVT could be a tolerant ecological niche for intracellular organisms because of the many specialized immune adaptations of the decidua to allow for tolerance of fetal tissues.⁷ For example due to the lack of classical MHC class I molecules infected EVT cannot present microbial antigens present in their cytoplasm.

Host Defenses

That being said, the maternal-fetal interface has defense mechanisms that limit infection by intracellular organisms. This has probably been most closely characterized in experimental models with *L. monocytogenes*. In pregnant guinea pigs only one in a million bacteria injected intravenously are able to colonize the maternal-fetal interface.¹⁸ Furthermore, EVT have the ability to eliminate approximately 80% of intracellular *L. monocytogenes* within 24 hours.¹⁹ However, the bacterial load can at times overcome host defenses such that the organism can no longer be eliminated and the placenta becomes a nidus for infection, leading to continuous seeding and reseeding of the maternal organs and the fetus until the placenta and the products of conception are expelled, or the mother dies.¹⁸ Further research is needed to better understand the molecular and cellular mechanisms of host defenses against pathogens at the maternal-fetal interface, and how these are breached.

Breaching the Barrier

Pathogens vary in their ability to elicit host responses and hence disease expression. *Mycobacterium tuberculosis*, for example, may not leave a significant inflammatory signature, and may be able to co-exist with the pregnancy rarely causing fetal damage or vertical transmission because of their slow replication rate and their ability to live for long periods of time inside of macrophages. Other organisms, like the obligate intracellular Cytomegalovirus, tend to evoke host cell apoptosis and cause a great deal of tissue necrosis, inflammation, and edema in placental tissues, leading to abruption and fetal loss.¹⁶ Damage to SYN by Cytomegalovirus opens the door to secondary invaders like Herpes simplex virus, which might not cause damage on its own.¹⁹ In general, coinfection by more than one organism is probably worth remembering as a possible mechanism by which damage to SYN might be permissive to infection by a second organism. Another example is placental infection with *Plasmodium falciparum*, the parasite that causes malaria. Infected erythrocytes are sequestered in the intervillous spaces, adhere to SYN, and cause SYN damage.²⁰ This damage might explain increased rates of vertical transmission of HIV-1²¹ in pregnant women with who are coinfecting.

Looking Ahead

We hope that this discussion will serve to increase awareness among OB/GYN clinicians that all maternal illnesses during pregnancy should be regarded with suspicion, because host immune cells can carry intracellular organisms to the placenta, and breach the placental barrier. Antibiotic therapy of infections such as typhoid fever and listeriosis might have to be extended beyond the 2 weeks recommended for non-pregnant individuals especially if infection occurs during the first trimester.^{22,23}

We do not know if histological examination of the placenta or bacterial cultures are the most important investigations in cases of fetal loss, preterm labor, or abruption. We do hope, however, that this discussion might stimulate some clinical research into culture-independent methods of diagnosing infection by intracellular organisms at the maternal-fetal interface, be

it by quantitative PCR for pathogen DNA, immunohistochemistry, electron microscopy, or some other method.

Acknowledgments

Sources of financial support: A.I.B. is supported by the US National Institutes of Health (R01AI084928), and a Burroughs Wellcome Fund Investigator in the Pathogenesis of Infectious Disease Award (41259).

References

1. Vigliani MB, Bakardjiev AI. First trimester typhoid Fever with vertical transmission of salmonella typhi, an intracellular organism. *Case reports in medicine*. 2013; 2013:973297. [PubMed: 24459469]
2. Robbins JR, Bakardjiev AI. Pathogens and the placental fortress. *Current Opinion in Microbiology*. 2012; 15(1):36–43. [PubMed: 22169833]
3. Daniel Givens M, Marley MSD. Infectious causes of embryonic and fetal mortality. *Theriogenology*. 70(3):270–285. [PubMed: 18502494]
4. Drevets DA, Jelinek TA, Freitag NE. *Listeria monocytogenes*-infected phagocytes can initiate central nervous system infection in mice. *Infect Immun*. 2001 Mar; 69(3):1344–1350. [PubMed: 11179297]
5. Vazquez-Torres A, Jones-Carson J, Baumler AJ, et al. Extraintestinal dissemination of *Salmonella* by CD18-expressing phagocytes. *Nature*. 1999 Oct 21; 401(6755):804–808. [PubMed: 10548107]
6. Krishnan L, Nguyen T, McComb S. From mice to women: the conundrum of immunity to infection during pregnancy. *Journal of reproductive immunology*. 2013 Mar; 97(1):62–73. [PubMed: 23432873]
7. Trundley A, Moffett A. Human uterine leukocytes and pregnancy. *Tissue Antigens*. 2004; 63(1):1–12. [PubMed: 14651517]
8. Olding L, Philipson L. Two cases of listeriosis in the newborn, associated with placental infection. *Acta pathologica et microbiologica Scandinavica*. 1960; 48:24–30. [PubMed: 14428619]
9. Driscoll SG, Gorbach A, Feldman D. Congenital listeriosis: diagnosis from placental studies. *Oncologia*. 1962 Aug; 20:216–220. [PubMed: 13887884]
10. Yamazaki K, Price JT, Altshuler G. A placental view of the diagnosis and pathogenesis of congenital listeriosis. *Am J Obstet Gynecol*. 1977 Nov 15; 129(6):703–705. [PubMed: 411374]
11. Stout MJ, Conlon B, Landeau M, et al. Identification of intracellular bacteria in the basal plate of the human placenta in term and preterm gestations. *Am J Obstet Gynecol*. 2013 Mar; 208(3):226 e221–227 e221. [PubMed: 23333552]
12. Bakardjiev AI, Stacy BA, Portnoy DA. Growth of *Listeria monocytogenes* in the guinea pig placenta and role of cell-to-cell spread in fetal infection. *J Infect Dis*. 2005 Jun 1; 191(11):1889–1897. [PubMed: 15871123]
13. Robbins JR, Skrzypczynska KM, Zeldovich VB, Kapidzie M, Bakardjiev AI. Placental syncytiotrophoblast constitutes a major barrier to vertical transmission of *Listeria monocytogenes*. *PLoS Pathog*. 2010 Jan; 6(1):e1000732. [PubMed: 20107601]
14. Robbins JR, Zeldovich VB, Poukchanski A, Boothroyd JC, Bakardjiev AI. Tissue Barriers of the Human Placenta to Infection with *Toxoplasma gondii*. *Infection and Immunity*. 2012 Jan 1; 80(1):418–428. 2012. [PubMed: 22083708]
15. Koi H, Zhang J, Makrigiannakis A, et al. Syncytiotrophoblast is a barrier to maternal-fetal transmission of herpes simplex virus. *Biol Reprod*. 2002 Nov; 67(5):1572–1579. [PubMed: 12390890]
16. Fisher S, Genbacev O, Maidji E, Pereira L. Human Cytomegalovirus Infection of Placental Cytotrophoblasts In Vitro and In Utero: Implications for Transmission and Pathogenesis. *Journal of Virology*. 2000 Aug 1; 74(15):6808–6820. 2000. [PubMed: 10888620]

17. Nancy P, Tagliani E, Tay CS, Asp P, Levy DE, Erlebacher A. Chemokine gene silencing in decidual stromal cells limits T cell access to the maternal-fetal interface. *Science (New York NY)*. 2012 Jun 8; 336(6086):1317–1321.
18. Bakardjiev AI, Theriot JA, Portnoy DA. *Listeria monocytogenes* traffics from maternal organs to the placenta and back. *PLoS Pathog*. 2006 Jun.2(6):e66. [PubMed: 16846254]
19. Zeldovich VB, Robbins JR, Kapidzic M, Lauer P, Bakardjiev AI. Invasive extravillous trophoblasts restrict intracellular growth and spread of *Listeria monocytogenes*. *PLoS Pathog*. 2011; 7(3):e1002005. [PubMed: 21408203]
20. Weisblum Y, Panet A, Haimov-Kochman R, Wolf DG. Models of vertical cytomegalovirus (CMV) transmission and pathogenesis. *Seminars in immunopathology*. 2014 Nov; 36(6):615–625. [PubMed: 25291972]
21. Crocker IP, Tanner OM, Myers JE, Bulmer JN, Walraven G, Baker PN. Syncytiotrophoblast Degradation and the Pathophysiology of the Malaria-infected Placenta. *Placenta*. 25(4):273–282. [PubMed: 15028419]
22. Bulterys PL, Chao A, Dalai SC, et al. Placental Malaria and Mother-to-Child Transmission of Human Immunodeficiency Virus-1 in Rural Rwanda. *The American journal of tropical medicine and hygiene*. 2011 Aug 1; 85(2):202–206. 2011. [PubMed: 21813835]
23. Chan BT, Hohmann E, Barshak MB, Pukkila-Worley R. Treatment of listeriosis in first trimester of pregnancy. *Emerging infectious diseases*. 2013 May; 19(5):839–841. [PubMed: 23697599]
24. Lamont RF, Sobel J, Mazaki-Tovi S, et al. Listeriosis in human pregnancy: a systematic review. *Journal of perinatal medicine*. 2011 May; 39(3):227–236. [PubMed: 21517700]

Figure 1. Structure of the human placenta

(A) Maternal blood from spiral arteries in the decidua flows into the intervillous space where it surrounds thousands of fetally derived chorionic villi.

(B) Villi are freely floating in maternal blood, or invade the decidua to form anchoring villi. The entire villous surface is covered with a continuous layer of multinucleate syncytiotrophoblast (SYN), which is the major fetal surface in contact with maternal blood. The apical side of the SYN consists of profuse, branched microvilli, which provide abundant surface area for gas and nutrient exchange between mother and fetus. The SYN is undergirded by cytotrophoblasts (sCTB), which are separated from fetal capillaries in the villous stroma by a basement membrane. Some cytotrophoblasts leave the basement membrane and differentiate along the invasive pathway to form anchoring villi. Columns of unpolarized cytotrophoblasts attach to and then penetrate the uterine wall where they give rise to extravillous cytotrophoblasts (EVT).

(Figure adapted with permission from Robbins et al. 2010)²

Table 1Placental invaders (adapted with permission from Robbins et al. 2010²)

Pathogen (Illness)	Type	Lifestyle	Cells infected
<i>Brucella</i> spp. (Brucellosis) ^{a, b, c, d, e}	Bacteria	Facultative	Leukocytes, epithelial
<i>Coxiella burnetii</i> (Q fever) ^{c, e, h}	Bacteria	Obligate	Leukocytes
<i>Listeria monocytogenes</i> (Listeriosis) ^{a, b, c, d, g, h 2423}	Bacteria	Facultative	Epithelial, phagocytes
<i>Mycobacterium tuberculosis</i> (TB) ^{d, g}	Bacteria	Facultative	Leukocytes
<i>Treponema pallidum</i> (Syphilis) ^{b, c, f, g}	Bacteria	Facultative	Unknown
<i>Leishmania</i> spp. (Leishmaniasis) ^h	Parasite	Obligate	Leukocytes
<i>Plasmodium falciparum</i> (Malaria) ^{b, c, e, g, h}	Parasite	Obligate	Erythrocytes, hepatocytes
<i>Toxoplasma gondii</i> (Toxoplasmosis) ^{a, b, d, g}	Parasite	Obligate	All nucleated cells
<i>Trypanosoma</i> spp. (Chagas disease, African sleeping sickness) ^{e, g}	Parasite	Facultative	Epi/endothelial
Cytomegalovirus (CMV) ^{a, b, d, g}	Virus	Obligate	Leukocytes, trophoblasts
Lymphocytic choriomeningitis virus (LCMV) ^{a, b, f, g25,2624,25}	Virus	Obligate	Leukocytes
Parvovirus B19 ^{a, b, c, d, f}	Virus	Obligate	Hematopoietic, endothelial
Rubella virus (German measles) ^g	Virus	Obligate	Many
Varicella zoster virus (Chicken pox) ^{g, h}	Virus	Obligate	Leukocytes, neurons, epithelial

^a first trimester fetal death;^b second trimester fetal death;^c stillbirth;^d preterm labor;^e intrauterine growth restriction;^f fetal hydrops;^g severe neonatal infection;^h increased severity of maternal disease.

Note: Many other intracellular organisms including *Babesia* spp., Coxsackie B virus, Japanese encephalomyelitis virus, *Leptospira* spp., *Cryptococcus neoformans*, *Pasteurella*, *Shigella*, *Salmonella typhi*, *Wucheria bancrofti*, *Candida* spp., *Campylobacter*, and many gingival bacteria including *Fusobacterium nucleatum* merit further study because of human case reports and/or animal studies.

Epstein-Barr virus, Hepatitis B virus, HIV, and HSV are known to cause perinatal infections via exposure to maternal secretions and blood, but rarely via vertical transmission across the placenta.