Published in final edited form as:

Trends Immunol. 2016 October ; 37(10): 659–667. doi:10.1016/j.it.2016.08.008.

Balancing Tolerance or Allergy to Food Proteins

Paul J Bryce, PhD

Associate Professor, Division of Allergy-Immunology, Department of Medicine, Northwestern
Feinberg School of Medicine, 240 E. Huron, Chicago, IL 60611, FAX: 312 503 0078

Paul J Bryce: p-bryce@northwestern.edu

Abstract

Dietary proteins usually induce immune tolerance, and yet may trigger life-threatening immune responses in the case of food allergy. The associated type 2 immunity, linked with specific IgE production and activation of mast cells and basophils, is well understood, but the mechanisms related to preventing food allergy are still being deciphered. Recent insights into the mechanisms that regulate oral tolerance and dietary antigen sampling has revealed unique regulatory events that occur during early life and into adulthood. Drawing from both recent clinical and experimental discoveries, this article focuses on the current evidence for how several key stages of life present mechanistic points that might participate in tipping the balance between food protein tolerance or allergy.

Increasing Prevalence of Food Allergies

In spite of its increasing prevalence and recognition, food allergy is clearly not a new problem. In 1928 Albert Rowe elegantly outlined the clinical history of a patient with food allergy and his successful experience with an elimination diet [1]. In stressing the importance for considering family history of atopy, the inconsistencies of skin testing, the psychological aspects of food aversions being discounted as “whims or fancies” by physicians, and several key foods to be eliminated, Rowe imparts opinions that seem far ahead of their time. The last decade has been an incredibly important window in helping to shine a spotlight on food allergy as a disease. An extensive wealth of studies has helped to highlight the limited treatment options for food allergy, escalating prevalence, continued difficulties in diagnosis, and the sociological impact on health and well-being. In contrast, our mechanistic understanding of the processes that limit most of us from exhibiting immunological reactions to foods has remained largely unclear. Recent breakthroughs in this area have finally allowed us to glimpse at the sophisticated and complex events that allow us to partake in something as simple as eating.

The key immunological mechanisms thought to contribute to the allergic sensitization responses (particularly dominated by allergen-specific IgE) and the processes of reactivity

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

(particularly the most severe forms, including anaphylaxis) have been reviewed previously [2]. Instead, this review examines the evidence for how the immune system exerts balance over our responses to dietary antigens during several key stages of development, drawing significantly from epidemiological studies of food allergy. The disruption of this balance, through relatively unknown causes, is likely responsible for the sensitization, and ultimately the reactivity to food antigens. It is interesting to note that these two events that are seemingly independent since, somewhat surprisingly, food sensitization remains much more common than food allergy.

Neonatal and Early-Life Influences on Food Allergy Development

The proteins in foods represent an important nutrient component that is transmitted from mother to embryo. A recent study of twenty healthy women during two distinct stages of pregnancy described the detection of several major food allergens in amniotic fluid [3], suggesting that our first experience of dietary allergens occurs in utero. Due to the need to suppress maternal immunity towards paternal antigens, the fetal microenvironment represents a uniquely specialized site of tolerance but how this contributes to establishing immune tolerance to foods is unknown. Cord blood has provided an opportunity to interrogate this maternal-offspring interface and has demonstrated intriguing findings. In a recent prospective study that examined the cord blood immune profiles from newborns, it was found that individuals that would ultimately develop food allergies presented with a higher ratio of monocytes to CD4⁺ T cells and a lowered numbers of regulatory T cells (Treg) [4]. Functionally, the CD14⁺ monocytes isolated from these cords also displayed a heightened innate inflammatory response (IL-1 β , IL-6 and tumor necrosis factor (TNF)) when stimulated by the bacterial ligand LPS, suggesting an intrinsic alteration in their activation potential. The authors connected this with the disruption of tolerance and priming towards Th2 and allergy by demonstrating that these cytokines modulated IL-2 production, suppressing Treg induction and instead promoting an IL-4 expressing T cell that bore similarities to Th2 cells. This potential for extremely early life events to influence food allergy has also been supported by circumstantial clinical evidence in adult patients who underwent cord blood transplantation for hematological disorders and developed new-onset food allergy within a year of transplant, a condition referred to as “Transplant-acquired food allergy” [5, 6]. One caveat to these conclusions is that differences in Tregs from cord blood have already been demonstrated between atopic (individuals with evidence for existing IgE responses) and non-atopic mothers and associate with atopic dermatitis [7, 8]. These differences might thus relate to atopy and allergy in general, as opposed to dietary tolerance or food allergy specifically.

While human studies did not directly address the role of specific antigens, Gerhold et al. utilized a well-established murine model of allergic airway inflammation to ovalbumin (OVA) to examine the effects of antigen exposure throughout pregnancy and studied the subsequent capacity for the offspring to become sensitized [9]. While the pups from non-exposed mothers generated robust antigen-specific IgE responses and developed airway inflammation, those from the exposed mothers were significantly protected. This protection was associated with the development of Tregs expressing the master regulator FoxP3 (which can be derived from both the thymus (tTregs) and the periphery (pTregs)). FoxP3 deficient

mice suffer from a variety of spontaneous inflammatory and autoimmune diseases in which both tTregs and pTregs contribute to suppression, including failure of oral tolerance and development of food allergy [70]. Similarly, maternal exposure to peanut during pregnancy and lactation was reported to reduce the development of peanut-specific IgE in the offspring, but only upon low dose exposure [10]. This effect seems dramatically different if the mother is already sensitized or undergoes sensitization during pregnancy, with several studies describing heightened sensitization in the offspring [11–13], and epigenetic alteration of the *il4* gene locus [11]. Such epigenetic modifications and their associations with food allergy have been extensively studied and were reviewed recently [14]

In the earliest days of life, breastfeeding represents another potential exposure to maternal dietary antigens. There has been extensive literature describing the detection of food allergens in breast milk, including several major peanut allergens [15–17]. In mice, administration of human breast milk that contained peanut allergens before weaning diminished the efficiency of subsequent sensitization attempts, suggesting the induction of an oral tolerance response [17]. Furthermore, the study of a limited panel of immune mediators present in breast milk demonstrated that reduced levels of IL-1 β and IL-6 were associated with subsequent food allergy development in children [18]. These two cytokines are associated with the induction of a Th17 response, that might help counteract Th2 priming and allergy development [19]. High IL-1 β in breast milk has also been shown to reduce the risks for development of atopic dermatitis [20]. However, the contribution of breast milk-associated influences as a protective or sensitizing influence over food allergy has remained controversial, particularly since several population-based studies have yielded modest or no associations between the choice of formula versus breastmilk feeding and subsequent food allergy development [21–23]. A recent study on peanut exposure during pregnancy and lactation in mice also concluded that there was no effect on the development of oral tolerance versus peanut allergy in the pups [24]. Mechanistically, this might not be surprising if we consider the important contribution of Tregs to the development of oral tolerance, as previously reviewed [25]. It has been shown in mice that FoxP3⁺ thymocytes are delayed in their appearance versus their CD25⁺ FoxP3[–] counterparts, with neonates requiring several weeks for FoxP3⁺ cells to reach full competency of numbers [26, 27]. Interestingly, this window coincides with the weaning age of a mouse and the introduction of solid dietary constituents. However, the origin of these Tregs during pup development remains unclear. While the thymus remains an important site of their development, the activation of CD4⁺ T cells in the periphery when TGF β is present promotes FoxP3 expression, dependent on the intronic FoxP3 enhancer conserved non-coding sequence 1 (CNS1), and differentiation into induced or peripheral Tregs (pTregs) [28]. These pTreg cells have been shown to be critical both for the control of intestinal microbiota composition and the suppression of allergic responses [29]. In particular, pTregs prevented the development of elevated IgE levels as the mice aged.

While it is unclear whether breastfeeding directly initiates protective tolerance in infants, transfer of maternal secretory IgA antibodies through breast milk is thought to be an important source of early protection against infection in newborns prior to the development of their own intestinal IgA-secreting cells [30]. In mice, maternal secretory IgA was demonstrated to be important in the establishment and composition of the intestinal

microbiota and to limit inflammation in a model of dextran sodium sulfate induced colonic damage [31]. In addition to the role of IgA during infection, it is likely to also be associated with food allergies. Indeed, a clinical study demonstrated that mothers who avoided cow's milk during breastfeeding had lower levels of specific IgA, and that this was associated with the development of cow's milk allergy in their children [32].

In addition to these antigen-specific immune processes, there also exists substantial evidence for a contribution of dietary factors on immune skewing and this is particularly evident for vitamin D and food allergy. It has been shown that high vitamin D levels in either mothers or cord blood associated with an increased risk for developing food allergy [33]. Suggesting the importance of a balance, vitamin D deficiency is also associated with increased risk of sensitivity towards food allergens [34]. The links between vitamin D biology and Tregs have been extensively dissected and recently reviewed [35].

Early Dietary Allergen Exposures Can Promote Tolerance

While the role of prenatal exposure to specific antigens has not been extensively studied in humans, recent findings from early introduction of peanuts into the diet of children have shed light on the role of early life exposure. Building from previous studies that had correlated protection from peanut allergy with early consumption [36], the "Learning Early about Peanut Allergy" (LEAP) study was designed to assess whether regular consumption of peanut could influence the progression towards peanut allergy in a group of high-risk children [37]. In this study, the risk factors were defined as severe eczema, preexisting food allergy to egg, or both. This highlights the important phenomenon of the "atopic march", in which preexisting allergic disease in an individual significantly increases their risk for developing further allergies [38]. The LEAP study achieved remarkable outcomes from early introduction of peanut, with an incidence of 17.2% in the avoidance group but only 3.2% in the consumption group. While primarily designed to test the efficacy of early introduction, the study did also provide insights into some immunological differences between the patient groups. Interestingly, both did show some development of peanut-specific IgE over the sixty months of the study but high levels of IgE were more common in the avoidance group, while peanut-specific IgG4 levels were higher in the consumption group. The potential for IgG4 being a protective antibody relates in part to its inability to bind complement or to form immune complexes [39]. The ratio of IgG4 to IgE has indeed been used as a mechanistic determinant associating with allergy protection in the LEAP study, as well as a number of studies on successful immunotherapy trials [40–42]. A recent follow-up study of the LEAP trial participants twelve months later demonstrated continued protection and maintained IgG4/IgE ratio against peanut allergens [43]. While these findings are exciting from the position of therapy and control of peanut allergy in the setting of high risk individuals, it is important to note that the majority of healthy people do not carry food allergen specific IgG4 or IgE antibodies and so this represents a protective mechanism in sensitized individuals, rather than a general mechanism of oral tolerance. Additionally, an expansion of this early introduction approach to six allergens (peanut, cooked hens egg, cow's milk, sesame, white fish and wheat) in the "Enquiring about Tolerance (EAT)" trial yielded mixed results, with significant benefit seen for peanut and egg allergy in patients who adhered to the treatment protocol but no benefits for the other allergens. [44] In light of these facts, it

seems more likely that early consumption acts to limit a pre-existing aberrant immune response, rather than resets the immune system to the tolerance response observed in healthy individuals. It also remains to be determined if the protective effects of early introduction are possible to all potential allergens and whether IgG4 is the underlying key mechanism of protection or whether further mechanisms are at play.

Despite the potential for these early-life events that might contribute to tolerance to dietary proteins, some individuals progress from sensitization to food allergy. The recent interests in food allergy are driven in part by the significant increases in prevalence that have been seen in recent years. Current estimates suggest that food allergy has nearly doubled in prevalence and now affects up to eight percent of school-aged children and five percent of adults in the United States [45]. The underlying cause of this increase remains unknown. Furthermore, treatment consists mainly of careful avoidance of the trigger foods, leading to a tremendous impact on quality of life [46]. Association studies have helped provide clues to some factors that are likely to have underlying mechanistic importance in tipping the balance from tolerance to allergy. In particular, strong associations between food allergy and preexisting or coincident atopic dermatitis have been seen over and over again, with food allergy affecting around fifteen percent of atopic dermatitis patients [47]. Patients with mutations in the skin matrix protein filaggrin have an intrinsic defect in their epidermal barrier and a strong predisposition to atopic dermatitis [48], but also to peanut allergy [49, 50]. Interestingly, this sensitization associated with levels of peanut dust in the homes of the children [50]. This finding supports the idea of a priming of the immune system to food allergens occurring through cutaneous exposures derived from within our environment. For asthma in particular, the hygiene hypothesis, whereby early life exposures to lipopolysaccharide might protect from allergy-associated immune priming, has been supported from a number of avenues [51]. However, in the case of food allergy, there is little evidence for protection via such mechanisms. Instead, an alternative hypothesis has been proposed in which the early ingestion of dietary allergens promotes tolerance while cutaneous exposures drive sensitization [52]. Food allergy would then occur after early avoidance of food and exposure through a disrupted skin barrier (for example, in the presence of filaggrin mutations mentioned above). In experimental models also, the skin has been demonstrated to be a significant site of sensitization [53–55]. Mechanistically, this sensitization requires the Th2-priming effects of thymic stromal lymphopoietin (TSLP) and its effects on basophils [54]. This epicutaneous priming route seems sufficient to exert intestinal influences, as shown by a significant expansion of mast cells within the jejunum and the fact that the allergen-specific IgE induced during skin exposure was necessary for anaphylaxis to occur on oral challenge [55]. Conversely, oral exposure to allergen has also been shown to protect mice from skin allergic immune responses in a similar atopic dermatitis model [56]. Taken together, the evidence seems to implicate the skin as a route through which food allergic reactivity can be initiated, while oral exposure is generally associated with tolerance.

Evolution of Food Allergies: from Childhood to Adulthood

Development of Natural Tolerance

Clinically, it is well established that food allergy can be transient, and that many children eventually ‘outgrow’ their food allergies, a phenomenon termed “natural tolerance”. This occurs at different rates for different food allergens: studies have shown that sixty to seventy five percent of those with cows milk allergy outgrow by the age of five years [57], while this number is only around fifty percent for egg [58] and twenty two percent for peanut [59]. For cow’s milk, natural tolerance was shown to be associated with increased IgG4/IgE ratios [60]. Recent data from our group demonstrated a significant increase in Tregs in the blood of children who had acquired natural tolerance of egg or peanut, compared to either healthy control or food allergic children [61]. Critically, IL-10-expressing Tregs, including the Tr1 subset of pTregs that are generally FoxP3 negative [62], were increased upon relevant allergen stimulation in natural tolerance patients but not in allergic individuals. Importantly, this type of outgrowth seems to reflect a true conversion from reactivity to life-long tolerance while many attempts to induce tolerance by immunotherapy approaches have predominantly resulted in desensitization and return of reactivity on cessation of treatment [63].

Adult-Onset Food Allergy

In the timeline of food allergy development, most of the focus has been on the early-life events and initiation or prevention of food allergy in infants. The LEAP study has helped to emphasize how failure to support early dietary exposure to food allergens might have facilitated a rise in food allergy due to a lack in the engagement of oral tolerance mechanisms. However, food allergy is also seen in adults, despite an extensive history of dietary intake of foods they subsequently become reactive to. To this point, we reported on 171 cases of physician-diagnosed food allergy that developed in adulthood, with the average age of diagnosis being thirty one years old [64]. The foods represented in these adult-onset food allergy patients were similar to those observed in childhood food allergy, including fish, nuts, soy, milk, eggs and peanuts. This raises an intriguing immunological question regarding the sustainability of oral tolerance and factors that might overcome its effects and drive sensitization and reactivity. It is well known that IgE-associated sensitization to foods is significantly more common than the frequency of people who exhibit food reactivity [65–67], a significant problem in the diagnosis of food allergy. It remains to be determined if adult-onset food allergy reflects acquisition of sensitization and reactivity or reactivity in the face of prior sensitization. Adult-onset food allergy also emphasizes the point that food allergy is not only due to failures to initiate tolerance but that oral tolerance to dietary antigens is a process that seemingly requires maintenance and can be reverted.

Interactions between Responses to Dietary Antigens and the Microbiota

The evidence from food allergy that is outlined above has begun to provide new avenues of investigation into the processes that balance our responses to dietary antigens. The immune system in the gastrointestinal tract in particular is capable of both profound inflammatory responses (e.g. towards pathogenic microbes) but also tolerance. This tolerance has been

recently brought to the forefront in the context of the recognition and responses to the microbiota that coexists within our intestinal tract, but several studies have connected this to dietary tolerance as well. Much of the immune mediators implicated in early-life aspects of food allergy and discussed above have also been implicated in shaping interactions with the intestinal microflora. For example, IgA in breast milk was demonstrated to alter the microbiome composition and to dramatically shift the flora that persisted into adulthood in mice [31]. This might be important in the context of generating efficient tolerance responses within the intestine since it has been clearly demonstrated that certain species of bacteria, particularly a specific collection of toxin and virulence factor lacking strains in the Clostridia class, are critical for the generation of FoxP3⁺ Tregs within the colon [68, 69]. Importantly, germ-free mice or neonates who undergo antibiotic depletion of their commensal flora have been shown to have a heightened susceptibility to food allergy, attributed to alterations in IL-22 that were induced by Clostridia [70]. In mice with an engineered gain-of-function mutation in the IL-4 receptor α chain and who exhibit a heightened susceptibility to oral sensitization versus tolerance, a profoundly altered microbiome signature was observed and this was sufficient to impart susceptibility when transferred into germ-free wild-type mice [71]. This has important implications, since it supports the conclusion that the personalized genetic landscape of an individual might be a contributing factor in shaping the balance of their own commensal microbiota, and that both the genetic landscape and the microbiota may contribute to allergic susceptibility. The mechanism for this susceptibility in IL-4R α mutant mice was attributed to deficiencies in Tregs numbers and functions. Subsequent studies using this mutant mouse have described a reprogramming towards a Th2-like cell that fails in its ability to maintain oral tolerance [72], perhaps through a cross-talk with type 2 innate lymphoid cells (ILC2) [73].

While the microbiota appears to play a critical role in the balance between intestinal tolerance and sensitization, recent discoveries have helped shine a spotlight on the important role that dietary proteins also play. A study of intestinal biology during development showed that the introduction of solid foods led to massive changes in bacterial composition, with a transition to a more stable community, but also to increased levels of short chain fatty acids, vitamin biosynthesis, carbohydrate utilization [74]. More specifically, a recent study focused on the direct influences of dietary antigens in intestinal homeostasis and showed a previously underappreciated biology [75]. By utilizing germ-free mice that were maintained on an elemental diet – an amino-acid based diet that is devoid of proteins and is utilized clinically in some food allergy patients [76] – they were able to effectively eliminate exposure to dietary antigens and to dissect the alterations seen in the intestinal homeostasis. Their findings elegantly demonstrate a bifurcation of the intestinal pTreg communities and the mechanisms needed for their initiation, with colonic pTregs being driven by microbiota and those in the small intestine being driven by dietary antigen exposure. The cells at these two distinct locations could also be discriminated from each other based on the expression of the transcription factor retinoid-related orphan receptor gamma t (ROR γ t), potentially suggesting differences in developmental cues or functional properties. While ROR γ t is critical for pro-inflammatory Th17 responses, the importance of ROR γ t expressing cells in intestinal tolerance, both innate and adaptive cell types, has increasingly become evident and was recently reviewed [77]. The study also demonstrates an alteration in the frequencies of

specialized tolerogenic dendritic cells (CD103+CD11b+), which have been shown to interact with macrophages at intestinal gap junctions to establish oral tolerance [78]. Mechanistically, these findings align with recent reports these cells are regulated by high-fiber diet and short-chain fatty acids produced by some microbial species and are required to protect against food allergy in mice [79]. While dietary proteins were required for small intestinal pTreg development and introduction of solid food was sufficient to induce their development, questions remain, particularly in how this relates to clinical tolerance in humans. Somewhat surprisingly, mice that did not develop these pTregs still did not exhibit sensitization or reactivity upon introduction of standard chow. Instead, the adoptive transfer of antigen-specific T cells was utilized to elicit allergen-specific inflammatory responses and these were enhanced in the absence of pTregs. One conclusion from this is that dietary-induced pTregs in the small intestine are critical for suppressing food allergy in sensitized individuals, a state of control that is observed in many people who possess evidence of sensitization to food allergens but no apparent reactivity [67]. However, the underlying mechanisms of oral tolerance that prevent the initiation of sensitization in the majority of individuals remain unclear.

Concluding Remarks

Our understanding of the mechanisms that balance tolerance versus food allergy are beginning to provide insights into how the intestinal immune system is influenced by a complex process of life exposures. Three key recent studies spotlight this

- Differences in monocyte responses from the cord blood of children destined to become food allergic implicates in utero and maternal influences [4].
- Successful treatment of susceptible children with early introduction of peanut consumption implicates early-life and diet-driven protective processes [37].
- Diet protein-induced pTreg induction within the small intestine defines a previously unappreciated direct mechanism through which diet alters immune balance [75].

These findings implicate both innate balances but also the importance of acquired protection.

Several questions remain in our understanding of the mechanisms working towards or away from food allergy (see Outstanding Questions). In particular, the differences between those who undergo conversion seems particularly important. A better understanding of the events driving natural tolerance acquisition are likely to guide therapeutic studies and the development of novel treatments while the events leading to loss of tolerance in some adults is perplexing. Are these adults failing to maintain diet-associated pTregs or is sensitization due to some non-specific failure in barriers? How do dietary processes interact with microbiota-driven signals over the course of food allergy development or protection?

Outstanding Questions

- What are the mechanisms driving natural tolerance acquisition?
- What overcomes tolerance in some adults and promotes food allergy?
- Diet-associated p Tregs are important for controlling inflammation post-sensitization, but most individuals do not get sensitized at all. What are the mechanisms that prevent sensitization?
- How do dietary processes interact with microbiota-driven signals over the course of food allergy development or protection?

Certainly, these recent developments represent the beginning of a new age in our clinical and mechanistic understanding of food allergies; one that Albert Rowe most likely would not have foreseen.

References

1. Rowe AH. Abdominal Food Allergy: Its History, Symptomatology, Diagnosis and Treatment. *Cal West Med.* 1928; 29:317–22. [PubMed: 18740859]
2. Johnston LK, Chien KB, Bryce PJ. The immunology of food allergy. *J Immunol.* 2014; 192:2529–34. [PubMed: 24610821]
3. Pastor-Vargas C, Maroto AS, Diaz-Perales A, Villalba M, Esteban V, Ruiz-Ramos M, de Alba MR, Vivanco F, Cuesta-Herranz J. Detection of major food allergens in amniotic fluid: initial allergenic encounter during pregnancy. *Pediatr Allergy Immunol.* 2016; doi: 10.1111/pai.12608
4. Zhang Y, Collier F, Naselli G, Saffery R, Tang ML, Allen KJ, Ponsonby AL, Harrison LC, Vuillermin P, Group BISI. Cord blood monocyte-derived inflammatory cytokines suppress IL-2 and induce nonclassic “T(H)2-type” immunity associated with development of food allergy. *Sci Transl Med.* 2016; 8:321ra8.
5. Mori T, Kato J, Sakurai M, Hashimoto N, Kohashi S, Hashida R, Saburi M, Kikuchi T, Yamane Y, Hoshino K, Okamoto S. New-onset food allergy following cord blood transplantation in adult patients. *Bone Marrow Transplant.* 2016; 51:295–6. [PubMed: 26457912]
6. Feliu J, Clay J, Raj K, Barber L, Devlia V, Shaw B, Pagliuca A, Mufti G. Transplant-acquired food allergy (TAFA) following cord blood stem cell transplantation in two adult patients with haematological malignancies. *Br J Haematol.* 2014; 167:426–8. [PubMed: 24962133]
7. Hinz D, Bauer M, Roder S, Olek S, Huehn J, Sack U, Borte M, Simon JC, Lehmann I, Herberth G, group Ls. Cord blood Tregs with stable FOXP3 expression are influenced by prenatal environment and associated with atopic dermatitis at the age of one year. *Allergy.* 2012; 67:380–9. [PubMed: 22187950]
8. Schaub B, Liu J, Hoppler S, Haug S, Sattler C, Lluís A, Illi S, von Mutius E. Impairment of T-regulatory cells in cord blood of atopic mothers. *J Allergy Clin Immunol.* 2008; 121:1491–9. 99 e1–13. [PubMed: 18539197]
9. Gerhold K, Avagyan A, Reichert E, Seib C, Van DV, Luger EO, Hutloff A, Hamelmann E. Prenatal allergen exposures prevent allergen-induced sensitization and airway inflammation in young mice. *Allergy.* 2012; 67:353–61. [PubMed: 22229690]
10. Lopez-Exposito I, Song Y, Jarvinen KM, Srivastava K, Li XM. Maternal peanut exposure during pregnancy and lactation reduces peanut allergy risk in offspring. *J Allergy Clin Immunol.* 2009; 124:1039–46. [PubMed: 19895992]
11. Song Y, Liu C, Hui Y, Srivastava K, Zhou Z, Chen J, Miller RL, Finkelman FD, Li XM. Maternal allergy increases susceptibility to offspring allergy in association with TH2-biased epigenetic alterations in a mouse model of peanut allergy. *J Allergy Clin Immunol.* 2014; 134:1339–45. e7. [PubMed: 25441650]

12. Scholl I, Ackermann U, Ozdemir C, Blumer N, Dicke T, Sel S, Sel S, Wegmann M, Szalai K, Knittelfelder R, Untersmayr E, Scheiner O, Garn H, Jensen-Jarolim E, Renz H. Anti-ulcer treatment during pregnancy induces food allergy in mouse mothers and a Th2-bias in their offspring. *FASEB J*. 2007; 21:1264–70. [PubMed: 17227952]
13. Herz U, Joachim R, Ahrens B, Scheffold A, Radbruch A, Renz H. Allergic sensitization and allergen exposure during pregnancy favor the development of atopy in the neonate. *Int Arch Allergy Immunol*. 2001; 124:193–6. [PubMed: 11306966]
14. Martino DJ, Saffery R, Allen KJ, Prescott SL. Epigenetic modifications: mechanisms of disease and biomarkers of food allergy. *Curr Opin Immunol*. 2016; 42:9–15. [PubMed: 27218660]
15. Schocker F, Baumert J, Kull S, Petersen A, Becker WM, Jappe U. Prospective investigation on the transfer of Ara h 2, the most potent peanut allergen, in human breast milk. *Pediatr Allergy Immunol*. 2016; 27:348–55. [PubMed: 26842773]
16. Vadas P, Wai Y, Burks W, Perelman B. Detection of peanut allergens in breast milk of lactating women. *JAMA*. 2001; 285:1746–8. [PubMed: 11277829]
17. Bernard H, Ah-Leung S, Drumare MF, Feraudet-Tarisse C, Verhasselt V, Wal JM, Creminon C, Adel-Patient K. Peanut allergens are rapidly transferred in human breast milk and can prevent sensitization in mice. *Allergy*. 2014; 69:888–97. [PubMed: 24773443]
18. Jarvinen KM, Suarez-Farinas M, Savilahti E, Sampson HA, Berin MC. Immune factors in breast milk related to infant milk allergy are independent of maternal atopy. *J Allergy Clin Immunol*. 2015; 135:1390–3. e1–6. [PubMed: 25533649]
19. Chen Z, O'Shea JJ. Th17 cells: a new fate for differentiating helper T cells. *Immunol Res*. 2008; 41:87–102. [PubMed: 18172584]
20. Jepsen AA, Chawes BL, Carson CG, Schoos AM, Thysen AH, Waage J, Brix S, Bisgaard H. High breast milk IL-1 β level is associated with reduced risk of childhood eczema. *Clin Exp Allergy*. 2016; doi: 10.1111/cea.12770
21. Goldsmith AJ, Koplin JJ, Lowe AJ, Tang ML, Matheson MC, Robinson M, Peters R, Dharmage SC, Allen KJ. Formula and breast feeding in infant food allergy: A population-based study. *J Paediatr Child Health*. 2016; 52:377–84. [PubMed: 27145499]
22. Lucciolli S, Zhang Y, Verrill L, Ramos-Valle M, Kwegyir-Afful E. Infant feeding practices and reported food allergies at 6 years of age. *Pediatrics*. 2014; 134(Suppl 1):S21–8. [PubMed: 25183751]
23. Osborn DA, Sinn J. Formulas containing hydrolysed protein for prevention of allergy and food intolerance in infants. *Cochrane Database Syst Rev*. 2003;CD003664. [PubMed: 14583987]
24. Jarvinen KM, Westfall J, De Jesus M, Mantis NJ, Carroll JA, Metzger DW, Sampson HA, Berin MC. Role of Maternal Dietary Peanut Exposure in Development of Food Allergy and Oral Tolerance. *PLoS One*. 2015; 10:e0143855. [PubMed: 26656505]
25. Pabst O, Mowat AM. Oral tolerance to food protein. *Mucosal Immunol*. 2012; 5:232–9. [PubMed: 22318493]
26. Fontenot JD, Dooley JL, Farr AG, Rudensky AY. Developmental regulation of Foxp3 expression during ontogeny. *J Exp Med*. 2005; 202:901–6. [PubMed: 16203863]
27. Burchill MA, Yang J, Vang KB, Moon JJ, Chu HH, Lio CW, Vegoe AL, Hsieh CS, Jenkins MK, Farrar MA. Linked T cell receptor and cytokine signaling govern the development of the regulatory T cell repertoire. *Immunity*. 2008; 28:112–21. [PubMed: 18199418]
28. Chen W, Jin W, Hardegen N, Lei KJ, Li L, Marinos N, McGrady G, Wahl SM. Conversion of peripheral CD4+CD25– naive T cells to CD4+CD25+ regulatory T cells by TGF- β induction of transcription factor Foxp3. *J Exp Med*. 2003; 198:1875–86. [PubMed: 14676299]
29. Josefowicz SZ, Niec RE, Kim HY, Treuting P, Chinen T, Zheng Y, Umetsu DT, Rudensky AY. Extrathymically generated regulatory T cells control mucosal TH2 inflammation. *Nature*. 2012; 482:395–9. [PubMed: 22318520]
30. Gustafson CE, Higbee D, Yeckes AR, Wilson CC, De Zoeten EF, Jedlicka P, Janoff EN. Limited expression of APRIL and its receptors prior to intestinal IgA plasma cell development during human infancy. *Mucosal Immunol*. 2014; 7:467–77. [PubMed: 24045575]
31. Rogier EW, Frantz AL, Bruno ME, Wedlund L, Cohen DA, Stromberg AJ, Kaetzel CS. Secretory antibodies in breast milk promote long-term intestinal homeostasis by regulating the gut

microbiota and host gene expression. *Proc Natl Acad Sci U S A*. 2014; 111:3074–9. [PubMed: 24569806]

32. Jarvinen KM, Westfall JE, Seppo MS, James AK, Tsuang AJ, Feustel PJ, Sampson HA, Berin C. Role of maternal elimination diets and human milk IgA in the development of cow's milk allergy in the infants. *Clin Exp Allergy*. 2014; 44:69–78. [PubMed: 24164317]
33. Weisse K, Winkler S, Hirche F, Herberth G, Hinz D, Bauer M, Roder S, Rolle-Kampczyk U, von Bergen M, Olek S, Sack U, Richter T, Diez U, Borte M, Stangl GI, Lehmann I. Maternal and newborn vitamin D status and its impact on food allergy development in the German LINA cohort study. *Allergy*. 2013; 68:220–8. [PubMed: 23253182]
34. Baek JH, Shin YH, Chung IH, Kim HJ, Yoo EG, Yoon JW, Jee HM, Chang YE, Han MY. The link between serum vitamin D level, sensitization to food allergens, and the severity of atopic dermatitis in infancy. *J Pediatr*. 2014; 165:849–54. e1. [PubMed: 25108543]
35. Zeng H, Chi H. Metabolic control of regulatory T cell development and function. *Trends Immunol*. 2015; 36:3–12. [PubMed: 25248463]
36. Du Toit G, Katz Y, Sasieni P, Meshier D, Maleki SJ, Fisher HR, Fox AT, Turcanu V, Amir T, Zadik-Mnuhin G, Cohen A, Livne I, Lack G. Early consumption of peanuts in infancy is associated with a low prevalence of peanut allergy. *J Allergy Clin Immunol*. 2008; 122:984–91. [PubMed: 19000582]
37. Du Toit G, Roberts G, Sayre PH, Bahnson HT, Radulovic S, Santos AF, Brough HA, Phippard D, Basting M, Feeney M, Turcanu V, Sever ML, Gomez Lorenzo M, Plaut M, Lack G, Team LS. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med*. 2015; 372:803–13. [PubMed: 25705822]
38. Simpson EL, Eichenfield LF, Ellis CN, Mancini AJ, Paller AS. Current issues in atopic comorbidities and preventing the atopic march. *Semin Cutan Med Surg*. 2012; 31:S6–9. [PubMed: 23021784]
39. Davies AM, Sutton BJ. Human IgG4: a structural perspective. *Immunol Rev*. 2015; 268:139–59. [PubMed: 26497518]
40. Wright BL, Kulis M, Orgel KA, Burks AW, Dawson P, Henning AK, Jones SM, Wood RA, Sicherer SH, Lindblad RW, Stablein D, Leung DY, Vickery BP, Sampson HA. Consortium of Food Allergy R, Component-resolved analysis of IgA, IgE, and IgG4 during egg OIT identifies markers associated with sustained unresponsiveness. *Allergy*. 2016; doi: 10.1111/all.12895
41. Wood RA, Togias A, Wildfire J, Visness CM, Matsui EC, Gruchalla R, Hershey G, Liu AH, O'Connor GT, Pongracic JA, Zoratti E, Little F, Granada M, Kennedy S, Durham SR, Shamji MH, Busse WW. Development of cockroach immunotherapy by the Inner-City Asthma Consortium. *J Allergy Clin Immunol*. 2014; 133:846–52. e6. [PubMed: 24184147]
42. Caubet JC, Bencharitiwong R, Moshier E, Godbold JH, Sampson HA, Nowak-Wegrzyn A. Significance of ovomucoid- and ovalbumin-specific IgE/IgG(4) ratios in egg allergy. *J Allergy Clin Immunol*. 2012; 129:739–47. [PubMed: 22277199]
43. Du Toit G, Sayre PH, Roberts G, Sever ML, Lawson K, Bahnson HT, Brough HA, Santos AF, Harris KM, Radulovic S, Basting M, Turcanu V, Plaut M, Lack G. Immune Tolerance Network L-OST, Effect of Avoidance on Peanut Allergy after Early Peanut Consumption. *N Engl J Med*. 2016; 374:1435–43. [PubMed: 26942922]
44. Perkin MR, Logan K, Tseng A, Raji B, Ayis S, Peacock J, Brough H, Marrs T, Radulovic S, Craven J, Flohr C, Lack G, Team EATS. Randomized Trial of Introduction of Allergenic Foods in Breast-Fed Infants. *N Engl J Med*. 2016; 374:1733–43. [PubMed: 26943128]
45. Sicherer SH, Sampson HA. Food allergy: Epidemiology, pathogenesis, diagnosis, and treatment. *J Allergy Clin Immunol*. 2014; 133:291–307. quiz 08. [PubMed: 24388012]
46. Greenhawt M. Food allergy quality of life and living with food allergy. *Curr Opin Allergy Clin Immunol*. 2016; 16:284–90. [PubMed: 27070333]
47. Nguyen TA, Leonard SA, Eichenfield LF. An Update on Pediatric Atopic Dermatitis and Food Allergies. *J Pediatr*. 2015; 167:752–6. [PubMed: 26118932]
48. McAleer MA, Irvine AD. The multifunctional role of filaggrin in allergic skin disease. *J Allergy Clin Immunol*. 2013; 131:280–91. [PubMed: 23374260]

49. Asai Y, Greenwood C, Hull PR, Alizadehfar R, Ben-Shoshan M, Brown SJ, Campbell L, Michel DL, Bussieres J, Rousseau F, Fujiwara TM, Morgan K, Irvine AD, McLean WH, Clarke A. Filaggrin gene mutation associations with peanut allergy persist despite variations in peanut allergy diagnostic criteria or asthma status. *J Allergy Clin Immunol.* 2013; 132:239–42. [PubMed: 23684069]
50. Brough HA, Simpson A, Makinson K, Hankinson J, Brown S, Douiri A, Belgrave DC, Penagos M, Stephens AC, McLean WH, Turcanu V, Nicolaou N, Custovic A, Lack G. Peanut allergy: effect of environmental peanut exposure in children with filaggrin loss-of-function mutations. *J Allergy Clin Immunol.* 2014; 134:867–75. e1. [PubMed: 25282568]
51. von Mutius E. Allergies, infections and the hygiene hypothesis--the epidemiological evidence. *Immunobiology.* 2007; 212:433–9. [PubMed: 17544828]
52. Lack G. Epidemiologic risks for food allergy. *J Allergy Clin Immunol.* 2008; 121:1331–6. [PubMed: 18539191]
53. Marsella R. Experimental model for peanut allergy by epicutaneous sensitization in atopic beagle dogs. *Exp Dermatol.* 2015; 24:711–2. [PubMed: 26014917]
54. Noti M, Kim BS, Siracusa MC, Rak GD, Kubo M, Moghaddam AE, Sattentau QA, Comeau MR, Spergel JM, Artis D. Exposure to food allergens through inflamed skin promotes intestinal food allergy through the thymic stromal lymphopoietin-basophil axis. *J Allergy Clin Immunol.* 2014; 133:1390–9. 99 e1–6. [PubMed: 24560412]
55. Bartnikas LM, Gurish MF, Burton OT, Leisten S, Janssen E, Oettgen HC, Beaupre J, Lewis CN, Austen KF, Schulte S, Hornick JL, Geha RS, Oyoshi MK. Epicutaneous sensitization results in IgE-dependent intestinal mast cell expansion and food-induced anaphylaxis. *J Allergy Clin Immunol.* 2013; 131:451–60. e1–6. [PubMed: 23374269]
56. Baek JO, Roh JY, Jung Y. Oral tolerance inhibits atopic dermatitis-like type 2 inflammation in mice by modulating immune microenvironments. *Allergy.* 2016
57. Saarinen KM, Pelkonen AS, Makela MJ, Savilahti E. Clinical course and prognosis of cow's milk allergy are dependent on milk-specific IgE status. *J Allergy Clin Immunol.* 2005; 116:869–75. [PubMed: 16210063]
58. Peters RL, Dharmage SC, Gurrin LC, Koplin JJ, Ponsonby AL, Lowe AJ, Tang ML, Tey D, Robinson M, Hill D, Czech H, Thiele L, Osborne NJ, Allen KJ. HealthNuts s. The natural history and clinical predictors of egg allergy in the first 2 years of life: a prospective, population-based cohort study. *J Allergy Clin Immunol.* 2014; 133:485–91. [PubMed: 24373356]
59. Peters RL, Allen KJ, Dharmage SC, Koplin JJ, Dang T, Tilbrook KP, Lowe A, Tang ML, Gurrin LC, HealthNuts S. Natural history of peanut allergy and predictors of resolution in the first 4 years of life: A population-based assessment. *J Allergy Clin Immunol.* 2015; 135:1257–66. e1–2. [PubMed: 25725989]
60. Savilahti EM, Rantanen V, Lin JS, Karinen S, Saarinen KM, Goldis M, Makela MJ, Hautaniemi S, Savilahti E, Sampson HA. Early recovery from cow's milk allergy is associated with decreasing IgE and increasing IgG4 binding to cow's milk epitopes. *J Allergy Clin Immunol.* 2010; 125:1315–21. e9. [PubMed: 20462631]
61. Qamar N, Fishbein AB, Erickson KA, Cai M, Szychlinski C, Bryce PJ, Schleimer RP, Fuleihan RL, Singh AM. Naturally occurring tolerance acquisition to foods in previously allergic children is characterized by antigen specificity and associated with increased subsets of regulatory T cells. *Clin Exp Allergy.* 2015; 45:1663–72. [PubMed: 25989379]
62. Roncarolo MG, Gregori S, Bacchetta R, Battaglia M. Tr1 cells and the counter-regulation of immunity: natural mechanisms and therapeutic applications. *Curr Top Microbiol Immunol.* 2014; 380:39–68. [PubMed: 25004813]
63. Burbank AJ, Burks W. Food specific oral immunotherapy: a potential treatment for food allergy. *Expert Rev Gastroenterol Hepatol.* 2015; 9:1147–59. [PubMed: 26145713]
64. Kamdar TA, Peterson S, Lau CH, Saltoun CA, Gupta RS, Bryce PJ. Prevalence and characteristics of adult-onset food allergy. *J Allergy Clin Immunol Pract.* 2015; 3:114–5. e1. [PubMed: 25577631]
65. Mahesh PA, Wong GW, Ogorodova L, Potts J, Leung TF, Fedorova O, Holla AD, Fernandez-Rivas M, Clare Mills EN, Kummeling I, Versteeg SA, van Ree R, Yazdanbakhsh M, Burney P.

- Prevalence of food sensitization and probable food allergy among adults in India: the EuroPrevall INCO study. *Allergy*. 2016; 71:1010–9. [PubMed: 27297800]
66. Burney PG, Potts J, Kummeling I, Mills EN, Clausen M, Dubakiene R, Barreales L, Fernandez-Perez C, Fernandez-Rivas M, Le TM, Knulst AC, Kowalski ML, Lidholm J, Ballmer-Weber BK, Braun-Fahlander C, Mustakov T, Kralimarkova T, Popov T, Sakellariou A, Papadopoulos NG, Versteeg SA, Zuidmeer L, Akkerdaas JH, Hoffmann-Sommergruber K, van Ree R. The prevalence and distribution of food sensitization in European adults. *Allergy*. 2014; 69:365–71. [PubMed: 24372074]
 67. McGowan EC, Peng RD, Salo PM, Zeldin DC, Keet CA. Changes in Food-Specific IgE Over Time in the National Health and Nutrition Examination Survey (NHANES). *J Allergy Clin Immunol Pract*. 2016; 4:713–20. [PubMed: 27133095]
 68. Atarashi K, Tanoue T, Oshima K, Suda W, Nagano Y, Nishikawa H, Fukuda S, Saito T, Narushima S, Hase K, Kim S, Fritz JV, Wilmes P, Ueha S, Matsushima K, Ohno H, Olle B, Sakaguchi S, Taniguchi T, Morita H, Hattori M, Honda K. Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. *Nature*. 2013; 500:232–6. [PubMed: 23842501]
 69. Tanoue T, Atarashi K, Honda K. Development and maintenance of intestinal regulatory T cells. *Nat Rev Immunol*. 2016; 16:295–309. [PubMed: 27087661]
 70. Stefka AT, Feehley T, Tripathi P, Qiu J, McCoy K, Mazmanian SK, Tjota MY, Seo GY, Cao S, Theriault BR, Antonopoulos DA, Zhou L, Chang EB, Fu YX, Nagler CR. Commensal bacteria protect against food allergen sensitization. *Proc Natl Acad Sci U S A*. 2014; 111:13145–50. [PubMed: 25157157]
 71. Noval Rivas M, Burton OT, Wise P, Zhang YQ, Hobson SA, Garcia Lloret M, Chehoud C, Kuczynski J, DeSantis T, Warrington J, Hyde ER, Petrosino JF, Gerber GK, Bry L, Oettgen HC, Mazmanian SK, Chatila TA. A microbiota signature associated with experimental food allergy promotes allergic sensitization and anaphylaxis. *J Allergy Clin Immunol*. 2013; 131:201–12. [PubMed: 23201093]
 72. Noval Rivas M, Burton OT, Wise P, Charbonnier LM, Georgiev P, Oettgen HC, Rachid R, Chatila TA. Regulatory T cell reprogramming toward a Th2-cell-like lineage impairs oral tolerance and promotes food allergy. *Immunity*. 2015; 42:512–23. [PubMed: 25769611]
 73. Noval Rivas M, Burton OT, Oettgen HC, Chatila T. IL-4 production by group 2 innate lymphoid cells promotes food allergy by blocking regulatory T-cell function. *J Allergy Clin Immunol*. 2016; doi: 10.1016/j.jaci.2016.02.030
 74. Koenig JE, Spor A, Scalfone N, Fricker AD, Stombaugh J, Knight R, Angenent LT, Ley RE. Succession of microbial consortia in the developing infant gut microbiome. *Proc Natl Acad Sci U S A*. 2011; 108(Suppl 1):4578–85. [PubMed: 20668239]
 75. Kim KS, Hong SW, Han D, Yi J, Jung J, Yang BG, Lee JY, Lee M, Surh CD. Dietary antigens limit mucosal immunity by inducing regulatory T cells in the small intestine. *Science*. 2016; 351:858–63. [PubMed: 26822607]
 76. Gonsalves N, Kagalwalla AF. Dietary treatment of eosinophilic esophagitis. *Gastroenterol Clin North Am*. 2014; 43:375–83. [PubMed: 24813522]
 77. Ohnmacht C. Tolerance to the Intestinal Microbiota Mediated by ROR(gamma)(+) Cells. *Trends Immunol*. 2016; 37:477–86. [PubMed: 27255270]
 78. Mazzini E, Massimiliano L, Penna G, Rescigno M. Oral tolerance can be established via gap junction transfer of fed antigens from CX3CR1(+) macrophages to CD103(+) dendritic cells. *Immunity*. 2014; 40:248–61. [PubMed: 24462723]
 79. Tan J, McKenzie C, Vuillermin PJ, Govere G, Vinuesa CG, Mebius RE, Macia L, Mackay CR. Dietary Fiber and Bacterial SCFA Enhance Oral Tolerance and Protect against Food Allergy through Diverse Cellular Pathways. *Cell Rep*. 2016; 15:2809–24. [PubMed: 27332875]

Trends Box

Food allergy has become an important clinical problem as its prevalence increases across the world.

Successful clinical trials on early introduction of foods have demonstrated the importance of early oral exposure in prevention.

The sampling of dietary proteins by the intestinal immune system imparts specialized and localized tolerance processes that occur independently from microbe-driven responses.

Figure 1. Factors Influencing Tolerance and Allergy to Dietary Antigens Across Development
 Throughout life, a progression of positive and negative factors might contribute to the fate of immune responsiveness to dietary antigens. In childhood, early events related to establishment of oral tolerance and control of antigen priming processes protect from becoming sensitized. Later in life, natural tolerance might develop in some food allergic individuals while some adults develop de novo food allergy. Tolerogenic factors are pictured in green, while factors involved in allergy development are pictured in red. Abbreviations used: TSLP, thymic stromal lymphopoietin, Treg, regulatory T cell