Published in final edited form as:

J Am Chem Soc. 2016 September 7; 138(35): 11360–11367. doi:10.1021/jacs.6b07221.

Iron(II)-Catalyzed Intermolecular Aminofluorination of Unfunctionalized Olefins Using Fluoride Ion

Deng-Fu Lu[†], Cheng-Liang Zhu[†], Jeffrey D. Sears, and Hao Xu^{*}

Department of Chemistry, Georgia State University, 100 Piedmont Avenue SE, Atlanta, Georgia 30303, United States

Abstract

We herein report a new catalytic method for intermolecular olefin aminofluorination using earth-abundant iron catalysts and nucleophilic fluoride ion. This method tolerates a broad range of unfunctionalized olefins, especially nonstyrenyl olefins that are incompatible with existing olefin aminofluorination methods. This new iron-catalyzed process directly converts readily available olefins to internal vicinal fluoro carbamates with high regioselectivity (*N* vs *F*), many of which are difficult to prepare using known methods. Preliminary mechanistic studies demonstrate that it is possible to exert asymmetric induction using chiral iron catalysts and that both an iron-nitrenoid and carbocation species may be reactive intermediates.

Graphical Abstract

INTRODUCTION

Selective incorporation of fluorine into organic molecules is an emerging area with increasing importance in organic synthesis because fluorine atoms are present in numerous pharmaceuticals and biological probes.¹ In particular, intermolecular olefin functionalization via fluorine atom transfer represents a convenient approach to synthesize high-value building blocks with fluorine-containing stereogenic centers.² Among a variety of olefin fluorination

^{*}Corresponding Author: hxu@gsu.edu.

[†]Author Contributions

D.-F.L. and C.-L.Z. contributed equally.

Notes

The authors declare no competing financial interest.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.6b07221.

Experimental procedure, characterization data for all new compounds, selected NMR spectra and HPLC traces (PDF)

si_002.cif for compound **6** (CIF)

si_003.cif for Fe(**L1**)₂(BF₄)₂ (CIF)

si_004.cif for compound **7d** (CIF)

si_005.cif for compound **11a** (CIF)

reactions, the *intermolecular amino-fluorination of unfunctionalized olefins* is of great value since it can directly convert simple olefins to vicinal fluoro primary amines which are important structural motifs in medicinal chemistry.³ However, development of this catalytic reaction has been challenging; only a few methods were reported, and most of them are exclusively compatible with styrenyl olefins.⁴

Amid these discoveries, Stavber reported Ritter-type electrophilic olefin vicinal fluoro acetamidation in acetonitrile using electrophilic fluorination reagent Accufluor NFTh, through β -fluorocarbenium intermediates, to afford terminal fluorides; however, the substrate scope was predominantly limited to styrenyl olefins.^{4a} Liu reported palladium-catalyzed styrene aminofluorination using electrophilic NFSI, which affords terminal sulfonamido fluorides;^{4b} the same group later documented a related styrene aminofluorination using AgF that provides internal sulfonamido fluorides.^{4c}

The existing intermolecular olefin aminofluorination methods are mechanistically illuminating; however, there still exist significant gaps for this important transformation. First, there has yet to be developed a method for catalytic amino-fluorination of *a broad range of unfunctionalized olefins* (especially for nonstyrenyl olefins). Next, the feasibility of asymmetric catalysis for intermolecular aminofluorination of unfunctionalized olefins has not yet been demonstrated.⁵ Additionally, the use of earth-abundant iron catalysts to promote olefin aminofluorination with simple nucleophilic fluoride ion is desirable,⁶ especially considering potential applications in ¹⁸F PET imaging; however, *iron-catalyzed intermolecular olefin fluorination using nucleophilic fluoride remains unreported*,⁷ presumably because of the high BDE of the Fe–F bond,⁸ the low solubility of iron fluoride species, and low nucleophilicity of solvated fluoride ion in organic solvents.⁹

Herein, we report an iron-catalyzed intermolecular amino-fluorination method for a broad range of unfunctionalized olefins, including nonstyrenyl olefins, using fluoride ion (Scheme 1). This new reaction readily affords a variety of synthetically valuable vicinal fluoro carbamates with high regioselectivity (*N* vs *F*), many of which are difficult to prepare using existing methods.¹⁰ We further demonstrate that it is possible to modulate the enantioselectivity for aminofluorination of cyclic olefins. Additionally, preliminary mechanistic studies suggest a possible new olefin aminofluorination pathway in which both an iron-nitrenoid and carbocation species may be involved in a stepwise nitrogen and fluorine atom transfer.

RESULTS AND DISCUSSION

We previously reported an iron-catalyzed method for *intra-molecular* olefin aminofluorination using fluoride ion in which iron(II)–bidentate ligand complexes catalyze the reaction in the presence of Et₃N·3HF (Scheme 2A);¹¹ however, this type of catalyst was ineffective to catalyze the *intermolecular* olefin aminofluorination because of lack of reactivity (Scheme 2B). We also reported an iron-catalyzed method for *intermolecular olefin aminohydroxylation* using iron(II)–tridentate ligand complexes;¹² however, these catalysts were unable to promote the *intermolecular olefin aminofluorination* in the presence of Et₃N·3HF, possibly due to catalyst deactivation (Scheme 2B). We therefore explored a range

of catalysts, oxidants, and fluorination reagents to search for a *stable yet reactive* catalyst in the presence of the nucleophilic fluoride ion.

To determine the possible cause of the aforementioned catalyst deactivation (Scheme 2B), we exposed an *in situ* generated $\text{Fe}(\text{BF}_4)_2\text{-L1}$ catalyst (1:1 ratio), previously discovered for styrene aminohydroxylation, to an intra-molecular aminofluorination condition in the absence of any olefin or oxidant (Scheme 3). Insoluble FeF_2 and a soluble iron complex were obtained, and X-ray crystallographic analysis revealed its structure as $\text{Fe}(\text{L1})_2(\text{BF}_4)_2$, which, surprisingly, is inactive for styrene aminohydroxylation.¹³ This result suggests that fluoride species may disrupt the $\text{Fe}(\text{BF}_4)_2\text{-L1}$ complex through iron sequestering and therefore lead to the inactive $\text{Fe}(\text{L1})_2(\text{BF}_4)_2$ catalyst formation.

During the discovery of the aforementioned iron-catalyzed intramolecular olefin aminofluorination, we identified that electrophilic XtalFluor-E¹⁴ effectively suppresses the competing olefin aminohydroxylation, presumably by sequestering the benzoate generated during the N–O bond cleavage (eq 1).^{11,14}

(1)

We also noted, through NMR titration experiments, that XtalFluor-E and $\text{Et}_3\text{N}\cdot 3\text{HF}$ ¹⁵ may form dynamic complexes (Scheme 4),¹⁶ which could attenuate the concentration of $\text{Et}_3\text{N}\cdot 3\text{HF}$ and therefore significantly slow down the putative fluoride-induced catalyst deactivation.

To test this hypothesis, we selected styrene **1** as a model substrate for catalyst discovery (Table 1). We first observed that the $\text{Fe}(\text{NTf}_2)_2\text{-L1}$ complex, a catalyst for styrene amino-oxygenation, led to inefficient aminofluorination when a 1:1 mixture of XtalFluor-E and $\text{Et}_3\text{N}\cdot 3\text{HF}$ was used: vicinal amino fluoride **3** was obtained in a low yield (16%) together with the amino-oxygenation product **4** (entry 1). Notably, there was no reaction in the absence of an iron catalyst; both **1** and **2a** were stable toward XtalFluor-E and $\text{Et}_3\text{N}\cdot 3\text{HF}$, and they were fully recovered (entry 2).

We also observed a higher yield of **3** when the XtalFluor-E/ $\text{Et}_3\text{N}\cdot 3\text{HF}$ ratio was increased to 1.6:1 (entry 3, XtalFluor-E (2.5 equiv), $\text{Et}_3\text{N}\cdot 3\text{HF}$ (1.5 equiv), 42% yield). Suspecting that an extra equivalent of XtalFluor-E may sequester the carboxylate generated during the N–O bond cleavage, we replaced **2a** with **2b**, from which we expected a more nucleophilic benzoate to be generated. A similar yield of **3** and a better mass balance were observed (entry 4). Interestingly, a 2:1 mixture of XtalFluor-E/ $\text{Et}_3\text{N}\cdot 3\text{HF}$ resulted in a decreased yield of **3** and isolation of imidazoline **5**, presumably through MeCN (cosolvent) incorporation

(entry 5). Notably, no desired product was observed in the absence of Et₃N·3HF, and only **5** was isolated (entry 6).

In order to attenuate the neighboring carbamate group's participation, we switched **2b** to more electrophilic **2c** and **2d** and observed further improved yield of **3** (entries 7–8, 46–73% yield). Furthermore, an excess amount of both XtalFluor-E and Et₃N·3HF is beneficial to the yield of **3** (entry 9, 81% yield), and the Fe(NTf₂)₂–**L2** complex provides essentially the same yield (entry 10, 78% yield). To our surprise, Fe(**L1**)–(MeCN)(H₂O)₂(BF₄)₂ (**6**), a crystalline solid that was easily prepared and handled outside of a glovebox (Figure 1), was found to effectively catalyze styrene aminofluorination in the absence of the MeCN cosolvent; **3** was isolated in an excellent yield without using an excess amount of fluorination reagent (entry 11, 88%).¹³

To better understand the structure–reactivity relationship of fluoride donors, we explored a range of variations using other nucleophilic fluorination reagents under the optimized conditions (Table 2). First, replacements of Et₃N·3HF with Et₃N·HF, Et₃N·2HF, TBAF·(tBuOH)₄, KF, or AgF all lead to unproductive decomposition of **2d** (entry 1).¹⁷ Next, replacement of XtalFluor-E with XtalFluor-M also affords **3d**, albeit in a lower yield (entry 2, 65% yield). Furthermore, using benzoyl fluoride instead of Et₃N·3HF/XtalFluor-E does not afford the desired product **3**, but rather leads to benzylation of **2d** (entry 3). Furthermore, using DFI,¹⁸ PhenoFluor,¹⁹ or DAST,²⁰ reagents that can presumably sequester benzoates, leads to unproductive decomposition of **2d** (entries 4–5). Surprisingly, the DAST/XtalFluor-E (1:1) combination can lead to full conversion of **1** and generation of amino fluoride **3d** (entry 6, 51% yield). This result corroborates the proposal that Et₃N·3HF and XtalFluor-E may form dynamic complexes, which are compatible with the iron catalyst (Scheme 4).

Upon discovering the optimal catalytic conditions, we explored the substrate scope of this new reaction (Table 3). First, a range of styrenes with various electronic properties are compatible with this method, and they were converted to internal amino fluorides with excellent yields (entries 1a–1d). Additionally, ene-yne and α-methylstyrene are both excellent substrates (entries 2–3).

Next, we evaluated olefins that are incompatible with existing aminofluorination methods. An allyl silane with a labile C–Si bond in the presence of fluoride ion was converted to an internal amino fluoride, which has a nitrogen, fluorine, and silicon-based triad with excellent yield and regioselectivity (entry 4, 82% yield). A silyl dienol and an ethyl dienoate, both with labile functional groups, were also tolerated by this method to produce amino allylic fluorides in reasonable yields (entries 5–6). Isoprene aminofluorination provided a pair of vicinal amino allylic fluorides with an excellent yield and good regioselectivity (entry 7, 84% yield). Furthermore, a linear aliphatic diene was converted to both 1,2- and 1,4-amino fluorides with excellent combined yield (entry 8, 91% yield).

We subsequently explored indene aminofluorination using the Fe(NTf₂)₂–**L1** complex: an *anti*-2-amino fluoride was isolated with excellent regioselectivity and significant diastereoselectivity (entry 9, 72% yield, dr: 6.2:1).²³ Moreover, we evaluated isolated

olefins, including 1,1-disubstituted and monosubstituted olefins, both of which are significantly less reactive. With increased catalyst loading and the amount of fluorination reagents, both of them were converted to internal amino fluorides with reasonable yields and high regioselectivity (entries 10–11, 62–71% yield). *Notably, this new method enables the synthesis of a range of structurally diverse vicinal amino fluorides for the first time (described in entries 4, 5, 7, 10, and 11).*

The furnished vicinal fluoro carbamates, both from styrenyl and aliphatic olefins, such as **7** and **9**, can be easily derivatized to *N*-Boc protected vicinal fluoro amines **8** and **10** in excellent yields under mild conditions (Scheme 5).¹³

With the success of this aminofluorination method for simple unfunctionalized olefins, we further explored this new reaction with complex olefins, including myrcene which contains both a trisubstituted olefin and a conjugated diene, and (–)-β-pinene (Scheme 6). The aminofluorination of myrcene readily affords both a standard 1,2-addition product **11b** and a 1,6-addition product **11a** presumably through cyclization. We also note that the nitrogen atom was selectively transferred to the same terminal positions in both **11a** and **11b**. To our surprise, the aminofluorination of (–)-β-pinene does not afford any 1,2-addition product, but rather the same 1,6-addition product **11a** presumably through rearrangement. Notably, complete stereochemistry retention is observed in this aminofluorination–rearrangement cascade.²⁴

Fluorinated amphetamines are of great interest to biological studies of the central nervous system; therefore, we evaluated the aminofluorination of *trans*-β-methylstyrene (Scheme 7).²⁵ Interestingly, although the Fe(BF₄)₂–**L1** complex **6** promotes undesirable C–H abstraction, a less sterically hindered Fe(NTf₂)₂–**L2** complex catalyzes efficient aminofluorination to afford **12** (Scheme 7, 72% yield, dr: 2.0:1).¹³

In order to demonstrate the potential practicality of this reaction, we explored the aminofluorination of indene and isoprene on a large scale (Scheme 8). To our pleasure, both catalytic reactions can be scaled up to gram scale with consistent yield and regio- and stereoselectivity.¹³

The catalytic enantioselective aminofluorination of unfunctionalized olefins has been an unsolved synthetic challenge;²⁶ therefore, we explored the feasibility of achieving asymmetric induction using chiral iron catalysts. As a first step, we studied asymmetric indene aminofluorination as a proof of concept, which may provide valuable mechanistic insights (Table 4). The Fe(NTf₂)₂–chiral ligand **L3** complex catalyzes indene aminofluorination with **2d** and affords *anti*-2-amino fluoride **7d** with significant ee (entry 1, 60% ee). Although less sterically hindered acyloxyl carbamates (**2b** and **2c**) lead to further increased ee for indene aminofluorination, both reactions evidently suffer from lower reactivity (entries 2–3, 78–83% ee). In order to achieve enhanced reactivity, the benzoyl activating group in **2c** was switched to 3,5-bis(trifluoromethyl)-benzoyl group in **2e**. To our surprise, a dramatically decreased ee was observed (entry 4, 55% ee). Fortunately, indene aminofluorination with acyloxyl carbamate **2f**, which has a 2,4-dichloro-benzoyl activating group, provides **7** with a synthetically useful yield and high ee (entry 5, 59% yield, 84% ee).

Other chiral ligands, including tridentate **L4** and **L5**, were also evaluated, and they all provided inferior results.

These results also provide valuable mechanistic insight. The acyloxyl moiety on **2** proves crucial for stereochemical communication in the iron-catalyzed indene aminofluorination (Table 4, entries 3–5). This observation suggests that the benzoate generated during N–O bond cleavage may be involved in the ee-determining step. Additionally, since the same ee was observed for both diastereomers of **7**, it is likely that the C–N bond forming step is ee-determining.

In order to formulate a mechanistic model, we carried out several control experiments to gain more insight into both the C–N and C–F bond forming steps (Scheme 9). First, two analogues of **2** in which the *N*–H moiety was replaced with either the *N*–Me or *N*–Ac groups were subjected to standard reaction conditions (Scheme 9, eq 2). Neither was reactive toward the aminofluorination, which suggests that the N–H moiety in **2** is critical for its activation.

Next, the aminofluorination of a cyclopropyl-substituted olefin **15** afforded only the ring-opening products **16** and **17** (Scheme 9, eq 3). This result, together with the observed 1,6-mycene aminofluorination, suggests that the reaction may proceed through a stepwise process and that a carbo-radical species may be an intermediate. To obtain more mechanistic details beyond the C–N bond forming step, we further evaluated a 1,1-disubstituted olefin **18** and observed both 1,2-amino fluoride **19** and 1,3-aminofluoride **20** which is presumably obtained through 1,2-hydride shift from a carbocation (Scheme 9, eq 4). This result suggests that a carbocation may also be involved in the reaction pathway.

Furthermore, benzoylated morpholine **21** was identified as a side product in styrene aminofluorination using Et₃N·3HF/ XtalFluor-M (Scheme 9, eq 5). We suspected that **21** might be derived from the interaction between XtalFluor-M and the benzoic acid that was generated during the iron-catalyzed N–O bond cleavage; therefore, a control experiment between benzoic acid and XtalFluor-M was conducted, from which **21** was indeed isolated (Scheme 9, eq 6). These results suggest that XtalFluor-E or XtalFluor-M sequesters benzoic acids to promote efficient fluorine atom transfer.

On the basis of a range of mechanistic evidence presented in Scheme 9, a mechanistic working hypothesis for the iron-catalyzed olefin aminofluorination is proposed in Scheme 10. First, the iron catalyst may reductively cleave the N–O bond in **2**, which can possibly be converted to an iron-nitrenoid **A**. The iron-nitrenoid may initiate a radical addition with olefin **1** to afford a carbo-radical species **B**. Subsequently, the high-valent iron intermediate may further oxidize the radical, presumably through single electron transfer (SET),²⁷ to generate a carbocation **C1**. When the concentration of a suitable fluoride donor is low, **C1** may be efficiently intercepted by the neighboring carbamate group to afford amino-oxygenation product **4**.

However, in the presence of a suitable nucleophilic fluoride and benzoate sequestering reagent (e.g., Et₃N·3HF/XtalFluor-E), carbocation **C1** may be rapidly converted to **C2**,

which can be captured and converted to amino fluoride **3**; therefore, the aminofluorination pathway will override the competing amino-oxygenation pathway. Since oxazoline **4** may also be converted to carbocation **C1** under reaction conditions which are strongly Lewis acidic, it is possible that this additional pathway may also operate, for a certain type of substrates, to afford the olefin aminofluorination product **3**.

CONCLUSION

In conclusion, we reported a catalytic method for intermolecular olefin aminofluorination using earth-abundant iron catalysts and nucleophilic fluoride ion. This method tolerates a broad range of unfunctionalized olefins, including nonstyrenyl olefins that are incompatible with existing aminofluorination methods. It also readily affords a variety of synthetically valuable vicinal fluoro carbamates with high regioselectivity. Notably, many of them are difficult to prepare using existing methods. We systematically described the catalytic discovery, structure–reactivity relationship studies of fluoride donors, substrate scope and limitation of this method, derivatization of vicinal fluoro carbamates under mild conditions, practical procedures for gram-scale olefin aminofluorination, chiral catalyst discovery for asymmetric olefin aminofluorination, and preliminary mechanistic studies. These studies demonstrate that it is feasible for asymmetric induction using chiral iron catalysts for aminofluorination of unfunctionalized olefins, in which both an iron nitrenoid and a carbocation species may be possible reactive intermediates. Our current efforts focus on both mechanistic understanding and synthetic applications of this new reaction.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institutes of Health (GM110382) and Georgia State University. H.X. is an Alfred P. Sloan Research Fellow. We also thank a referee for insightful comments and suggestions.

References

1. For a selected reference of fluorine in medicinal chemistry, see: Müller K, Faeh C, Diederich F. *Science*. 2007; 317:1881. [PubMed: 17901324]
2. For iron-mediated intermolecular olefin hydrofluorination, see: Barker TJ, Boger DL. *J Am Chem Soc*. 2012; 134:13588. [PubMed: 22860624] For recent references of catalytic olefin difluorination, see: Banik SM, Medley JW, Jacobsen EN. *J Am Chem Soc*. 2016; 138:5000. [PubMed: 27046019] Banik SM, Medley JW, Jacobsen EN. *Science*. 2016; 353:51. [PubMed: 27365443] Molnár IG, Gilmour R. *J Am Chem Soc*. 2016; 138:5004. [PubMed: 26978593] For selected references of styrene hydrofluorination, see: Emer E, Pfeifer L, Brown JM, Gouverneur V. *Angew Chem, Int Ed*. 2014; 53:4181. Wilger DJ, Grandjean JMM, Lammert TR, Nicewicz DA. *Nat Chem*. 2014; 6:720. [PubMed: 25054943] For olefin azido-fluorination, see: Li Z, Zhang C, Zhu L, Liu C, Li C. *Org Chem Front*. 2014; 1:100. For styrene fluorosulfonylation, see: Yuan Z, Wang HY, Mu X, Chen P, Guo YL, Liu G. *J Am Chem Soc*. 2015; 137:2468. [PubMed: 25649748]
3. For a selected reference of vicinal fluoro amines in medicinal chemistry, see: Hagmann WK. *J Med Chem*. 2008; 51:4359. [PubMed: 18570365]
4. Stavber S, Pec n TS, Papež M, Zupan M. *Chem Commun*. 1996:2247. In this report, the substrate scope was predominantly limited to styrenyl olefins with two exceptions. First, tetramethylethylene.

Second, 1-octene was converted to an inseparable mixture of 1,2- and 1,3-terminal fluorides. For recent references related to intermolecular styrene aminofluorination, see: Qiu S, Xu T, Zhou J, Guo Y, Liu G. *J Am Chem Soc.* 2010; 132:2856. [PubMed: 20148557] Zhang H, Song Y, Zhao J, Zhang J, Zhang Q. *Angew Chem, Int Ed.* 2014; 53:11079. Saavedra-Olavarria J, Arteaga GC, López JJ, Pérez EG. *Chem Commun.* 2015; 51:3379. Chen P, Liu G. *Eur J Org Chem.* 2015; 2015:4295. For a recent review of intramolecular olefin aminofluorination, see: Serguchev YA, Ponomarenko MV, Ignat'ev NV. *J Fluorine Chem.* 2016; 185:1.

5. For a reference of asymmetric fluoro aminooxylation of enals using chiral diphenyl prolinol-type of organocatalysts, see: Appayee C, Brenner-Moyer SE. *Org Lett.* 2010; 12:3356. [PubMed: 20575576]
6. The aminofluorination of terminal olefins using nucleophilic fluorination reagents affords internal amino fluorides with the inverse regioselectivity obtained using electrophilic fluorination reagents.
7. For iron-mediated olefin hydrofluorination with electrophilic Selectfluor, see ref 2a. For iron-catalyzed benzylic C–H fluorination with Selectfluor, see: Bloom S, Pitts CR, Woltornist R, Griswold A, Holl MG, Lectka T. *Org Lett.* 2013; 15:1722. [PubMed: 23527764] For other selected examples of radical fluorination that are not promoted by iron catalysts, see: Rueda-Becerril M, Chatalova Sazepin C, Leung JCT, Okbinoglu T, Kennepohl P, Paquin J-F, Sammis GM. *J Am Chem Soc.* 2012; 134:4026. [PubMed: 22320293] Liu W, Huang X, Cheng MJ, Nielsen RJ, Goddard WA, Groves JT. *Science.* 2012; 337:1322. [PubMed: 22984066] Xia JB, Zhu C, Chen C. *J Am Chem Soc.* 2013; 135:17494. [PubMed: 24180320] Zhang C, Li Z, Zhu L, Yu L, Wang Z, Li C. *J Am Chem Soc.* 2013; 135:14082. [PubMed: 24025164] Ventre S, Petronijevic FR, MacMillan DWC. *J Am Chem Soc.* 2015; 137:5654. [PubMed: 25881929]
8. BDE for Fe–F is 447 kJ/mol, while BDEs for Fe–Cl and Fe–Br are 336 and 243 kJ/mol, respectively. See: Luo Y-R. *Comprehensive Handbook of Chemical Bond Energies.* CRC Press Boca Raton, FL 2007:803.
9. The nonsolvated fluoride ion is exceptionally nucleophilic; for a reference, see: Sun H, DiMaggio SG. *J Am Chem Soc.* 2005; 127:2050. [PubMed: 15713075]
10. For existing vicinal fluoro acetamide and sulfonamide synthesis from olefins, see ref 4. For selected examples of fluoro amine synthesis from other precursors, see: Duthion BR, Pardo DG, Cossy J. *Org Lett.* 2010; 12:4620. [PubMed: 20849096] Schulte ML, Lindsley CW. *Org Lett.* 2011; 13:5684. [PubMed: 21942742] Wade TN. *J Org Chem.* 1980; 45:5328. Kalow JA, Schmitt DE, Doyle AG. *J Org Chem.* 2012; 77:4177. [PubMed: 22489747] Sladojevich F, Arlow SI, Tang P, Ritter T. *J Am Chem Soc.* 2013; 135:2470. [PubMed: 23397884] Davies SG, Fletcher AM, Frost AB, Roberts PM, Thomson JE. *Org Lett.* 2015; 17:2254. [PubMed: 25871478] Zhang Q, Stockdale DP, Mixdorf JC, Topczewski JJ, Nguyen HM. *J Am Chem Soc.* 2015; 137:11912. [PubMed: 26348569]
11. Lu DF, Liu GS, Zhu CL, Yuan B, Xu H. *Org Lett.* 2014; 16:2912. [PubMed: 24829034] For related iron-catalyzed enantioselective olefin aminochlorination and diastereoselective aminobromination, see: Zhu CL, Tian JS, Gu ZY, Xing GW, Xu H. *Chem Sci.* 2015; 6:3044. [PubMed: 26807211] Tian JS, Zhu CL, Chen YR, Xu H. *Synthesis.* 2015; 47:1709. [PubMed: 26806984]
12. Lu DF, Zhu CL, Jia ZX, Xu H. *J Am Chem Soc.* 2014; 136:13186. [PubMed: 25166591] For a related iron-catalyzed intramolecular olefin aminohydroxylation, see: Liu GS, Zhang YQ, Yuan YA, Xu H. *J Am Chem Soc.* 2013; 135:3343. [PubMed: 23402638]
13. For experimental details, see the Supporting Information.
14. For synthesis of XtalFluor-E, see: L'Heureux A, Beaulieu F, Bennett C, Bill DR, Clayton S, LaFlamme F, Mirmehrabi M, Tadayon S, Tovell D, Couturier M. *J Org Chem.* 2010; 75:3401. [PubMed: 20405933] XtalFluor-E is hygroscopic; see Supporting Information for correct handling and storage. XtalFluor-E has no effect on the catalytic activity of the iron–bi-dentate ligand complexes in the intramolecular olefin aminofluorination. Full conversion of the substrate was observed in the absence of XtalFluor-E (eq 1).
15. For a reference describing the structure of Et₃N·3HF, see: Wiechert D, Mootz D, Franz R, Siegemund G. *Chem - Eur J.* 1998; 4:1043.
16. XtalFluor-E is prepared using DAST and BF₃·Et₂O. For experimental details, see the Supporting Information. For reactivity of these complexes, see entry 11 of Table 1 and entry 6 of Table 2.

17. Acyloxyl carbamate **2d** decomposes under weakly basic conditions. We suspect that the reaction medium may be slightly basic due to solvation of fluoride ion with deleterious moisture in these reagents.
18. Hayashi H, Sonoda H, Fukumura K, Nagata T. Chem Commun. 2002:1618.
19. Tang P, Wang W, Ritter T. J Am Chem Soc. 2011; 133:11482. [PubMed: 21736304]
20. Middleton WJ. J Org Chem. 1975; 40:574.
21. An excess amount of XtalFluor-E/Et₃N·3HF was applied to suppress the rapid competing olefin aminohydroxylation.
22. An additional amount of XtalFluor-E/Et₃N·3HF was used to convert oxazolines to corresponding amino fluorides.
23. For a diastereoselective indene aminofluorination to afford 1-amino fluoride, which is the regioisomer of **7**, see ref 4e.
24. The ee for (–)-β-pinene is 92%, and the ee for **11a** is 91%. For experimental details, see the Supporting Information.
25. For a selected example of fluorinated amphetamine synthesis, see: Cresswell AJ, Davies SG, Lee JA, Roberts PM, Russell AJ, Thomson JE, Tyte MJ. Org Lett. 2010; 12:2936. [PubMed: 20509635]
26. For iron-catalyzed asymmetric intramolecular olefin amino-fluorination, see ref 11. For a recent review of asymmetric fluorocyclization, see: Wolstenhulme JR, Gouverneur V. Acc Chem Res. 2014; 47:3560. [PubMed: 25379791] For other selected examples of asymmetric fluorination, see: Beeson TD, MacMillan DWC. J Am Chem Soc. 2005; 127:8826. [PubMed: 15954790] Kalow JA, Doyle AG. J Am Chem Soc. 2010; 132:3268. [PubMed: 20163118] Rauniyar V, Lackner AD, Hamilton GL, Toste FD. Science. 2011; 334:1681. [PubMed: 22194571] For asymmetric olefin difluorination, see ref 2b–d. For asymmetric fluoro aminooxylation of enals using organocatalysis, see ref 5. For a recent review of asymmetric olefin aminohalogenation, see: Chemler SR, Bovino MT. ACS Catal. 2013; 3:1076. [PubMed: 23828735]
27. (a) Kharasch MS, Sosnovsky G. J Am Chem Soc. 1958; 80:756.(b) Kochi JK. Science. 1967; 155:415. [PubMed: 17737550]

Figure 1.
Crystal structure of $\text{Fe}(\text{L1})(\text{MeCN})(\text{H}_2\text{O})_2(\text{BF}_4)_2$ (**6**).

Scheme 1.

Iron-Catalyzed Intermolecular Olefin Aminofluorination Using Fluoride Ion

A) previous work: intramolecular olefin aminofluorination

B) early attempts for intermolecular olefin aminofluorination

Scheme 2.

Iron-Catalyzed Intramolecular Olefin Aminofluorination and Early Attempts for Intermolecular Olefin Aminofluorination

Scheme 3.

Experiments To Explore the Putative Fluoride-Induced Catalyst Deactivation

Scheme 4.

Proposed Complex Formation from $\text{Et}_3\text{N} \cdot 3\text{HF}$ and XtalFluor-E

Scheme 5.

Derivatization of Olefin Aminofluorination Products to *N*-Boc-Protected Vicinal Fluoro Amines

Scheme 6.
Iron-Catalyzed Aminofluorination of Myrcene and (-)- β -Pinene

Scheme 7. Iron-Catalyzed Aminofluorination of *trans*- β -Methyl-styrene

^aReaction condition: Fe(NTf₂)₂ (15 mol %), L2 (15 mol %), XtalFluor-E (2.5 equiv)/Et₃N·3HF (1.5 equiv), 4 Å MS, CH₂Cl₂/MeCN, -15 °C, 1 h; then, another portion of XtalFluor-E (1.5 equiv)/Et₃N·3HF (1.5 equiv), 0 °C, 3 h.

Scheme 8.

Iron-Catalyzed Gram-Scale Aminofluorination of Indene and Isoprene

Scheme 9. Control Experiments To Probe for Mechanistic Insight

^a**6** (15 mol %), XtalFluor-E (2.5 equiv)/Et₃N·3HF (1.5 equiv), -15 °C, CH₂Cl₂. ^b**6** (15 mol %), XtalFluor-E (4.0 equiv)/Et₃N·3HF (2.4 equiv), -40 °C, CH₂Cl₂. ^c**6** (15 mol %), XtalFluor-E (3.0 equiv)/Et₃N·3HF (1.8 equiv), -35 to 0 °C, CH₂Cl₂. R^1 : 2,4-Cl₂-benzoyl, R^2 : CF₃CH₂, R^3 : (CF₃)₂CH.

Scheme 10.
 Mechanistic Working Hypothesis of the Iron-Catalyzed Intermolecular Olefin
 Aminofluorination Using Fluoride Ion

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

The yield of **5** is 8–12%.

No desired product observed, and **5** (42%) was isolated.

Preformed complex **6** was used as the catalyst in CH₂Cl₂ in the absence of MeCN; –15 °C, 2 h; the yield of **5** is less than 5%.

Table 2

Structure–Reactivity Relationship Studies of Nucleophilic Fluorination Reagents

entry ^a	variation from the std conditions	observations
1	replace Et ₃ N·3HF with Et ₃ N·HF, Et ₃ N·2HF, TBAF(^t BuOH) ₄ , or AgF	decomposition of 2d , no desired product
2	replace XtalFluor-E with XtalFluor-M	3d (65%) with full conversion
3	replace Et ₃ N·3HF/XtalFluor-E with benzoyl fluoride	3d (<5%) with 66% conversion ^b
4	replace Et ₃ N·3HF/XtalFluor-E with DFI or PhenoFluor	decomposition of 2d , no desired product
5	replace Et ₃ N·3HF/XtalFluor-E with DAST	decomposition of 2d , no desired product
6	replace Et ₃ N·3HF/XtalFluor-E (1.5 equiv/2.5 equiv) with DAST/XtalFluor-E (1.5 equiv/ 1.0 equiv)	3d (51%) with full conversion

^aReactions were carried out under N₂ at -15 °C in 2 h, and 4 Å activated molecular sieves (powder) were used to remove deleterious moisture, unless stated otherwise.

^bAmino-oxygenation product **4d** (42%) and benzoylated **2d** (24%) were isolated.

Table 3

Substrate Scope for the Iron-Catalyzed Olefin Aminofluorination

$ \begin{array}{c} \text{R}^1 \text{---} \text{C} = \text{C} \text{---} \text{R}^3 \\ \\ \text{R}^2 \end{array} + 2\text{d} \xrightarrow[\text{CH}_2\text{Cl}_2, -30 \text{ to } -15^\circ\text{C}^a]{\begin{array}{c} \text{iron(II) catalyst} \\ (10 \text{ mol } \%) \\ \text{Et}_3\text{N} \cdot 3\text{HF} (1.5 \text{ equiv}) \\ \text{XtalFluor-E} (2.5 \text{ equiv}) \end{array}} \begin{array}{c} \text{F} \text{---} \text{C} \text{---} \text{C} \text{---} \text{R}^3 \\ \quad \\ \text{R}^2 \quad \text{NHCO}_2\text{R}^4 \end{array} $ <p>R¹, R², R³: alkyl and aryl groups; R⁴: (CF₃)₂CH</p>				
entry ^a	olefins	reaction time	vincinal fluoro carbamates	yield
1				
		2h	a: R=H	88%
		4h	b: R=Me ^b	82%
		3h	c: R=Br	87%
		4h	d: R=CO ₂ Me ^{b,c}	81%
2		4h		75%
3 ^b		3h		83%
4		5h		82%
5 ^{c,d}		3h		65%
6 ^{e,f}		4h		65%

entry ^a	olefins	reaction time	vicinal fluoro carbamates	yield
7 ^d		6h	 + (6.7 : 1)	84%
8 ^b		6h	 + (2.7 : 1)	91%
9 ^{c,g}		6h	 dr. 6.2 : 1	72%
10 ^{b,c}		3h		71%
11 ^{f,h}		2h		62%

^aReactions were carried out under N₂ with preformed iron(II)-**L1** catalysts (10 mol %), unless stated otherwise.

^bXtalFluor-E (3.0 equiv)/Et₃N·3HF (1.8 equiv).²¹

^cCatalyst loading: 15 mol %.

^dXtalFluor-E (4.0 equiv)/Et₃N·3HF (2.4 equiv).²¹

^e Catalyst loading: 20 mol %.

^f Additional XtalFluor-E (2.5 equiv)/Et₃N·3HF (2.5 equiv) were added after 1 h.²²

^g XtalFluor-E (5.0 equiv)/Et₃N·3HF (3.0 equiv).²¹

^h Catalyst loading: 30 mol %, XtalFluor-E (1.5 equiv)/Et₃N·3HF (0.9 equiv).

Table 4

Catalytic Asymmetric Indene Aminofluorination

entry ^a	2	ligand	yield ^b	dr ^c	ee (major/minor) ^d
1	2d	L3	48%	5.2:1	60%/60%
2	2b	L3	38%	2.5:1	78%/77%
3	2c	L3	35%	2.4:1	83%/82%
4	2e	L3	55%	2.2:1	55%/55%
5	2f	L3	59%	3.0:1	84%/84%
6	2f	L4	32%	1.8:1	36%/37%
7	2f	L5	39%	1.9:1	32%/30%

^a Unless stated otherwise, the reactions were carried out under a nitrogen atmosphere with 4 Å molecular sieves.

^b Isolated yield.

^c dr was determined by ¹H NMR.

^d Enantiomeric excess (ee) was measured by HPLC with chiral columns; the absolute stereochemistry was determined by X-ray crystallographic analysis of **7d** (R: (CF₃)₂CH).