Published in final edited form as:

Health Aff (Millwood). 2016 August 1; 35(8): 1382–1390. doi:10.1377/hlthaff.2015.1635.

Physician Participation In ACOs Is Lower In Places With Vulnerable Populations Compared To More Affluent Communities

Laura C. Yasaitis, PhD,

Leonard Davis Institute of Health Economics, University of Pennsylvania, Center for Health Equity Research and Promotion, Philadelphia VA Medical

William Pajeroski, BA,

Leonard Davis Institute of Health Economics, University of Pennsylvania, Center for Health Equity Research and Promotion, Philadelphia VA Medical

Daniel Polsky, PhD, and

Perelman School of Medicine, University of Pennsylvania; Leonard Davis Institute of Health Economics, University of Pennsylvania, Center for Health Equity Research and Promotion, Philadelphia VA Medical

Rachel M. Werner, MD, PhD

Perelman School of Medicine, University of Pennsylvania; Leonard Davis Institute of Health Economics, University of Pennsylvania, Center for Health Equity Research and Promotion, Philadelphia VA Medical

Abstract

Early evidence suggested that accountable care organizations (ACOs) could improve health care quality while constraining costs, and ACOs are expanding throughout the United States. However, if disadvantaged patients have unequal access to physicians who participate in ACOs, that expansion may exacerbate health care disparities. We examined the relationship between physician participation in both Medicare and commercial ACOs across the country and the sociodemographic characteristics of their likely patient populations. Physician participation in ACOs varied widely across hospital referral regions, from nearly 0 percent to over 85 percent. After we adjusted for individual physician and practice characteristics, we found that physicians who practiced in ZIP Code Tabulation Areas where a higher percentage of the population was black, living in poverty, uninsured, or disabled or had less than a high school education—compared to other areas—had significantly lower rates of ACO participation than other physicians. Our findings suggest that vulnerable populations may not have as great access as other groups to physicians participating in ACOs, which could exacerbate existing disparities in health care quality.

Keywords

Health Reform; Disparities; Physicians

An accountable care organization (ACO) is a network of doctors and hospitals whose members share financial and medical responsibility for providing coordinated care to patients, with the goals of improving care quality and limiting unnecessary spending. ACOs are becoming increasingly common in the United States: At least 744 of them have been formed since 2011. Between Medicare and the commercial (or private) sector, an estimated 23.5 million Americans are now being served by an ACO.[1] Early evidence from both Medicare and commercial ACO initiatives shows that ACOs have succeeded in improving the quality of care and patient experience and, in some cases, constraining costs.[2-7]

While enthusiasm for ACOs is growing, some stakeholders warn of possible unintended consequences. In particular, there are concerns that ACOs may worsen existing disparities in health care quality.[8-10] Historically, the care of vulnerable patients has been concentrated among relatively few providers, who tend to have fewer financial and health care resources and perform worse on most traditional quality metrics, compared to [please provide].[8-14] If ACO participation is linked to improved care quality, and vulnerable patients have less access than other patients to physicians participating in ACOs, existing disparities may increase. Indeed, early evidence suggests that ACOs tend to be located in areas of the country with lower poverty rates than [please provide],[15] and that patients covered under early Medicare ACO contracts had higher incomes and were less likely to be black, covered under Medicaid, or disabled, compared to [please provide].[16]

These early findings are only part of the story. Previous work has focused on the earliest ACOs (those established by 2012) and, in some cases, only Medicare ACOs.[15,16] Early participants in ACOs and Medicare ACOs may vary systematically from later participants and those in commercial ACOs, and thus it is unknown how generalizable these early findings are. More important, it is unknown whether physicians serving more vulnerable patients are less likely to participate in an ACO within an ACO region. If this were the case, it might suggest that such physicians are being systematically excluded from ACOs.

Evidence is needed to better understand how differences in physician ACO participation may affect health care disparities. We used a national data set of US physicians to examine whether a physician's ACO participation is related to sociodemographic factors of local populations both nationally and within small geographic areas.

Study Data And Methods

Data

The unit of analysis in our study was the physician. We used a telephone-verified database of office-based US physicians that included information about their participation in an ACO (public or commercial). The database is continuously updated by SK&A, a private marketing company.

These data have been shown to be a nearly complete sample frame of all office-based US physicians.[17] However, we investigated their completeness by using National Provider Identifiers and last names to match them to data in the Physician and Other Supplier Public Use File of the Centers for Medicare and Medicaid Services (CMS).

We found that 81.6 percent of physicians in the Medicare file were also in the SK&A data. Compared to physicians in both data sets, those who were only in the Medicare file treated fewer unique beneficiaries annually (359 versus 500), received less in annual Medicare payments (\$57,467.39 versus \$137,895.40), and submitted fewer claims annually (1,074 versus 3,631). Physicians who were only in the Medicare file were evenly distributed geographically; in no state did the SK&A data match less than 70 percent of Medicare physicians. Of the physicians in the SK&A data, 83.4 percent were also in the Medicare file. Previous comparisons of SK&A data with those in the American Medical Association's Physician Masterfile and in the American Community Survey found similar totals of physicians and selection across broad specialties.[18]

We identified each physician's ZIP Code Tabulation Area and used summary statistics for that area from the American Community Survey for the period 2009-13 as a proxy for the sociodemographic characteristics of the population that the physician was likely to care for. These areas overlap substantially with ZIP codes and are identical in the case of many residents. For physicians with multiple practice locations, we used the location with the highest patient volume reported by SK&A.

Variables

Our main dependent variable in all analyses was whether or not a physician participated in an ACO. Our main independent variables were the following sociodemographic characteristics at the level of the ZIP Code Tabulation Area: the percentages of the population that had less than a high school education and that were black, Hispanic, living in poverty, uninsured, or disabled.

We included the following physician-level variables: sex, number of physicians in the practice, whether or not the practice was multispecialty, rurality of the practice location, and medical specialty (internal medicine, primary care, surgery, or other).[19] We calculated the number of physicians in the practice by using the largest organizational level present in the data, whether that was integrated health system, physician group, or physician office.[17] We defined rurality of the practice location using Rural-Urban Commuting Area codes, classified into four categories.[20] We summarized the characteristics of the physicians overall and by whether or not they participated in an ACO.

We estimated physician ACO participation at the level of the hospital referral region (HRR) because these 306 regions reflect markets for tertiary hospital care and are representative of all health care markets.[21] In addition, preliminary analyses at the level of the ZIP Code Tabulation Area revealed that variations in ACO penetration largely followed HRR boundaries. Each HRR is a collection of unique ZIP Code Tabulation Areas. Each physician was assigned to an HRR based on the area in which he or she practiced. We then determined the percentages of physicians within each HRR who participated in an ACO.

We included HRR-level health system variables to describe health resources in HRRs with varying physician ACO participation rates. These variables included the supplies of primary care and medical specialist physicians and of acute care hospital beds; Medicare beneficiary primary care visit rates; and the percentage of Medicare beneficiaries with diabetes who received a hemoglobin A1c test.[22]

We sorted HRRs into quartiles based on physicians' ACO participation and weighted by population, so that each quartile represented roughly a quarter of the US population. We then described the population and health system characteristics across quartiles, which we used as independent variables in regression analyses.

Regression Analyses

We performed multivariate linear regressions to examine the association between population-level sociodemographic characteristics of physicians' practice locations and rates of physician ACO participation, after adjusting for physician and practice characteristics. Our dependent variable was ACO participation reported by physicians to [please provide]. We used sociodemographic characteristics of physician ZIP Code Tabulation Areas as our main independent variables, to test for the association of physician ACO participation with those characteristics. To represent these variables, we grouped the areas into quartiles for each sociodemographic characteristic. Each independent sociodemographic variable was an indicator of the quartile of the physician's ZIP Code Tabulation Area for that sociodemographic measure. The mean and distribution of the area characteristics within each quartile are available in the online Appendix.[23]

Each set of sociodemographic indicators was examined in a separate regression. In all regressions, we included the physician- and practice-level covariates described above to address potential confounding of the relationship between local population characteristics and ACO participation.

Each regression was run first without and then with HRR fixed effects (that is, a dummy variable for each HRR). Regressions without HRR fixed effects estimated the probability of physician participation in ACOs nationally. Adding HRR fixed effects controlled for regional characteristics, such as HRR-level income or population demographic characteristics, and regressions with those fixed effects estimated the probability of physician participation conditional on those characteristics. Thus, if ACOs are more likely to be in high-income HRRs than in low-income ones, but all physicians participated equally within each HRR regardless of within-HRR variations in income, we would expect to find that ZIP Code Tabulation Area sociodemographic characteristics had a significant effect on ACO participation in a regression without HRR fixed effects, but no effect in a regression with the effects. We also performed the regressions after limiting the sample to primary care physicians, as this group of physicians is most central to the ACO model.[24]

To summarize the associations between various population characteristics in a physician's practice location and his or her likelihood of participation, we used multivariate models to predict the percentage of physicians participating in an ACO in each quartile of ZIP Code Tabulation Area population sociodemographic characteristics.

Limitations

Our study had several limitations. First, the physician data that we used included the vast majority of practicing physicians, but not all of them. However, based on analyses of Medicare physicians who were and were not successfully matched with physicians in our primary data source, we found that the physicians in our study were geographically representative and more clinically active than those for whom we did not have information about ACO participation.

Second, we used general population characteristics of the ZIP Code Tabulation Area where each physician's practice was located as a proxy for the patient population he or she was likely to care for. We were unable to study populations specifically attributed to ACOs or to the physicians in our sample. For some subspecialties, patients travel substantial distances to receive care, and as a result these characteristics are a noisy representation of physicians' likely patient panels. However, sensitivity analyses among the subset of physicians for whom we had Medicare data supported the validity of our measures.

Finally, our physician data were self-reported. Moreover, the data identified only one ACO for each physician and did not specify if the ACO was commercial or sponsored by Medicare or Medicaid.

Study Results

Our study included 521,543 US physicians. Of those physicians, 25.9 percent participated in an ACO (Exhibit 1). ACO participation was more common among female physicians than among males; physicians in large and multispecialty practices, compared to other practices; and primary care physicians, compared to those in other specialties. Rates of ACO participation were highest in the Northeast, mid-Atlantic region, upper Midwest, and West (Exhibit 2).

Compared to HRRs with higher physician participation in ACOs, in HRRs with lower participation, residents were more likely to have less than a high school education and to be living in poverty, black, or disabled (Exhibit 3). The largest percentages of Hispanics were found in HRRs with neither the lowest nor the highest physician ACO participation.

HRRs with higher physician participation also tended to have a greater supply of all types of physicians, but a lower supply of acute care hospital beds, compared to HRRs with lower participation. In addition, among Medicare beneficiaries, visits to primary care physicians and rates of hemoglobin A1c testing were slightly higher in HRRs with greater physician ACO participation. Full results on characteristics of HRRs by ACO participation rate are available in the Appendix.[23]

Nationally, we found that physician ACO participation was inversely related to the percentage of the population that was black, living in poverty, uninsured, or disabled or that had less than a high school education (Exhibit 4). There was no consistent pattern of participation across levels of Hispanic population. The largest differences in participation were across quartiles of the proportion of population that was black, with 30.8 percent of all

physicians (35.7 percent of primary care physicians) participating in an ACO in ZIP Code Tabulation Areas with the lowest quartile of black population, compared to 22.9 percent of all physicians (26.4 percent of primary care physicians) in the highest quartile. For both groups of physicians, the difference between those two quartiles was significant ($p < 0.001$). Full regression results are available in the Appendix.[23]

When we adjusted for HRR fixed effects, we found persistent differences in ACO participation rates, which suggest within-HRR variation across quartiles of population sociodemographic characteristics (Exhibit 5). In this analysis, the percentages of physicians participating in ACOs were still significantly different across quartiles of ZIP Code Tabulation Areas within HRRs, but the magnitude of the differences was smaller than was the case with the national estimates.

Discussion

We found that physician participation in ACOs varied significantly with local sociodemographic characteristics of the population. Physicians practicing in ZIP Code Tabulation Areas with more blacks, higher poverty, more uninsured or disabled residents, or more people with less than a high school education were less likely to participate in ACOs than their counterparts practicing in areas whose populations of such vulnerable groups were smaller.

Physicians serving populations with relatively high rates of disadvantaged people could be excluded from emerging ACOs in two ways. First, ACOs may be more likely to form in regions where the overall patient population is more affluent, compared to regions with less affluent populations. Second, emerging ACOs may be more likely to contract with local physicians who serve more affluent patients than with physicians whose patient populations are less affluent, thus excluding disadvantaged patients.

Our study confirmed that ACOs are more likely to be formed in regions of the country whose populations have higher socioeconomic status and a smaller percentage of blacks, compared to regions whose populations have lower socioeconomic status and a larger percentage of blacks.[15] However, we also found that within the regions where an ACO is located, physicians are more likely to participate if their practice is located in a ZIP Code Tabulation Area with a higher socioeconomic status and smaller percentage of blacks, since limiting our analyses to within-HRR differences diminished these associations but did not eliminate them.

Our results suggest that vulnerable patients have less access to physicians who participate in ACOs and thus less access to the potential benefits of ACOs, compared to other patients. These findings are consistent with early analyses of regions where ACOs began operating and analyses of Medicare patients who were and were not attributed to some of the first public ACO contracts.[15,16] These findings raise concerns that ACOs may increase existing disparities between the quality of health care received by vulnerable populations and the quality of care received by other groups.

As discussed above, studies have shown that ACOs increase the quality of care. However, ACOs also aim to contain costs, and it is possible that ACO participation may encourage providers to stint on necessary care. If that is the case, vulnerable populations would be somewhat sheltered from exposure to such physicians.

While previous studies have examined areas where ACO contracts have been initiated,[15] physician rates of participation may be a more important indicator than the simple presence of an ACO in a region, since patients can be attributed to an ACO only through their physicians. Additionally, given the leadership role that physicians have played in early ACOs,[25,26] it is important to examine possible factors that may prevent some physicians from embracing the ACO model.

There are several possible mechanisms for lower rates of ACO participation among physicians serving vulnerable populations, compared to other physicians. Given that it may be hard to meet some benchmarks for quality of care among hard-to-treat, vulnerable populations,[27-29] ACOs may be less likely to locate in regions with these populations than elsewhere. At the same time, even within areas with ACOs, the organizations may exclude physicians likely to care for vulnerable patients in an effort to ensure a population that will make it possible to achieve high scores on specific quality measures.

Rates of ACO participation may also be lower among physicians serving vulnerable populations, compared to other physicians, because of physician choice. Individual physicians or physician groups may choose to hold off on joining an ACO if they deem it would be too difficult to achieve specific quality and spending goals with their patient populations. Finally, physician leaders of early ACOs reported that limited capital was a major hurdle to entering an ACO contract.[26] Groups serving vulnerable populations may find such hurdles insurmountable.

Our findings suggest several potential policy interventions to prevent the widening of health care disparities as ACOs expand. First, given that previous work[26] has shown limited capital to be a major concern of physician leaders in starting an ACO, providing physician groups that serve vulnerable populations with additional incentives to form an ACO or assistance with start-up costs could ameliorate these disparities. The CMS ACO Investment Model is a new policy intervention to encourage new ACOs to form in rural and underserved areas by addressing their lack of adequate access to capital. Our findings suggest that this new model should help address barriers, but future versions of this program may need to enhance incentives for physician participation in ACOs to attract participation from the physicians in the most underserved parts of these areas. In addition, risk-adjusting ACO-linked quality indicators for patients' sociodemographic characteristics, or perhaps rewarding improvements over time, may encourage providers with vulnerable patient panels to pursue involvement in ACO contracts.[30]

Despite the fact that our study was limited by our inability to assess which patients were directly attributed to ACOs, the results of sensitivity analyses were consistent with our main findings. Specifically, using publicly available Medicare patient-panel data, we found that physicians who participated in ACOs saw significantly fewer black beneficiaries than

physicians who did not participate in ACOs (6.3 percent versus 6.9 percent among all physicians, and 5.8 percent versus 6.7 percent among primary care physicians; $p < 0.01$). Additionally, patient panels of primary care physicians tend to closely approximate the population of the ZIP Code Tabulation Areas in which they practice because patients seek primary care services close to home. This is supported by the fact that our results were slightly stronger when we limited our sample to primary care physicians.

Finally, because of our relatively short study period, we could not test whether rates of ACO participation were increasing over time among physicians who serve vulnerable populations. In particular, ongoing ACO expansion in Medicaid and federally qualified health centers may alleviate some potential disparities.[30] However, in conjunction with previous studies covering periods through 2012, our study—which extended through 2013—found persistent trends in ACO location and physician participation in ACOs with respect to population sociodemographic characteristics. Additionally, even if such expansion occurs, the quality improvements documented for early ACOs—which were likely achieved among relatively affluent patient populations—may differ as ACOs expand to care for more diverse populations.

Conclusion

Overall, our results suggest that current patterns of physician participation in ACOs risk exacerbating disparities in the quality of care received between vulnerable populations and other groups. Greater care is warranted in creating policies to encourage the development of ACOs and physician participation in them in areas with vulnerable populations. Additionally, any changes in the care and health outcomes of these populations should be monitored to assess the potential effects of ACO implementation on health care disparities.

Acknowledgment

This study was supported by grants from the National Institute on Aging (Grant Nos. K24-AG047908 and PO1-AG19783).

Bios for 2015-1635_Yasaitis

Bio 1: Laura C. Yasaitis (yasaitis@mail.med.upenn.edu) is a postdoctoral research fellow at the Perelman School of Medicine at the University of Pennsylvania, in Philadelphia.

Bio 2: William Pajeroski is a doctoral student at the Leonard Davis Institute of Health Economics at the University of Pennsylvania.

Bio 3: Daniel E. Polsky is executive director of and the Robert D. Eilers Professor in Health Care Management and Economics at the Leonard Davis Institute of Health Economics at the University of Pennsylvania.

Bio 4: Rachel M. Werner is an assistant professor of medicine at the University of Pennsylvania.

Notes

1. Muhlestein, D. Continued growth of public and private accountable care organizations. Health Affairs Blog [blog on the Internet]. 2013 Feb 19 [cited 2016 Jun 14]. Available from: <http://healthaffairs.org/blog/2013/02/19/continued-growth-of-public-and-private-accountable-care-organizations/>
2. McWilliams JM, Chernew ME, Landon BE, Schwartz AL. Performance differences in year 1 of Pioneer Accountable Care Organizations. *N Engl J Med*. 2015; 372(20):1927–36. [PubMed: 25875195]
3. McWilliams JM, Landon BE, Chernew ME, Zaslavsky AM. Changes in patients' experiences in Medicare accountable care organizations. *N Engl J Med*. 2014; 371(18):1715–24. [PubMed: 25354105]
4. McWilliams JM, Landon BE, Chernew ME. Changes in health care spending and quality for Medicare beneficiaries associated with a commercial ACO contract. *JAMA*. 2013; 310(8):829–36. [PubMed: 23982369]
5. Song Z, Rose S, Safran DG, Landon BE, Day MP, Chernew ME. Changes in health care spending and quality 4 years into global payment. *N Engl J Med*. 2014; 371(18):1704–14. [PubMed: 25354104]
6. Song Z, Safran DG, Landon BE, Landrum MB, He Y, Mechanic RE, et al. The “Alternative Quality Contract,” based on a global budget, lowered medical spending and improved quality. *Health Aff (Millwood)*. 2012; 31(8):1885–94. [PubMed: 22786651]
7. Nyweide DJ, Lee W, Cuerdon TT, Pham HH, Cox M, Rajkumar R, et al. Association of Pioneer Accountable Care Organizations vs traditional Medicare fee for service with spending, utilization, and patient experience. *JAMA*. 2015; 313(21):2152–61. [PubMed: 25938875]
8. Pollack CE, Armstrong K. Accountable care organizations and health care disparities. *JAMA*. 2011; 305(16):1706–7. [PubMed: 21521853]
9. Anderson RE, Ayanian JZ, Zaslavsky AM, McWilliams JM. Quality of care and racial disparities in Medicare among potential ACOs. *J Gen Intern Med*. 2014; 29(9):1296–304. [PubMed: 24879050]
10. Lewis VA, Larson BK, McClurg AB, Boswell RG, Fisher ES. The promise and peril of accountable care for vulnerable populations: a framework for overcoming obstacles. *Health Aff (Millwood)*. 2012; 31(8):1777–85. [PubMed: 22869656]
11. Bach PB, Pham HH, Schrag D, Tate RC, Hargraves JL. Primary care physicians who treat blacks and whites. *N Engl J Med*. 2004; 351(6):575–84. [PubMed: 15295050]
12. Jha AK, Orav EJ, Li Z, Epstein AM. Concentration and quality of hospitals that care for elderly black patients. *Arch Intern Med*. 2007; 167(11):1177–82. [PubMed: 17563027]
13. Werner RM, Goldman LE, Dudley RA. Comparison of change in quality of care between safety-net and non-safety-net hospitals. *JAMA*. 2008; 299(18):2180–87. [PubMed: 18477785]
14. Joynt KE, Orav EJ, Jha AK. Thirty-day readmission rates for Medicare beneficiaries by race and site of care. *JAMA*. 2011; 305(7):675–81. [PubMed: 21325183]
15. Lewis VA, Colla CH, Carluzzo KL, Kler SE, Fisher ES. Accountable care organizations in the United States: market and demographic factors associated with formation. *Health Serv Res*. 2013; 48(6 Pt 1):1840–58. [PubMed: 24117222]
16. Epstein AM, Jha AK, Orav EJ, Liebman DL, Audet AM, Zezza MA, et al. Analysis of early accountable care organizations defines patient, structural, cost, and quality-of-care characteristics. *Health Aff (Millwood)*. 2014; 33(1):95–102. [PubMed: 24395940]
17. Dunn A, Shapiro AH. Do physicians possess market power? *J Law Econ*. 2014; 57(1):159–93.
18. Gresenz CR, Auerbach DI, Duarte F. Opportunities and challenges in supply-side simulation: physician-based models. *Health Serv Res*. 2013; 48(2 Pt 2):696–712. [PubMed: 23347041]
19. Primary care includes family practice, general internal medicine, geriatrics, and pediatrics. Medical specialties include internal medicine specialties such as cardiology, neurology, endocrinology, and rheumatology. Surgery includes general surgery and subspecialties such as orthopedic and vascular surgery and neurosurgery. Hospital specialties are the remaining physician specialties, including anesthesiology, emergency medicine, radiology, and pathology.

20. Hart, G. Temporary ZIP RUCA 3.10 file ccess page [Internet]. University of North Dakota Center for Rural Health; Grand Forks (ND): 2014 Aug 4 [cited 2016 Jun 14]. Available from: <https://ruralhealth.und.edu/ruca>
21. Dartmouth Institute for Health Policy and Clinical Practice. Dartmouth atlas of health care: data by region [Internet]. Dartmouth Institute; Lebanon (NH): c 2015 [cited 2016 Jun 14]. Available from: <http://www.dartmouthatlas.org/data/region/>
22. Dartmouth Institute for Health Policy and Clinical Practice. Dartmouth atlas of health care: downloads [Internet]. Dartmouth Institute; Lebanon (NH): c 2016 [cited 2016 Jun 14]. Available from: <http://www.dartmouthatlas.org/tools/downloads.aspx>
23. To access the Appendix, click on the Appendix link in the box to the right of the article online.
24. McClellan M, McKethan AN, Lewis JL, Roski J, Fisher ES. A national strategy to put accountable care into practice. *Health Aff (Millwood)*. 2010; 29(5):982–90. [PubMed: 20439895]
25. Shortell SM, McClellan SR, Ramsay PP, Casalino LP, Ryan AM, Copeland KR. Physician practice participation in accountable care organizations: the emergence of the unicorn. *Health Serv Res*. 2014; 49(5):1519–36. [PubMed: 24628449]
26. Colla CH, Lewis VA, Shortell SM, Fisher ES. First national survey of ACOs finds that physicians are playing strong leadership and ownership roles. *Health Aff (Millwood)*. 2014; 33(6):964–71. [PubMed: 24889945]
27. Reschovsky JD, O'Malley AS. Do primary care physicians treating minority patients report problems delivering high-quality care? *Health Aff (Millwood)*. 2008; 27(3):w222–31. [PubMed: 18430747]
28. Hong CS, Atlas SJ, Chang Y, Subramanian S, Ashburner JM, Barry MJ, et al. Relationship between patient panel characteristics and primary care physician clinical performance rankings. *JAMA*. 2010; 304(10):1107–13. [PubMed: 20823437]
29. Zaslavsky AM, Hochheimer JN, Schneider EC, Cleary PD, Seidman JJ, McGlynn EA, et al. Impact of sociodemographic case mix on the HEDIS measures of health plan quality. *Med Care*. 2000; 38(10):981–92. [PubMed: 11021671]
30. Casalino LP, Elster A, Eisenberg A, Lewis E, Montgomery J, Ramos D. Will pay-for-performance and quality reporting affect health care disparities? *Health Aff (Millwood)*. 2007; 26(3):w405–14. [PubMed: 17426053]
31. American Hospital Association. Accountable care organizations: findings from the survey of care systems and payment [Internet]. AHA; Chicago (IL): [cited 2016 Jun 14]. Available from: <http://www.aha.org/content/14/14aug-acocharts.pdf>

Exhibit 2. Percentages of physicians participating in an accountable care organization in [please provide], by hospital referral region

SOURCE Authors' analysis of SK&A physician data for 2013. NOTES Hospital referral regions were divided into roughly equal population-weighted quartiles. Not populated is [please provide]. Quartile 1 (lowest) is 0.4–12.7 percent. Quartile 2 is 12.8–22.1 percent. Quartile 3 is 22.6–31.2 percent. Quartile 4 (highest) is 31.3–87.1 percent.

Values represent the percent of physicians in each HRR that reported participating in any ACO. Ranges are defined by dividing HRRs into roughly equal population-weighted quartiles. The range of percent of physicians participating in an ACO in each quartile is as follows: Very Low: 0.4 – 12.7%; Low: 12.8 – 22.1%; Medium: 22.6 – 31.2%; High: 31.3 – 87.1%.

Source: Authors' calculations from SK&A physician data (2013).

Exhibit 3. Percentages of hospital referral region populations with selected sociodemographic characteristics, by geographic access to physicians participating in an accountable care organization in [please provide]

SOURCE Authors' analysis of SK&A physician data for 2013 and American Community Survey data for 2009–13. NOTES As rates of physician participation in an accountable care organization (ACO) increase, the unadjusted percentages of the hospital referral region (HRR) population with less than a high school education and is living in poverty, uninsured, or disabled decreases. Rates of ACO participation are the proportions of physicians in each HRR who reported participating in any ACO. HRRs were divided into quartiles weighted by the total resident population. Quartile 1 (lowest) is 0.4–12.7 percent. Quartile 2 is 12.8–22.1 percent. Quartile 3 is 22.6–31.2 percent. Quartile 4 (highest) is 31.3–87.1 percent.

As physician ACO participation increases, the proportion of the resident population that is black, has less than a HS education, is living in poverty, uninsured, or disabled decreases. ACO participation was defined at the HRR level by determining the total proportion of physicians in the HRR who report participating in any ACO. HRRs were then divided into quartiles weighted by the total resident population. The range of percent of physicians participating in an ACO in each quartile is as follows: Very Low: 0.4 – 12.7%; Low: 12.8 – 22.1%; Medium: 22.6 – 31.2%; High: 31.3 – 87.1%. The sociodemographic statistics reported represent the unadjusted percent of the HRRs' total resident population defined by each of the sociodemographic indicators.

Source: Authors' calculations from SK&A physician data (2013) and American Community Survey data (2009-2013).

		Hispanic	Less than	Uninsured	Living in P	Black	Disabled
Quartile of physician participation	Very Low	0.13796	0.17614	0.1485	0.15943	0.15257	0.13384
	Low	0.19495	0.17517	0.1646	0.1561	0.14009	0.11943
	Medium	0.22401	0.15814	0.15482	0.14357	0.1375	0.10472
	High	0.10527	0.15856	0.11602	0.13976	0.1148	0.11942

Exhibit 1

Characteristics of office-based US physicians in [please provide]

	Participating in an ACO		Not participating in an ACO		All	
	No.	%	No.	%	No.	%
All physicians	135,026	25.9	386,517	74.1	521,543	100
Sex						
Male	92,116	68.2	283,480	73.3	375,596	72.0
Female	42,910	31.8	103,037	26.7	145,947	28.0
Number of physicians in practice ^a						
Less than 10	20,811	15.4	199,896	51.7	220,707	42.3
10–100	19,590	14.5	89,137	23.1	108,727	20.8
More than 100	94,625	70.1	97,484	25.2	192,109	36.8
Multispecialty practice						
No	88,816	65.8	298,050	77.1	386,866	74.2
Yes	46,210	34.2	88,467	22.9	134,677	25.8
Primary practice rurality ^b						
Urban	127,548	94.5	356,829	92.3	484,377	92.9
Large rural city or town	4,807	3.6	19,740	5.1	24,547	4.7
Small rural town	1,986	1.5	7,750	2.0	9,736	1.9
Isolated small rural town	685	0.5	2,198	0.6	2,883	0.6
Medical specialty ^c						
Internal medicine	35,837	26.5	110,145	28.5	145,982	28.0
Primary care	52,876	39.2	114,877	29.7	167,753	32.2
Surgery	26,627	19.7	93,722	24.3	120,349	23.1
Other	19,686	14.6	67,773	17.5	87,459	16.8

SOURCE Authors' analysis of SK&A physician data for 2013. NOTE ACO is accountable care organization.

^aTotal number of physicians in the largest unique medical practice to which the physician could be assigned.^bBased on the Rural Urban Commuting Area (RUCA) of the ZIP code of the physician's primary practice. RUCA scores were grouped into four categories that captured the predominant commuting patterns of each ZIP code's residents.^cFor details about the specialties, see Note 19 in text.

Exhibit 4

Predicted percentages of all physicians and primary care physicians participating in an accountable care organization in [please provide], by quartiles of demographic characteristics of the ZIP Code Tabulation Area in which they practice

	Quartile			
Percent of population that is or has:	1 (lowest)	2	3	4 (highest)
Black				
All physicians	30.8	26.7 ***	24.7 ***	22.9 ***
Primary care physicians	35.7	32.7 ***	31.3 ***	26.4 ***
Hispanic				
All physicians	24.1	27.8 ****	26.5 ****	24.2 ****
Primary care physicians	29.0	32.5 ****	32.6	31.9 **
Living in poverty				
All physicians	28.9	26.2 ****	25.5 ****	23.2 ****
Primary care physicians	35.5	31.6 ****	30.7 **	28.0 ****
Uninsured				
All physicians	27.4	27.7	24.9 ****	22.1 ****
Primary care physicians	33.3	31.7 ****	30.4 ****	29.8 ****
Less than a high school education				
All physicians	26.4	26.8 **	24.5 ****	25.2 ****
Primary care physicians	31.6	32.9 ****	31.1 ****	29.9 ****
Disabled				
All physicians	28.5	26.7 ****	24.4 ****	23.2 ****
Primary care physicians	35.4	33.3 ****	29.6 ****	27.5 ****

SOURCE Authors' analysis of SK&A physician data for 2013 and American Community Survey data for 2009–13. NOTES Significance refers to the difference between each estimate and that of the next lowest quartile. Estimates were derived from a multiple regression model that predicted physicians' participation in any accountable care organization, adjusted for sex, medical specialty, number of physicians in the practice, and rurality of the practice location. Demographic estimates were derived from American Community Survey data, with ZIP Code Tabulation Areas divided into population-weighted quartiles. The quartiles were treated as a categorical variable; indicators were used to estimate the change in participation associated with each quartile, with the lowest quartile as the reference group.

**
 $p < 0.05$

 $p < 0.01$

 $p < 0.001$

Exhibit 5

Predicted percentages of all physicians and primary care physicians participating in an accountable care organization in [please provide], by quartiles of demographic composition of the ZIP Code Tabulation Area in which they practice, adjusted for hospital referral region fixed effects

	Quartile			
Percent of population that is or has:	1 (lowest)	2	3	4 (highest)
Black				
All physicians	27.5	25.9 ****	25.1 ****	25.5 **
Primary care physicians	33.1	31.8 ****	31.5	29.7 ****
Hispanic				
All physicians	25.6	27.3 ****	25.8 ****	24.6 ****
Primary care physicians	32.0	33.2 ****	31.1 ****	29.1 ****
Living in poverty				
All physicians	27.1	26.2 ****	25.7 ***	24.7 ****
Primary care physicians	34.0	31.7 ****	30.9 ***	29.3 ****
Uninsured				
All physicians	25.1	27.0 ****	26.6 ***	24.7 ****
Primary care physicians	32.1	31.8	31.4	30.3 ***
Less than a high school education				
All physicians	26.1	26.3	24.9 ****	26.0 ****
Primary care physicians	31.7	32.3 **	31.1 ****	30.7
Disabled				
All physicians	27.6	26.0 ****	24.9 ****	24.7
Primary care physicians	33.3	32.7 **	30.4 ****	29.5 ***

SOURCE Authors' analysis of SK&A physician data for 2013 and American Community Survey data for 2009–13. NOTES Adjusting for hospital referral region made it possible to estimate differences in participation across ZIP Code Tabulation Areas with different sociodemographic characteristics located within the same region. Significance refers to the difference between each estimate and that of the next lowest quartile. Estimates were derived from a multiple regression model described in the Notes to Exhibit 4. Demographic estimates were derived from American Community Survey data, with ZIP Code Tabulation Areas divided into population-weighted quartiles. These quartiles were treated as a categorical variable; indicators were used to estimate the change in participation associated with each quartile, with the lowest quartile as the reference group.

**
 $p < 0.05$

 $p < 0.01$

 $p < 0.001$