Editorial

The number seven is a magical figure. The cranium has seven bones, the orbit has seven bones and the nasal framework has seven components. The seven facial bones are hollow containing air spaces inside them and there are seven muscles which control the movements of the eye which offer us perfect binocular vision.

The cranial bones have sutures which are quite different from the rest of the bones. They are held together by fibrous joints which are either simple end to end (buttoned) e.g. median palatine suture or bevelled parieto-temporal or serrated sagittal suture depending upon the stress they are supposed to handle and are universally present at their designated places. Whether they are locally controlled or there is a remote control system regulated by some 'nano-mechanism, remains to be investigated. The serrated sutures are also not simple, V-shaped serrations but have a complexity of having small and large indentations without any names.

The number of serrations, streaks or convolutions is also to be studied, to find out whether it is only their size or also the number which increases with advancing age. Is it possible to slide horizontally the two separated parietal bones to interlock them or this is not possible due to the convolutions having obliquity for better strength to the suture.

Some sutures fail to grow, either due to some failure or deficiency or remote control mechanism and cause synostosis and lead to craniofacial abnormalities which need to be managed well in time. The genesis is obscure and an early identification and research is called for, so that either this could be prevented or treated, as soon as the diagnosis is established.

The bony growth is at right angle to the suture line, but in cases of craniosynostosis no growth at right angle to the suture takes place and hence this dimension of the skull remains short while other sutures contribute to the growth of the skull. The exact location of the growth centres need to be established.

The children born with cleft lip and palate invariably have problems with occlusion. Some sort of chip or device or a dynamic expansion plate simulating distraction osteogenesis which should be user friendly and may easily be fabricated, needs to be devised to stimulate the growth of the maxilla on the affected side, which will go a long way in correcting occlusion and obviate the need for orthognathic surgery in these children.

The craniofacial clefts from zero to fourteen are established clinical entities with characteristic soft tissue and/or bony abnormalities which are quite consistent. Obviously, there is some sort of temporary arrest of development and then there is normal development beyond it. These clefts have a radially progressive mechanism below and above the level of the eye. What is it, that causes these ecto-mesodermal clefts locally or through a remote control, nano-mechanism is something worth investigating to prevent the occurrence of these clefts. This would be a real breakthrough and go a long way in helping the society at large.

Prof. Ramesh Chandra M.S. (Gen. Surg.), M.S. (Plastic Surg.),
FRCS (Eng.)
KG's Medical College, Lucknow, India

E-mail address: chandradocorsclinic@gmail.com (R. Chandra).