Insights into Interspecific Hybridization Events in Allotetraploid Cotton Formation from Characterization of a Gene-Regulating Leaf Shape

Lijing Chang, Lei Fang, Yajuan Zhu, Huaitong Wu, Zhiyuan Zhang, Chunxiao Liu, Xinghe Li, and Tianzhen Zhang¹

State Key Laboratory of Crop Genetics and Germplasm Enhancement, Cotton Hybrid R & D Engineering Center (the Ministry of Education), College of Agriculture, Nanjing Agricultural University, 210095, China

ABSTRACT The morphology of cotton leaves varies considerably. Phenotypes, including okra, sea-island, super-okra, and broad leaf, are controlled by a multiple allele locus, L_2 . Okra leaf (L_2^o) is an incomplete mutation that alters leaf shape by increasing the length of lobes with deeper sinuses. Using a map-based cloning strategy, we cloned the L_2 locus gene, which encodes a *LATE MERISTEM IDENTITY 1* (*LMI1*)-like transcription factor (*GhOKRA*). Silencing *GhOKRA* leads to a change in phenotype from okra to broad leaf. Overexpression of *GhOKRA* in *Arabidopsis thaliana* greatly increases the degree of the leaf lobes and changes the leaf shape. Premature termination of translation in *GhOKRA* results in the production of broad leaves. The sequences of *OKRA* from diploid progenitor D-genome species, and wild races and domesticated allotetraploid cottons in *Gossypium hirsutum* show that a premature termination mutation occurred before and after the formation of tetraploid cotton, respectively. This study provides genomic insights into the two interspecific hybridization events: one produced the present broad leaf and another formed okra leaf phenotype with complete *OKRA*, that occurred during allotetraploid cotton formation.

KEYWORDS cotton; leaf shape; HD-ZIP I transcription factor; interspecific hybridization; evolution

THE cotton genus *Gossypium* includes ~45 diploid ($2n = 2x = 26$) and five allotetraploid ($2n = 4x = 52$) species (Fryxell 1992; Wendel *et al.* 2009). Diploid *Gossypium* species fall into eight different genome groups (A–G, plus K) based on the observations of meiotic pairing behavior (Fryxell 1979; Endrizzi *et al.* 1985; Percival *et al.* 1999). Allotetraploids originated from interspecific hybridization between diploid species in the New World (Wendel 1989; Wendel *et al.* 1995). *Gossypium herbaceum* or *G. arboreum* and *G. raimondii* are generally regarded as the best extants of the A- and D-subgenome progenitors, respectively (Endrizzi *et al.* 1985; Wendel *et al.* 1995; Zhao *et al.* 1998).

The length of the leaf lobes is highly variable in the *Gossypium* genus, but the predominant form in cultivated cottons is the broad leaf type. The lobe of the leaf varies in shape from broadly triangular to lanceolate or even willow leaf. Okra (L_2^o), sea-island (L_2^e), super-okra (L_2^s), sub okra (L_2^u), and broad leaf (L_2) are members of a multiple allelic series that exist at the L_2 locus in chromosome (Chr) D1 (or Chr 15) (Stephens 1945). Okra leaf shape solely found in the D subgenome in tetraploid cotton is incomplete dominant inheritance and alters leaf shape by increasing the length of lobes with deeper sinuses. Broad leaf cotton has wider leaf lobes and less indentation between the major lobes than the okra types. Broad leaf (L_2) is recessive to okra leaf (L_2^o) (Stephens 1945). Cotton with the okra leaf phenotype has a potential resistance to insect pests and drought. The okra leaf type also has a greater CO₂ exchange rate, water use efficiency, and a lower stomatal conductance compared to normal (broad) leaf type of cotton (Pettigrew *et al.* 1993). It also provides less favorable microenvironmental conditions for the habitat of *Bemisia tabaci*, leading to fewer adults and immatures (eggs and nymphs) because of the more open

Copyright © 2016 by the Genetics Society of America

doi: 10.1534/genetics.116.193086

Manuscript received June 24, 2016; accepted for publication August 9, 2016; published Early Online August 24, 2016.

Supplemental material is available online at www.genetics.org/lookup/suppl/doi:10.1534/genetics.116.193086/-/DC1.

¹Corresponding author: State Key Laboratory of Crop Genetics and Germplasm Enhancement, Cotton Hybrid R & D Engineering Center (the Ministry of Education), College of Agriculture, Nanjing Agricultural University, Nanjing 210095, China. E-mail: cotton@njau.edu.cn

canopy, as evidenced by the higher leaf perimeter-to-leaf area ratio (Chu *et al.* 2002). Okra leaf traits can be used as a source of biotic and abiotic stress resistance. Many extra-long staple cotton (*G. barbadense*) cultivars have the L_2^e sea-island genotype, which is similar to L_2^u and heterozygous okra (L_2^o/l_2) in phenotype (Meredith 1984). Okra leaf Upland cotton was grown in Australia on a major scale, accounting for between 40 and 60% of Australian seed sales between 1987 and 1993 - (Stiller and Wilson 2014). The Commonwealth Scientific and Industrial Research Organization has released many okra leaf cultivars, including Siokra1-4 and Siokra S-324. Okra leaf cultivars such as Biaoza 1A have also been developed for a range of circumstances, such as short season production systems and dryland production systems in Xinjiang, China (Ma *et al.* 2001).

L_2^o was mapped to a 5.4 cM interval with molecular markers, and two *LATE MERISTEM IDENTITY 1* (*LM1*)-like genes have been identified as candidates for okra leaf (Andres *et al.* 2014). Using a combination of targeted association analysis, F_2 population-based fine mapping, and comparative sequencing of orthologs, Zhu *et al.* (2016) further identified one *LM1*-like gene to be a candidate gene underlying the okra leaf trait in *G. hirsutum* and named as *GhOKRA*, which was a class I homeodomain leucine zipper (HD-ZIP I) transcription factor. The homologous gene of *LM1*, *REDUCED COMPLEXITY* (*RCO*), was reported to sculpt developing leaflets by repressing growth at their flanks in *Cardamine hirsute* (Vlad *et al.* 2014). In the present research, we used a chromosomal segment introgression line (CSIL) IL-15-5-1 with sea-island leaves (Figure 1A) and map-based isolated the HD-ZIP I transcription factor, which is responsible for okra leaf development in *Gossypium*. Virus-induced gene silencing (VIGS) assay and transgenic *Arabidopsis* studies confirmed that *GhOKRA* regulates okra leaf development in *Gossypium*. Sequence analysis of *OKRA* in D-genome diploid species and *G. hirsutum* races further provides genomic insights into the two interspecific hybridization events that occurred during allotetraploid cotton formation.

Materials and Methods

Plant materials and genotyping

TM-1 is a standard genetic line of Upland cotton and has broad leaves (Kohel *et al.* 1970). T586 is a *G. hirsutum* multiple gene marker line and has okra leaves (Fryxell 1984; Endrizzi *et al.* 1985). Hai7124 is a commercial Sea-island *Verticillium*-resistant cultivar with sea-island leaves. Materials used in our research are summarized in Supplemental Material, Table S1. Eighty-nine accessions from seven races of *G. hirsutum* were listed in Table S2, provided by the Institute of Cotton Research of the Chinese Academy of Agricultural Sciences (CAAS). IL-15-5-1 was derived from a CSIL with sea-island leaves that carried the Hai7124 segment from chromosome D1 in the background of the recurrent parent TM-1 (Wang *et al.* 2012). Two F_2 mapping populations of 770 and 508 individuals were developed from a cross between TM-1 as the female parent and IL-15-5-1 as the male

parent. Their F_1 had an intermediate phenotype. In the first mapping population, (IL-15-5-1 $\times$ TM-1) F_3 produced 176 homozygous sea-island, 392 segregation for sea-island and broad leaf, and 202 homozygous broad leaf family lines. And the second mapping population contained 112 homozygous sea-island individuals from (IL-15-5-1 $\times$ TM-1) F_2 . All the materials and mapping populations were grown at the Jiangpu Breeding Station with normal field practices and in the greenhouses of Nanjing Agriculture University. Fresh leaves were used as the source of genomic DNA, and leaf primordia of different leaf shape varieties were harvested for total RNA extraction. All samples were immediately frozen in liquid nitrogen after collection and stored at -70° .

Cloning of the L_2 gene

The genome sequence of L_2 was amplified from Super okra, T586, TM-1, and Hai7124 plants and from the other materials listed in Table S1. The gene-specific PCR primer pairs used are described in Table S3. The complementary DNA (cDNA) of *GhOKRA* was cloned from the leaf primordium of super-okra cotton by reverse transcription-polymerase chain reaction (RT-PCR). Amplification products produced using ExTaq DNA Polymerase (TaKaRa Bio, Shiga, Japan) were cloned into the pMD19-T vector (TaKaRa Bio) for sequencing by Jinsite Biotechnology (Nanjing, China). The A-subgenome and D-subgenome sequences were distinguished by alignment with the homologous genes in *G. raimondii* (Paterson *et al.* 2012) and *G. arboreum* (Li *et al.* 2014).

Phylogenetic analysis of *OKRA* sequences

Alignment of the sequences of *OKRA* from different cotton materials (Table S1) was performed with ClustalX (1.83) (Thompson *et al.* 1997) and CLC sequence view 6. A phylogenetic tree was constructed by the maximum likelihood (ML) method in MEGA 6.06 (www.megasoftware.net) and the bootstrap test of phylogeny was performed with 1000 replications (Tamura *et al.* 2013).

Quantitative RT-PCR analysis

Total RNA was extracted from cotton leaf primordia and *A. thaliana* rosette leaves using the Biospin Plant Total RNA Extraction Kit (BioFlux, cat. no. BSC65S1). Leaf primordia at squaring stage were used for detecting the expression difference between species. Leaf primordia at seedling stage were used in the VIGS experiment. First-strand cDNA was generated using TransScript One-Step Genomic DNA Removal and cDNA Synthesis SuperMix (TransGen Biotech, Beijing, China, cat. no. AT311) kits according to the manufacturer's instructions. The cotton *Histone 3* gene (His3, GenBank accession no. AF024716) and *Actin2* (Act2, At3g18780) of *A. thaliana* were used as reference genes. The sequences of the quantitative RT-PCR (qRT-PCR) primers are listed in Table S3. qRT-PCR products were quantified by the ABI 7500 Real Time System (Applied Biosystems, Foster City, CA) and the Light Cycler Fast Start DNA Master SYBR Green I kit (Roche, Basel, Switzerland) according to the manufacturer's instructions.

Figure 1 Map-based cloning of the leaf morphology gene L_2^e . (A) The leaf phenotype of parents used for mapping: TM-1 and IL-15-5-1. From left to right: Third leaf, second leaf, and first leaf in the top to represent mature to young leaves. Bar, 5 cm. (B) L_2^e was first mapped on the D1 chromosome between the markers JESPR152 and NAU3040 using an F_2 population. The genetic distance between the markers was 6.91 cM. L_2^e was further fine mapped to a region between markers JESPR152 and H1776 using 112 dominant individuals. A 183-kb region containing 18 putative ORFs was obtained by mapping this region with sequences in *G. raimondii* Chr 02. The homologous gene of *AtLMI1* is shown at the red arrow.

VIGS assay

The cDNA fragment of *GhOKRA* was cloned from the leaf primordium of super-okra cotton. Gene-specific primers (Table S3) and anchored oligo (dT)₁₈ primers were used to obtain a cDNA fragment for 3'-rapid amplification of cDNA ends (3'-RACE). A 273-bp fragment from the 3' end of *GhOKRA* cDNA was amplified from the super-okra cotton. The product was cloned into *EcoRI*-*XbaI*-digested pTRV2, generating a pTRV2-*GhOKRA* construct. The vectors, pTRV1 and pTRV2-*GhOKRA*, were introduced into *Agrobacterium tumefaciens* strain GV3101. More than 15 super-okra individuals were infiltrated with a 1:1 mixture of *Agrobacterium* carrying pTRV1 and pTRV2 as a negative control, or pTRV1 and pTRV2-*GhOKRA*, separately. The primers used for the construction of VIGS vectors are listed in Table S3. The VIGS assay was carried out independently three times, according to methods described previously (Liu *et al.* 2002).

Subcellular localization of *GhOKRA*

The coding region of the *GhOKRA* gene was amplified from the super-okra cotton and inserted into GFP vector pBinGFP4 (Liu *et al.* 2014) to generate the construct pBin-35S::*GhOKRA*-GFP4. The construct was transformed into *A. tumefaciens* strain GV3101 using methods described previously (Sparkes *et al.* 2006). The subcellular localization of the OKRA-GFP4 fusion protein was observed with a confocal laser scanning microscope (Zeiss LSM 780, Berlin, Germany).

A. thaliana transformation

The fragments of truncated *GhOKRA*^M from TM-1, *GrOKRA*^M from *G. raimondii*, and complete *GhOKRA* from super-okra cotton were obtained using primers (Table S3) and recombined into the pBI121 plasmid using a ClonExpress II One Step Cloning Kit (C112-01, Vazyme Biotech, Nanjing, China) to construct the expression vectors pBI121-35S::*GhOKRA*/*GhOKRA*^M/*GrOKRA*^M used for plant transformation. *GhOKRA*^M

and *GrOKRA*^M represented the genome sequence of premature termination *GhOKRA* in TM-1 and *GrOKRA* in *G. raimondii*, respectively. The complete *GhOKRA* fragment was cloned from the cDNA of super-okra cotton. Vectors were introduced into *A. tumefaciens* strain GV3101 for transformation of *A. thaliana* using the floral dip method (Clough and Bent 1998). Transgenic plants were tested by kanamycin selection marker and PCR analysis of NPTII and 35S-*GhOKRA*.

Data availability

Table S1 contains detailed descriptions of materials used in this research. Table S2 contains accessions from seven races of *G. hirsutum* and genotypes for each accession. Table S3 contains all of the primers designed in this research.

Results

Mapping of the okra leaf gene

IL-15-5-1 is a CSIL derived from a cross between *G. barbadense* cv. Hai7124 with a sea-island leaf and the recurrent parent *G. hirsutum* accession TM-1 (Figure 1A). Although it is phenotypically a sea-island leaf plant (Wang *et al.* 2012), the CSIL line is a near-isogenic line to TM-1. IL-15-5-1 was crossed as a male parent with TM-1 to produce a mapping population. Their F_1 had an intermediate phenotype and (IL-15-5-1 $\times$ TM-1) F_3 produced 176 homozygous sea-island, 392 segregation for sea-island and broad leaf, and 202 homozygous broad leaf family lines, consistent with the segregation ratio of 1:2:1 ($\chi^2 = 2.01 < \chi^2_{0.05, 2} = 5.99$, Table S4). Using microsatellites or simple sequence repeats (SSRs) distributed on Chr 1D, the L_2^e gene was anchored in the introgression segment and linked with SSRs, JESPR152 and NAU3040 (Table S3). To isolate the L_2^e gene, we reproduced (IL-15-5-1 $\times$ TM-1) F_2 , consisting of 508 individuals for the fine mapping of L_2^e (Table S4). We screened out 112 homozygous sea-island individuals from (IL-15-5-1 $\times$ TM-1) F_2 to narrow down

Figure 2 Analysis of genome sequences and transcript levels of *OKRA* in leaves of different shapes. (A) Different leaf shapes of cotton. From left to right: broad (TM-1), sea-island (Hai7124), okra (T586), super-okra (Super okra). (B) qRT-PCR analysis of *OKRA* gene expression in leaves of different shapes. The expression of *OKRA* was consistent with the phenotype of the leaf lobes. (C) Genome sequence analysis of *OKRA*. One 8-bp deletion and one 1-bp deletion were located in the TM-1 genome sequence at positions 742 and 861 bp from the ATG start codon, respectively. In this super-okra line, the single-nucleotide “G” labeled by red frame was not deleted at the C terminus. Data represent the means $\pm$ SDs of three replicated experiments. Different letters indicate significant differences between columns at $P < 0.05$.

the L_2^e locus between markers JESPR152 and NAU3040, using other SSRs (Table S3). The L_2^e gene was delimited in a 183-kb region flanked by markers JESPR152 and H1776, based on the genome sequences of *G. raimondii* (Paterson *et al.* 2012) and *G. hirsutum* (Zhang *et al.* 2015) (Figure 1B).

Cloning of the okra leaf gene

In the 183-kb region, 18 open reading frames (ORFs) were predicted through annotation of the genome sequence of *G. raimondii* Chr 02 (Table S5). qRT-PCR of ORFs revealed that the expression of ORF17 in leaf primordium at squaring stage was consistently different between the super-okra leaf (Super okra), okra leaf (T586), sea-island leaf (Hai7124), and broad leaf (TM-1) (Figure 2, A and B and Figure S1). This gene was not expressed detectably in the broad leaf (TM-1), but its expression was significantly higher in the super-okra leaf (Super okra) than in the okra leaf (T586) and sea-island leaf (H7124). This gene is annotated as a homologous gene of *LATE MERISTEM IDENTITY 1* (*LM11*) of *Arabidopsis* and is an HD-ZIP I transcription factor. This HD-ZIP I transcription factor likely controlling the deeply lobed leaf shape in *Gossypium* is the same as the gene *GhOKRA* identified by Zhu *et al.* (2016).

We cloned the genomic sequences of *OKRA* in Chr 1D (referred as *OKRA-1D*) from different varieties of leaf including Super okra, T586, Hai7124, and TM-1. Multiple sequence alignment revealed that two single-nucleotide transitions at positions 185 and 268 bp from the ATG start codon were detected, along with one 8-bp deletion and one 1-bp deletion at positions 742 and 861 bp from the ATG start codon, respectively, in TM-1 (Figure 2C and Figure S2 and Table S1). In our super-okra line, we did not observe a single-nucleotide deletion at the C terminus that caused the super-okra leaf phenotype reported by Zhu *et al.* (2016). It is supposed that the broad leaf is mutated from the wild-type (WT) okra leaf, due to a frameshift mutation. *OKRA*-GFP was located in the nucleus, which was consistent with its putative function as a transcription factor (Figure S3).

GhOKRA controls cotton okra leaf development

To explore whether *GhOKRA* participates in leaf morphogenesis or not, we cloned 3'-end fragments from *GhOKRA* of super-okra cotton into VIGS pTRV2 vectors (Liu *et al.* 2002) and constructed a pTRV2-*GhOKRA* construct (Figure 3A). At 45 days after the induction of TRV-mediated gene silencing, the leaf of all super-okra cotton infiltrated with pTRV2-*GhOKRA* changed from deeply lobed okra to palmate (Figure 3, B and C). qRT-PCR analysis showed that the *GhOKRA* transcript was down-regulated in pTRV2-*GhOKRA* VIGS lines as compared with controls (Figure 3D). These results demonstrate that *GhOKRA* regulates leaf shape development.

In *G. hirsutum*, there are three homologous genes of *AtLM11* that include *GhOKRA*. The other two homologs were not related to leaf development by VIGS assay (data not shown), suggesting their function has diverged.

Overexpression of *GhOKRA* changes *A. thaliana* leaf shape

To determine whether premature termination or low expression of *GhOKRA* in TM-1 changes leaf development, we developed transgenic *A. thaliana* lines that overexpressed *GhOKRA^M* (a genomic sequence of premature termination of *GhOKRA* in TM-1), *GrOKRA^M* (a genomic sequence of *GrOKRA* in *G. raimondii*), and complete *GhOKRA* (encoding region of cDNA from super-okra cotton) under control of the 35S promoter (Pro35S::*GhOKRA*) in Col-0. The expression of *GhOKRA* and *GhOKRA^M* increased significantly in these transgenic plants (Figure 4A). The rosette leaf blades in transgenic plants overexpressing *GhOKRA* became narrow, abaxially curled, and divided into lobes as compared to the WT in the T1 generation (Figure 4, B and C), and the petiole was longer than that in the WT (Figure 4, D and E). There was no phenotypic difference between transgenic plants overexpressing *GhOKRA^M*, *GrOKRA^M*, and the WT (Figure 4, B and F). These results show that premature termination of translation of *GhOKRA* derived from an 8-bp deletion made this gene nonfunctional in TM-1. This result further verifies

Figure 3 VIGS phenotypes and suppression of the endogenous transcripts. (A) Schematic representation of the structure of pTRV2-GhOKRA containing the 3' cDNA in a 273-bp length of GhOKRA. (B and C) Leaf shape phenotypes: Super okra, pTRV2, pTRV2-GhOKRA, and TM-1. The leaves of plants with pTRV2-GhOKRA transformed to a broad shape but the leaves of plants with pTRV2 did not change. pTRV2 was used as a negative control. (D) qRT-PCR analysis was used to examine the transcript levels of GhOKRA in leaf primordia of plants infected with pTRV2 and pTRV2-GhOKRA. The expression of GhOKRA was significantly silenced by VIGS in plants infected with pTRV2-GhOKRA. Data represent the means $\pm$ SDs of three replicated experiments. Asterisks indicate a significant difference at * $P < 0.05$ and ** $P < 0.01$.

that GhOKRA controls leaf shape in cotton in the same way as in *Arabidopsis*.

Evolution of leaf shape in D-genome diploid species

We found two GhOKRA homologs, *Thecc1EG022391t1* and *CGD0020808*, in cacao (*Theobroma cacao*) (Argout *et al.* 2011). Of the sequenced species, cacao is most closely related to the cotton genus (Paterson *et al.* 2012). The multiple sequence alignments indicated that no such 1- and 8-bp deletions occur in cacao (Figure S4). There exist 13–14 D-genome diploid species (Wendel *et al.* 2009). *G. raimondii* (D₅) is generally regarded as the best extant of the D-subgenome progenitor (Endrizzi *et al.* 1985; Wendel *et al.* 1995; Zhao *et al.* 1998). Leaf shapes in D-genome diploid species are variable from the broad or unlobed leaf in *G. aridum*, *G. davidsonii*, and *G. raimondii*, to the deeply lobed leaf in *G. thurberi* and *G. trilobum*. The leaf shape of *G. gossypoides* is more or less deeply three lobed. We isolated the sequences

Figure 4 Analysis of truncated GhOKRA by overexpression in *A. thaliana*. (A) Relative expression levels of GhOKRA in transgenic plants. The expression levels of GhOKRA and GhOKRA^M in leaves from three transgenic individual plants were all significantly higher than that in the WT. (B) Rosette of *A. thaliana* WT (Col-0). (C) Rosette of *A. thaliana* expressing GhOKRA. (D and E) Leaf phenotype of WT and transgenic plants expressing GhOKRA. (F) Rosette of *A. thaliana* expressing GhOKRA^M. Transgenic plants overexpressing GhOKRA showed narrow, abaxially curled and divided leaves compared with the WT, while transgenic plants overexpressing GhOKRA^M showed no differences compared with the WT. Premature termination led to the nonfunction of GhOKRA in TM-1. Data represent the means $\pm$ SDs of three replicated experiments. Asterisks indicate a significant difference at * $P < 0.05$ and ** $P < 0.01$. Bar, 1 cm.

of OKRA from these six D-genome diploid species mentioned above and found that there were remarkable differences in sequences in D-genome diploid species, especially at the position of the termination codon (Figure S5). Sequence analysis showed that the 1-bp deletion present in TM-1 was also found in *G. raimondii* (Figure S6), but the 8-bp deletion in TM-1 was absent from all D-genome species. There was an 18-bp deletion in *G. gossypoides* and a transition mutation (T/C) delaying the termination in *G. davidsonii* (Figure S5).

Figure 5 Multiple sequence alignment of *GhOKRA* in seven races of *G. hirsutum*. (A) Genome sequences of *GhOKRA* from some accessions randomly selected from 89 accessions from all seven races were isolated and classified into three sequence types based on the 1-bp deletion and 8-bp deletion at positions 742 and 861 bp from the ATG start codon, respectively, in TM-1. Five accessions from *morrilli*, *palmeri*, and *marie-galante* races had sequences that were consistent with Super okra (type 1). Three accessions had the same sequence type as *G. raimondii* containing the 1-bp deletion (type 2). The remaining five accessions had the same sequence type as TM-1 (type 3). (B) The two interspecific hybridizations that occurred during allotetraploid cotton formation. One hybridization event occurred between extant progenitors of *G. herbaceum* or *G. arboreum* and *G. raimondii*. Another hybridization event occurred between *G. herbaceum* or *G. arboreum* and the WT or nascent *G. raimondii*, which had an okra leaf phenotype with complete OKRA.

Phylogenetic analysis showed that the OKRA sequence in *G. raimondii* was closest to that in tetraploid cotton (Figure S6), confirming that *G. raimondii* (D₅) is the best extant of the D-subgenome progenitor.

The evolution of leaf shape in allotetraploid cottons

Allotetraploid cotton originated from hybridization event(s) between an extant A-genome progenitor of *G. herbaceum* (A₁) or *G. arboreum* (A₂) and a D-genome progenitor, *G. raimondii* (D₅), 1–1.5 million years ago (MYA) (Endrizzi *et al.* 1985; Wendel *et al.* 1995; Zhao *et al.* 1998; Zhang *et al.* 2015). However, the number of times hybridization events occurred to produce allotetraploid cottons remains to be explored. Based mainly on their morphologies and distinct geographical distributions, one wild and six domesticated races of *G. hirsutum* have been identified (Hutchinson 1951). We isolated *GhOKRA* genome sequences from 89 accessions of all seven races with variable leaf shapes to evaluate the evolution of leaf shape in tetraploid cotton (Figure 5A and Figure S7). There were three sequence types in the *GhOKRA* genome sequence among the 89 accessions (Table S2). Seventeen accessions from *morrilli*, *palmeri*, *latifolium*, and *marie-galante* races had sequences that were completely consistent with okra leaf cultivated species (type 1). Twelve accessions from *punctatum*, *yucatanen*, *palmeri*, *marie-galante*, and *latifolium* races had the same mutated sequence as three accessions from *G. raimondii*, containing 1-bp deletion (type 2), suggesting that the broad leaf may originate from diploid progenitor cotton *G. raimondii*, although we can not exclude the possibility that the same mutated sequence 1-bp deletion occurred after

the formation of tetraploid cottons. The remaining accessions from *richmondii*, *latifolium*, *morrilli*, *punctatum*, *palmeri*, and *marie-galante* races had the same sequence as TM-1 and modern broad leaf cultivars, containing both the 8-bp and 1-bp deletions at positions 742 and 861 bp from the ATG start codon, respectively (type 3). These results revealed that another mutation (an 8-bp deletion) occurred to develop the broad leaf phenotype after the formation of tetraploid cotton.

Discussion

The tetraploid was spawned from the interspecific hybridization of an A-genome species resembling an African (*G. herbaceum*) or Asian (*G. arboreum*) species and a D-genome species resembling extant *G. raimondii* (Wendel 1989). Under long-term human selection of a wide range of morphological and physiological traits, the two tetraploid species, *G. hirsutum* and *G. barbadense*, have been domesticated in South and Central America, respectively. Twelve D-genome diploid species are endemic to western Mexico, indicating this area is the center of diversity of the D genome. However, *G. raimondii* is presently distributed only in Peru. The 1-bp deletion in OKRA present in wild race species and modern cotton cultivars was found only in *G. raimondii*, indicating that the mutation in tetraploid cottons originated from the D-diploid progenitor species *G. raimondii*. We identified three types of *GhOKRA* in 89 accessions from seven wild and domesticated races of tetraploid Upland cotton and found that another 8-bp deletion mutation occurred after the formation of tetraploid cotton. From the evolution of leaf

shape in allotetraploid cottons, we can reasonably deduce that allotetraploid cottons may originate from at least two interspecific hybridizations (Figure 5B). One hybridization event occurred between extant progenitors of *G. herbaceum* or *G. arboreum* and *G. raimondii* in which a 1-bp “G” deletion had already occurred before the formation of tetraploid cotton and produced the present broad leaf tetraploid cottons. Although a premature termination mutation with 1-bp deletions at positions 861 bp could occur after the tetraploid cotton formation, the possibility of such a premature termination mutation occurred at the same position in the tetraploid cotton and *G. raimondii* would be very low. Another hybridization event occurred between *G. herbaceum* or *G. arboreum* and the WT or nascent *G. raimondii*, which had an okra leaf phenotype with complete *OKRA*, to form the okra leaf tetraploid cottons. The phenotype is validated with *GhOKRA* haplotypes in the wild and domesticated races of tetraploid Upland cotton. After the formation of tetraploid cotton, a further 8-bp deletion occurred in the tetraploid cottons, originating from already broad leaf tetraploid cottons, and resulting in the gene *L₂*, as found in TM-1.

The phenotypic differences in leaf shape are controlled by a single multiple allele series in tetraploid cotton (Stephens 1945). Leaf shape in the D-genome diploid is highly variable. In the present research, using a map-based cloning strategy, we cloned the *L₂* locus genes using a sea-island okra leaf (*L₂^e*) CSIL, IL-15-5-1 (Wang *et al.* 2012), and found that both 1- and 8-bp deletions cause premature termination of translation and the nonfunction of *OKRA* and lead to broad leaf development in *Gossypium*. Super-okra leaves had a higher expression of *OKRA* than other okra leaves at the transcriptional level. Although the remarkable morphological variation between okra (*L₂^o*), sea-island (*L₂^e*), super-okra (*L₂^s*), and sub-okra (*L₂^u*) types are attributed to the same *L₂* multiple allele locus (Stephens 1945), their ORFs in this *OKRA* did not show any differences in their sequences. It remains to be explored whether it is epigenetics, a *cis*-regulatory mutation at this locus, or a causative mutation on a distant gene exerting its influence on *OKRA* that has caused such varied expression in these different okra leaf lines with the same ORF sequences.

Acknowledgments

This work was financially supported in part by grants from the National Natural Science Foundation of China (31330058), National Key R & D Program for Crop Breeding (2016YFD0100300), and the Jiangsu Province Collaborative Innovation Center-Modern Crop Production (JCIC-MCP) project. We thank the Medium-Term Gene Bank of Cotton in China and the Institute of Cotton Research of CAAS for providing some of the cotton seeds used in the present study. The authors declare no competing financial interests.

Author contributions: T.Z. designed the research; L.C., L.F., and Y.Z. performed the research; T.Z., L.C., and L.F. analyzed all data and wrote the manuscript; Y.Z., H.W.,

and X.L. constructed mapping populations; H.W. designed SSR primers; L.C. and Z.Z. conducted subcellular localization; L.C., L.F., and C.L. analyzed the evolution; and all authors discussed results and commented on the manuscript.

Literature Cited

- Andres, R. J., D. T. Bowman, B. Kaur, and V. Kuruparthi, 2014 Mapping and genomic targeting of the major leaf shape gene (*L*) in Upland cotton (*Gossypium hirsutum* L.). *Theor. Appl. Genet.* 127: 167–177.
- Argout, X., J. Salse, J. M. Aury, M. J. Guiltinan, G. Droc *et al.*, 2011 The genome of *Theobroma cacao*. *Nat. Genet.* 43: 101–108.
- Chu, C. C., E. T. Natwick, and T. J. Henneberry, 2002 Bemisia tabaci (Homoptera: Aleyrodidae) biotype B colonization on okra- and normal-leaf upland cotton strains and cultivars. *J. Econ. Entomol.* 95: 733–738.
- Clough, S. J., and A. F. Bent, 1998 Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* 16: 735–743.
- Endrizzi, J. E., E. L. Turcotte, and R. J. Kohel, 1985 Genetics, cytology, and evolution of *Gossypium*. *Adv. Genet.* 23: 271–375.
- Fryxell, P. A., 1979 *The Natural History of the Cotton Tribe*. Texas A&M University Press, College Station, TX.
- Fryxell, P. A., 1984 Taxonomy and germplasm resources. pp. 27–57 in *Cotton*, edited by R. J. Kohel, and C. F. Lewis. American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America, Madison, WI.
- Fryxell, P. A., 1992 A revised taxonomic interpretation of *Gossypium* L. (Malvaceae). *Rheedea* 2: 108–165.
- Hutchinson, J. B., 1951 Intra-specific differentiation in *Gossypium hirsutum*. *Heredity* 5: 161–193.
- Kohel, R., T. R. Richmond, and C. F. Lewis, 1970 Texas Marker1. Description of a genetic standard for *Gossypium hirsutum* L. *Crop Sci.* 10: 670–673.
- Li, F., G. Fan, K. Wang, F. Sun, Y. Yuan *et al.*, 2014 Genome sequence of the cultivated cotton *Gossypium arboreum*. *Nat. Genet.* 46: 567–572.
- Liu, T., T. Song, X. Zhang, H. Yuan, L. Su *et al.*, 2014 Unconventionally secreted effectors of two filamentous pathogens target plant salicylate biosynthesis. *Nat. Commun.* 5: 4686.
- Liu, Y., M. Schiff, and S. P. Dinesh-Kumar, 2002 Virus-induced gene silencing in tomato. *Plant J.* 31: 777–786.
- Ma, Q., X. Yang, J. Liu, Z. Wang, T. Xing *et al.*, 2001 Hybrid cotton Biaoza A1. *China Cotton* 28: 23–24.
- Meredith, W. R., 1984 Influence of leaf morphology on lint yield of cotton-enhancement by the sub okra trait. *Crop Sci.* 24: 855–857.
- Paterson, A. H., J. F. Wendel, H. Gundlach, H. Guo, J. Jenkins *et al.*, 2012 Repeated polyploidization of *Gossypium* genomes and the evolution of spinnable cotton fibres. *Nature* 492: 423–427.
- Percival, A. E., J. F. Wendel, and J. M. Stewart, 1999 Taxonomy and germplasm resources, pp. 33–63 in *Cotton: Origin, History, Technology, and Production*, edited by C. W. Smith, and J. T. Cothren. John Wiley and Sons, New York.
- Pettigrew, W. T., J. J. Heitholt, and K. C. Vaughn, 1993 Gas-exchange differences and comparative anatomy among cotton leaf-type isolines. *Crop Sci.* 33: 1295–1299.
- Sparkes, I. A., J. Runions, A. Kearns, and C. Hawes, 2006 Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. *Nat. Protoc.* 1: 2019–2025.
- Stephens, S. G., 1945 A genetic survey of leaf shape in New World cottons: a problem in critical identification of alleles. *J. Genet.* 56: 313–330.

- Tamura, K., G. Stecher, D. Peterson, A. Filipski, and S. Kumar, 2013 MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* 30: 2725–2729.
- Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G. Higgins, 1997 The CLUSTAL X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 25: 4876–4882.
- Vlad, D., D. Kierzkowski, M. I. Rast, F. Vuolo, R. Dello Ioio *et al.*, 2014 Leaf shape evolution through duplication, regulatory diversification, and loss of a homeobox gene. *Science* 343: 780–783.
- Wang, P., Y. J. Zhu, X. L. Song, Z. B. Cao, Y. Z. Ding *et al.*, 2012 Inheritance of long staple fiber quality traits of *Gossypium barbadense* in *G. hirsutum* background using CSILs. *Theor. Appl. Genet.* 124: 1415–1428.
- Wendel, J. F., 1989 New world tetraploid cottons contain old world cytoplasm. *Proc. Natl. Acad. Sci. USA* 86: 4132–4136.
- Wendel, J. F., A. Schnabel, and T. Seelanan, 1995 An unusual ribosomal DNA sequence from *Gossypium gossypoides* reveals ancient, cryptic, intergenomic introgression. *Mol. Phylogenet. Evol.* 4: 298–313.
- Wendel, J. F., C. Brubaker, I. Alvarez, R. Cronn, and J. Stewart, 2009 Evolution and natural history of the cotton genus, pp. 3–22 in *Genetics and Genomics of Cotton*, edited by H. A. Paterson. Springer-Verlag, New York.
- Stiller, W. N., and I. W. Wilson, 2014 Australian cotton germplasm resources, pp. 1–34 in *World Cotton Germplasm Resources*, edited by I. Y. Abdurakhmonov. InTech, Rijeka, Croatia.
- Zhang, T., Y. Hu, W. Jiang, L. Fang, X. Guan *et al.*, 2015 Sequencing of allotetraploid cotton (*Gossypium hirsutum* L. acc. TM-1) provides a resource for fiber improvement. *Nat. Biotechnol.* 33: 531–537.
- Zhao, X. P., Y. Si, R. E. Hanson, C. F. Crane, H. J. Price *et al.*, 1998 Dispersed repetitive DNA has spread to new genomes since polyploid formation in cotton. *Genome Res.* 8: 479–492.
- Zhu, Q. H., J. Zhang, D. Liu, W. Stiller, Z. Zhang *et al.*, 2016 Integrated mapping and characterization of the gene underlying the okra leaf trait in *Gossypium hirsutum* L. *J. Exp. Bot.* 67: 763–774.

Communicating editor: S. Poethig

Figure S1 Expression levels of ORFs from different leaf shape varieties in the mapping region.

The expression of ORFs in the overlapped mapping region with previous study was detected in different leaf shape varieties. The overlapped region contained 13 ORFs (ORF6-ORF18) including predicted candidate gene ORF17. ORF16 and ORF18 were not expressed detectably in all varieties. ORF14 and ORF15 were homologous genes encoding pyruvate kinase family protein with high sequence similarity, whose expressions were detected together. Among the 13 ORFs, except ORF 17 (*OKRA*), the expression level of other ORFs in TM-1 (broad), Hai7124 (sea-island), T586 (okra), Super okra (super okra) was not consistent with the leaf shape. Data represent the means $\pm$ SDs of three replicated experiments.

```

      *      20      *      40      *      60      *      80      *      100      *
Hai17124 : ATGGATTGGGATGGACACCATTCGACCCCTTATTTCACGACTAGAACCTTCCTTAACTCCCTTCTAACATAAATTATAATCAATATCCAGTAAATATATATTTATATATAGT : 115
T586 : ATGGATTGGGATGGACACCATTCGACCCCTTATTTCACGACTAGAACCTTCCTTAACTCCCTTCTAACATAAATTATAATCAATATCCAGTAAATATATATTTATATATAGT : 115
Super-okra : ATGGATTGGGATGGACACCATTCGACCCCTTATTTCACGACTAGAACCTTCCTTAACTCCCTTCTAACATAAATTATAATCAATATCCAGTAAATATATATTTATATATAGT : 115
TM-1 : ATGGATTGGGATGGACACCATTCGACCCCTTATTTCACGACTAGAACCTTCCTTAACTCCCTTCTAACATAAATTATAATCAATATCCAGTAAATATATATTTATATATAGT : 115

      120      *      140      *      160      *      180      *      200      *      220      *
Hai17124 : CCTTTATTTCTTGTCTTCCACTTCTTTATTTCTAACTTTTGTTCAGAGGTGTGGAGGACATGACAAATCAAGGATTTGAAGAAGCTTGCAGATGGTTCAGATT : 230
T586 : CCTTTATTTCTTGTCTTCCACTTCTTTATTTCTAACTTTTGTTCAGAGGTGTGGAGGACATGACAAATCAAGGATTTGAAGAAGCTTGCAGATGGTTCAGATT : 230
Super-okra : CCTTTATTTCTTGTCTTCCACTTCTTTATTTCTAACTTTTGTTCAGAGGTGTGGAGGACATGACAAATCAAGGATTTGAAGAAGCTTGCAGATGGTTCAGATT : 230
TM-1 : CCTTTATTTCTTGTCTTCCACTTCTTTATTTCTAACTTTTGTTCAGAGGTGTGGAGGACATGACAAATCAAGGATTTGAAGAAGCTTGCAGATGGTTCAGATT : 230

      240      *      260      *      280      *      300      *      320      *      340      *
Hai17124 : TGAATATGAACGGCTTCATGAACATGGTAAAGGTAAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAAGACAATAAGTTGGA : 345
T586 : TGAATATGAACGGCTTCATGAACATGGTAAAGGTAAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAAGACAATAAGTTGGA : 345
Super-okra : TGAATATGAACGGCTTCATGAACATGGTAAAGGTAAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAAGACAATAAGTTGGA : 345
TM-1 : TGAATATGAACGGCTTCATGAACATGGTAAAGGTAAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAAGACAATAAGTTGGA : 345

      *      360      *      380      *      400      *      420      *      440      *      460      *
Hai17124 : TCCCGACAGGAAATGAAGCTTCCAGGAACCTTGAAGCTTCAACCAAGACAATCGCTGTCTGGTTCCAAACAGGCGTGTAGATGGAAGCTTAACAGCTTGAACACTCATAT : 460
T586 : TCCCGACAGGAAATGAAGCTTCCAGGAACCTTGAAGCTTCAACCAAGACAATCGCTGTCTGGTTCCAAACAGGCGTGTAGATGGAAGCTTAACAGCTTGAACACTCATAT : 460
Super-okra : TCCCGACAGGAAATGAAGCTTCCAGGAACCTTGAAGCTTCAACCAAGACAATCGCTGTCTGGTTCCAAACAGGCGTGTAGATGGAAGCTTAACAGCTTGAACACTCATAT : 460
TM-1 : TCCCGACAGGAAATGAAGCTTCCAGGAACCTTGAAGCTTCAACCAAGACAATCGCTGTCTGGTTCCAAACAGGCGTGTAGATGGAAGCTTAACAGCTTGAACACTCATAT : 460

      *      480      *      500      *      520      *      540      *      560      *
Hai17124 : AATACGCTTAAACATGAGTATGATATTATCTCCATTGAAAAGCAAAGCTACAAGACGAGGTAGCCTCTATATTCAAAACCTATTTTTTTATCCCTTCTCTCGCTCTCTATA : 575
T586 : AATACGCTTAAACATGAGTATGATATTATCTCCATTGAAAAGCAAAGCTACAAGACGAGGTAGCCTCTATATTCAAAACCTATTTTTTTATCCCTTCTCTCGCTCTCTATA : 575
Super-okra : AATACGCTTAAACATGAGTATGATATTATCTCCATTGAAAAGCAAAGCTACAAGACGAGGTAGCCTCTATATTCAAAACCTATTTTTTTATCCCTTCTCTCGCTCTCTATA : 575
TM-1 : AATACGCTTAAACATGAGTATGATATTATCTCCATTGAAAAGCAAAGCTACAAGACGAGGTAGCCTCTATATTCAAAACCTATTTTTTTATCCCTTCTCTCGCTCTCTATA : 575

      580      *      600      *      620      *      640      *      660      *      680      *
Hai17124 : TATCTTTACGTGGAACTTGGTGCACTGTTTTATAAAACGAATATGCTTGTGTCTTATTGTTGAGGTTTCCCATTTTGTATACACACATACATACATATATATATTAT : 690
T586 : TATCTTTACGTGGAACTTGGTGCACTGTTTTATAAAACGAATATGCTTGTGTCTTATTGTTGAGGTTTCCCATTTTGTATACACACATACATACATATATATATTAT : 690
Super-okra : TATCTTTACGTGGAACTTGGTGCACTGTTTTATAAAACGAATATGCTTGTGTCTTATTGTTGAGGTTTCCCATTTTGTATACACACATACATACATATATATATTAT : 690
TM-1 : TATCTTTACGTGGAACTTGGTGCACTGTTTTATAAAACGAATATGCTTGTGTCTTATTGTTGAGGTTTCCCATTTTGTATACACACATACATACATATATATATTAT : 690

      700      *      720      *      740      *      760      *      780      *      800      *
Hai17124 : ACTGTACTCTCATACCTTTTACGCAAGTGAATGAAAGGAATGCTAGTGGACAGCGCACAAGGAACCAAGTCTCCACGGTTTACAAGGAAATCTCCGCGGAGAGATCAT : 805
T586 : ACTGTACTCTCATACCTTTTACGCAAGTGAATGAAAGGAATGCTAGTGGACAGCGCACAAGGAACCAAGTCTCCACGGTTTACAAGGAAATCTCCGCGGAGAGATCAT : 805
Super-okra : ACTGTACTCTCATACCTTTTACGCAAGTGAATGAAAGGAATGCTAGTGGACAGCGCACAAGGAACCAAGTCTCCACGGTTTACAAGGAAATCTCCGCGGAGAGATCAT : 805
TM-1 : ACTGTACTCTCATACCTTTTACGCAAGTGAATGAAAGGAATGCTAGTGGACAGCGCACAAGGAACCAAGTCTCCACGGTTTACAAGGAAATCTCCGCGGAGAGATCAT : 797
      ACTGTACTCTCATACCTTTTACGCAAGTGAATGAAAGGAATGCTAGTggaaacagCGCACAAGGAACCAAGTCTCCACGGTTTACAAGGAAATCTCCGCGGAGAGATCAT

      *      820      *      840      *      860      *      880      *      900      *      920      *
Hai17124 : CGAAAGTAGCTCGATTTCGGAGCTCGAACAGACGAGTATAGCAGGAACGACTATATCCGATAGTTGAATGTAACATATTTCAATGAGGATGAGAATAACCCAGTTCCACT : 920
T586 : CGAAAGTAGCTCGATTTCGGAGCTCGAACAGACGAGTATAGCAGGAACGACTATATCCGATAGTTGAATGTAACATATTTCAATGAGGATGAGAATAACCCAGTTCCACT : 920
Super-okra : CGAAAGTAGCTCGATTTCGGAGCTCGAACAGACGAGTATAGCAGGAACGACTATATCCGATAGTTGAATGTAACATATTTCAATGAGGATGAGAATAACCCAGTTCCACT : 920
TM-1 : CGAAAGTAGCTCGATTTCGGAGCTCGAACAGACGAGTATAGCAGGAACGACTATATCCGATAGTTGAATGTAACATATTTCAATGAGGATGAGAATAACCCAGTTCCACT : 911
      CGAAAGTAGCTCGATTTCGGAGCTCGAACAGACGAGTATAGCAGGAACGACTATgATTCGATAGTTGAATGTAACATATTTCAATGAGGATGAGAATAACCCAGTTCCACT

      *      940      *
Hai17124 : CACTACTGGGATATTCAACTCCCTTCTATCCCTAA : 956
T586 : CACTACTGGGATATTCAACTCCCTTCTATCCCTAA : 956
Super-okra : CACTACTGGGATATTCAACTCCCTTCTATCCCTAA : 956
TM-1 : CACTACTGGGATATTCAACTCCCTTCTATCCCTAA : 947
      CACTACTGGGATATTCAACTCCCTTCTATCCCTAA

```

Figure S2 Alignment of *OKRA* gene sequences from Super okra, T586, Hai7124, and TM-1. The black shadow indicates exon, the gray shadow indicates SNP sites and deletions. The sequences without shadow are introns. The blue shadow labels termination codon in TM-1. The 1-bp deletion and 8-bp deletion at the positions 742bp and 861bp from the ATG start codon are indicated by blue arrows.

Figure S3 Subcellular localization of *GhOKRA*. Subcellular localization of *GhOKRA* was obtained by transient expression of *GhOKRA* fused to the pBinGFP4-35S::GFP vector in tobacco leaf cells. The pBinGFP4-35S::GFP construct was used as the control. Merged overlay of bright field and GFP fluorescence images; Bright field images of tobacco leaf epidermis cell; Scale bar=50 μm .

```

      *      20      *      40      *      60      *      80      *      100      *
G. raimondii : ATGGATTGGGATGGCACCAATTCGACCTTTATTTTCGACCTAGAACCTTCTCTTAACCTTCCTTCTAACTATAATATATCAATATCCAGAAAGGTGTGGAGGACATGACACATC : 115
TM-1 : ATGGATTGGGATGGCACCAATTCGACCTTTATTTTCGACCTAGAACCTTCTCTTAACCTTCCTTCTAACTATAATATATCAATATCCAGAAAGGTGTGGAGGACATGACACATC : 115
Super okra : ATGGATTGGGATGGCACCAATTCGACCTTTATTTTCGACCTAGAACCTTCTCTTAACCTTCCTTCTAACTATAATATATCAATATCCAGAAAGGTGTGGAGGACATGACACATC : 115
T. cacao_1 : ATGGATTGGGATGGCACCTTAAGGACCTTTGTTTCGACCTAGAACCTTCTCTTAACCTTCCTAACACTATAATCATGATCAATATCCAG-----GTATGGGATGAGGATC : 109
T. cacao_2 : -----ATGGGATGAGGATC : 16

atggatgg atggcacc t cg ccttt ttcc cgac agaaccttctcttaacttcc t aactataa tat atcaatatccag GGA ATGA ATC

      120      *      140      *      160      *      180      *      200      *      220      *
G. raimondii : PAGGAATTCAGGAGCGTGGCATGGGTTGGTTCCAGATTGGAATTCAGAACGCTTCATGAACAATGGTAAGGTAAACACACAGAGAGAAGAGATTGACAAATGATCAGTTGGA : 230
TM-1 : PAGGAATTCAGGAGCGTGGCATGGGTTGGTTCCAGATTGGAATTCAGAACGCTTCATGAACAATGGTAAGGTAAACACACAGAGAGAAGAGATTGACAAATGATCAGTTGGA : 230
Super okra : PAGGAATTCAGGAGCGTGGCATGGGTTGGTTCCAGATTGGAATTCAGAACGCTTCATGAACAATGGTAAGGTAAACACACAGAGAGAAGAGATTGACAAATGATCAGTTGGA : 230
T. cacao_1 : CAGGACATCTGAGCGATTTTCATGGGTTGGTTCCAGCTTTGACAGAAACAGCT---ACAACAATC---AAG-----AAGAGAAGAGAGATTGACAAATGATCAGTTGGA : 209
T. cacao_2 : CAGGACATCTGAGCGATTTTCATGGGTTGGTTCCAGCTTTGACAGAAACAGCT---ACAACAATC---AAG-----AAGAGAAGAGAGATTGACAAATGATCAGTTGGA : 116

AGGA T G GA GC G ATGGGTTGGTTCCAG TTG A A GAAC GCT AACAAT AAG A AAGAAGAAGAG TTGACAA TGATCAATT GA

      240      *      260      *      280      *      300      *      320      *      340      *
G. raimondii : TTCCTTGGAGAGGAGTTTCCAGGAACAGATTAAGTTGGTCCGACAGAGAAATGAAGCTTTCCAGGAACTTGGACTTCAACCAAGCAAAATGGCTCTGGTTCCAAAACAGG : 345
TM-1 : TTCCTTGGAGAGGAGTTTCCAGGAACAGATTAAGTTGGTCCGACAGAGAAATGAAGCTTTCCAGGAACTTGGACTTCAACCAAGCAAAATGGCTCTGGTTCCAAAACAGG : 345
Super okra : TTCCTTGGAGAGGAGTTTCCAGGAACAGATTAAGTTGGTCCGACAGAGAAATGAAGCTTTCCAGGAACTTGGACTTCAACCAAGCAAAATGGCTCTGGTTCCAAAACAGG : 345
T. cacao_1 : TTCCTTGGAGAGGAGTTTCCAGGAACAGATTAAGTTGGTCCGACAGAGAAATGAAGCTTTCCAGGAACTTGGACTTCAACCAAGCAAAATGGCTCTGGTTCCAAAACAGG : 324
T. cacao_2 : TTCCTTGGAGAGGAGTTTCCAGGAACAGATTAAGTTGGTCCGACAGAGAAATGAAGCTTTCCAGGAACTTGGACTTCAACCAAGCAAAATGGCTCTGGTTCCAAAACAGG : 231

TC TTGGA AGGAGTTTCCA GAAGAGA AAGTTGGA CC GACAGAGAAATGAAGCTTTC A GGA CTGG CT CAACCAAG CAAAT GC GCTCTGGTTCCAAAACAGG

      360      *      380      *      400      *      420      *      440      *      460      *
G. raimondii : CGTGCTAGATGGAAGCTTAACAGCTTGACCTCTATATATATCGCTTAAACAGGAGTATGATTTATCTCCATTGAAAGCAAGCAAGCTCAAGAGAGAGGTGATGGAATTGAAG : 460
TM-1 : CGTGCTAGATGGAAGCTTAACAGCTTGACCTCTATATATATCGCTTAAACAGGAGTATGATTTATCTCCATTGAAAGCAAGCAAGCTCAAGAGAGAGGTGATGGAATTGAAG : 460
Super okra : CGTGCTAGATGGAAGCTTAACAGCTTGACCTCTATATATATCGCTTAAACAGGAGTATGATTTATCTCCATTGAAAGCAAGCAAGCTCAAGAGAGAGGTGATGGAATTGAAG : 460
T. cacao_1 : CGTGCTAGATGGAAGCTTAACAGCTTGACCTCTATATATATCGCTTAAACAGGAGTATGATTTATCTCCAGGGAAGCAAGCAAGCTCAAGAGAGAGGTGATGGAATTGAAG : 439
T. cacao_2 : CGTGCTAGATGGAAGCTTAACAGCTTGACCTCTATATATATCGCTTAAACAGGAGTATGATTTATCTCCAGGGAAGCAAGCAAGCTCAAGAGAGAGGTGATGGAATTGAAG : 346

CGTGCTAGATGGAAGCTTAACAGCTTGACCTCTATATATATCGCTTAAACAGGAGTATGATTTATCTCCA GA AAGCA AAGCT CAAGA GAGGT ATG AATTGAAG

      480      *      500      *      520      *      540      *      560      *
G. raimondii : CBAATGCTATGGAAACAGCGACAGAGGAACCGAGCTCCACAGCTTTACAGGAAATCTCCGGGGAGAGAGATCTCGAAAGCACTTGATTCTGCTCAACCAAGAGATATAC : 575
TM-1 : CBAATGCTATGGAAACAGCGACAGAGGAACCGAGCTCCACAGCTTTACAGGAAATCTCCGGGGAGAGAGATCTCGAAAGCACTTGATTCTGCTCAACCAAGAGATATAC : 534
Super okra : CBAATGCTATGGAAACAGCGACAGAGGAACCGAGCTCCACAGCTTTACAGGAAATCTCCGGGGAGAGAGATCTCGAAAGCACTTGATTCTGCTCAACCAAGAGATATAC : 575
T. cacao_1 : CBAATGCTATGGAAACAGCGACAGAGGAACCGAGCTCCACAGCTTTACAGGAAATCTCCGGGGAGAGAGATCTCGAAAGCACTTGATTCTGCTCAACCAAGAGATATAC : 554
T. cacao_2 : CBAATGCTATGGAAACAGCGACAGAGGAACCGAGCTCCACAGCTTTACAGGAAATCTCCGGGGAGAGAGATCTCGAAAGCACTTGATTCTGCTCAACCAAGAGATATAC : 461

G ATGCTA ggaacagGC AC A GAACC AG CTCCAC G TTACA GA AT TCCGG G AGAGA C TCGAAG A ctc attcg gctc aacaag c ag tag

      580      *      600      *      620      *      640      *      660      *      680      *
G. raimondii : AGCAAGGAGATATATCGGATG : 597
TM-1 : ----- : -
Super okra : AGCAAGGAGATATATCGGATGTTGAATGTAAATATTTTCAATGAGGATGAGATTAACCCAGTTTCACACTACTACTGGGATATTCAGTCTCTCTTTATCCCTAA : 684
T. cacao_1 : AGCAAGGAGATATATCGGATGTTGAATGTAACTATGTTTCAATGAGGATGAGATTAACCCAGTTTCACACTACTACTGGGATGTTGAGGTGCTCTCTTTATCCCTAA : 663
T. cacao_2 : AGCAAGGAGATATATCGGATGTTGAATGTAACTATGTTTCAATGAGGATGAGATTAACCCAGTTTCACACTACTACTGGGATGTTGAGGTGCTCTCTTTATCCCTAA : 570

aq aaac ac at atc gat g

```

Figure S4 Sequence alignment between *OKRA* of *Gossypium* and homologs of *T. cacao*. CDS sequences of *OKRA* in *G. raimondii*, TM-1, Super okra and homologs of *T. cacao* (*T. cacao_1*, Thecc1EG022391t1; *T. cacao_2*, CGD0020808) were aligned. The 1-bp deletion and 8-bp deletion appeared in *Gossypium* are indicated by blue arrows.

Figure S6 Phylogenetic analysis and sequence alignment analysis of *OKRA* in D-genome diploid species. Phylogenetic analysis of aligned genome sequences of *OKRA* in *G. aridum*, *G. davidsonii*, *G. raimondii*, *G. thurberi*, *G. trilobum* and *G. gossypoides*. Tetraploid cotton was phylogenetically close to *G. raimondii*. Sequence alignment showed that the 1-bp deletion present in TM-1 was also found in *G. raimondii* and the 8-bp deletion in TM-1 was absent from all D-genome species. Phylogenetic analysis was done using the ML method with 1,000 resampling replicates.

		20	40	60	80	
morrl18	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	T	ATGCTCCCCCAACAGT	80		
palmeri11	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	T	ATGCTCCCCCAACAGT	80		
palmeri8	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	T	ATGCTCCCCCAACAGT	80		
palmeri5	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	T	ATGCTCCCCCAACAGT	80		
Superokra_Dt	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	T	ATGCTCCCCCAACAGT	80		
marie-galante6	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	T	ATGCTCCCCCAACAGT	80		
richmondii4	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
latifolium98B	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
palmeri4	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
TM-1_Dt	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
marie-galante4	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
morrl11	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
punctatum4	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
latifolium191	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
yucatum9	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCATTGTTG	TCATGCTGCC	CCCCCAACAGT	80		
G.raimondii	ATTAACAGACAAGTCCACCCTCTCTCCATCTCCCTTTAAATACACAGTTAAGGCAT	G	TTGTCATGCTCCCCCAACAGT	80		
Conservation						
		100	120	140	160	
morrl18	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
palmeri11	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
palmeri8	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
palmeri5	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
Superokra_Dt	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
marie-galante6	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
richmondii4	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
latifolium98B	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
palmeri4	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
TM-1_Dt	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
marie-galante4	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
morrl11	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
punctatum4	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
latifolium191	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
yucatum9	CAACCCCTTCACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTCACTCTTTGG	160				
G.raimondii	CAACCCCTTACAACAGTCAACCCCTTTTCAGTTCTGGTCTGCTCCCCATTGCTATGCTACTTTAGGCTTACTCTTGG	160				
Conservation						
		180	200	220	240	
morrl18	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
palmeri11	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
palmeri8	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
palmeri5	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
Superokra_Dt	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
marie-galante6	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
richmondii4	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
latifolium98B	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
palmeri4	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
TM-1_Dt	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
marie-galante4	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
morrl11	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
punctatum4	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	240				
latifolium191	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
yucatum9	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
G.raimondii	GCAGC - TTCGTTTCAGTTCTGGTGGTTCCATTTATATATACAGATATACTCGCTTTCTTCATTCAAAGAAATGGATTGGGA	239				
Conservation						
		260	280	300	320	
morrl18	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
palmeri11	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
palmeri8	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
palmeri5	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
Superokra_Dt	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
marie-galante6	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
richmondii4	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
latifolium98B	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
palmeri4	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
TM-1_Dt	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
marie-galante4	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
morrl11	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
punctatum4	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	320				
latifolium191	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
yucatum9	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
G.raimondii	TGGCACCATTTCGACCCCTTTATTTACGACTAGAACCTTCTCTTAACCTTCCCTTCTAACTATAATTATAATCAATATCCAG	319				
Conservation						

		340	360	380	400	
morrll18	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
palmeri11	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
palmeri8	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
palmeri5	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
Superokra_Dt	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
marie-galante6	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
richmondii4	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
latifolium98B	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
palmeri4	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
TM-1_Dt	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
marie-galante4	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
morrll1	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
punctatum4	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					400
latifolium191	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
yucatum9	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
G.raimondii	GTAATATATATATTTATATATAGTCCCTTTATTTTCTTGTTCTTCCACTTTCCTTTATTTCTAACTTTTTGTTTCAGAAGG					399
Conservation	<div></div>					
		420	440	460	480	
morrll18	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
palmeri11	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
palmeri8	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
palmeri5	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
Superokra_Dt	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
marie-galante6	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
richmondii4	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
latifolium98B	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
palmeri4	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
TM-1_Dt	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
marie-galante4	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
morrll1	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
punctatum4	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					480
latifolium191	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
yucatum9	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
G.raimondii	TGTGGAGGACATGACAATCAAGGATTGGAAGAAGCTGGCAATGGGTTGGTCCAGATTGAATATGAACGGCTTCATGA					479
Conservation	<div></div>					
		500	520	540	560	
morrll18	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGTTCGTTGGAAGGAGTTTCCAAGAA					559
palmeri11	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTCGTTGGAAGGAGTTTCCAAGAA					559
palmeri8	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
palmeri5	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
Superokra_Dt	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
marie-galante6	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
richmondii4	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
latifolium98B	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
palmeri4	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
TM-1_Dt	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
marie-galante4	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
morrll1	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
punctatum4	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					560
latifolium191	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
yucatum9	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
G.raimondii	ACAATGGTAAAGGTAACAACAACAAGAAGAAGAGATTGACAAGTGATCAGTTAGATTTCGTTGGAAGGAGTTTCCAAGAA					559
Conservation	<div></div>					
		580	600	620	640	
morrll18	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
palmeri11	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
palmeri8	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
palmeri5	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
Superokra_Dt	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
marie-galante6	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
richmondii4	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
latifolium98B	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
palmeri4	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
TM-1_Dt	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
marie-galante4	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
morrll1	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
punctatum4	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					640
latifolium191	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
yucatum9	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
G.raimondii	GAGAATAAGTTGGATCCCGACAGGAAAAATGAAGCTTTCCAAGGAACTTGGACTTCAACCAAGACAAATCGCTGTCTGGTT					639
Conservation	<div></div>					

	660	680	700	720	
morrll18	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
palmeri11	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
palmeri8	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
palmeri5	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
Superokra_Dt	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
marie-galante6	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
richmondii4	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
latifolium98B	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
palmeri4	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
TM-1_Dt	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
marie-galante4	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
morrll1	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
punctatum4	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	720
latifolium191	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
yucatatum9	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
G.raimondii	CCAAAACAGGCGTGCTAGATG	GAAAGCTAAACAGCTTGA	AACTCATATAATACGCTTAA	ACATGAGTATGATATTATCT	719
Conservation
	740	760	780	800	
morrll18	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
palmeri11	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
palmeri8	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
palmeri5	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
Superokra_Dt	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
marie-galante6	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
richmondii4	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	798
latifolium98B	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
palmeri4	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
TM-1_Dt	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
marie-galante4	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
morrll1	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
punctatum4	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	800
latifolium191	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
yucatatum9	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
G.raimondii	CCATTGAAAAGCAAAGCTACA	AGACGAGGTTAGCCTCTATA	TTCAAAACCTCTATTTTTT	TATCCCTTCTCTCGCTCTC	799
Conservation
	820	840	860	880	
morrll18	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
palmeri11	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
palmeri8	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
palmeri5	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
Superokra_Dt	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
marie-galante6	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
richmondii4	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	878
latifolium98B	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
palmeri4	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
TM-1_Dt	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
marie-galante4	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
morrll1	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
punctatum4	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	880
latifolium191	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
yucatatum9	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
G.raimondii	TATATATCTTTACGTGGAAC	TGGTGCACGTGTTTTATAA	ACGAATATGCTTGTGTGCT	CTATTGTTGAGGTTTCCCCAT	879
Conservation
	900	920	940	960	
morrll18	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
palmeri11	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
palmeri8	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
palmeri5	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
Superokra_Dt	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
marie-galante6	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
richmondii4	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	956
latifolium98B	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
palmeri4	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
TM-1_Dt	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
marie-galante4	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
morrll1	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
punctatum4	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	958
latifolium191	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
yucatatum9	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
G.raimondii	TTTGTTTATACACACATACAT	ACATATAATATATAT - -	TTATACTGTACTCTCATAC	CTTTTACGCAGGTGATGGA	957
Conservation

		1,300	1,320	1,340	1,360	
morrl118	TTTGTGGAGAAATT	T	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1357		
palmer11	TTTGTGGAGAAATT	T	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1357		
palmer8	TTTGTGGAGAAATT	T	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1357		
palmer5	TTTGTGGAGAAATT	T	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1357		
Superokra_Dt	TTTGTGGAGAAATT	T	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1357		
marie-galante6	TTTGTGGAGAAATT	T	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1357		
richmondii4	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGA	1347		
latifolium98B	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1348		
palmer4	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1348		
TM-1_Dt	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1348		
marie-galante4	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1348		
morrl11	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1348		
punctatum4	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1357		
latifolium191	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1356		
yucatum9	TTT	A	TTGGAGAAATTGACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1356		
G.raimondii	TTTGTGGAGAAATT	G	ACGTAACATTAATGTATCAAAGGCTGCATCAGTGAGTATTGGATCAAAAGATTGAAGTACATTT	1358		
Conservation
		1,380				
morrl118	TGCTTGCTTACTACGGCTGAATAG	1381				
palmer11	TGCTTGCTTACTACGGCTGAATAG	1381				
palmer8	TGCTTGCTTACTACGGCTGAATAG	1381				
palmer5	TGCTTGCTTACTACGGCTGAATAG	1381				
Superokra_Dt	TGCTTGCTTACTACGGCTGAATAG	1381				
marie-galante6	TGCTTGCTTACTACGGCTGAATAG	1381				
richmondii4	TGCTTGCTTACTACGGCTGAATAG	1371				
latifolium98B	TGCTTGCTTACTACGGCTGAATAG	1372				
palmer4	TGCTTGCTTACTACGGCTGAATAG	1372				
TM-1_Dt	TGCTTGCTTACTACGGCTGAATAG	1372				
marie-galante4	TGCTTGCTTACTACGGCTGAATAG	1372				
morrl11	TGCTTGCTTACTACGGCTGAATAG	1372				
punctatum4	TGCTTGCTTACTACGGCTGAATAG	1381				
latifolium191	TGCTTGCTTACTACGGCTGAATAG	1380				
yucatum9	TGCTTGCTTACTACGGCTGAATAG	1380				
G.raimondii	TGCTTGCTTACTACGGCTGAATAG	1382				
Conservation

Table S1 Phenotypes and their origins of materials used in the present research.

Materials	Leaf Shape	Origin
TM-1	Normal	USDA-ARS. Col. Station, TX, USA
Hai7124	Sea-island	Commercial Sea island
T586	okra	USDA-ARS. Col. Station, TX, USA
IL-15-5-1	Sea-island	Nanjing Agricultural University, China
Super Okra	super okra	National Medium-term Gene Bank of Cotton in China
<i>G. aridum</i>	unlobed	Institute of Cotton Research of CAAS
<i>G. davidsonii</i>	unlobed	Nanjing Agricultural University, China
<i>G. raimondii</i>	unlobed	Nanjing Agricultural University, China
<i>G. thurberi</i>	deeply lobed	Nanjing Agricultural University, China
<i>G. trilobum</i>	deeply lobed	Institute of Cotton Research of CAAS
<i>G. gossypoides</i>	broad	Nanjing Agricultural University, China

Table S2 Sequence analysis of *GhOKRA* in 89 accessions from seven races of *G. hirsutum*.

Accession	Acc. number ^a	Sequence type
Latifolium 1	TX-2	3
Latifolium 15	TX-36	3
Latifolium 39	TX-69	3
Latifolium 45	TX-76	3
Latifolium 46	TX-77	3
Latifolium 48	TX-79	3
Latifolium 50	TX-81	3
Latifolium 53	TX-85	3
Latifolium 57	TX-89	3
Latifolium 58	TX-90	1
Latifolium 60	TX-93	3
Latifolium 63	TX-97	3
Latifolium 65	TX-100	3
Latifolium 67	TX-102	3
Latifolium 71	TX-106	3
Latifolium 73	TX-108	3
Latifolium 74	TX-109	3
Latifolium 77	TX-116	3
Latifolium 79	TX-118	3
Latifolium 80	TX-119	3
Latifolium 88	TX-143	3
Latifolium 92	TX-155	3
Latifolium 93	TX-156	3
Latifolium 95	TX-158	3
Latifolium 98B	TX-162	3
Latifolium 100	TX-164	3
Latifolium 105	TX-173	3
Latifolium 109	TX-177	3
Latifolium 161	TX-242	3
Latifolium 168	TX-252	3
Latifolium 178	TX-482	3
Latifolium 191	TX-16	2
Marie-Galante 2	TX-141	3
Marie-Galante 4	TX-367	3
Marie-Galante 5	TX-368	1
Marie-Galante 6	TX-370	1
Marie-Galante 7	TX-371	2
Marie-Galante 12	TX-817	1
Marie-Galante 19	TX-839	2
Marie-Galante 20	TX-840	1
Marie-Galante 25	TX-849	3
Marie-Galante 27	TX-853	1
Marie-Galante 30	TX-865	1
Marie-Galante 32	TX-867	1
Marie-Galante 56	TX-1139	1

Marie-Galante 57	TX-1141	3
Marie-Galante 58	TX-1142	2
Marie-Galante 60	TX-1144	3
Marie-Galante 62	TX-1191	1
Marie-Galante 64	TX-1203	1
Marie-Galante 66	TX-1205	3
Marie-Galante 71	TX-1297	2
Marie-Galante 73	TX-1606	3
Marie-Galante 116	TX-2304	3
Morrilli 1	TX-125	3
Morrilli 5	TX-130	3
Morrilli 10	TX-136	1
Morrilli 11	TX-137	3
Morrilli 18	TX-191	1
Morrilli 20	TX-194	3
Morrilli 25	TX-257	3
Morrilli 27	TX-260	3
Morrilli 32	TX-278	3
Morrilli 36	TX-284	3
Morrilli 37	TX-285	1
Morrilli 38	TX-286	3
Morrilli 39	TX-292	3
Paimerii 1	TX-1	2
Paimerii 4	TX-11	3
Paimerii 5	TX-51	1
Paimerii 6	TX-303	3
Paimerii 8	TX-1530	1
Paimerii 11	TX-2513	1
Punctatum 4	TX-28	2
Punctatum 5	TX-44	3
Punctatum 9	TX-115	3
Punctatum 14	TX-231	3
Punctatum 15	TX-251	3
Punctatum 20	TX-487	2
Punctatum 24	TX-997	2
Punctatum 31A	TX-1244	3
Richmondii 4	TX-19	3
Richmondii 6	TX-256	3
Richmondii 8	TX-461	3
Richmondii 10	TX-463	3
Richmondii 11	TX-1101	3
Yucatanense 1	TX-1039	2
Yucatanense 6A	TX-2096	2
Yucatanense 9	TX-2094	2
Latifolium 1	TX-2	3
Latifolium 15	TX-36	3
Latifolium 39	TX-69	3
Latifolium 45	TX-76	3

^a The number of the Texas race collected in the USA.

Table S3 Primers used in this study.

Primer name	Forward primer 5'-3'	Reverse primer 5'-3'	Purpose
H2035	AAGTGATATCGGTGGTACTTCT	GGTATCCCATGATTTATAGCC	fine mapping
H1960	CGTCGTATTTGTCGTACTTCT	CATTTATCGGTAGACTCATCG	fine mapping
H1933	GGAAAGAAAGAAAGGAAGGA	GCTTTTATTTCTGCTGCTCT	fine mapping
H1857	CTGGCCTTCTCATTTATCTG	CTAATCGTATCTGGTCAGCAA	fine mapping
H1851	CAATACAGGAAGACGTGAGTT	GTATCATGATGTCTTGGGTTC	fine mapping
H1851	CAATACAGGAAGACGTGAGTT	GTATCATGATGTCTTGGGTTC	fine mapping
H1792	GGGAAAAAGAGAGTGAATGAG	GCAGCGATAAATGTAAAAGC	fine mapping
H1776	TCACAATAACAAGGTGAGTG	CCGGTATATTCGAGTTTCATT	fine mapping
H1768	GGAAGTGTGTCATCCATT	GGTTGGAGAAGAATATGGTAAG	fine mapping
H1758	GACCGAGACATTGAAGTTAGA	GCCTTTTACGTGAAGAGTTC	fine mapping
JESPR152	GATGCACCAGATCCTTTTATTAG	GGTACATCGGAATCACAGTG	fine mapping
NAU3040	AAAAGCCGTGCTAACTGAAT	AATCACAAACACCCGAAACT	fine mapping
G1	ATTAACAGACAAGTCCACCCTC	CTATTCAGCCGTAGTAAGCAAG	genome sequence of <i>OKRA</i>
G2	ACTTTAGGCTCACTCTTTGGG	CAAATTGTACTGAAACGAGGA	cDNA sequence of <i>GhOK3</i>
S1	GGGGTACCATGGATTGGGATGGCACCATTG	TCCCCCGGGGGGATAAGAAGGGAGTTGAATAT	Subcellular localization
R1-1	GAATTGAAAGGAATGCTAGTGG	TGTTTTAGGGATAAGAAGGGAG	3'RACE of <i>GhOKRA</i>
R1-2	GGCACCATTGACCCCTTATTT	ATTGTCCATGTCCTCCACACCT	3'RACE of <i>GhOKRA</i>
V1	CGGAATTCCGAAGATCGCCGGAAACAACCA TTTGGAGAGAACACGGGGGACTCTAGAATGGA	GCTCTAGAAGTTCAAGCTCTTGATCCAATA TTGAACGATCGGGGAAATTCGAGCTCTTAGGG	VIGS- <i>GhOKRA</i> transformation of <i>A. thaliana</i>
T1	TTGGGATGGCACCATTTCGA TTTGGAGAGAACACGGGGGACTCTAGAATGGA	ATAAGAAGGGAGTTGAAT TTGAACGATCGGGGAAATTCGAGCTCCTACTTT	(<i>GhOKRA</i>) transformation of <i>A. thaliana</i>
T2	TTGGGATGGCACCATTTCGA TTTGGAGAGAACACGGGGGACTCTAGAATGGA	CGATGATCTCTCCGCCG TTGAACGATCGGGGAAATTCGAGCTCCTATCGG	(<i>GhOKRA</i> ^M) transformation of <i>A. thaliana</i>
T3	TTGGGATGGCACCATTTCGA	ATATAGTCGTTTCC	(<i>GrOKRA</i> ^M)
293/294	GAGGCTATTCGGCTATGACTG	TAGAAGGCGATGCGCTGCGA	NPTII

Table S4. Segregation of different leaf shapes in mapping populations.

Mapping populations	Total	Okra leaf	Normal leaf	Intermediate leaf	Chi-square test
(IL-15-5-1×TM-1)F ₂	770	176	202	392	2.01
(IL-15-5-1×TM-1)F ₂	508	112	115	281	5.77
Total	1278	288	317	673	

χ^2 test for goodness-of-fit at 0.05 significance level ($\chi^2_{0.05,2}=5.99$)

Table S5 Annotated gene sequences in the genomic region of *G. raimondii* Chr. 02 delineated by markers closely linked to the leaf shape gene and their homology with *Arabidopsis thaliana* genes.

ORF	<i>G. raimondii</i> gene ID	Physical location	<i>Arabidopsis</i> gene ID	Protein homolog	<i>Arabidopsis thaliana</i> gene name
ORF1	Gorai.002G242400.1	60675284-60676284	AT3G44735.1	ATPSK3,PSK1,PSK3	PHYTOSULFOKINE 3 PRECURSOR
ORF2	Gorai.002G242500.1	60679931-60680949	AT3G44735.1	ATPSK3,PSK1,PSK3	PHYTOSULFOKINE 3 PRECURSOR
ORF3	Gorai.002G242600.1	60681030-60684351	AT3G23620.1		Ribosomal RNA processing Brix domain protein
ORF4	Gorai.002G242700.1	60684967-60690830	AT3G52940.1	ELL1,FK,HYD2	Ergosterol biosynthesis ERG4/ERG24 family
ORF5	Gorai.002G242800.1	60691443-60699456	AT5G03740.1	HD2C,HDT3	histone deacetylase 2C
ORF6	Gorai.002G242900.1	60700190-60718341	AT3G52940.2	ELL1,FK,HYD2	Ergosterol biosynthesis ERG4/ERG24 family
ORF7	Gorai.002G243000.1	60721265-60724366	AT5G03740.1	HD2C,HDT3	histone deacetylase 2C
ORF8	Gorai.002G243100.1	60725536-60729527	AT2G36530.1	ENO2,LOS2	Enolase
ORF9	Gorai.002G243200.1	60733892-60734923	AT3G52960.1		Thioredoxin superfamily protein
ORF10	Gorai.002G243300.1	60734924-60737571	AT5G12100.1		pentatricopeptide (PPR) repeat-containing protein
ORF11	Gorai.002G243400.1	60739751-60745494	AT5G03760.1	ATCSLA09,ATCSLA9 CSLA09,CSLA9,RAT4	Nucleotide-diphospho-sugar transferases superfamily protein
ORF12	Gorai.002G243500.1	60763514-60767171	AT3G23160.1		Protein of unknown function (DUF668)
ORF13	Gorai.002G243600.1	60775425-60777957	AT2G36570.1		Leucine-rich repeat protein kinase family protein
ORF14	Gorai.002G243700.1	60784258-60789349	AT3G52990.1		Pyruvate kinase family protein
ORF15	Gorai.002G243800.1	60793342-60798916	AT3G52990.1		Pyruvate kinase family protein
ORF16	Gorai.002G243900.1	60802515-60804472	AT3G10040.1		sequence-specific DNA binding transcription factors
ORF17	Gorai.002G244000.1	60816695-60817565	AT5G03790.1	ATHB51,HB51,LMI1	homeobox 51
ORF18	Gorai.002G244100.1	60818675-60821101	AT5G60900.1	RLK1	receptor-like protein kinase 1