CASE REPORT

Coping with a life event in bipolar disorder: ambulatory measurement, signalling and early treatment

Stefan E Knapen,¹ Rixt F Riemersma-van der Lek,² Bartholomeus C M Haarman,² Robert A Schoevers¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bcr-2016-216123>)

¹Department of Psychiatry, University of Groningen, University Medical Center Groningen, Research School of Behavioural and Cognitive Neurosciences (BCN), Interdisciplinary Center for Psychopathology and Emotion regulation (ICPE), Groningen, The Netherlands

²Department of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

Correspondence to Stefan Erik Knapen, stefanknapen@gmail.com

Accepted 27 September 2016

SUMMARY

Disruption of the biological rhythm in patients with bipolar disorder is a known risk factor for a switch in mood. This case study describes how modern techniques using ambulatory assessment of sleep parameters can help in signalling a mood switch and start early treatment. We studied a 40-year-old woman with bipolar disorder experiencing a life event while wearing an actigraph to measure sleep-wake parameters. The night after the life event the woman had sleep later and shorter sleep duration. Adequate response of both the woman and the treating psychiatrist resulted in two normal nights with the use of 1 mg lorazepam, possibly preventing further mood disturbances. Ambulatory assessment of the biological rhythm can function as an add-on to regular signalling plans for prevention of episodes in patients with bipolar disorder. More research should be conducted to validate clinical applicability, proper protocols and to understand underlying mechanisms.

BACKGROUND

Modern techniques such as ambulatory assessment of physiological parameters may be helpful to monitor and signal potential elicitors of new episodes in patients with a mental disorder. In bipolar disorder disruptions of the biorhythm such as a change in sleep are known risk factors for a switch in mood.¹ We describe a case where ambulatory measurement could have functioned as a signal for early treatment.

CASE PRESENTATION

A 40-year-old woman is in treatment for bipolar disorder type I since 8 years. The bipolar disorder was diagnosed by a psychiatrist from the department of psychiatry of the University Medical Center Groningen according to the Diagnostic and Statistical Manual of Mental Disorders Fourth Edition criteria. Since diagnosis and treatment she was admitted once to the psychiatric department for a depressive episode. She visited her treating psychiatrist around every 6 weeks. Furthermore she experienced subclinical mood swings once or twice a year, particularly after stressful life events. These subclinical mood swings persisted after different psychological treatments (cognitive-behavioural therapy, interpersonal psychotherapy and individual conversations). She was treated with lithium, 600 mg a day, with a blood level of 0.56 mmol/L,

after recent dosage lowering because of a developing renal insufficiency, a known side effect of lithium treatment. As the renal insufficiency continued after dosage decrease of lithium the patient switched to lamotrigine as mood stabiliser. This medication change happened after the presented events.

The patient participated in the study 'Circadian timing systems in bipolar disorder' in the Netherlands from 15 July 2014 to 22 July 2014.² The aim of the study was to identify variations in circadian rest-activity rhythms in-vivo. As part of the study she wore an actigraph (the Actiwatch 2, Philips Respironics, validated for sleep analysis), a small wristband that can measure activity and light and calculates sleep timing, duration and quality of sleep.^{3 4} During the study the patient had no insight in her own actigraphy data. During this period she experienced a serious life event, the loss of a friend by an accident.

The news of the loss came to her in the afternoon. The following night she immediately had a change in her sleep pattern, went to bed ~2 hours later than usual and slept around 1.5 hours less than usual.

TREATMENT

The day after the life event, she contacted her psychiatrist to discuss the event and its emotional impact, because she knew from earlier experience that this might destabilise her. The psychiatrist prescribed 1 mg lorazepam for two consecutive nights, knowing that sleep loss can provoke a mood episode. Using lorazepam restored a stable rhythm, with regular in bed and out of bed times and having a night of about 7.5 hours.

OUTCOME AND FOLLOW-UP

After two nights of lorazepam use, she slept at her regular time, going to bed at 23:14 and waking up at 8:10, without the use of lorazepam. Although she was sad, her mood remained euthymic afterwards.

See [figure 1](#) for a visualisation of the rest-activity rhythm in an actogram and online supplementary information for average sleep time.

DISCUSSION

A major goal of bipolar disorder treatment is the prevention of manic or depressive episodes by early

CrossMark

To cite: Knapen SE, Riemersma-van der Lek RF, Haarman BCM, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-216123

Figure 1 Actogram of 8 days. Black bars represent activity per minute. The dark blue overlay indicates sleep time, while light blue overlay shows time in bed, without being asleep. The red blocks under the activity represent time awake. Red blocks during a sleep period indicate moments of being awake. Single asterisk marks means a life event, and double asterisks mark the intake of 1 mg lorazepam. On the third day there was a life event, coped with two nights of lorazepam use.

signalling and treatment.⁵ In this patient, an episode was probably prevented by adequate signalling of an increased risk and the use of a benzodiazepine. Two factors were important to achieve this, the insight of the patient into her disease and the adequate use of medication to maintain a regular sleeping pattern. Although there were no formal psychotherapeutic interventions from the psychiatrist, getting a brief supportive reaction from her psychiatrist could have been helpful as well. A strong, direct effect on sleep with the same strength as one might expect from lorazepam is unlikely.

As part of the study, sleep and activity were also measured by an actiwatch during this period. Apart from recognising the

potential risk of this emotional event herself, the objective measure of sleep could also have been helpful to alert the patient. A monitoring system could inform both the patient and treating health personnel. For many years, the use of early alarm-systems, such as a signalling plan, has been advocated and tested, but modern technology ('wearable technology') now makes objective measurements very accessible to patients and practitioners.^{6,7}

It is clear that there is an interplay between sleep and the onset of mood episodes in bipolar disorder. The onset of mania is often preceded by changes in sleep patterns, and the antidepressant effect of sleep deprivation on relieving

depression suggests a similar mechanism.⁸ With this in mind, continuous sleep measurement in patients with bipolar disorder could help to prevent full-blown episodes by early signalling of changes in these patterns. Future studies are needed to see if this would outperform patients' own early recognition of an upcoming mood episode and be helpful to prevent new episodes. Current commercially available actigraphs may show the result of sleep-wake measurement on a person's smartphone, but lack the clinical validity needed to apply them to patients.⁹ Further research is therefore needed, in the form of longitudinal studies to determine what early changes in physiological patterns precede new episodes, and in the form of randomised controlled trials examining the benefit of feedback of such changes to patients and clinicians in order to intervene early.

Learning points

- ▶ Switches in mood in patients with bipolar disorder might be triggered with changes in sleep-wake parameters.
- ▶ Changes in sleep-wake parameters can be monitored by ambulatory assessment methods such as actigraphy.
- ▶ Monitoring sleep-wake parameters can be helpful in preventing mood switches in patients with bipolar disorder.

Contributors SEK and RFRL analysed the case and wrote the initial paper. BCMH and RAS contributed to the paper. All authors read and approved the final manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Jackson A, Cavanagh J, Scott J. A systematic review of manic and depressive prodromes. *J Affect Disord* 2003;74:209–17.
- 2 Verkooijen S, van Bergen AH, Knapen SE, *et al*. An actigraphy study investigating sleep in bipolar I patients, unaffected siblings and controls. *J Affect Disord* 2016.
- 3 Kaplan KA, Talbot LS, Gruber J, *et al*. Evaluating sleep in bipolar disorder: comparison between actigraphy, polysomnography, and sleep diary. *Bipolar Disord* 2012;14:870–9.
- 4 Kushida CA, Chang A, Gadkary C, *et al*. Comparison of actigraphic, polysomnographic, and subjective assessment of sleep parameters in sleep-disordered patients. *Sleep Med* 2001;2:389–96.
- 5 National Collaborating Centre for Mental Health. Bipolar Disorder: The Assessment and Management of Bipolar Disorder in Adults, Children and Young People in Primary and Secondary Care. Leicester UK: The British Psychological Society (2014).
- 6 Perry A, Tarrier N, Morriss R, *et al*. Randomised controlled trial of efficacy of teaching patients with bipolar disorder to identify early symptoms of relapse and obtain treatment. *BMJ* 1999;318:149–53.
- 7 Bos FM, Schoevers RA, aan het Rot M. Experience sampling and ecological momentary assessment studies in psychopharmacology: a systematic review. *Eur Neuropsychopharmacol* 2015;25:1853–64.
- 8 Wehr T, Wirz-Justice A, Goodwin F. Phase advance of the circadian sleep-wake cycle as an antidepressant. *Science* 1979;206:710–13.
- 9 Kooiman TJM, Dontje ML, Sprenger SR, *et al*. Reliability and validity of ten consumer activity trackers. *BMC Sports Sci Med Rehabil* 2015;7:24.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow