Published in final edited form as:

Proteins. 2016 September ; 84(9): 1333. doi:10.1002/prot.25098.

Structure alignment of membrane proteins: Accuracy of available tools and a consensus strategy

Marcus Stamm¹ and Lucy R. Forrest^{1,2,*}

¹Computational Structural Biology Group, Max Planck Institute of Biophysics, Frankfurt Am Main, Germany ²Computational Structural Biology Section, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Maryland

The following sentence on page 1722 is incorrect: Using this clustering approach, we identified 159 α -helical protein chains in 37 families corresponding to 354 protein pairs (Supporting Information Table S1), plus 68 β -barrels in 8 families corresponding to 319 alignments (Supporting Information Table S2), for a total of 673 alignments.

The correct sentence should read: Using this clustering approach, we identified 152 α -helical protein chains in 40 families corresponding to 354 protein pairs (Supporting Information Table S1), plus 68 β -barrels in 8 families corresponding to 319 alignments (Supporting Information Table S2), for a total of 673 alignments.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

*Correspondence to: Lucy R. Forrest, Computational Structural Biology Section, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, USA. lucy.forrest@nih.gov.