Published in final edited form as:

Bio Protoc. ; 5(14): . doi:10.21769/BioProtoc.1530.

Large-scale Phenotypic Profiling of Gene Deletion Mutants in *Candida glabrata*

Fabian Istel[#], Tobias Schwarzmüller[#], Michael Tscherner, and Karl Kuchler^{*}

¹Medical University Vienna, Max F. Perutz Laboratories, Department of Medical Biochemistry, Vienna, Austria

[#] These authors contributed equally to this work.

Abstract

Here, we describe a method enabling the phenotypic profiling of genome-scale deletion collections of fungal mutants to detect phenotypes for various stress conditions. These stress conditions include among many others antifungal drug susceptibility, temperature-induced and osmotic as well as heavy metal or oxidative stress. The protocol was extensively used to phenotype a collection of gene deletion mutants in the human fungal pathogen *Candida glabrata* (*C. glabrata*) (Schwarzmüller *et al.*, 2014).

Materials and Reagents

1. *C. glabrata* strain (ATCC2001, clinical isolates)
2. Sterile water (double distilled)
3. Bacto™ peptone (BD Biosciences, catalog number: 211820)
4. Bacto™ yeast extract (BD Biosciences, catalog number: 212720)
5. Bacto™ agar (BD Biosciences, catalog number: 214030)
6. Difco™ yeast nitrogen base (YNB) (BD Biosciences, catalog number: 233520)
7. Glucose (Merck KGaA, catalog number: 108337)
8. Adenine (Sigma-Aldrich, catalog number: A8626)
9. L-arginine (Sigma-Aldrich, catalog number: A5006)
10. L-tyrosine (Sigma-Aldrich, catalog number: T3754)
11. L-isoleucine (Sigma-Aldrich, catalog number: I2752)
12. L-phenylalanine (Sigma-Aldrich, catalog number: P2126)
13. L-glutamic acid (Sigma-Aldrich, catalog number: G1251)
14. L-aspartic acid (Sigma-Aldrich, catalog number: A9256)

^{*}For correspondence: karl.kuchler@meduniwien.ac.at.

15. L-threonine (Sigma-Aldrich, catalog number: T8625)
16. L-serine (Sigma-Aldrich, catalog number: S4500)
17. L-valine (Sigma-Aldrich, catalog number: V0500)
18. L-methionine (Sigma-Aldrich, catalog number: M9625)
19. Glycerol (Sigma-Aldrich, catalog number: G5516)
20. YPD media (see Recipes)
21. Solid YPD media (see Recipes)
22. 2x YPD media (see Recipes)
23. SC media (see Recipes)
24. Solid SC media (see Recipes)
25. Amino acid mix (see Recipes)
26. 15% glycerol solution (see Recipes)

Equipment

1. Square plates (PlusPlate) (40 ml/plate) (Singer Instruments, catalog number: PLU-001)
2. Pin Pad (RePad 96 Long) (Singer Instruments, catalog number: REP-001)
3. 96-well plates (200 µl/well) (Starlab, catalog number: CC76827596)
4. Petri dishes (92 mm) (SARSTEDT AG, catalog number: 82.1473)
5. Culture tubes (Starlab, catalog number: I14850810)
6. Yeast replica robot (Singer Instruments, catalog number: RoToR HDA)
7. Plate reader (Victor³V) (PerkinElmer, catalog number: 1420040)
8. 48-well replica stamp (V&P Scientific, catalog number: VP 408H)
9. 96-well replica stamp (V&P Scientific, catalog number: VP 407)
10. Library copier (guide for stamps) (V&P Scientific, model: VP 381)
11. Plate mixer (Eppendorf, catalog number: 5353000014)
12. Rotary shaker for culture flasks (innova44) (New Brunswick Scientific, catalog number: M1282-0002)
13. Incubator (Heraeus Instruments, catalog number: B6120)

Procedure

- A. Arraying and storage

1. For arraying deletion strains in the 96-well format see the Note section. Arrayed strains can be suspended in 15% glycerol and stored frozen at minus 70 °C in 96-well plates.
 2. Frozen stocks can be either handled by manual replicating tools or by an automated system such as the RoToR HDA by Singer Instruments. The latter offers a higher reproducibility and reduces the risk of contamination when compared to manual systems. Therefore, this protocol for large-scale screening was optimized for the work with the Singer HDA RoToR replica-plating robot.
- B.** Initial large-scale screening (on solid YPD media)
1. For initial pilot screenings, prepare square plates containing the screening compounds and control plates lacking compounds as shown in Figure 1. The concentration of the compound for the solid screening plate is usually twice the concentration of the IC₅₀. However, the exact concentration should be determined prior to the large-scale screening by plating control strains.
 - a. To exclude replica-plating errors, at least two plates should be prepared for each condition (stress-inducing compound, stress-inducing media, temperature, *etc.*).
 - b. Source plates can be either liquid media (96-well microtiter plates) or solid media (square plates).
 2. Dry plates on a clean bench for 10 min, for light-sensitive compounds turn off lights.
 - a. Make sure that no droplets of condensed water are on the plates or lids before using them.
 - b. Label plates before spotting strains.
When using a Singer robot use the following settings in the ROTOR HDA software:
 - c. For solid source plates activate the “offset” function, to avoid pins taking cells from exactly the same spot.
 - d. If liquid source plates are used, enable the “wet mix” option in the software menu

(diameter: 1 mm; speed: 3 mm/sec; cycles: 1; Travel: 2D). Alternatively, liquid source plates can be shaken on a MixMate for 15 sec at 750 rpm.

- e. If the so-called RePads are used for more than one target plate (“Recycle Mode”), the “Revisit Source” option should be enabled as well.

3. After plating, turn plates upside down as soon as spots have dried. In case of solid source plates, the target plates can be turned right away. Incubate plates for 72 h at 30 °C without shaking and inspect colony growth. Pictures should be taken of every plate at 24, 48 and 72 h. Visually compare the growth of each mutant strains and controls under stress and without stress with the growth of wild type strain(s). Important points that should be considered are shown in Figures 2 and 3.

C. Serial dilution spot assays (solid SC media)

1. For refined screening, prepare at least two SC media plates for each condition. Prior to their use, dry them on a clean bench. Grow overnight cultures of sensitive and wild type strains from the 96-well screening in SC medium at 30 °C with shaking at 200 rpm (200 rpm are used for growth in culture tubes; a total amount of 5 ml for each culture is sufficient). Dilute cultures into fresh SC medium resulting in an OD₆₀₀ of ≈ 0.2 and regrow them to an OD₆₀₀ of ≈ 1.0 . Prepare 1:10 serial dilutions of deletions strains in sterile water in a 96-well plate, starting with an OD₆₀₀ of 0.1.
2. Spotting uses a 48-well metal pin replicator, also referred to as “frogger” or “hopper”.
 - a. Sterilize the tool by submerging the pins in EtOH and brief flaming over a bunsen burner. Repeat this process three times and let the pins cool down.
 - b. Stir the serial dilution in the 96-well plate using the replicator and retract the pins from the wells without touching the wall of the wells.
 - c. A droplet of cell suspension will remain on each tip of the replicator.

- Europe PMC Funders Author Manuscripts

Europe PMC Funders Author Manuscripts

- Europe PMC Funders Author Manuscripts

Europe PMC Funders Author Manuscripts

- Europe PMC Funders Author Manuscripts

2. To validate the link between a genotype and a phenotype more experiments are necessary. These experiments include the complementation of deletion strains and the deletion of a single gene in different backgrounds.

Notes

1. Plate layouts

When designing the layout several aspects should be considered.

Outer spots grow faster.

Outer spots are framed red in Figure 2 and show a faster growth when compared to spots in inner rows and columns.

This phenomenon may hamper comparison of growth phenotypes of deletion strains to the growth of a wild type strain. In the worst case, this may lead to missing relevant phenotype. This problem can be solved by filling all outer rows / columns with a dummy strain (*e.g.* the wild type). Therefore, all strains that are to be screened are surrounded on all sides and experience the same growth conditions. Of note, this procedure yields only 60 usable spots. Alternatively, a 96-well plate can be spotted in quadruplicate for each deletion in the 384-well format using the Singer HDA RoToR. This method will result in each deletion strain being spotted 4 times on the agar plate as illustrated in Figure 3.

Include empty spots for controls

To compare results, at least the background strain should be spotted on each plate. Since this strain is usually derived from a wild type strain, the wild type should also be included. Moreover, it is very useful to include empty spots on each plate and to add additional positive and negative control strains for a specific screening. For instance, it can be very helpful to temporarily add a known hypersensitive strain for a specific screening as positive control [for an azole screening a *pdr1* (CAGL0A00451g) strain is a great control].
2. Liquid source plates are preferable to solid source plates, since the reproducibility is higher when using liquid sources plates. We experienced that the amount of cells can be inconsistent when using solid source plates. Liquid source plates can be shaken prior to plating, resulting in a very homogenous source cell suspension.
3. There are various standardized protocols available for the microdilution assays to quantify drug susceptibilities including those of the Clinical and Laboratory Standards Institute (Fothergill *et al.*, 2011) or the European Committee on Antimicrobial Susceptibility Testing (Arendrup *et al.*, 2012).

4. A video of the Singer RoToR in use is available at: <https://www.youtube.com/watch?v=1JabqneqXPE>.
5. Stocks for water insoluble compounds can be prepared in the appropriate solvent (*e.g.* DMSO). However, the dilution should be done in water. Otherwise, the wells would contain up to 50% of solvent.

Recipes

-
1. YPD media (yeast extract peptone dextrose)

25 g/L	Bacto™ peptone
12.5 g/L	Bacto™ yeast extract
2%	glucose
 2. Solid YPD media

25 g/L	Bacto™ peptone
12.5 g/L	Bacto™ yeast extract
2%	glucose
2%	agarose
 3. 2x YPD media

50 g/L	Bacto™ peptone
25 g/L	Bacto™ yeast extract
4%	glucose
 4. SC media (synthetic complete)

3.4 g/L	Difco™ YNB
2.9 g/L	amino acid mix
2 %	glucose
 5. Solid SC media

3.4 g/L	Difco™ YNB
2.9 g/L	amino acid mix
2 %	glucose
2 %	agarose
 6. Amino acid mix (g per 11.7 g total mix)

0.4 g	adenine
0.2 g	L-arginine
0.3 g	L-tyrosine
0.3 g	L-isoleucine
0.5 g	L-phenylalanine
1.0 g	L-glutamic acid
1.0 g	L-aspartic acid
2.0 g	L-threonine
4.0 g	L-serine
1.5 g	L-valine
0.5 g	L-methionine
 7. 15% glycerol solution

150 ml/L glycerol

850 ml/L water

Acknowledgements

This work was supported by the Austrian Science Foundation FWF through the ERA-Net Pathogenomics project FunPath (FWF-API-0125), and in part by grants from the Christian Doppler Society, the FP7 EC project FUNGITECT, the Marie-Curie ITN ImResFun (MC-ITN-606786) and the FWF Project FWF-P25333 „Chromatin“ to KK.

References

1. Arendrup MC, Cuenca-Estrella M, Lass-Flörl C, Hope W, Eucast A. EUCAST technical note on the EUCAST definitive document EDef 7.2: method for the determination of broth dilution minimum inhibitory concentrations of antifungal agents for yeasts EDef 7.2 (EUCAST-AFST). Clin Microbiol Infect. 2012; 18(7):E246–247. [PubMed: 22563750]
2. Fothergill AW. Antifungal susceptibility testing: clinical laboratory and standards institute (CLSI) methods. Interactions of Yeasts, Moulds, and Antifungal Agents. 2011:65–74.
3. Schwarzmüller T, Ma B, Hiller E, Istel F, Tscherner M, Brunke S, Ames L, Firon A, Green B, Cabral V, Marcet-Houben M, et al. Systematic phenotyping of a large-scale *Candida glabrata* deletion collection reveals novel antifungal tolerance genes. PLoS Pathog. 2014; 10(6):e1004211. [PubMed: 24945925]

Figure 1. Screening process-from large scale to single strains.

Putative hits (purple, blue, pink) are identified by comparing growth of strains on a control plate (YPD only) with a plate supplemented with screening compounds. In addition, deletion strains must be compared to the corresponding wild types (first two rows). These strains are then used for a refined screening in liquid assays through serial dilution on solid media to confirm primary phenotypes.

Figure 2. Faster growth of outer spots

Figure 3. 96-well to 384-well format.

The 384-well format (right panel) offers two distinct advantages when compared to the 96-well format (left panel). Only the outer spots grow faster, so every original spot has at least one spot now that grows normally. In addition, it can be easier to identify growth phenotypes due to the uniform appearance of growth patterns.

Serial dilution of cells**Table 1**

OD₆₀₀	Dilution	Cells/ml
0.1	1:1	3×10^6
0.01	1:10	3×10^5
0.001	1:100	3×10^4
0.0001	1:1,000	3×10^3