Published in final edited form as:

Acad Pediatr. 2016 ; 16(8): 734–741. doi:10.1016/j.acap.2016.04.012.

Effect of Medication Label Units of Measure on Parent Choice of Dosing Tool: A Randomized Experiment

H. Shonna Yin, MD, MS^{a,b}, Ruth M. Parker, MD^c, Lee M. Sanders, MD, MPH^d, Benard P. Dreyer, MD^a, Alan Mendelsohn, MD^a, Stacy Bailey, PhD, MPH^e, Deesha A. Patel, MS^f, Jessica J. Jimenez, BA^a, Kwang-Youn A. Kim, PhD^g, Kara Jacobson, MPH^h, Laurie Hedlund, MA^f, Rosa Landa, BA^d, Leslie Maness, BA^a, Purvi Tailor Raythatha, BSc^a, Terri McFadden, MDⁱ, and Michael S. Wolf, PhD, MPH^f

H. Shonna Yin: yinh02@med.nyu.edu; Ruth M. Parker: rpark01@emory.edu; Lee M. Sanders: leesanders@stanford.edu; Benard P. Dreyer: benard.dreyer@nyumc.org; Alan Mendelsohn: alm5@nyu.edu; Stacy Bailey: scbailey@unc.edu; Deesha A. Patel: deesha.patel@northwestern.edu; Jessica J. Jimenez: jessica.jimenez@nyumc.org; Kwang-Youn A. Kim: kykim@northwestern.edu; Kara Jacobson: kljacob@emory.edu; Laurie Hedlund: l-hedlund@northwestern.edu; Rosa Landa: rlanda@stanford.edu; Leslie Maness: ldmaness@bu.edu; Purvi Tailor Raythatha: purvi.tailor@nyumc.org; Terri McFadden: tmcfadd@emory.edu; Michael S. Wolf: mswolf@northwestern.edu

^aPediatrics, NYU School of Medicine - Bellevue Hospital, 550 First Avenue, New York, NY, 10016, United States

^bPopulation Health, NYU School of Medicine, 550 First Avenue, New York, NY, 10016, United States

^cMedicine, Emory University School of Medicine, 1518 Clifton Road, Atlanta, GA, 30322, United States

Corresponding author: H4. Shonna Yin, MD, MS, New York University School of Medicine, Department of Pediatrics, 550 First Avenue, NBV 8S4-11, New York, NY 10016; Telephone: (646) 501-4284, Fax: (212) 263-8172; yinh02@med.nyu.edu.

Conflict of Interest: The authors have no conflicts of interest to report.

Preliminary findings from this study were presented in part at the 2015 Pediatric Academic Societies meeting (April 25, 2015 in San Diego, CA), as well as at two meetings of the CDC Prevention of Overdoses & Treatment Errors in Children Taskforce (PROTECT) initiative (November 13, 2014 and November 4, 2015 in Atlanta, GA).

Financial Disclosure: Drs. Bailey, Parker, and Wolf, and Ms. Jacobson, have served as a consultant to, and received grant funding, from Merck, Sharp and Dohme for work unrelated to this study.

CONTRIBUTOR STATEMENT

Dr. Yin and Ms. Patel had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. The authors made contributions as delineated below.

Study concept and design: Yin, Wolf, Parker, Sanders, Dreyer, Mendelsohn, Bailey, Patel, Kim.

Acquisition, analysis, or interpretation of data: Yin, Wolf, Parker, Sanders, Dreyer, Mendelsohn, Bailey, Patel, Jimenez, Kim, Jacobson, Landa, Maness, Tailor Raythatha, Hedlund, McFadden.

Critical revision of the manuscript for important intellectual content: Yin, Wolf, Parker, Sanders, Dreyer, Mendelsohn, Bailey, Patel, Jimenez, Kim, Jacobson, Landa, Maness, Tailor Raythatha, Hedlund, McFadden.

Statistical analysis: Patel, Kim, Yin, Wolf, Dreyer, Mendelsohn.

Obtained funding: Yin, Wolf, Parker, Sanders, Dreyer, Mendelsohn, Bailey, Kim.

Administrative, technical, or material support: Yin, Patel, Jimenez, Kim, Jacobson, Landa, Maness, Tailor Raythatha, Hedlund, McFadden.

Study supervision: Yin, Wolf, Parker, Sanders, Dreyer, Mendelsohn, Bailey, Jacobson, Patel. All authors have approved the final manuscript as submitted.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

^dPediatrics, Stanford University School of Medicine, 117 Encina Commons, Stanford, CA, CA, 94305, United States

^ePharmaceutical Outcomes and Policy, UNC Eshelman School of Pharmacy, 2204 Kerr Hall, Chapel Hill, NC, 27599, United States

^fDivision of General Internal Medicine and Geriatrics, Northwestern University Feinberg School of Medicine, 750 N. Lake Shore Drive, Chicago, IL, 60611, United States

^gPreventive Medicine, Northwestern University Feinberg School of Medicine, 750 N. Lake Shore Drive, Chicago, IL, 60611, United States

^hRollins School of Public Health Emory University, 1518 Clifton Road, Atlanta, GA, 30322, United States

ⁱPediatrics, Emory University School of Medicine, 1518 Clifton Road, Atlanta, GA, USA, 30322

Abstract

OBJECTIVE—Some experts recommend eliminating “teaspoon” and “tablespoon” terms from pediatric medication dosing instructions, as they may inadvertently encourage use of nonstandard tools (i.e. kitchen spoons), which are associated with dosing errors. We examined whether use of “teaspoon” or “tsp” on prescription labels affects parents’ choice of dosing tools, and the role of health literacy and language.

METHODS—Analysis of data collected as part of a controlled experiment (SAFE Rx for Kids study), which randomized English/Spanish-speaking parents (n=2110) of children 8 years old to 1 of 5 groups which varied in unit of measurement pairings on medication labels/dosing tools. Outcome assessed was parent self-reported choice of dosing tool. Parent health literacy measured using the Newest Vital Sign.

RESULTS—77.0% had limited health literacy (36.0% low, 41.0% marginal); 35.0% completed assessments in Spanish. Overall, 27.7% who viewed labels containing either “tsp” or “teaspoon” units (alone or with “mL”) chose nonstandard dosing tools (i.e. kitchen teaspoon, kitchen tablespoon), compared to 8.3% who viewed “mL”-only labels (AOR=4.4[95% CI: 3.3–5.8]). Odds varied based on whether “teaspoon” was spelled out or abbreviated (“teaspoon”-alone: AOR=5.3[3.8–7.3]); “teaspoon” with mL: AOR=4.7[3.3–6.5]; “tsp” with mL (AOR=3.3[2.4–4.7]); p<0.001)). Similar findings were noted across health literacy and language groups.

CONCLUSIONS—Use of teaspoon units (“teaspoon” or “tsp”) on prescription labels is associated with increased likelihood of parent choice of nonstandard dosing tools. Future studies may be helpful to examine the real-world impact of eliminating teaspoon units from medication labels, and identify additional strategies to promote the safe use of pediatric liquid medications.

Keywords

medication errors; dosing errors; health literacy; ambulatory care; health communication; injury prevention

INTRODUCTION

Standardizing labels on pediatric liquid medications has been proposed as a way to reduce dosing errors in the United States (US).^{1–7} The Centers for Disease Control and Prevention (CDC), through its PROTECT initiative, along with the American Academy of Pediatrics (AAP), have recently begun to advocate for “milliliter”- or “mL”-only dosing, and the elimination of “teaspoon” and “tablespoon” terms on medication labels.^{8–10} While one study showed that parents are able to dose more accurately using mL instructions, a significant proportion of parents continued to make errors even with mL dosing.¹⁰ Concerns about moving to the metric-only labeling of medications and dosing tools in the US, which has historically relied on a non-metric spoon-based system^{8,9,11}, have led some to call for additional research to better delineate the benefits of a move to an mL-exclusive system.

In this context, some medical providers and public health professionals have suggested that “teaspoon” and “tablespoon” labels may inadvertently cause dosing errors by encouraging parent use of nonstandard dosing tools to measure medications such as kitchen spoons, which vary widely in size and shape.^{10,12–14} There is longstanding recognition that kitchen spoons should not be used for dosing pediatric medications.^{12,15,16} In fact, since 1975, the AAP has recommended the use of standard dosing tools such as oral syringes, droppers, and dosing cups, which have markings to guide parents in the accurate dosing of medications.¹² The US Food and Drug Administration (FDA) recommends inclusion of standard tools with over-the-counter liquid medications.³

In this study, we sought to examine the degree to which parents’ choice of medication dosing tools is affected by the unit of measurement present on a medication label, in particular the impact of teaspoon terms (“teaspoon”, “tsp”). We hypothesized that presence of a teaspoon unit on the label, whether spelled out or abbreviated, would increase the likelihood of a parent inappropriately choosing a nonstandard tool (i.e. kitchen spoon) to administer medications. We also sought to examine differences in impact by parent health literacy and language, as those with low health literacy and limited English proficiency have been previously found to be at greater risk for misunderstanding drug labels and subsequent medication errors.^{10,13,17–20}

METHODS

Participants, Recruitment, Randomization

Data for this study were collected as part of a randomized controlled experiment to examine the degree to which specific attributes of medication labels and dosing tools affect parent dosing errors (SAFE Rx for Kids study). Subjects were enrolled from pediatric outpatient clinics affiliated with New York University (Bellevue Hospital Center), Stanford University (Gardner Packard Children’s Health Care Center), and Emory University (Children’s Hospital of Atlanta (CHOA) at Hughes Spalding) Schools of Medicine. These clinics serve predominantly low income families (at least 70% enrolled in Medicaid or the Women Infants and Children (WIC) supplemental nutrition program). Northwestern University served as the data management site; no subjects were enrolled at this site. Institutional review board approval was obtained from each partner site (Emory, New York University, Northwestern,

Stanford University Schools of Medicine) and CHOA, as well as the Research Review Committee of Bellevue Hospital Center.

Parents/legal guardians were consecutively approached by trained research assistants to determine eligibility. Those who met inclusion criteria were English or Spanish-speaking parents or legal guardians (18 years of age) who had a child 8 years of age that was presenting for care in the pediatric clinic. Parents also had to be primarily responsible for administering medications to their child, and could not have participated in a previous medication dosing study. Parents were excluded from participation if they had: 1) visual acuity worse than 20/50 (Rosenbaum Pocket Screener), 2) uncorrectable hearing impairment, or 3) if they or their child were too ill to participate. Written, informed consent was obtained from parents prior to participating in the study.

Once parents were enrolled, they were randomized to one of five groups which varied by the pairing of units of measurement used on the medication bottle label and on the dosing tool (Table 1; Figure 1). Randomization was blocked by site, in sets of 100 (20 per group for each of the 5 groups); a random number generator was used. Unit label/dosing tool combinations were selected to represent commonly seen standard practices. Consistent with pharmacy guidelines, teaspoon units on English language medication labels were translated for Spanish-language parents.^{21,22} For the purposes of this analysis, which only involved what was present on the label, Groups 1 and 4 were collapsed into one group which had “mL-only” labels. Group 2 had both mL and a “tsp” abbreviation. Group 3 had mL and “teaspoon” spelled out, and Group 5 had “teaspoon” units alone.

Parents were also randomized to receive medication labels with 3 different dose amounts (2.5, 5, 7.5 mL) and as part of the larger study, were asked to measure those amounts using 3 different dosing tools (2 types of oral syringes, and 1 dosing cup) (total of 9 trials).

Assessments

Assessments were performed immediately following subject enrollment. Trained research assistants conducted interviews in English or Spanish, as per caregiver preference. The assessment involving choice of nonstandard dosing tool was conducted first; a structured survey was used subsequently to assess sociodemographic characteristics and health literacy. A gift card (\$20) was provided to study subjects as a nominal incentive.

Choice of Dosing Tool

At the beginning of the dosing assessment, parents were shown the first medication bottle label they were randomized to and asked “If you had all of these dosing tools at home, which of these would let you give your child the correct amount of medicine?” Parents were shown a response card with the following dosing tool names, accompanied by a photograph of the tool (kitchen teaspoon, kitchen tablespoon, dosing spoon, measuring spoon, dosing cup, dropper, oral syringe) (Figure 2); parents were able to select more than one. In addition, parents were asked if they would choose any other tool not shown on the card, with responses recorded verbatim. Parents who selected kitchen teaspoon or kitchen tablespoon were considered to have chosen a “nonstandard” dosing tool.

Health Literacy

Parent health literacy was assessed using the Newest Vital Sign (NVS),²³ which is validated in English and Spanish. A score of 0 or 1 was considered to be “low” health literacy, reflecting a high likelihood of limited literacy; 2 or 3, “marginal” health literacy, reflecting possible limited literacy; and 4 to 6, adequate health literacy.

Sociodemographic Data and Child Health Status

The following sociodemographic data were collected: child age, gender; parent age, gender, relationship to child, marital status, income, country of birth, race/ethnicity, language, educational attainment. Child’s chronic disease status and associated medication use were assessed using questions from the Children With Special Health Care Needs screener, which were adapted to assess chronic disease and medicine use in any child in the household.²⁴

Statistical Analyses

Statistical analyses were performed using SAS software version (9.4, (SAS Institute, Inc, Cary, North Carolina). Descriptive statistics were calculated for each variable. Chi-square analyses were conducted to compare parent characteristics between randomization groups for categorical variables, while ANOVA and Kruskal-Wallis tests were used to compare differences between groups for continuous measurements. Similar analyses were performed to compare families who did and did not enroll in the study.

Chi-square analyses were used to compare nonstandard dosing tool choice within those groups where a teaspoon unit was present on the label (Groups 2, 3, 5) versus those groups where only mL was present on the label (Groups 1 and 4). Analyses were also used to compare nonstandard dosing tool choice by each group which had a teaspoon unit present on the label individually (Groups 2, 3, 5). Multiple logistic regression was performed, controlling for key study variables of label language and dose amount, which were established *a priori*, and those characteristics found to be statistically different between randomization groups (i.e. health literacy). Stratified analyses and interaction tests were also performed by health literacy and by language.

RESULTS

Between August 26, 2013 and December 18, 2014, a total of 3116 parents were assessed for eligibility (Figure 3). Of these, 816 were not eligible based on one or more criteria, leaving 2300 eligible to complete the study. Ultimately, 2110 parents went on to enroll in the study and underwent randomization (74 parents ran out of time after they were assessed for eligibility and 116 refused to participate). Characteristics did not differ between subjects enrolled in the study and those who were eligible but did not enroll ($p>0.05$ for all characteristics). Two thousand and ninety-six parents completed the assessment of dosing tool preference and are included in this study analysis.

Characteristics of the study sample are shown in Table 2. Over half of parents were Hispanic, and a third Black, non-Hispanic. Nearly 80% of parents had an annual household

income <\$40,000. The majority of parents fell into the lowest 2 levels of health literacy (36.0% low, 41.0% marginal, 23.0% adequate).

Choice of Dosing Tool

Parents who received medication labels containing “tsp” or “teaspoon” units (Groups 2, 3, and 5) were more likely to choose a nonstandard dosing tool compared to those parents who received medication labels containing only “mL” units (Groups 1 and 4) (27.7 vs. 8.3%, $p<0.001$; adjusted odds ratio (AOR)=4.4[95% CI 3.3–5.8]) (see Table 3); the odds of choosing a nonstandard tool varied based on whether “teaspoon” was spelled out or abbreviated on the label (“teaspoon”-alone: AOR=5.3[3.8–7.3]; “teaspoon” with mL: AOR=4.7[3.3–6.5]; “tsp” with mL: AOR=3.4[2.4–4.7]). There was no statistically significant difference between parents who received labels with “teaspoon” spelled out with and without mL present (Group 3 vs. 5). Odds of choosing a kitchen spoon was statistically significantly lower when the “tsp” abbreviation was present compared to when the “teaspoon” was spelled out (Group 2 vs. 3: $p=0.04$; Group 2 vs. 5: $p=0.005$). There was no difference by site for group on tool preference.

Health Literacy and Choice of Nonstandard Dosing Tool

Parents with low health literacy had a higher odds of choosing a nonstandard dosing tool compared to parents with adequate health literacy (25.0% vs. 18.1%; AOR=1.6 [1.2–2.2]) (see Table 4). There were no significant interactions between label type and health literacy ($p=0.2$).

Language and Choice of Nonstandard Dosing Tool

There was no statistically significant difference in choice of nonstandard dosing tool by language ($p=0.1$) (see Table 4). In addition, no significant interactions were found between label type and language ($p=0.6$).

DISCUSSION

This study is the first to examine the impact of medication label terms (units of measurement) on parents’ choice of dosing tools. We found that labels with “tsp” or “teaspoon” were associated with an increased likelihood of a parent choosing a nonstandard kitchen spoon, compared to when metric ‘mL’ units were used. Odds of choosing a nonstandard tool were higher when the word “teaspoon” was spelled out, compared to when it was abbreviated (“tsp”). These findings remained consistent across parent health literacy and language groups, as well as across sites.

Among parents who were shown labels with “tsp” or “teaspoon” – nearly a third considered using a kitchen spoon, compared with less than 10% of parents who were shown labels with “milliliter” or “mL.” This 20% decrease in absolute risk is likely to be significant, as it is well-established that use of nonstandard spoons increases the likelihood of parent dosing errors;^{15,16} use of standard tools, such as oral syringes and droppers, which contain markings to help parents dose accurately, is endorsed by numerous organizations, including the AAP and FDA.^{3,9,12,25}

Some findings may appear intuitive. Choosing a kitchen spoon is an obvious and logical choice when presented with the word “teaspoon” or its abbreviation. Within a traditionally non-metric environment, many US parents may be unaware of the official metric equivalent for one teaspoon (5 mL). A parent’s choice of dosing tool is likely to be driven by a desire to seek concordance between the prescribed or recommended dose and selection of a tool which would allow for measurement of that dose.

Unexpectedly, the precise language on the label had an impact on parent choice of nonstandard tools. When the word “teaspoon” was spelled out rather than abbreviated (“tsp”), parents were more likely to choose a nonstandard kitchen spoon. This may be because the “tsp” abbreviation may be less familiar than words like “tablespoon” or “teaspoon”. The abbreviation might also be more likely to direct parents to think about the “tsp” units on standard dosing tools. Also of interest, a label with “mL” alongside “teaspoon” did not reduce parent’s likelihood of choosing a kitchen spoon. This further suggests a parent’s propensity to seek out the familiar.

Interestingly, there was a lack of significant effect modification by parent literacy or language. While those with low health literacy were slightly more likely to choose a nonstandard dosing tool, parents in each health literacy group had similar odds of choosing a kitchen spoon when presented with labels with teaspoon units vs. “mL”-only labels. Still, even with “mL”-only labels, more than 1 in 10 parents with low health literacy chose a kitchen spoon. Interestingly, parents’ English language proficiency was not associated with choice of nonstandard kitchen spoons. Perhaps, parents whose primary language is Spanish may have prior knowledge and orientation to metric measurements, particularly if they immigrated from one of the majority of nations that use metric standards. Nonetheless, parents across language levels benefited from avoidance of teaspoon units on labels.

Notably, our study was conducted in three geographically distinct sites, and there was no difference in the impact of group on tool preference by site. These findings suggest that avoidance of teaspoons on labels is likely to benefit parents across the US.

This study has limitations common for a cross-sectional analysis of data from a controlled experiment. We used a “mock” medication label for a hypothetical child; our findings therefore might not reflect the actual dosing tool choices parents would make at home when medications are prescribed to their own children. Dosing tool choices were purposefully shown as photographs, with markings not clearly identifiable, as we sought to determine whether parents would choose a specific dosing tool type; we did not want parents to be distracted by the individual markings of each tool. Also, images of kitchen spoons were the first ones on the page, followed by a range of standardized dosing tools, which may have affected which choices parents selected. There are also limitations resulting from the study parameters we used. We restricted testing to 4 commonly seen label variations. Other variations (e.g. “tsp” alone, tablespoon units) were not assessed, and only one label design format was used. Furthermore, this study was conducted with parents who speak English and Spanish and who brought their children to university-affiliated pediatric clinics which serve predominantly low income families; results may not be generalizable. We note, however, that our sample included a disproportionate number of lower literate parents, many

of who were from more medically vulnerable communities. We purposefully targeted these patients given their higher risk for misunderstanding medication instructions and unintentionally misusing medications.

CONCLUSION

Four decades later, despite recommendations from the AAP that parents not use nonstandard kitchen spoons to dose,¹² parents still commonly consider the use of nonstandard spoons to administer medications. Our study findings show that the units of measurement used on prescription medication labels heavily influences parent choice of liquid medication dosing tool; when teaspoon units were present on the label, parents were significantly more likely to choose a nonstandard kitchen spoon. The recently proposed AAP and CDC-endorsed mL-exclusive system seeks to limit the use of terms like “teaspoon” and “tablespoon”.⁸ We found in our study, however, that even with “mL”-only labels, nearly 1 in 10 parents still considered using nonstandard tools, indicating that additional strategies might be needed to further reduce kitchen spoon use and familiarize parents with proper dosing. Health literacy-informed communication strategies such as demonstration and teachback/showback using recommended dosing tools would help providers reinforce the idea that standardized dosing tools are preferable to kitchen spoons.^{26–28} Developing processes and policies in both clinical settings and pharmacies to support the provision of dosing tools to families would further reinforce this concept.^{26,27} Because the US has a longstanding reliance on non-metric units, a move to an “mL”-only medication label will likely require a public health campaign to ensure that a move towards eliminating that which is familiar does not drive further disparities, especially among parents with low health literacy.

Acknowledgments

Funding Source: Supported by the National Institutes of Health (NIH)/National Institute of Child Health and Human Development (R01HD070864). Dr. Yin is supported by HRSA Grant 12-191-1077- Academic Administrative Units in Primary Care. Dr. Sanders is supported by FDA CERSI grant (UCSF-Stanford CERSI Award #13).

We thank our research staff, and the staff of the pediatric outpatient clinics at Bellevue Hospital Center, Gardner Packard Children’s Health Care Center, and Children’s Hospital of Atlanta at Hughes Spaulding, for their support.

References

1. Institute of Medicine. Standardizing Medication Labels: Confusing Patients Less, Workshop Summary. Washington, DC: Roundtable on Health Literacy Board on Population Health and Public Health Practice; 2008.
2. Institute of Medicine. Preventing Medication Errors. Washington, DC: The National Academies Press; 2006.
3. U.S. Department of Health and Human Services. Guidance for industry: dosage delivery devices for orally ingested OTC liquid drug products. Silver Spring, MD: Food and Drug Administration Center for Drug Evaluation and Research (CDER); 2011.
4. Wallace LS, Keenum AJ, DeVoe JE. Characteristics of container labeling in a sample of commonly prescribed children’s oral medications. *Res Soc Adm Pharm*. 2010; 6:272–279.
5. Kaushal R, Jaggi T, Walsh K, Fortescue EB, Bates DW. Pediatric medication errors: what do we know? what gaps remain? *Ambul Pediatr*. 2004; 4(1):73–81. [PubMed: 14731088]

6. Yin HS, Wolf MS, Dreyer BP, Sanders LM, Parker RM. Evaluation of consistency in dosing directions and measuring devices for pediatric nonprescription liquid medications. *JAMA*. 2010; 304(23):2595–2602. [PubMed: 21119074]
7. DeWalt DA. Ensuring safe and effective use of medication and health care: perfecting the dismount. *JAMA*. 2010; 304(23):2641–2642. [PubMed: 21119075]
8. Paul IM, Yin HS. Out with teaspoons, in with metric units: pediatricians urged to prescribe liquid medications in mLs only. *AAP News*. 2012; 33(3):10.
9. Paul IM, Neville K, Galinkin JL, et al. Metric units and the preferred dosing of orally administered liquid medications. *Pediatrics*. 2015; 135(4):784–787.
10. Yin HS, Dreyer BP, Ugboaja DC, et al. Unit of measurement used and parent medication dosing errors. *Pediatrics*. 2014; 134(2):e354–e361. [PubMed: 25022742]
11. McQueen MJ. Conversion to SI units: the Canadian experience. *JAMA*. 1986; 256(21):3001–3002. [PubMed: 3773219]
12. Yaffe SJ, Bierman CW, Cann HM, et al. Inaccuracies in administering liquid medication. *Pediatrics*. 1975; 56(2):327–328. [PubMed: 1161381]
13. Yin HS, Mendelsohn AL, Wolf MS, et al. Parents' medication administration errors: role of dosing instruments and health literacy. *Arch Pediatr Adolesc Med*. 2010; 164(2):181–186. [PubMed: 20124148]
14. Wolf MS, Davis TC, Curtis LM, et al. Effect of standardized, patient-centered label instructions to improve comprehension of prescription drug use. *Med Care*. 2011; 49(1):96–100. [PubMed: 21150800]
15. Yin HS, Dreyer BP, Foltin GL, van Schaick L, Mendelsohn AL. Association of low caregiver health literacy with reported use of nonstandardized dosing instruments and lack of knowledge of weight-based dosing. *Ambul Pediatr*. 2007; 7(4):292–298. [PubMed: 17660100]
16. Madlon-Kay DJ, Mosch FS. Liquid medication dosing errors. *J Fam Pract*. 2000; 49(8):741–744. [PubMed: 10947142]
17. Levya M, Sharif I, Ozuah PO. Health literacy among Spanish-speaking Latino parents with limited English proficiency. *Ambulat Pediatr*. 2005; 5(1):56–59.
18. Bailey SC, Agarwal N, Sleath B, Gumusoglu S, Wolf MS. Improving drug labeling and counseling for limited English proficient adults. *J Health Care Poor Underserved*. 2011; 22(4):1131–1143. [PubMed: 22080698]
19. Masland MC, Kang SH, Ma Y. Association between limited English proficiency and understanding prescription labels among five ethnic groups in California. *Ethn Health*. 2011; 16(2):125–144. [PubMed: 21491287]
20. Bailey SC, Pandit AU, Yin S, et al. Predictors of misunderstanding pediatric liquid medication instructions. *Fam Med*. 2009; 41(10):715–721. [PubMed: 19882395]
21. U.S. Pharmacopeial Convention. USP 36 - NF 31. Prescription Container Labeling. Vol USP 362012
22. New York State Education Department Office of the Professions. Article 137, Pharmacy. Interpretation and translation requirements for prescription drugs and standardized medication labeling. Vol §68292015
23. Weiss BD, Mays MZ, Martz W, et al. Quick assessment of literacy in primary care: the newest vital sign. *Ann Fam Med*. 2005; 3(6):514–522. [PubMed: 16338915]
24. Bethell CD, Read D, Stein REK, Blumberg SJ, Wells N, Newacheck PW. Identifying children with special health care needs: development and evaluation of a short screening instrument. *Ambul Pediatr*. 2002; 2(1):38–48. [PubMed: 11888437]
25. National Council for Prescription Drug Programs. NCPDP Recommendations and Guidance for Standardizing the Dosing Designations on Prescription Container Labels of Oral Liquid Medications. Scottsdale, AZ: 2014.
26. Yin HS, Dreyer BP, Moreira HA, et al. Liquid medication dosing errors in children: role of provider counseling strategies. *Academic pediatrics*. 2014; 14(3):262–270. [PubMed: 24767779]
27. Yin HS, Jay M, Maness L, Zabar S, Kalet A. Health literacy: an educationally sensitive patient outcome. *Journal of general internal medicine*. 2015; 30(9):1363–1368. [PubMed: 26173523]

28. Yin HS, Dreyer BP, van Schaick L, Foltin GL, Dinglas C, Mendelsohn AL. Randomized controlled trial of a pictogram-based intervention to reduce liquid medication dosing errors and improve adherence among caregivers of young children. *Archives of pediatrics & adolescent medicine*. 2008; 162(9):814–822. [PubMed: 18762597]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

WHAT'S NEW

Parents who received prescription labels with teaspoon units had over 4 times the odds of choosing a nonstandard kitchen spoon. Findings support avoidance of teaspoon terms on labels, with benefits likely to be seen across health literacy and language groups.

CARLOS HERNANDEZ 444 Main St., Chicago, IL 60611		4/1/13
Amoxicillin 250 mg / 5 mL Take 5 mL (1 teaspoon) by mouth in the morning and at night for 10 days. Take for ear infection.		IMPORTANT: Finish all of this medicine (unless your doctor tells you to stop).
Rx: 0664978-5527 Do not use after: 4/15/13 Amount: 100 mL No refills Provider: Shonna Yin, MD		Pink liquid
		CITY PHARMACY 10 E. Wabash Chicago, IL 60601 (312) 555-5555

FIGURE 1.
Example of Medication Label (Group 3)

FIGURE 2.
Dosing Tool Card Shown to Parents

FIGURE 3.
Study Enrollment Flow Chart

^aRan out of time after signing consent

TABLE 1

Comparison of Randomization Group Characteristics

Group	Unit(s) Used on Medication Bottle Label	Example of how 5 mL or 1 tsp amount displayed on Label	
		English	Spanish
1&4	mL	5 mL	5 mL
2	mL, tsp	5 mL (1 tsp)	5 mL (1 cda)
3	mL, teaspoon	5 mL (1 teaspoon)	5 mL (1 cucharadita)
5	teaspoon	1 teaspoon	1 cucharadita

^aFull study involves variations in pairing of unit on label and dosing tool. Findings from this study involved only the bottle label; Groups 1 and 4 were exposed to the same label.

TABLE 2

Characteristics of Study Population, by Randomization Group (n=2096)

Label unit	Entire Population	Groups 1 & 4	Group 2	Group 3	Group 5	p-value
		mL	mL/tsp	mL/teaspoon	teaspoon	
		N=834	N=423	N=421	N=418	
	Mean (SD) or n(%)	Mean (SD) or n(%)	Mean (SD) or n(%)	Mean (SD) or n(%)	Mean (SD) or n(%)	
Child Characteristics						
Age, mean (SD), y	2.1 (2.2)	2.1 (2.2)	2.3 (2.2)	2.0 (2.2)	1.9 (2.1)	0.1
Gender, n(%)	983 (46.9)	385 (46.2)	209 (49.4)	196 (46.6)	193 (46.2)	0.7
Female						
Chronic medical problem treated with medication, n (%) ^a	353 (17.3)	151 (18.6)	64 (15.5)	64 (15.7)	74 (18.2)	0.4
Parent Characteristics						
Age, mean (SD), y	29.7 (7.3)	29.8 (7.3)	30.3 (7.5)	29.1 (7.1)	29.5 (7.5)	0.2
Relationship to Child, n (%)						
Mother	1876 (89.5)	750 (89.9)	382 (90.3)	377 (89.5)	367 (87.8)	0.6
Income, n (%)						
<\$10,000	497 (23.7)	217 (26.0)	99 (23.4)	85 (20.2)	96 (23.0)	0.02
\$10,000–\$19,999	553 (26.4)	201 (24.1)	114 (27.0)	143 (34.0)	95 (22.7)	
\$20,000–\$39,999	584 (27.9)	221 (26.5)	122 (28.8)	107 (25.4)	134 (32.1)	
\$40,000	255 (12.2)	111 (13.3)	47 (11.1)	48 (11.4)	49 (11.7)	
Unknown/Missing	207 (9.9)	84 (10.1)	41 (9.7)	38 (9.0)	44 (10.5)	
Country of birth: Non-US Born, n (%) ^b	1027 (49.4)	403 (48.6)	228 (54.2)	200 (48.2)	196 (47.3)	0.2
Race/Ethnicity, n (%) ^c						
Hispanic	1136 (54.7)	465 (56.2)	225 (53.6)	223 (54.0)	223 (53.9)	0.8
Non-Hispanic						

Label unit	Entire Population		Groups 1 & 4		Group 2		Group 3		Group 5		p-value
			mL		mL/tsp		mL/teaspoon		teaspoon		
			N=834		N=423		N=421		N=418		
	Mean (SD) or n(%)	Mean (SD) or n(%)	Mean (SD) or n(%)	Mean (SD) or n(%)	Mean (SD) or n(%)	Mean (SD) or n(%)	Mean (SD) or n(%)				
White, non-Hispanic	79 (3.8)	32 (3.9)	14 (3.3)	15 (3.6)	18 (4.3)						
Black, non-Hispanic	695 (33.5)	275 (33.2)	146 (34.8)	134 (32.4)	140 (33.8)						
Other, non-Hispanic	165 (8.0)	56 (6.8)	35 (8.3)	41 (9.9)	33 (8.0)						
Language Spanish, n (%) ^d	733 (35.0)	302 (36.2)	157 (37.1)	133 (31.6)	141 (33.7)					0.3	
Education, n (%) ^e											
Less than HS Graduate	633 (30.5)	258 (31.2)	136 (32.4)	116 (28.0)	123 (29.7)					0.7	
HS Graduate or Equivalent	674 (32.5)	266 (32.2)	140 (33.3)	138 (33.3)	130 (31.4)						
Higher than HS Graduate	769 (37.0)	303 (36.6)	144 (34.3)	161 (38.8)	161 (38.9)						
Health Literacy, n (%) ^f											
Inadequate Literacy	741 (36.0)	306 (37.4)	139 (33.4)	142 (34.6)	154 (37.2)					0.02	
Marginal Literacy	843 (41.0)	341 (41.7)	191 (45.9)	167 (40.7)	144 (34.8)						
Adequate Literacy	474 (23.0)	171 (20.9)	86 (20.7)	101 (24.6)	116 (28.0)						

^aMissing for 59 children overall (22 in groups 1 and 4, 11 in group 2, 14 in group 3, and 12 in group 5).

^bMissing for 17 parents (5 in groups 1 and 4, 2 in group 2, 6 in group 3, and 4 in group 5).

^cMissing for 21 parents (6 in groups 1 and 4, 3 in group2, 8 in group3, and 4 in group 4).

^dLanguage of survey administration.

^eMissing for 20 parents (7 in groups 1 and 4, 3 in group 2, 6 in group 3, 4 in group 5).

^fHealth literacy measured using Newest Vital Sign (NVS). Data missing for 38 subjects who did not complete the NVS (16 in groups 1 and 4, 7 in group 2, 11 in group 3, and 4 in group 5).

TABLE 3
Parent Choice of Nonstandard Tool by Randomization Group, Health Literacy, and Language (n=2096)

Group #	Label Unit(s)	% parents choosing non-standard tool ^a		AOR ^{b,c}	95% CI	p-value
		n (%)	p-value			
1&4	mL	69 (8.3)	<0.001	1.0	Ref	Ref
2	mL/tsp	94 (22.2)		3.4	2.4–4.7	<0.001
3	mL/teaspoon	122 (29.0)		4.7	3.3–6.5	<0.001
5	teaspoon	133 (31.8)		5.3	3.8–7.3	<0.001
<i>Health Literacy^d</i>						
Low		185 (25.0)	<0.001	1.6	1.2–2.2	0.004
Marginal		138 (16.4)		0.9	0.7–1.3	0.7
Adequate		86 (18.1)		1.0	Ref	Ref
<i>Language</i>						
English		256 (18.8)	0.07	1.0	Ref	Ref
Spanish		162 (22.1)		1.1	0.9–1.5	0.3

^aChi square.

^bFull model adjusting for group, dose amount, language, health literacy.

^cStatistically significant differences found for Group 2 vs. 3 [AOR (95% CI): 0.7 (0.5–0.99), p=0.04] and Group 2 vs. 5 [AOR (95% CI): 0.6 (0.5–0.9), p=0.005]; No statistically significant difference found between Group 3 vs. 5 [AOR (95% CI): 0.9 (0.7–1.2), p=0.4].

^dHealth literacy assessed using the Newest Vital Sign (NVS).

TABLE 4

Parent Choice of Nonstandard Tool by Randomization Group, Stratified by Health Literacy and Language

		% parents choosing non-standard tool		AOR ^b	95% CI	p-value
		n (%)	p-value ^a			
By Health Literacy ^{c,d}						
Low Health Literacy (n=741)						
Group #	Label Unit(s)					
1&4	mL	41 (13.4)	<0.001	1.0	Ref	Ref
2	mL/tsp	38 (27.3)		2.5	1.5–4.1	<0.001
3	mL/teaspoon	50 (35.2)		3.7	2.3–6.0	<0.001
5	teaspoon	56 (36.4)		3.8	2.4–6.1	<0.001
Marginal Health Literacy (n=843)						
Group #	Label Unit(s)					
1&4	mL	17 (5.0)	<0.001	1.0	Ref	Ref
2	mL/tsp	43 (22.5)		5.5	3.0–10.0	<0.001
3	mL/teaspoon	40 (24.0)		6.1	3.3–11.1	<0.001
5	teaspoon	38 (26.4)		6.8	3.7–12.5	<0.001
Adequate Health Literacy (n=474)						
Group #	Label Unit(s)					
1&4	mL	10 (5.9)	<0.001	1.0	Ref	Ref
2	mL/tsp	12 (14.0)		2.7	1.1–6.4	0.03
3	mL/teaspoon	27 (26.7)		5.7	2.6–12.5	<0.001
5	teaspoon	37 (31.9)		7.7	3.6–16.3	<0.001
By Language ^e						
English (n=1363)						
Group #	Label Unit(s)					
1&4	mL	35 (6.6)	<0.001	1.0	Ref	Ref
2	mL/tsp	58 (21.8)		4.1	2.6–6.5	<0.001
3	mL/teaspoon	81 (28.1)		5.5	3.5–8.5	<0.001

		% parents choosing non-standard tool		AOR ^b	95% CI	p-value
		n (%)	p-value ^a			
5	teaspoon	82 (29.6)		6.0	3.9–9.3	<0.001
Spanish (n=733)						
Group #	Label/ Unit(s)					
1&4	mL	34 (11.3)	<0.001	1.0	Ref	Ref
2	mL/tsp	36 (22.9)		2.5	1.5–4.3	<0.001
3	mL/teaspoon	41 (30.8)		3.7	2.2–6.2	<0.001
5	teaspoon	51 (36.2)		4.6	2.7–7.5	<0.001

^a Chi square.

^b Models by health literacy adjusting for group, dose amount, language; Models for language adjusting for group, dose amount, health literacy.

^c Health literacy assessed using the Newest Vital Sign (NVS).

^d Group by health literacy interaction non-significant (p=0.2).

^e Group by language interaction non-significant (p=0.6)