Published in final edited form as:

Arch Biochem Biophys. 2016 June 15; 600: 23–32. doi:10.1016/j.abb.2016.04.006.

Tropomyosin-Binding Properties Modulate Competition Between Tropomodulin Isoforms

Mert Colpan[#], Natalia A. Moroz, Kevin T. Gray, Dillon A. Cooper, Christian A. Diaz, and Alla S. Kostyukova[#]

Voiland School of Chemical Engineering and Bioengineering, Washington State University, Pullman, WA, 99164-6515, United States

Abstract

The formation and fine-tuning of cytoskeleton in cells are governed by proteins that influence actin filament dynamics. Tropomodulin (Tmod) regulates the length of actin filaments by capping the pointed ends in a tropomyosin (TM)-dependent manner. Tmod1, Tmod2 and Tmod3 are associated with the cytoskeleton of non-muscle cells and their expression has distinct consequences on cell morphology. To understand the molecular basis of differences in the function and localization of Tmod isoforms in a cell, we compared the actin filament-binding abilities of Tmod1, Tmod2 and Tmod3 in the presence of Tpm3.1, a non-muscle TM isoform. Tmod3 displayed preferential binding to actin filaments when competing with other isoforms. Mutating the second or both TM-binding sites of Tmod3 destroyed its preferential binding. Our findings clarify how Tmod1, Tmod2 and Tmod3 compete for binding actin filaments. Different binding mechanisms and strengths of Tmod isoforms for Tpm3.1 contribute to their divergent functional capabilities.

Keywords

tropomodulin; tropomyosin; actin; cytoskeleton

INTRODUCTION

Actin filaments are essential components of the cytoskeleton in cells; they facilitate various processes, such as cell migration, determination of cell shape and muscle contraction (for reviews, see (Pollard and Cooper 2009; Gunning et al. 2015a)). Actin filaments (F-actin) are formed by polymerization of globular actin (G-actin). Actin filaments have two distinct ends: a fast-growing (barbed) end and a slow-growing (pointed) end. Numerous actin-binding proteins regulate the organization of actin filaments (for review, see (dos Remedios et al. 2003; Hansen and Kwiatkowski 2013)). Tropomodulin (Tmod), a 40-kDa tropomyosin

[#]Corresponding authors, Address correspondence to: Phone number: +1-509-335-1888, Fax number: +1-509-335-4806, mert.colpan@wsu.edu; alla.kostyukova@wsu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

(TM)-binding protein, caps the pointed ends of actin filaments (Weber et al. 1994; Weber et al. 1999). Tmods regulate actin dynamics in muscle and non-muscle cells by preventing the polymerization and depolymerization of actin filaments from the pointed ends (for reviews, see (Yamashiro et al. 2012; Colpan et al. 2013; Fath 2013)).

Tmod has two structurally different domains: a disordered N-terminal domain and a compact C-terminal leucine-rich repeat (LRR) domain that consists of a repeated pattern of α -helices and β -strands (Kostyukova et al. 2000; Fujisawa et al. 2001; Kostyukova et al. 2001; Krieger et al. 2002). The N-terminal domain of Tmod has two TM-binding sites and one actin-binding site (Fowler et al. 2003; Greenfield et al. 2005; Kostyukova et al. 2005; Kostyukova et al. 2006). The N-terminal domain of one Tmod molecule binds two molecules of TM at the pointed end (Kostyukova et al. 2006; Kostyukova et al. 2007). TM is a coiled-coil protein that binds along the actin filaments in a head-to-tail fashion (for review, see (Gunning et al. 2015b)). The two TM-binding sites of Tmods have different affinities for TM isoforms (Uversky et al. 2011).

The C-terminal domain of Tmod contains a second actin-binding site (Gregorio et al. 1995; Fowler et al. 2003; Kostyukova and Hitchcock-DeGregori 2004; Rao et al. 2014). In addition to actin capping, Tmods can sequester actin monomers and nucleate actin polymerization (Fischer et al. 2006; Yamashiro et al. 2010).

Three Tmod isoforms, Tmod1, Tmod2 and Tmod3, are components of the cytoskeleton of brain cells (Cox and Zoghbi 2000; Cox et al. 2003). Tmod1 and Tmod2 have been shown to differently modulate formation of appendages in model cell lines used to study neuritogenesis (Fath et al. 2011; Moroz et al. 2013a; Guillaud et al. 2014). Tmod2 expression level changes in the brain of rats during epilepsy or cerebral ischemia (Iwazaki et al. 2006; Chen et al. 2007). Deletion of Tmod2 results in behavioral changes such as impaired memory and learning, hyperactivity and reduced sensorimotor gating in mice (Cox et al. 2003).

Tmod isoforms share some cellular compartments. Their structural similarity allows one isoform to compensate for the absence of another. For example, Tmod1 was found to be upregulated after Tmod2 knockout in mice (Cox et al. 2003). Tmod1 expression increased to compensate for Tmod2 knockdown in neuroblastoma N2a cells (Fath et al. 2011). When Tmod1 was knocked out, Tmod3 has been shown to replace it in skeletal muscle cells (Gokhin et al. 2010). Tmod3 is normally not found in red blood cells, but its expression was detected in mouse erythrocytes after Tmod1 knockout (Moyer et al. 2010). Although a substitution mechanism exists for Tmods, replacement of one isoform with another does not restore normal cell physiology and function (Gokhin et al. 2010; Moyer et al. 2010). Molecular differences between isoforms seem to attribute distinctive functions to Tmods, which affect cellular development and morphology. However, the molecular basis behind these differences in their functions is not very clear.

The purpose of this work was to compare the TM-dependent actin-binding abilities of Tmod1, Tmod2, Tmod3 and study their competition for binding at the pointed ends of actin filaments in the presence of Tpm3.1 (previously TM5NM1, see (Geeves et al. 2015) for the

new TM nomenclature), a non-muscle TM isoform that is involved in neuritogenesis (Fath et al. 2010) and cell elasticity (Jalilian et al. 2015). Based on data obtained from pyrene-actin polymerization assay, co-sedimentation, cross-linking and non-denaturing PAGE experiments, we propose that cooperative binding of Tmod3 to Tpm3.1 provides an advantage for Tmod3 to bind preferentially to the Tpm3.1-coated actin filaments when in competition with other Tmod isoforms.

EXPERIMENTAL PROCEDURES

Protein sequences

Sequences of Tmod1, Tmod2 and Tmod3 were downloaded from UniProt, Ensembl or NCBI for comparison of their amino acid sequences. The accession numbers for the analyzed Tmod1, Tmod2, Tmod3 isoforms are NP_990358.1, NP_001033799.1 and NP_058659.1, respectively.

Plasmid Construction

The plasmids for Tmod1; pET(His)Tmod1 and pET(His)Tmod1₁₋₃₄₄ (Kostyukova et al. 2000) were generated previously. The plasmids for Tmod2; pET(His)Tmod2 and Tmod3; pET(His)Tmod3 were used as templates to obtain pET(His)Tmod2₁₋₃₄₆ and pET(His)Tmod3₁₋₃₄₇, respectively. pET(His)Tmod3₁₋₃₄₇ was then used as a template to construct pET(His)Tmod3₁₋₃₄₇[L29E], pET(His)Tmod3₁₋₃₄₇[L134D] and pET(His)Tmod3₁₋₃₄₇[L29E/L134D]. For site-directed mutagenesis, the plasmids encoding Tmods were amplified by PCR using a set of two complementary oligonucleotides and *Pfu Turbo* DNA polymerase (Agilent Technologies, USA). The original template plasmid was digested with *DpnI* (New England Biolabs), and the mutated plasmid was transformed into *Escherichia coli* (max efficiency DH5α). All designed oligonucleotides were synthesized by Integrated DNA Technologies, Inc. (Coralville, IA). Introduced mutations were confirmed by DNA sequencing at GENEWIZ, Inc. (South Plainfield, NJ).

Tmod2₁₋₃₄₆ was generated using an oligonucleotide with the sequence: 5'-CTG GTT CGA AAG AAG AGA GTT TAA GGA GAC AGG AGG-3'. Tmod3₁₋₃₄₇ was generated using an oligonucleotide with the sequence: 5'-C AAG AGG CGA ATT TAA GGA GAC CAC CAG TAG CTC G-3'. The L29E mutation was introduced to Tmod3₁₋₃₄₇ using an oligonucleotide with the sequence: 5'-CTG TCC GAA TCA GAG GAG AAA CAG CTG GAG ACT GTC C-3'. The L134D mutation was introduced to Tmod3₁₋₃₄₇ using an oligonucleotide with the sequence: 5'-G CTG TGC GAC GAT GCA GCT ATT CTT GGG ATG C-3'. The construct for Tmod3₁₋₃₄₇[L29E] and the set of oligonucleotides for changing Leu-134 to Asp were used to generate Tmod3₁₋₃₄₇[L29E/L134D]. Mutated triplets in the sequences above are underlined.

Protein Expression and Purification

WT-Tmod1 and Tmod1₁₋₃₄₄ were expressed and purified as described in (Kostyukova and Hitchcock-DeGregori 2004). WT-Tmod2 and Tmod2₁₋₃₄₆ constructs were overexpressed in *E. coli* BL21 (DE3) using auto-induction medium according to the method described in (Studier 2005) and purified using Ni-NTA agarose and ion-exchange chromatography

according to the method described in (Moroz et al. 2013a). Tmod3 constructs (WT, truncated and mutants) were overexpressed in BL21 Rosetta™ 2(DE3)plysS Singles™ cells (Novagen) using LB media with 1% glucose, 0.1 mg/mL carbenicillin and 0.034 mg/mL chloramphenicol that was supplemented with 0.1 mM IPTG for induction at OD₆₀₀ = 0.6. The cells were grown for 5 hours after induction and pelleted at 8,000 RPM (Beckman-Coulter JA-10 Rotor), 4°C, 8 minutes. The pellets were resuspended in 20 mM Tris-HCl pH 7.5, 100 mM NaCl, 1 mM Pefabloc and 1 mM TLCK, sonicated and spun down at 16,000 RPM (Beckman-Coulter JA-17 Rotor), 4°C, 30 minutes. The pellet that contained Tmod3 in inclusion bodies was resuspended in 50 mM sodium phosphate pH 7.0, 150 mM NaCl and 10 mM Imidazole with 6 M Urea and solubilized overnight at 4°C on ice. The resuspension was centrifuged at 70,000 RPM (Beckman-Coulter MLA-80 Rotor), 4°C, 40 minutes and Tmod3 in the supernatant was purified using Ni-NTA agarose chromatography as in (Kostyukova et al. 2000) with denaturing conditions in the presence of 6 M urea. Purified fractions were combined for dialysis against 6 M Urea in 20 mM Tris-HCl, 150 mM NaCl, 1 mM EDTA and 1 mM DTT. Urea was removed and the protein refolded by buffer change from 6 M to 0 M Urea in 1 M steps using dialysis. Aggregates were removed by centrifugation at 70,000 RPM (Beckman-Coulter MLA-80 Rotor), 4°C, 1 hour before experiments.

Actin acetone powder was purified as described in (Pardee and Spudich 1982). Chicken skeletal G-actin was purified from acetone powder as described in (Kostyukova and Hitchcock-DeGregori 2004). Purified actin was used fresh in fluorescence experiments. Pyrene-iodoacetamide labeled G-actin was prepared (Kouyama and Mihashi 1981), labeling ratios were calculated according to (Cooper et al. 1983) and the labeled protein was stored in liquid nitrogen. Pyrene-actin was defrosted at 37°C and aggregates were removed by centrifugation at 100,000 RPM (Beckman-Coulter TLA-120.1 Rotor), 4°C, 1 hour before experiments.

The construct for rat γ -Tpm3.1 was a generous gift from Dr. Sarah Hitchcock-DeGregori (Rutgers University, Piscataway, NJ), and was expressed and purified as described previously (Hitchcock-DeGregori and Heald 1987; Hammell and Hitchcock-DeGregori 1996; Moraczewska et al. 1999) with modifications. Tpm3.1 construct was overexpressed in *E. coli* BL21 (DE3) using LB media with 0.1 mg/mL carbenicillin that was supplemented with 0.1 mM IPTG for induction at OD₆₀₀ = 0.6. Cells were grown for 5 hours after induction and pelleted at 8,000 RPM (Beckman-Coulter JA-10 Rotor), 4°C, 8 minutes. The pellet was resuspended in 50 mM Tris-HCl pH 7.5, 10 mM EDTA, 20% sucrose, and 5 mg/mL lysozyme, and frozen at -80°C for 10 minutes. After thawing at room temperature, 1 M NaCl was added to the resuspension drop wise, with stirring, and then sonicated for 4 minutes. The suspension was spun down at 50,000 RPM (Beckman-Coulter MLA-80 Rotor) at 4°C, 1 hour and the supernatant was immersed in boiling water for 5 minutes. After cooling for 45 minutes at room temperature, the boiled supernatant was spun at 17,000 RPM (Beckman-Coulter JA-17 Rotor), 4°C for 30 minutes. The supernatant was further fractionated by precipitation with (NH₄)₂SO₄. At 30% (NH₄)₂SO₄ saturation, the mixture was incubated for 20 minutes at 4°C and the precipitate was removed by centrifugation at 17,000 RPM (Beckman-Coulter JA-17 Rotor), 4°C, 30 minutes. The (NH₄)₂SO₄ concentration of the supernatant was further increased to 70% saturation and the mixture

was stirred overnight at 4°C. The precipitate contained Tpm3.1 and was pelleted at 17,000 RPM (Beckman-Coulter JA-17 Rotor), 4°C, 30 minutes. The pellet was resuspended in 20 mM Tris-HCl pH 7.5 and 0.5 mM DTT, and dialyzed against the same buffer overnight at 4°C.

The dialyzed Tpm3.1 was loaded onto a DE52 pre-swollen microgranular DEAE cellulose (Whatman) and was eluted by linear 0-1 M NaCl gradient. Fractions containing Tpm3.1 were combined and dialyzed against 1 mM sodium phosphate, pH 7.0, and 0.5 mM DTT for further hydroxyapatite chromatography. Tpm3.1 was then loaded onto HA Ultrigel column (Pall Life Technologies) and eluted by linear 1-200 mM sodium phosphate pH 7.0 gradient. Fractions containing Tpm3.1 were combined, dialyzed against 50 mM NH_4HCO_3 , lyophilized and resuspended in 20 mM Tris-HCl pH 7.5, 150 mM NaCl, 2 mM MgCl_2 and 0.5 mM DTT before experiments.

Recombinant human gelsolin was a generous gift from Dr. John H. Hartwig (Brigham and Women's Hospital & Hematology Division, Boston, MA). $\gamma\text{TM1bZip}$ was synthesized by the Tufts University Core Facility (Boston, MA). $\gamma\text{TM1bZip}$ is a chimeric peptide that contains 19 residues of $\gamma\text{Tpm3.1}$, encoded by exon 1b and the 18 C-terminal residues of the GCN4 leucine zipper domain, which helps to stabilize the coiled-coil structure of TM (Kostyukova et al. 2007). All peptide and protein concentrations were determined by difference method as described in (Kostyukova et al. 2007) and/or by BCA Protein Assay Kit (Pierce) according to manufacturer's instructions.

Titration of actin filaments with Tpm3.1

3 μM G-actin and 30 nM gelsolin were mixed and polymerized in centrifuge tubes with F-buffer (25 mM Imidazole, pH 7.0, 100 mM KCl, 2 mM MgCl_2 , 1 mM EGTA, 2 mM Tris-HCl, 0.2 mM CaCl_2 , 0.01% (v/v) NaN_3 , 0.2 mM ATP and 0.5 mM DTT) for 1 hour at room temperature to create short actin filaments capped at the barbed ends by gelsolin. Tpm3.1 was added to the filaments at different final concentrations (0-3 μM in 0.5 μM steps), stirred gently and incubated for 30 minutes. The filaments were pelleted at 70,000 RPM (Beckman-Coulter TLA-100 Rotor), room temperature for 30 minutes. The pellets were washed with F-buffer, and then solubilized in 8 M Urea at room temperature for 30 minutes. The pellet and supernatant samples were mixed with 2x Laemmli Sample Buffer (0.125 M Tris-HCl, pH 6.8, 20% glycerol, 4% SDS, 0.004% bromophenol blue and 10% 2-mercaptoethanol), heated and loaded on 12% acrylamide SDS-PAGE. Gels were stained with Coomassie Brilliant Blue R-250. The images were captured with Molecular Imager ChemiDoc XRS+ (Bio-Rad, Hercules, CA). Quantitative analysis for the background-corrected band densities of the pellet and supernatant samples was done with Image Lab™ Software (Bio-Rad, Hercules, CA). 1.5 μM was determined as the saturating concentration of Tpm3.1 for actin filaments. This concentration was used for the subsequent capping and co-sedimentation experiments with truncated Tmods.

Fluorescence Measurements

Actin polymerization was measured by the change in pyreneactin fluorescence using a PTI fluorometer (Lawrenceville, NJ) (excitation, 366 nm, and emission, 387 nm, with 2 nm slit).

To measure inhibition rates of actin polymerization at the pointed ends by Tmod isoforms, short filaments capped with gelsolin at their barbed ends were prepared by polymerization of 3 μ M G-actin in the presence of 30 nM gelsolin. Polymerization was monitored by the increase in fluorescence when the filaments were diluted 5-fold with G-actin (10% pyrenyl-actin) in F-buffer in the presence of 1.5 μ M Tpm3.1 and truncated Tmods at various concentrations (5, 7, 10, 20, 35, and 50 nM). After dilution, the final concentrations of F-actin and G-actin were 0.6 μ M and 1.1 μ M, respectively, according to (Kostyukova and Hitchcock-DeGregori 2004). Exponential growth curves to maximum were fitted to the polymerization data using SigmaPlot 12.0, and initial rates (R) were calculated as the first derivatives at time zero.

To measure actin nucleation, 1 μ M G-actin (10% pyrenyl-actin) was mixed with 50, 100 or 200 nM Tmod2 or Tmod3 (WT or truncated). Polymerization reactions were started by adding F-buffer. Spontaneous actin nucleation in the absence of Tmod was measured as control for each experiment.

Co-sedimentation of actin filaments in the presence of Tpm3.1 and truncated Tmods

Short gelsolin-capped actin filaments with 1.5 μ M Tpm3.1 were prepared as described above. 100 nM of Tmod1₁₋₃₄₄, Tmod2₁₋₃₄₆, Tmod3₁₋₃₄₇, Tmod3₁₋₃₄₇[L29E], Tmod3₁₋₃₄₇[L134D] or Tmod3₁₋₃₄₇[L29E/L134D] were added to the filaments individually or as a mixture for competition experiments using 5 μ M Tmod stock solutions prepared in G-buffer (2 mM Tris-HCl, pH 8.0, 0.2 mM CaCl₂, 0.01% (v/v) NaN₃, 0.2 mM ATP and 0.5 mM DTT). Actin filaments in the presence of Tpm3.1 were incubated with Tmods for 30 minutes at room temperature. The mixtures were then centrifuged at 70,000 RPM (Beckman-Coulter TLA-100 Rotor), room temperature, 30 minutes to separate Tmods bound to the filament in the pellet from unbound Tmods in the supernatant. The pellets were prepared for further SDS-PAGE as described above. Tpm3.1 and Tmods were centrifuged at 70,000 RPM (Beckman-Coulter TLA-100 Rotor), room temperature, 30 minutes prior to experiments to eliminate nonspecific sedimentation.

Western Blot

The primary antibodies used in Western Blot experiments included polyclonal rabbit antibodies against Tmod1 (Thermo Fisher Scientific, Waltham, MA), Tmod2 and Tmod3 (Pacific Immunology, Ramona, CA) custom made using Tmods that we purified. Optimal dilutions for each antibody were determined prior to experiments. The secondary antibodies used in the experiments were peroxidase-conjugated AffiniPure goat anti-Rabbit IgG (H+L) (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA).

12% acrylamide SDS-PAGE gels were run with samples of pellets from co-sedimentation experiments. SDS-gels were transferred to Immuno-Blot PVDF Membranes (0.2 μ m) for Protein Blotting (Bio-Rad, Hercules, CA) using Trans-Blot® Turbo™ Transfer System (Bio-Rad, Hercules, CA). After transfer, the membranes were blocked in 7% (w/v) nonfat milk in 1×PBS with 0.1% (v/v) Tween-20 (Sigma-Aldrich, St. Louis, MO) overnight at 4°C. Membranes were incubated with 2.5% (w/v) milk in 1×PBS with 0.1% (v/v) Tween-20, containing 1×50,000 dilution of the primary antibodies against Tmod1 or 1×25,000 dilution

of the primary antibodies against Tmod2 or Tmod3 for 1 hour. Unbound antibodies were washed off from membranes by shaking in 1×PBS with 0.1% (v/v) Tween-20 for 10 minutes, 5 times. Membranes were then incubated with 1×50,000 dilution of the secondary antibodies and washed using the same conditions as for primary antibodies. Membranes were finally incubated with Pierce ECL Western Blotting Substrate (Thermo Fisher Scientific, Waltham, MA) according to manufacturer's instructions for imaging. The images for blots were captured using Molecular Imager ChemiDoc™XRS+ System (Bio-Rad, Hercules, CA).

Densitometric analysis

Densitometric analysis of Western Blot membranes and Coomassie-stained gels was done using ImageJ Software (NIH). The background-corrected band densities for individual Tmods that were co-sedimented with actin filaments were taken as controls (100%). First, band densities of Tmods on the membranes were divided over the background-corrected band densities of actin in the gels from the corresponding samples. Second, for internal normalization the obtained values were divided by the value for the control to calculate the percentage of Tmod that co-sediments with actin filament when in competition (relative band density, experiment/control × 100). The bands on the membranes that were used for densitometric analysis were never allowed to reach signal saturation during image development. Sigma Plot 12.0 was used to generate bar graphs for relative band densities of co-sedimented Tmods and for statistical analysis (one-way ANOVA).

Nondenaturing-PAGE

Nondenaturing gel electrophoresis was performed using 9% polyacrylamide gels without SDS that were polymerized in the presence of 10% glycerol. 6 μM of Tmods were incubated with γTM1bZip peptide at 0-4 TM/Tmod molar ratio in 20 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM DTT and 1 mM EDTA. Native gel sample loading buffer (125 mM Tris-HCL, pH 6.8, 40% glycerol and 0.01% bromophenol blue) was added to samples before loading. Electrophoresis was run with constant 10 mA current on ice to prevent dissociation of complexes. After electrophoresis, gels were stained with Coomassie Brilliant Blue R-250 and imaged with Molecular Imager ChemiDoc™ XRS+ System (Bio-Rad, Hercules, CA). Quantitative analysis for the background-corrected band densities of the Tmod/TM complexes was done with Image Lab™ Software (Bio-Rad, Hercules, CA).

Cross-linking

For cross-linking Tpm3.1 to Tmod3₁₋₃₄₇, proteins were dialyzed against 5 mM HEPES, pH 7.5, 150 mM NaCl and 2 mM MgCl₂. 5 μM of Tpm3.1 was incubated with equimolar concentrations of Tmod3 at room temperature for 30 minutes. Cross-linking was initiated with 0.01% (v/v) glutaraldehyde at room temperature. After 30 minutes, the reaction was stopped by adding 100 mM Tris-HCl, pH 8.0. Samples were loaded on 15% SDS-PAGE gels. The gels were stained with Coomassie Brilliant Blue R-250 and scanned with Molecular Imager ChemiDoc™ XRS+ System (Bio-Rad, Hercules, CA). Quantitative analysis for the background-corrected band densities of Tmods was done with ImageJ Software (NIH). Sigma Plot 12.0 was used to generate bar graphs for percentages of Tmod3

bound or unbound to Tpm3.1 after cross-linking and for statistical analysis (one-way ANOVA).

RESULTS

Truncating the 5 C-terminal residues of Tmod2 and Tmod3 disrupts their nucleation ability

Tmods have isoform-dependent actin nucleating abilities; Tmod2 is a better nucleator of actin polymerization than Tmod3 and Tmod1 (Yamashiro et al. 2010). Nucleation by Tmods cause formation of new barbed ends, and as a result may confound the exact determination of their ability to inhibit actin polymerization at the pointed ends. To compare the pointed-end capping abilities of Tmod isoforms in the presence of Tpm3.1, we needed to disrupt the nucleating ability of Tmods without affecting their TM-dependent capping abilities.

It was previously shown that truncating the last 15 residues at the C-terminal domain of Tmod1 impaired its actin-nucleating ability, while not affecting its actin-capping ability in the presence of TM (Kostyukova and Hitchcock-DeGregori 2004). Alignment of the sequence of Tmod1 with Tmod2 and Tmod3 suggested that removing their 5 C-terminal amino acid residues should have the same effect (Figure 1). Therefore, we generated truncated proteins Tmod2₁₋₃₄₆ and Tmod3₁₋₃₄₇ to avoid the problems of distinguishing capping from nucleating and eliminating differences in nucleation abilities between Tmod isoforms.

The nucleating ability of the truncated Tmods was compared with full-length Tmods at 50, 100 and 200 nM concentrations using pyrene-actin nucleation assay. The full-length proteins demonstrated a concentration-dependent increase in the level of nucleation for actin polymerization (Figure 2A). No nucleation effect was observed for Tmod2₁₋₃₄₆ or Tmod3₁₋₃₄₇; there was no difference in polymerization of actin in the absence (spontaneous polymerization) or presence of Tmods (Figure 2). Therefore, truncated Tmod2 and Tmod3 lost their ability to nucleate actin polymerization and could be used for subsequent capping experiments.

Comparison of the capping ability of Tmod isoforms in the presence of Tpm3.1

To compare the capping abilities of truncated Tmod isoforms, we used pyrene-actin pointed end polymerization assay in the presence of Tpm3.1 (Figure 3A, B). Exponential growth curves were fit to the polymerization data and initial rates were calculated as derivatives at time 0 (Figure 3C). All isoforms reached full inhibition at 50 nM concentrations. Although the differences between the capping abilities of truncated Tmods were small, Tmod2 consistently demonstrated higher efficiency for capping at all concentrations.

Tmod3₁₋₃₄₇ co-sediments preferentially with actin filaments in the presence of Tpm3.1

To find if Tmod2 also binds better to actin filaments in the presence of other isoforms, co-sedimentation experiments were carried out. Tmods' binding to the pointed ends of actin filaments was studied in the presence of Tpm3.1. Binding and co-sedimentation of each isoform alone was used as a control. The filaments were co-sedimented by centrifugation and the pellets were used for subsequent Western Blot analysis.

Because of different specificities of antibodies against Tmods, the band densities quantified from the blots were not directly comparable between different Tmod isoforms. Thus for quantification purposes, we assumed individual Tmods co-sedimented in the control pellets to be 100%. To calculate the percentage of each Tmod isoform that co-sediments with actin filaments when in competition with other isoforms, the background-corrected band intensities obtained from competition experiments were divided over that of the corresponding control.

There were small decreases in amounts of Tmod3₁₋₃₄₇ co-sedimented with F-actin in the presence of Tmod1₁₋₃₄₄ or Tmod2₁₋₃₄₆. The amount of co-sedimented Tmod3₁₋₃₄₇ was slightly less in the presence of Tmod2₁₋₃₄₆ than that of Tmod1₁₋₃₄₄ (Figure 4 and Table 1). The amount of Tmod1₁₋₃₄₄ that co-sedimented with actin filaments decreased by ~60% and ~75% in the presence of Tmod2₁₋₃₄₆ or Tmod3₁₋₃₄₇, respectively. The amount of co-sedimented Tmod2₁₋₃₄₆ decreased by ~60% when Tmod2₁₋₃₄₆ was in competition with Tmod3₁₋₃₄₇. The amount of co-sedimented Tmod2₁₋₃₄₆ also went down when Tmod1₁₋₃₄₄ was present. However, we found that our polyclonal antibodies against Tmod2 cross-reacted with Tmod1 (Figure 4A). Therefore, the band obtained for the Tmod1-Tmod2 pair may contain either one of the two isoforms or their mixture and the actual decrease in the amount of co-sedimented Tmod2₁₋₃₄₆ may be even less than what was detected.

Surprisingly, while Tmod3₁₋₃₄₇ formed a weaker cap at the pointed ends compared to Tmod2₁₋₃₄₆, it co-sedimented preferentially with Tpm3.1-coated actin filaments in the presence of other Tmod isoforms.

The second TM-binding site is crucial for Tmod31-347's ability to outcompete other isoforms in binding to the pointed ends

We aimed to identify the reason for Tmod3₁₋₃₄₇'s preferential co-sedimentation with actin filaments in the presence of Tpm3.1. To study and compare the effects of individual TM-binding sites on Tmod3₁₋₃₄₇'s co-sedimentation, we generated Tmod3₁₋₃₄₇ with mutations in its first and second TM-binding sites. Previously it was shown that L27E and L131D mutations in Tmod1 disrupted the first and second TM-binding sites, respectively (Greenfield et al. 2005; Kostyukova et al. 2005). Homologous mutations in Tmod3, L29E and L134D, were introduced into Tmod3₁₋₃₄₇. Single and double mutants were purified. The mutants were tested using the co-sedimentation assay with actin filaments in the presence of Tpm3.1 and other Tmod isoforms. The amount of Tmod in the pellets was determined using isoform-specific antibodies (Figure 5 and Table 1).

Co-sedimentation of Tmod3₁₋₃₄₇[L29E] with Tpm3.1-coated actin filaments showed a similar trend to that of Tmod3₁₋₃₄₇; there was no statistically significant change in the amount of co-sedimented Tmod3₁₋₃₄₇[L29E] in the presence of Tmod1₁₋₃₄₄ compared to Tmod3₁₋₃₄₇[L29E] alone (Figure 5B). The amount of Tmod3₁₋₃₄₇[L29E] decreased in the presence of Tmod2₁₋₃₄₆. Consistent with these results, the amounts of Tmod1₁₋₃₄₄ or Tmod2₁₋₃₄₆ in the pellets in the presence of Tmod3₁₋₃₄₇[L29E] did not demonstrate a statistically significant change compared to the experiments with Tmod3₁₋₃₄₇.

Tmod3₁₋₃₄₇[L134D] demonstrated decreased binding to actin filaments in the presence of Tmod1₁₋₃₄₄ or Tmod2₁₋₃₄₆. This decrease was more pronounced in the presence of Tmod2₁₋₃₄₆ than in the presence of Tmod1₁₋₃₄₄. Consistently, the amounts of Tmod1₁₋₃₄₄ or Tmod2₁₋₃₄₆ in the pellets increased in the presence of Tmod3₁₋₃₄₇[L134D].

Tmod3₁₋₃₄₇[L29E/L134D] exhibited drastically decreased binding to actin filaments in the presence of Tmod1₁₋₃₄₄ or Tmod2₁₋₃₄₆. Correspondingly, the amount of Tmod1₁₋₃₄₄ or Tmod2₁₋₃₄₆ in the pellets increased further in the presence of Tmod3₁₋₃₄₇[L29E/L134D] compared to that of Tmod3₁₋₃₄₇[L134D].

Overall, the amounts of Tmod1₁₋₃₄₄ and Tmod2₁₋₃₄₆ in the pellets increased in the presence of Tmod3₁₋₃₄₇ only when Tmod3₁₋₃₄₇'s second or both TM-binding sites were disrupted.

Tmod3, but not Tmod1 or Tmod2, binds Tpm3.1 cooperatively

To compare the binding of Tmod isoforms to Tpm3.1, they were titrated with γ TM1bZip, a model peptide that encompasses the Tmod-binding site of Tpm3.1 (Kostyukova et al. 2007) and run on non-denaturing PAGE.

For all Tmod isoforms, the band for free Tmod disappeared and a new band appeared when γ TM1bZip was added (Figure 6A-C). This indicates the formation of γ TM1bZip/Tmod complexes. We quantified the band densities for the complexes to determine their stoichiometry. The Tmod1 band disappeared and the complex band for γ TM1bZip/Tmod1 reached saturation at ~1:1 molar ratio (Figure 6A, D). Therefore, Tmod1 molecule was able to bind only one γ TM1bZip molecule under these conditions. The complex bands for γ TM1bZip/Tmod2 and γ TM1bZip/Tmod3 reached saturation at ~2:1 molar ratio. Thus, Tmod2 and Tmod3 were able to bind two molecules of TM.

The band for free Tmod2 disappeared at 1:1 molar ratio (Figure 6B). However, the band density of the complex continued to increase until ~2:1 molar ratio (Figure 6D). This was in agreement with our previous study where one of Tmod2's two TM-binding sites is occupied by γ TM1bZip first, and after that the other site starts to bind γ TM1bZip (Moroz et al. 2013a). Unlike Tmod2, the band for free Tmod3 did not disappear completely until ~2:1 molar ratio (Figure 6C, D). These findings suggest that binding of Tmod3 to Tpm3.1 is cooperative, while there is no cooperativity in Tmod2's binding to Tpm3.1.

Tmod3₁₋₃₄₇[L29E] binds to Tpm3.1 stronger than Tmod3₁₋₃₄₇[L134D]

To compare the binding of the two TM-binding sites of Tmod3 independently to full-length Tpm3.1, we performed cross-linking experiments using Tpm3.1 and the TM-binding mutants of Tmod3. Tmod3₁₋₃₄₇ was used as a positive control. Individual proteins, as well as their mixtures with Tpm3.1 were cross-linked using glutaraldehyde.

TM molecule is a coiled-coil dimer and it can self-associate. Cross-linking by glutaraldehyde resulted in appearance of a band corresponding to the dimer (Figure 7A). With the addition of Tmod3, while the density of the dimer band decreased, the number and density of the higher molecular mass bands increased, which indicated complex formation between Tmod3 and Tpm3.1. Because of the high background from Tpm3.1 intramolecular

cross-linking and a large number of complex bands, we quantified the bands of free Tmods to estimate the amount of Tmods that was incorporated into complexes with Tpm3.1 (Figure 7B).

Approximately 60% of Tmod3₁₋₃₄₇ remained free after 30 minutes of cross-linking to Tpm3.1, whereas ~70% of Tmod3₁₋₃₄₇[L29E] and ~90% of Tmod3₁₋₃₄₇[L134D] was free under the same conditions. As expected, almost 100% of Tmod3₁₋₃₄₇[L29E/L134D] remained free. Based on these results, Tmod3₁₋₃₄₇[L29E] demonstrated more propensity to form a complex with Tpm3.1 than Tmod3₁₋₃₄₇[L134D]. This indicates that the second TM-binding site of Tmod3 binds Tpm3.1 stronger than its first site.

DISCUSSION

While Tmod isoforms can be located in the same cellular compartments, they regulate cellular morphology at different measures (Weber et al. 2007; Fath et al. 2011; Moroz et al. 2013a). In this study, we compared the pointed end-binding abilities of Tmod1, Tmod2 and Tmod3 to understand the molecular basis behind the differences in their function.

Previously the removal of the 28 C-terminal residues of Tmod3 (Tmod3₁₋₃₂₅) resulted in no detectable nucleation activity (Yamashiro et al. 2010). Tmod3₁₋₃₂₅ does not contain the 6th α -helix in the C-terminal domain, which was later shown to be a part of the second actin-binding site of Tmod1 (Rao et al. 2014). Here we showed a shorter, 5-residue truncation from the C-terminal end of Tmod2 and Tmod3 that destroyed their nucleation ability entirely. The truncations in this study were done on the unstructured C-terminal tail of Tmods (Kostyukova et al. 2000). Interestingly, this region was observed in the crystal structure of the second actin-binding site of Tmod1 with actin but it does not directly interact with actin (Rao et al. 2014). However, it immediately follows the second actin-binding site of Tmod1. Therefore this region should be crucial for the proper folding of a part of the second actin-binding site of Tmods, which is important for actin nucleation.

Tmod3 demonstrated different mechanism of binding TM. Both Tmod2 and Tmod3 were able to bind two molecules of γ TM1bZip. However, unlike Tmod2, Tmod3 bound γ TM1bZip in a cooperative fashion. Previously we found that the fragment for the first TM-binding site of Tmod1 has very low affinity for γ TM1bZip (Kostyukova et al. 2007). Here we showed that full-length Tmod1 binds only one molecule of γ TM1bZip under non-denaturing PAGE conditions. Consequently, the presence of the second TM-binding site in full-length Tmod1 does not strengthen interaction of its first TM-binding site with γ TM1bZip. This indicates that there is no cooperativity in Tmod1's binding to Tpm3.1. Conclusively, each Tmod isoform demonstrates a different mechanism for binding γ TM1bZip.

Tmod3 co-sedimented preferentially with Tpm3.1-coated actin filaments when in competition with Tmod1 and Tmod2. Mutating individual or both of the TM-binding sites of Tmod3 did not prevent its binding to Tpm3.1-coated actin filaments individually. Earlier it was reported that disrupting the two TM-binding sites of Tmod3 did not affect its ability to inhibit depolymerization of actin filaments in the presence of Tpm3.1. (Lewis et al. 2014).

Hence, actin-binding ability of Tmod3 may be significantly influential in its association with actin filaments.

In earlier studies TM was shown to increase Tmod's affinity for the pointed end ~1000-fold (Gregorio and Fowler 1995; Weber et al. 1999). A recent study demonstrated that Tmod1's capping ability for bare actin filaments was only ~4-fold weaker than for Tpm1.1-coated actin filaments (Rao et al. 2014). Based on our data, we suggest that actin binding by Tmods may significantly contribute for association at the pointed ends, and the extent of this contribution may change with respect to their affinities for different TM and actin isoforms.

Our cross-linking results suggest that the second TM-binding site of Tmod3 is crucial for its interaction with Tpm3.1. Recently Lewis et al. used Tmod3 with L29G and L134D mutations in blot overlay assays with Tpm3.1 and concluded that Tmod3 does not show a preference on its binding sites for interacting with Tpm3.1 (Lewis et al. 2014). Unlike L29E used in our study, the L29G mutation destroys the formation of the α -helix in the first TM-binding site. On the other hand, the L29E mutation only disrupts the hydrophobic surface in the first TM-binding site and therefore its effect is similar to the L134D mutation. The choices in different residues for introducing a mutation to the first TM-binding site of Tmod3 may be the reason for differences between our results and the published data. Alternatively, the nature of the experiments (blot overlay vs. cross-linking) may also have resulted in different findings. Our data obtained from co-sedimentation experiments support our cross-linking results. The second TM-binding site influences the preferential binding of Tmod3 at the pointed ends of actin filaments when in competition with other Tmod isoforms *in vitro*. We propose that the second TM-binding site of Tmod3 enables the other site to bind Tpm3.1 stronger.

Surprisingly Tmod2₁₋₃₄₆ capped actin filaments in the presence of Tpm3.1 better than Tmod1₁₋₃₄₄ and Tmod3₁₋₃₄₇, however, the difference was small. We did not observe a difference between the capping efficiencies of Tmod1₁₋₃₄₄ and Tmod3₁₋₃₄₇. This result correlates with a previous study, where full-length Tmod1 and Tmod3 (at concentrations < 10 nM) had comparable effects in inhibiting polymerization of actin filaments in the presence of Tpm3.1 (Lewis et al. 2014; Yamashiro et al. 2014).

Inhibition of actin polymerization by Tmod at the pointed end is a dynamic process (Littlefield et al. 2001). We hypothesize that Tmods may continuously associate and dissociate from the pointed end. On and off-rates of binding at the pointed ends should rely on multiple molecular interactions. The differences in the capping efficiencies of Tmod isoforms may not be as drastic considering the kinetics of capping. Nevertheless, the binding affinities of Tmod isoforms for actin and TM influence their localization at the pointed ends in cells (Tsukada et al. 2011; Moroz et al. 2013b). Since inhibition of actin polymerization can only take place once Tmods assemble at the pointed end, we suggest that the differences between the affinities of Tmod isoforms for their binding partners should confine them to different populations of TM-coated actin filaments in cells.

Altogether our findings show that there are substantial differences between the binding of Tmod isoforms to Tpm3.1 and Tpm3.1-coated actin filaments, which may account for the

differences in their cellular localization and modulation of pointed end dynamics. Our results suggest that there is a TM-dependent competition mechanism between Tmod isoforms for binding to the pointed end. Such mechanism may exist to fine-tune actin filament lengths in non-muscle cells. Tmod3 shows a preference for binding to Tpm3.1-decorated actin filaments in skeletal muscle (Gokhin and Fowler 2011) and adipocytes, and impairing both of its TM-binding sites reduces its interaction with Tpm3.1 in cells (Lim et al. 2015). Cooperative binding of Tmod3 to Tpm3.1 may influence its preferred binding to Tpm3.1 in cells, while other TM isoforms are also present in the environment. Studying mechanisms of Tmod isoforms' binding to other TM isoforms in future studies can reveal their different localization patterns, distinct populations and functions in the cells.

ACKNOWLEDGEMENTS

This work was supported by a National Institutes of Health grant GM081688 and startup funds to ASK. We would like to thank Dr. Sarah Hitchcock-DeGregori for providing the construct for Tpm3.1 expression and Kaylin Pan and Nina Yi for their assistance in optimizing the purification protocol for Tpm3.1.

REFERENCES

- Chen A, Liao WP, Lu Q, Wong WS, Wong PT. Upregulation of dihydropyrimidinase-related protein 2, spectrin alpha II chain, heat shock cognate protein 70 pseudogene 1 and tropomodulin 2 after focal cerebral ischemia in rats--a proteomics approach. *Neurochemistry international*. 2007; 50(7-8): 1078–1086. [PubMed: 17196711]
- Colpan M, Moroz NA, Kostyukova AS. Tropomodulins and tropomyosins: working as a team. *Journal of muscle research and cell motility*. 2013; 34(3-4):247–260. [PubMed: 23828180]
- Cooper JA, Walker SB, Pollard TD. Pyrene actin: documentation of the validity of a sensitive assay for actin polymerization. *Journal of muscle research and cell motility*. 1983; 4(2):253–262. [PubMed: 6863518]
- Cox PR, Fowler V, Xu B, Sweatt JD, Paylor R, Zoghbi HY. Mice lacking Tropomodulin-2 show enhanced long-term potentiation, hyperactivity, and deficits in learning and memory. *Molecular and cellular neurosciences*. 2003; 23(1):1–12. [PubMed: 12799133]
- Cox PR, Zoghbi HY. Sequencing, expression analysis, and mapping of three unique human tropomodulin genes and their mouse orthologs. *Genomics*. 2000; 63(1):97–107. [PubMed: 10662549]
- dos Remedios CG, Chhabra D, Kekic M, Dedova IV, Tsubakihara M, Berry DA, Nosworthy NJ. Actin binding proteins: regulation of cytoskeletal microfilaments. *Physiological reviews*. 2003; 83(2):433–473. [PubMed: 12663865]
- Fath T. Tropomodulins and tropomyosins - organizers of cellular microcompartments. *Biomolecular concepts*. 2013; 4(1):89–101. [PubMed: 25436568]
- Fath T, Agnes Chan YK, Vrhovski B, Clarke H, Curthoys N, Hook J, Lemckert F, Schevzov G, Tam P, Watson CM, Khoo PL, Gunning P. New aspects of tropomyosin-regulated neuritogenesis revealed by the deletion of Tm5NM1 and 2. *European journal of cell biology*. 2010; 89(7):489–498. [PubMed: 20223554]
- Fath T, Fischer RS, Dehmelt L, Halpain S, Fowler VM. Tropomodulins are negative regulators of neurite outgrowth. *European journal of cell biology*. 2011; 90(4):291–300. [PubMed: 21146252]
- Fischer RS, Yarmola EG, Weber KL, Speicher KD, Speicher DW, Bubb MR, Fowler VM. Tropomodulin 3 binds to actin monomers. *The Journal of biological chemistry*. 2006; 281(47): 36454–36465. [PubMed: 17012745]
- Fowler VM, Greenfield NJ, Moyer J. Tropomodulin contains two actin filament pointed end-capping domains. *The Journal of biological chemistry*. 2003; 278(41):40000–40009. [PubMed: 12860976]

- Fujisawa T, Kostyukova A, Maeda Y. The shapes and sizes of two domains of tropomodulin, the P-end-capping protein of actin-tropomyosin. *FEBS letters*. 2001; 498(1):67–71. [PubMed: 11389900]
- Geeves MA, Hitchcock-DeGregori SE, Gunning PW. A systematic nomenclature for mammalian tropomyosin isoforms. *Journal of muscle research and cell motility*. 2015; 36(2):147–153. [PubMed: 25369766]
- Gokhin DS, Fowler VM. Cytoplasmic gamma-actin and tropomodulin isoforms link to the sarcoplasmic reticulum in skeletal muscle fibers. *The Journal of cell biology*. 2011; 194(1):105–120. [PubMed: 21727195]
- Gokhin DS, Lewis RA, McKeown CR, Nowak RB, Kim NE, Littlefield RS, Lieber RL, Fowler VM. Tropomodulin isoforms regulate thin filament pointed-end capping and skeletal muscle physiology. *The Journal of cell biology*. 2010; 189(1):95–109. [PubMed: 20368620]
- Greenfield NJ, Kostyukova AS, Hitchcock-DeGregori SE. Structure and tropomyosin binding properties of the N-terminal capping domain of tropomodulin 1. *Biophysical journal*. 2005; 88(1):372–383. [PubMed: 15475586]
- Gregorio CC, Weber A, Bondad M, Pennise CR, Fowler VM. Requirement of pointed-end capping by tropomodulin to maintain actin filament length in embryonic chick cardiac myocytes. *Nature*. 1995; 377(6544):83–86. [PubMed: 7544875]
- Guillaud L, Gray KT, Moroz N, Pantazis C, Pate E, Kostyukova AS. Role of tropomodulin's leucine rich repeat domain in the formation of neurite-like processes. *Biochemistry*. 2014; 53(16):2689–2700. [PubMed: 24746171]
- Gunning PW, Ghoshdastider U, Whitaker S, Popp D, Robinson RC. The evolution of compositionally and functionally distinct actin filaments. *Journal of cell science*. 2015a; 128(11):2009–2019. [PubMed: 25788699]
- Gunning PW, Hardeman EC, Lappalainen P, Mulvihill DP. Tropomyosin - master regulator of actin filament function in the cytoskeleton. *Journal of cell science*. 2015b; 128(16):2965–2974. [PubMed: 26240174]
- Hammell RL, Hitchcock-DeGregori SE. Mapping the functional domains within the carboxyl terminus of alpha-tropomyosin encoded by the alternatively spliced ninth exon. *The Journal of biological chemistry*. 1996; 271(8):4236–4242. [PubMed: 8626768]
- Hansen MD, Kwiatkowski AV. Control of actin dynamics by allosteric regulation of actin binding proteins. *International review of cell and molecular biology*. 2013; 303:1–25. [PubMed: 23445807]
- Hitchcock-DeGregori SE, Heald RW. Altered actin and troponin binding of amino-terminal variants of chicken striated muscle alpha-tropomyosin expressed in *Escherichia coli*. *The Journal of biological chemistry*. 1987; 262(20):9730–9735. [PubMed: 2954961]
- Iwazaki T, McGregor IS, Matsumoto I. Protein expression profile in the striatum of acute methamphetamine-treated rats. *Brain research*. 2006; 1097(1):19–25. [PubMed: 16729985]
- Jalilian I, Heu C, Cheng H, Freittag H, Desouza M, Stehn JR, Bryce NS, Whan RM, Hardeman EC, Fath T, Schevzov G, Gunning PW. Cell Elasticity Is Regulated by the Tropomyosin Isoform Composition of the Actin Cytoskeleton. *Plos One*. 2015; 10(5)
- Kostyukova A, Maeda K, Yamauchi E, Krieger I, Maeda Y. Domain structure of tropomodulin: distinct properties of the N-terminal and C-terminal halves. *European journal of biochemistry / FEBS*. 2000; 267(21):6470–6475.
- Kostyukova AS, Choy A, Rapp BA. Tropomodulin binds two tropomyosins: a novel model for actin filament capping. *Biochemistry*. 2006; 45(39):12068–12075. [PubMed: 17002306]
- Kostyukova AS, Hitchcock-DeGregori SE. Effect of the structure of the N terminus of tropomyosin on tropomodulin function. *The Journal of biological chemistry*. 2004; 279(7):5066–5071. [PubMed: 14660556]
- Kostyukova AS, Hitchcock-DeGregori SE, Greenfield NJ. Molecular basis of tropomyosin binding to tropomodulin, an actin-capping protein. *Journal of molecular biology*. 2007; 372(3):608–618. [PubMed: 17706248]
- Kostyukova AS, Rapp BA, Choy A, Greenfield NJ, Hitchcock-DeGregori SE. Structural requirements of tropomodulin for tropomyosin binding and actin filament capping. *Biochemistry*. 2005; 44(12):4905–4910. [PubMed: 15779917]

- Kostyukova AS, Tiktopulo EI, Maeda Y. Folding properties of functional domains of tropomodulin. *Biophysical journal*. 2001; 81(1):345–351. [PubMed: 11423419]
- Kouyama T, Mihashi K. Fluorimetry study of N-(1-pyrenyl)iodoacetamide-labelled F-actin. Local structural change of actin protomer both on polymerization and on binding of heavy meromyosin. *European journal of biochemistry / FEBS*. 1981; 114(1):33–38.
- Krieger I, Kostyukova A, Yamashita A, Nitani Y, Maeda Y. Crystal structure of the C-terminal half of tropomodulin and structural basis of actin filament pointed-end capping. *Biophysical journal*. 2002; 83(5):2716–2725. [PubMed: 12414704]
- Lewis RA, Yamashiro S, Gokhin DS, Fowler VM. Functional Effects of Mutations in the Tropomyosin-Binding Sites of Tropomodulin1 and Tropomodulin3. *Cytoskeleton*. 2014; 71(7):395–411. [PubMed: 24922351]
- Lim CY, Bi XZ, Wu DH, Kim JB, Gunning PW, Hong WJ, Han WP. Tropomodulin3 is a novel Akt2 effector regulating insulin-stimulated GLUT4 exocytosis through cortical actin remodeling. *Nat Commun*. 2015; 6
- Littlefield R, Almenar-Queralt A, Fowler VM. Actin dynamics at pointed ends regulates thin filament length in striated muscle. *Nature cell biology*. 2001; 3(6):544–551. [PubMed: 11389438]
- Moraczewska J, Nicholson-Flynn K, Hitchcock-DeGregori SE. The ends of tropomyosin are major determinants of actin affinity and myosin subfragment 1-induced binding to F-actin in the open state. *Biochemistry*. 1999; 38(48):15885–15892. [PubMed: 10625454]
- Moroz N, Guillaud L, Desai B, Kostyukova AS. Mutations changing tropomodulin affinity for tropomyosin alter neurite formation and extension. *PeerJ*. 2013a; 1:e7. [PubMed: 23638401]
- Moroz NA, Novak SM, Azevedo R, Colpan M, Uversky VN, Gregorio CC, Kostyukova AS. Alteration of tropomyosin-binding properties of tropomodulin-1 affects its capping ability and localization in skeletal myocytes. *The Journal of biological chemistry*. 2013b; 288(7):4899–4907. [PubMed: 23271735]
- Moyer JD, Nowak RB, Kim NE, Larkin SK, Peters LL, Hartwig J, Kuypers FA, Fowler VM. Tropomodulin 1-null mice have a mild spherocytic elliptocytosis with appearance of tropomodulin 3 in red blood cells and disruption of the membrane skeleton. *Blood*. 2010; 116(14):2590–2599. [PubMed: 20585041]
- Pardee JD, Spudich JA. Purification of muscle actin. *Methods in enzymology*. 1982; 85(Pt B):164–181. [PubMed: 7121269]
- Pollard TD, Cooper JA. Actin, a central player in cell shape and movement. *Science*. 2009; 326(5957):1208–1212. [PubMed: 19965462]
- Rao JN, Madasu Y, Dominguez R. Mechanism of actin filament pointed-end capping by tropomodulin. *Science*. 2014; 345(6195):463–467. [PubMed: 25061212]
- Studier FW. Protein production by auto-induction in high density shaking cultures. *Protein expression and purification*. 2005; 41(1):207–234. [PubMed: 15915565]
- Tsukada T, Kotlyanskaya L, Huynh R, Desai B, Novak SM, Kajava AV, Gregorio CC, Kostyukova AS. Identification of residues within tropomodulin-1 responsible for its localization at the pointed ends of the actin filaments in cardiac myocytes. *The Journal of biological chemistry*. 2011; 286(3):2194–2204. [PubMed: 21078668]
- Uversky VN, Shah SP, Gritsyna Y, Hitchcock-DeGregori SE, Kostyukova AS. Systematic analysis of tropomodulin/tropomyosin interactions uncovers fine-tuned binding specificity of intrinsically disordered proteins. *Journal of molecular recognition : JMR*. 2011; 24(4):647–655. [PubMed: 21584876]
- Weber A, Pennise CR, Babcock GG, Fowler VM. Tropomodulin caps the pointed ends of actin filaments. *The Journal of cell biology*. 1994; 127(6 Pt 1):1627–1635. [PubMed: 7798317]
- Weber A, Pennise CR, Fowler VM. Tropomodulin increases the critical concentration of barbed end-capped actin filaments by converting ADP.P(i)-actin to ADP-actin at all pointed filament ends. *The Journal of biological chemistry*. 1999; 274(49):34637–34645. [PubMed: 10574928]
- Weber KL, Fischer RS, Fowler VM. Tmod3 regulates polarized epithelial cell morphology. *Journal of cell science*. 2007; 120(Pt 20):3625–3632. [PubMed: 17928307]

- Yamashiro S, Gokhin DS, Kimura S, Nowak RB, Fowler VM. Tropomodulins: pointed-end capping proteins that regulate actin filament architecture in diverse cell types. *Cytoskeleton*. 2012; 69(6): 337–370. [PubMed: 22488942]
- Yamashiro S, Gokhin DS, Sui Z, Bergeron SE, Rubenstein PA, Fowler VM. Differential actin-regulatory activities of Tropomodulin1 and Tropomodulin3 with diverse tropomyosin and actin isoforms. *The Journal of biological chemistry*. 2014; 289(17):11616–11629. [PubMed: 24644292]
- Yamashiro S, Speicher KD, Speicher DW, Fowler VM. Mammalian tropomodulins nucleate actin polymerization via their actin monomer binding and filament pointed end-capping activities. *The Journal of biological chemistry*. 2010; 285(43):33265–33280. [PubMed: 20650902]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- Tropomodulin (Tmod) isoforms demonstrate different Tpm3.1-binding modes.
- Tmod3 outcompetes other Tmod isoforms in binding to Tpm3.1-coated actin filaments.
- Second tropomyosin-binding site of Tmod3 is influential in its competition ability.

Figure 1. Schematic view of domain organization of Tmod

TM1 and TM2 are the first and second tropomyosin-binding sites, respectively. A1 and A2 are the N-terminal and C-terminal actin binding sites, respectively. LRR is the Leucine-Rich Repeat domain. Underlined: residues that were removed from Tmod1, Tmod2 and Tmod3 to terminate their nucleation ability.

Figure 2. Nucleating effects of Tmod2 and Tmod3 on actin polymerization

A: Concentration-dependent increase in the polymerization of 1 μ M actin in the presence of Tmod2 or Tmod3 (full-length or truncated). The increase in actin polymerization was calculated by dividing the fluorescence counts of actin in the presence of Tmods by that of actin alone (control) at $t = 40$ min. B and C: Representative fluorescence curves of actin polymerization in the presence of 100 nM Tmod2 or Tmod3, respectively. Control: spontaneous actin polymerization.

Figure 3. Tpm3.1-dependent pointed end capping by truncated Tmod isoforms

A and B: Time courses of pointed end elongation of gelsolin-capped actin filaments in the presence of Tpm3.1 (1.5 M) and truncated Tmods (5 or 50 nM) detected as an increase in fluorescence. C: Concentration-dependent inhibition of actin polymerization by truncated Tmods. Initial rates were calculated as the first derivatives at time zero after fitting. The inhibition of actin polymerization by truncated Tmods was calculated as $R_{\text{exp}}/R_{\text{control}}$. R_{control} denotes the rate of polymerization in the absence of Tpm3.1 and Tmod, and R_{exp} is the rate of polymerization in the presence of both Tpm3.1 and truncated Tmods. The relative

rate of actin polymerization in the presence of Tpm3.1 was 0.90 ± 0.01 (N=2). *Circle*, control; *triangle*, Tmod1₁₋₃₄₄; *square*, Tmod2₁₋₃₄₆; and *diamond*, Tmod3₁₋₃₄₇.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 4. Co-sedimentation of truncated Tmod isoforms with Tpm3.1-coated actin filaments

A: 100 nM truncated Tmod isoforms were added to gelsolin-capped actin filaments in the presence of Tpm3.1 (1.5 μ M), pelleted, ran on SDS-PAGE and transferred to nitrocellulose membranes for detection using antibodies against Tmods. Parallel Coomassie Blue-stained gels show loading. **0:** No Tmod, **1:** Tmod1₁₋₃₄₄, **2:** Tmod2₁₋₃₄₆, **3:** Tmod3₁₋₃₄₇. **B:** Relative percentages of Tmods in the pellets. Columns represent averages and error bars represent standard deviations (N=3). Asterisk indicates statistically significant groups ($p < 0.05$) determined by one-way ANOVA post-hoc Holm-Sidak test.

A **B**

Figure 5. Co-sedimentation of truncated Tmod isoforms with Tpm3.1-coated actin filaments

A: 100 nM truncated Tmod isoforms were added to gelsolin-capped actin filaments in the presence of Tpm3.1 (1.5 μ M), pelleted, ran on SDS-PAGE and transferred to nitrocellulose membranes for detection using antibodies against Tmods. Parallel Coomassie Blue-stained gels show loading. **1:** Tmod1₁₋₃₄₄, **2:** Tmod2₁₋₃₄₆, **3:** Tmod3₁₋₃₄₇, **3M1:** Tmod3₁₋₃₄₇[L29E], **3M2:** Tmod3₁₋₃₄₇[L134D], **3DM:** Tmod3₁₋₃₄₇[L29E/L134D]. B: Relative percentages of Tmods in the pellets. Columns represent averages and error bars represent standard deviations (N=3). Different letters (a, b, c) or asterisks above columns indicate statistically significant groups ($p < 0.05$) determined by one-way ANOVA post-hoc Holm-Sidak test. Note that the presence of L29E and/or L134D mutations changes the mobility of Tmod3 slightly on the SDS-PAGE.

Figure 6. Complex formation between Tmod isoforms and γ TM1bZip detected by non-denaturing PAGE

A-C: 9% acrylamide gels in the absence of SDS were used to assess the binding of 6 μ M of Tmod1-3 to γ TM1bZip at 0.0 to 4.0 TM/Tmod molar ratios. Arrowheads indicate free Tmod and arrows indicate γ TM1bZip/Tmod complex. Free γ TM1bZip does not enter the gel because of its high isoelectric point. D: Background-corrected band densities of γ TM1bZip/Tmod complexes were calculated from Coomassie Blue-stained non-denaturing PAGE gels and plotted over γ TM1bZip/Tmod molar ratios. Data are averages and error bars are standard deviations (N=3). A.U.: arbitrary units.

Figure 7. Cross-linking of Tpm3.1 with Tmod3

A: 5 μ M Tmod3 was cross-linked to equimolar concentration of Tpm3.1 using glutaraldehyde. Samples were run on 15% acrylamide SDS-PAGE gels. **3**: Tmod3₁₋₃₄₇, **3M1**: Tmod3₁₋₃₄₇[L29E], **3M2**: Tmod3₁₋₃₄₇[L134D], **3DM**: Tmod3₁₋₃₄₇[L29E/L134D]. B: Percent of Tmod3 that was free or in complex after cross-linking was calculated by taking the fraction of band density of free Tmod3 in the presence of Tpm3.1 over free Tmod3 in the absence of Tpm3.1. Columns are averages and error bars are the standard deviations (N=4). The differences between all groups were statistically significant (One-way ANOVA post-hoc Holm-Sidak test, $p < 0.05$).

Relative band densities of Tmods in the pellets after co-sedimenting with actin/Tpm3.1 filaments. Numbers are averages and errors are the standard deviations (N=3).

Table 1

	In the presence of:					
	Tmod1 _{1,344}	Tmod2 _{1,346}	Tmod3 _{1,347}	Tmod3 _{1,347} [L29E]	Tmod3 _{1,347} [L134D]	Tmod3 _{1,347} [L29E/L134D]
Tmodin pellet	Tmod1 _{1,344}	38±16%	26±6%	31±6%	60±9%	79±8%
	Tmod2 _{1,346}	100%	42±5%	42±6%	72±8%	91±4 %
	Tmod3 _{1,347}	81±3%	100%	-	-	-
	Tmod3 _{1,347} [L29E]	53±6%	-	89±14%	-	-
	Tmod3 _{1,347} [L134D]	29±3%	-	-	108±10%	-
	Tmod3 _{1,347} [L29E/L134D]	16±8%	-	-	-	80±18%