Published in final edited form as:

Bio Protoc. 2012 October 20; 2(20): .

Promoter Orientation of Prokaryotic Phase-variable Genes by PCR

Stacey L. Bateman and Patrick Seed*

Department of Molecular Genetics and Microbiology Center for Microbial Pathogenesis,
Department of Pediatrics, Duke University School of Medicine, Durham, NC, USA

Abstract

One major mechanism of phase variable gene expression in prokaryotes is through inversion of the promoter element for a gene or operon. This protocol describes how to detect the promoter orientation of a phase-variable gene by PCR. This protocol, including primer design, is specific to detection of the promoter orientations of *hyxR*, a LuxR-like response regulator in Extraintestinal Pathogenic *Escherichia coli* (ExPEC) isolates (Bateman and Seed, 2012); however, this protocol can be generalized to other organisms and genes to discriminate prokaryotic promoter inversions by PCR through size discrimination of the amplification products. Expression of *hyxR* is regulated through bidirectional phase inversion of the upstream promoter region mediated by a member of the family of site-specific tyrosine recombinases called Fim-like recombinases. The recombinases recognize inverted DNA repeat sequences flanking the promoter and produce a genomic rearrangement, orientating the promoter in favor or disfavor of gene expression.

Materials and Reagents

1. *Escherichia coli* (*E. coli*) isolate UTI893
2. Sterile distilled, deionized water (diH₂O)
3. Agarose, molecular biology grade, standard sieve
4. Ethidium Bromide, 1 mg/ml in distilled, deionized water (diH₂O) (or other agent to visualize DNA)
5. Taq polymerase (1 U/μl) with 10× NH₄ buffer (APEX Bioreserch Products, catalog number: 42–409)
6. 10 mM dNTP mix, PCR grade (Life Technologies, catalog number: 18427-013)
7. Tryptone
8. Yeast extract
9. NaCl
10. Tris base

*For correspondence: patrick.seed@duke.edu.

11. Boric acid
12. EDTA (pH 8.0)
13. SDS
14. Glycerol
15. Xylene cyanol
16. Bromophenol blue
17. hyxR phase-specific primers, 100 µM stock solution [Integrated DNA Technologies (IDT)]

5' – ACTGATAATAACCAGAGGCTTCTT – 3'

5' – CAGTGATTAACCTTTCGAACATATTG – 3'

5' – GCGAAAGTTAATCACTGGTATGACC – 3'

18. Tris-Borate-EDTA (TBE) (10× stock) (see Recipes)
19. 10× DNA Loading Dye (see Recipes)
20. Luria-Bertani broth (LB) culture medium (Sigma-Aldrich, catalog number: L3022-250G (see Recipes)
21. 2% TBE agarose gel with EtBr (see Recipes)
22. 20 ml 10× DNA Loading dye⁶ (xylene cyanol/bromophenol blue) (see Recipes)

Equipment

1. 37 °C Incubator for bacteria with aeration
2. Thermal cycler (Bio-Rad Laboratories)
3. Gel electrophoresis system (Owl Separation Systems)
4. UV Transilluminator with photo documentation (Bio-Rad Laboratories)

Procedure

1. Start 5 ml overnight culture of *E. coli* isolate UTI89 (Note 1) in LB at 37 °C with aeration. Culture can be started from a colony on an agar plate or from a –80 °C glycerol stock.
2. Setting up the phase-PCR reaction
 - a. Phase PCR was performed using a 1:1:1 mixture of 3 phase primers including 1) an anchor primer outside of the

¹UTI89 is a prototypic ExPEC cystitis isolate that was obtained from an adult patient with cystitis and has been well described in the literature (Mulvey, 1998). Primers for this protocol were designed to UTI89 genomic sequence, and while ExPEC isolates are similar, their genomic sequence may not be exact. It is recommended that researchers check that the primers listed in this protocol will work with other ExPEC isolates.

invertible region (primer 1); 2) a phase-specific primer for the OFF orientation (primer 2); and 3) a phase-specific primer for the ON orientation (primer 3).

Reaction mixture (Note 2)

1 μ l overnight bacterial culture (template)

2.5 μ l of 10 $\times$ APEX NH_4 buffer (1 $\times$ final concentration)

0.5 μ l of 1 unit/ μ l APEX Taq polymerase (0.5 unit final concentration)

1.25 μ l of 50 mM MgCl_2 (2.5 mM final concentration)

0.1 μ l of 100 μ M stock of each of 3 primers (0.4 μ M final concentration of each primer)

0.5 μ l of 10 mM dNTP mix (0.2 mM final concentration)

up to 25 μ l total volume with sterile dH_2O .

b. Thermal cycler program:

94°C for 3 min	☐	$\times 1$ cycle
94°C for 30 sec	}	$\times 30$ cycles
54°C for 30 sec		
72°C for 25 sec		

²All PCR reactions were performed using APEX Taq polymerase and accompanying buffers distributed by Genesee Scientific, Inc (San Diego, California, USA). Thermal cycler parameters were based on the manufacturer's recommendations. Taq polymerase or related thermostable polymerases would be expected to provide similar performance in this assay with minor modifications of the reactions.

72°C for 6 min × 1 cycle

Hold at 4°C

3. Make 50 ml volume of 2% agarose in 1× TBE (see below). Pour gel into casting apparatus.
4. Cover gel completely with 1× TBE (running buffer).
5. Mix 5–10 µl of PCR reaction with 0.5–1.0 µl 10× DNA loading dye (approx. 1× final concentration). Load into wells and run samples at 100–120 V for 45 min to 1 h.
6. Visualize and photograph using a UV gel transilluminator.
7. Mixed phase populations show two bands by PCR, corresponding to the OFF and ON promoter orientations. The OFF orientation gives a 196 bp amplicon vs. a 231 bp amplicon for the ON orientation.

Recipes

1. 1× LB
 Suspend 20 g in 1 L of distilled water
 Autoclave for 15 min at 121 °C
 Final concentration of components (Tryptone 10 g/L, yeast extract 5 g/L, NaCl 5 g/L)
2. 1 L 10× TBE buffer stock (Tris-borate-EDTA)
 108 g Tris base
 55 g Boric acid
 40 ml 0.5 M EDTA (pH 8.0)
 diH₂O to 1 L
 Stir until completely dissolved.
3. 1 L 1× TBE buffer (working) (Note 3)
 100 ml 10× TBE
 900 ml diH₂O
 Mix well
4. 2% TBE agarose gel with EtBr
 2 g agarose per 100 ml 1×

³1× TBE is what you will use to make your agarose gel as well as the buffer used to run the gel.

TBE Heat in microwave (approx. 2 min) until agarose is dissolved

Let stand at room temperature for 15–20 min to cool (you do not want to re-solidify)

Add EtBr to a final concentration of 0.1 µg/ml just prior to pouring gel.

5. 20 ml 10× DNA Loading dye (Note 4) (xylene cyanol/bromophenol blue)

Add 0.025 g each xylene cyanol and bromophenol blue

1.25 ml 10% SDS

12.5 ml 100% glycerol

6.25 ml H₂O

Mix thoroughly and store at room temperature

Acknowledgments

This protocol is adapted from Bateman and Seed (2012).

References

1. Bateman SL, Seed PC. Epigenetic regulation of the nitrosative stress response and intracellular macrophage survival by extraintestinal pathogenic *Escherichia coli*. *Mol Microbiol.* 2012; 83(5): 908–925. [PubMed: 22221182]
2. Mulvey MA, Lopez-Boado YS, Wilson CL, Roth R, Parks WC, Heuser J, Hultgren SJ. Induction and evasion of host defenses by type 1-piliated uropathogenic *Escherichia coli*. *Science.* 1998; 282(5393):1494–1497. [PubMed: 9822381]

⁴Any commercially available DNA loading dye is appropriate. The recipe given is only a home-made suggestion. Other recipes available use different tracking dyes and/or density agents.