Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/jor

Original Article

Short-term evaluation of arthroscopic outside-in repair of ulnar side TFCC tear with vertical mattress suture

Maysara Abdelhalim Bayoumy^{*}, Hesham A. Elkady, Hatem G. Said,
Amr El-Sayed, Waleed Riad Saleh

Orthopaedic and Traumatology Department, Assiut University Hospital, Assiut 71111, Egypt

ARTICLE INFO

Article history:

Received 13 August 2015

Accepted 11 October 2015

Available online 6 December 2015

Keywords:

TFCC

Wrist

Arthroscopy

ABSTRACT

Background: There has been controversy regarding the pathogenesis and treatment of TFCC tear. Different surgical techniques for the treatment of TFCC are prescribed. The purpose of this study was to evaluate the short-term outcomes of arthroscopic outside-in repair of the ulnar side TFCC tear with vertical mattress suture.

Methods: In this study, arthroscopic outside-in repair of the ulnar side TFCC tear was done. 37 patients with TFCC tear with failure of conservative treatment for a minimum of 6 months had surgery. Outcomes were assessed using modified Mayo wrist score (MMWS), the Disability of the Arm, Shoulder, and Hand (DASH), visual analog scale (VAS) for pain and satisfaction criteria, and patient-related wrist evaluation score (PRWE), in addition to the range of motion and grip strength.

Results: After arthroscopic surgery, overall satisfaction was extremely positive, and out of the 37 patients, 91.9% of the patients are satisfied. The mean score for pain improved from 7.6 to 2.9 points. The mean MMWS was improved from 62.1 to 91.2 points. DASH score also improved from 29.9 to 10.2 points and PRWE improved from 60 to 33 points. The ROM was improved from 85.8% to 92% of the normal side. The grip strength of the affected side improved from 82.5% to 89% of the normal side. All improvements are statistically significant ($P < 0.05$).

Conclusion: Arthroscopic outside-in repair of ulnar side TFCC tear with mattress suture is a reproducible method with a marked improvement in function within a short period.

© 2015 Prof. PK Surendran Memorial Education Foundation. Published by Elsevier, a division of Reed Elsevier India, Pvt. Ltd. All rights reserved.

^{*} Corresponding author.

E-mail address: mysarahalim2010@gmail.com (M.A. Bayoumy).

Abbreviations: TFCC, triangular fibrocartilage complex; NSAIDS, non-steroidal anti-inflammatory drugs; DRUJ, distal radio-ulnar joint; MMWS, modified Mayo wrist score; DASH, Disability of the Arm Shoulder and Hand; VAS, visual analog scale; PRWE, patient related wrist evaluation; ROM, range of motion; ECU, extensor carpi ulnaris.

<http://dx.doi.org/10.1016/j.jor.2015.10.013>

0972-978X/© 2015 Prof. PK Surendran Memorial Education Foundation. Published by Elsevier, a division of Reed Elsevier India, Pvt. Ltd. All rights reserved.

1. Introduction

Triangular fibrocartilage (TFC) injuries are increasingly recognized as a cause of ulnar-sided wrist pain. Palmer grouped these tears into either traumatic or degenerative with various subclassifications.¹

The traumatic lesions (class I) were grouped into four types. Type A is a central tear in the TFCC disk. Type B is peripheral tear from the ulnar side. Type C is a tear in the palmar side of the TFCC. Type D is a radial avulsion of the TFCC. Degenerative tears of the TFCC (type II) are caused by normal aging or arthritis.² The central area of fibrocartilage disk is avascular, and is therefore called the debridement zone (60% of TFCC). 10–40% of the peripheral area of the TFCC has an extensive blood supply, and is therefore called the repair zone. This vascularity enables the surgeon to effect arthroscopic suture repair of peripheral Palmer type IB and ID TFCC tears which yielded encouraging results in many studies.³

Conservative treatment for TFCC injury generally includes compresses, long arm splints or elastic bandages, and NSAIDs. Many patients experience relief of symptoms with such conservative treatment.³ When the response to conservative treatment is poor, and in cases of wide rupture of the TFCC, surgical treatment is recommended due to the risk of DRUJ instability.⁴

Acute type IB peripheral tears have been shown to be amenable to suture repair if they fail non-operative management. Both open and arthroscopic series have shown good improvement in pain, grip strength, and function with repair of peripheral tears.⁵

Several arthroscopic methods have been described, including inside-out, outside-in, and all-arthroscopic techniques.⁶

The “outside-in” repair, which involves piercing the TFCC via the ulnar side of the wrist, has been described by several authors.⁷ The “outside-in” techniques vary among authors with instrumentation and subtle surgical modifications.⁸

The purpose of this study was to evaluate the short-term outcomes of arthroscopic outside-in repair of the ulnar side TFCC tear with vertical mattress suture, using different validated scores. Authors also attempted to identify socio-demographic factors and that may be associated with the outcomes after arthroscopic surgery for type 1B TFCC tear.

2. Materials and methods

2.1. Patients

Our study consisted of 37 consecutive arthroscopic outside-in TFCC repair for type 1B TFCC tear performed in 37 patients between February 2011 and March 2013. There were 29 male (29 wrists, 78.4%) and 8 female (8 wrists, 21.6%). The average patient age at the time of surgery was 23.3 years (range: 18–34 years). 17 patients (45.9%) were involved in heavy manual labor, 11 patients (29.8%) were house wives, 5 patients (13.5%) were employed, and 4 patients (10.8) not working at the time of surgery. 26 (70.2%) of the 37 wrist involved the dominant arm. All of these TFCC tears were treated non-operatively before undergoing arthroscopic release. Non-operative measures

used were rest, activity modification, splint, physical therapy, non-steroidal anti-inflammatory medication, and corticosteroid injection. The mean duration of this treatment was 11.1 months (range: 5–36 months). Clinically, the predominant symptoms were pain that increased with ulnar deviation and grasping. On physical examination, patients consistently had point tenderness distal to the ulnar styloid as well as positive TFCC stress test and TFCC stress test with compression. A neutral rotation posteroanterior radiograph and a pronated grip view to detected ulna positive showed 21 wrists with ulna neutral and 13 wrists with ulna minus and 4 wrists with ulna positive. MRI detected the lesion in 33 of the affected wrists.

All patients underwent an examination and evaluation preoperative and at 1.5, 3, 6, 12, 18, and 24 months postoperatively. Patients' assessment consisted of evaluating the range of motion (ROM) and grip strength (grip strength data are reported as percentage of strength compared with the contralateral healthy wrist). Health-related quality of life outcomes were assessed using two different scores including the Disability modified Mayo wrist score (MMWS) of the Arm, Shoulder, and Hand (DASH), Visual Analog Scale (VAS), and Patient Reported Wrist Evaluation (PRWE).

2.2. Operative procedure

The operation is performed under general anesthesia. The patient is positioned supine with the shoulder abducted and the involved extremity on an arm board. A non-sterile tourniquet is used. The hand and forearm are draped freely after being prepared and 10–15 lbs of traction are applied to the long and ring fingers through the finger traps and placed into the wrist traction tower (Linvatec) (Fig. 1).

A standard 3–4 portal just distal to Lister's tubercle is used as the viewing portal for the 2.7-mm arthroscope throughout the TFCC repair process (Fig. 2). We routinely make a 6-R portal under needle localization just radial to the ECU as the working portal during the repair. After completing a standard diagnostic arthroscopy, the morphology of the TFCC tears had been closely examined. The tear pattern most amenable to the repair technique described here is type IB tear of the TFCC without any clinical or arthroscopic evidence of DRUJ instability (Fig. 3). Such a tear will often have a positive trampoline test or loss of elastic recoil when the articular disk is depressed with a probe, suggestive of loss of tension to the ulnar capsule.

Once the tear has been identified and clearly visualized with the assistance of a probe (Fig. 4), the scar tissue is removed with a synovial shaver, allowing for a new bleeding surface for new scar tissue formation.

2 cm longitudinal incision is made on the ulnar side of the wrist just volar to ECU. Blunt dissection with a right-angled clamp is used to identify and protect any branches of the dorsal sensory branch of the ulnar nerve within the field, and dissection is carried down to the retinaculum.

The instrument set for repair composed from 2 of 24 gauge needle is used for outside-in repair and 3-0 proline suture as a loop (as the set of the meniscus is of large caliber and can injure the disk). The following steps pertain to each repair suture placed. The typical tear will accommodate between 1 and 2 mattress stitches, with the most volar stitch placed first.

Fig. 1 – Traction applied to the wrist joint by wrist traction tower.

Fig. 2 – Viewing and working dorsal wrist portals.

With the arthroscope in the 3-4 portal, the first needle loaded with 2-0 PDS II polydioxanone monofilament suture inserted, starting approximately 1 cm proximal to the level of the ulnocarpal joint (Fig. 5). The needle is then angled distal and radial to pierce the articular disk several millimeters radial to the edge of the tear (Fig. 6). The 2-0 PDS is advanced into the joint, and then withdrawn with small artery forceps outside the joint through the 6R portal, and the 2nd straight needle loaded with a loop of 3-0 PDS is inserted starting at the level of the ulnocarpal joint (Fig. 7), and is angled directly radial to come through the disk in a point dorsal to the first needle, and the loop is advanced through the needle into the joint, and then withdrawn with small artery forceps outside the joint through the 6R portal. The 2-0 PDS is carefully passed through the loop. The loop is removed together with the 2-0 PDS from within the joint, creating the first mattress suture (Fig. 8). Both ends of each paired suture are sequentially rerouted such that the knot will lie directly on retinaculum with no interposed subcutaneous tissue or potential nerve branches and are then the knots are sequentially tied. Standard closure of the ulnar incision and arthroscopy portals is performed.

Postoperatively, the patient was placed in a short arm volar splint postoperatively for 4 weeks. Intermittent bracing with a

plastic brace was prescribed for another 4 weeks, and progressive gentle active motion of the wrist was allowed at this stage. Passive stretching exercise and active wrist rotation were prohibited until 3 months postoperatively.

Fig. 3 – Type 1B TFCC tear.

Fig. 4 – Probing of the TFCC tear.

Fig. 6 – 1st suture passage through the TFC radial to the site of the tear.

2.3. Statistical analysis

Statistical analysis was performed using the Student's paired t-test to compare between the preoperative and postoperative variables with the statistical significance set at $P < 0.05$. Statistical analysis was performed with SPSS version 19 (SPSS, Chicago, IL).

Fig. 5 – Passage of the 1st suture through the capsule at the level of ulnocarpal joint.

3. Results

According to the patient's reports, after surgery, 16 patients described themselves as completely asymptomatic, 18 were better or much better, and 3 remained the same. A highly statistically significant improvement was identified comparing preoperative with postoperative VAS for pain and PRWE ($P < 0.05$). The mean score for pain improved from 7.6 (range, 5–9) to 2.9 (range, 0–7). The total PRWE improved from 55 out of 150 (range, 35–72) to 33 out of 150 (range, 7–68).

The mean MMWS was improved from 62.1 (range 40–80) preoperatively to 91.2 (range, 75–100) postoperatively, and this improvement was statistically significant ($P < 0.05$). Evaluation of functional outcomes with the DASH score also revealed statistically significant improvement ($P < 0.05$). The mean preoperative score was 29.9 (range, 6.8–44.2), while the mean postoperative score was 10.2 (range, 0–31.8). The range of motion was improved from 85.8% (range, 70–100%) to 92% (range, 70–100%) of the normal side. The grip strength of the affected side improved from 82.5% (range, 60–100% of the normal side) to 89% (range, 75–100%). No significant associations were observed between age, gender, preoperative length of non-operative treatment, time to return to work, and final outcome ($P > 0.05$). Two complications were reported: one patient had dorsal ulnar nerve (DUN) neurapraxia, and the other one had weakness in extension of little finger (extensor digiti minimi irritation). Both of them resolved with non-operative treatment.

4. Discussion

TFCC injuries are common, especially in an active patient population. When non-operative modalities fail to resolve

Fig. 7 – Passage of the suture loop just proximal to the 1st suture.

symptoms associated with this injury, surgical management is indicated. A number of open, arthroscopic-assisted, and arthroscopic techniques have been described.⁵

Several series have shown good results of arthroscopic repair of peripheral TFCC tears with the outside-in technique.⁴ In a multicenter study of arthroscopic TFCC repairs with an average follow-up of 37 months, Corso et al.⁹ reported that 29 of 45 wrists were rated as excellent, 12 as good, and 4 as poor according to postoperative MMWS. Millants et al.¹⁰ in 2002 published an outcome study of 35 patients who had arthroscopic surgical repair. Patients were assessed using the VAS and DASH score. According to the DASH score, 29

patients had good results, 5 had fair results, and 1 had a poor result. Grip strength and wrist motion were not assessed. Haugstvedt et al.³ reported 70% good to excellent results by Mayo score in their cohort of 20 peripheral tears, with a 90% range of motion, and 83% grip strength in comparison with the uninjured side.

The three previously described studies critically examined the results of arthroscopic treatment of TFCC tears. These studies note the definitive value of arthroscopy with respect to diagnosis upon direct visualization, minimization of soft-tissue trauma, cosmesis, and quicker recovery.

Trumble et al.¹¹ reported that 22 patients treated with arthroscopic technique had a mean arc of motion at 36 months that was 86% of the contralateral side, and a grip strength that was approximately equal to 82% of the contralateral side.

Although the majority of patients received treatment within 4 months of injury, we cannot confirm this for all patients. Our study seems to confirm the results of the previous papers. The technique is easy and the required equipment is available in all the operating theaters. Since all the patients had longstanding therapy previously, and no other pathologies were disclosed, the correlation between the pathology (TFCC tear) and the symptoms (ulnar wrist pain) and the efficacy of the treatment seems to be obvious.

Conflicts of interest

The authors have none to declare.

Acknowledgements

We acknowledge Prof. Maher Elassal and Prof. Mohamed M. Elhamid for their support in collecting the data and preparing the photos of the article.

REFERENCES

1. Palmer AK. Triangular fibrocartilage complex lesions: a classification. *J Hand Surg Am.* 1989;14:594–606.
2. Infanger M, Grimm D. Meniscus and discus lesions of triangular fibrocartilage complex (TFCC): treatment by laser-assisted wrist arthroscopy. *J Plast Reconstr Aesthet Surg.* 2009;466–471.
3. Haugstvedt J, Husby T. Results of repair of peripheral tears in the triangular fibrocartilage complex using an arthroscopic suture technique. *Scand J Plast Reconstr Surg Hand Surg.* 1999; 33:439–447.
4. Wysocki RW, Ruch DS. Outside-in repair of TFCC tears. *Oper Tech Sports Med.* 2010;18:163–167.
5. Yao J. All-arthroscopic TFCC repair using fast-fix. *Oper Tech Sports Med.* 2010;18:168–172.
6. Chloros GD, Wiesler ER, Poehling GG. Current concepts in wrist arthroscopy. *Arthroscopy.* 2008;24:343–354.
7. Ruch DS, Papadonikolakis A. Arthroscopically assisted repair of peripheral triangular fibrocartilage complex tears: factors affecting outcome. *Arthroscopy.* 2005;21: 1126–1130.

Fig. 8 – Creating the 1st suture mattress.

8. Micucci CJ, Schmidt CC. Arthroscopic repair of ulnar-sided TFCC tear. *Oper Tech Orthop.* 2007;17:118-124.
9. Corso SJ, Savoie FH, Geissler WB, et al. Arthroscopic repair of peripheral avulsions of the triangular fibrocartilage complex of the wrist: a multicenter study. *Arthroscopy.* 1997;13:78-84.
10. Millants P, De Smet L, Van Ransbeeck H. Outcome study of arthroscopic suturing of ulnar avulsions of the triangular fibrocartilage complex of the wrist. *Chir Main.* 2002;21:298-300.
11. Trumble TE, Gilbert M, Vedder N. Arthroscopic repair of the triangular fibrocartilage complex. *Arthroscopy.* 1996;12:588-597.