The laparoscopic approach for repair of indirect inguinal hernias in infants and children

David Juang, Jason D. Fraser, George W. Holcomb III

Department of Surgery, Children's Mercy Hospital, Kansas City, MO, USA

Contributions: (I) Conception and design: GW Holcomb III; (II) Administrative support: All authors; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: None; (V) Data analysis and interpretation: None; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: George W. Holcomb III, MD, MBA. Surgeon-in-Chief/Senior Vice President, Department of Surgery, Children's Mercy Hospital, 2401 Gillham Road, Kansas City, MO 64108, USA. Email: gholcomb@cmh.edu.

Abstract: Repair of an indirect inguinal hernia is one of the most common operations performed around the world by pediatric surgeons. Until the last 15 years, most inguinal hernia repairs were performed using an inguinal crease incision and extraperitoneal ligation of the patent processes vaginalis. However, since 2000, the laparoscopic approach has gained popularity and there have been increasing descriptions about various techniques for laparoscopic hernia (LH) repair. At our institution, we have transitioned the majority of inguinal hernia repairs to the laparoscopic approach. In this article, we will describe the technique that is utilized at Children's Mercy Hospital in Kansas City, Missouri (USA) and express our thoughts on the current debate regarding laparoscopic versus open inguinal hernia repair in infants and children.

Keywords: Laparoscopy; indirect inguinal hernias; infants and children

Submitted Sep 12, 2016. Accepted for publication Sep 26, 2016.

doi: 10.21037/tp.2016.10.03

View this article at: <http://dx.doi.org/10.21037/tp.2016.10.03>

Repair of an indirect inguinal hernia is one of the most common operations performed around the world by pediatric surgeons. Until the last 15 years, most repairs of an indirect inguinal hernia were performed via an inguinal crease incision and an extraperitoneal repair. Since 2000, the laparoscopic approach has gained popularity, and there have been increasing descriptions about the laparoscopic technique. In fact, the majority of studies published in the last 20 years have focused on the laparoscopic approach. There have been a number of case series describing an author's or institution's experience with the laparoscopic approach as well as several meta-analyses comparing the laparoscopic to the open technique (1-14). At our institution, we have transitioned the majority of inguinal hernia repairs to the laparoscopic approach. In this article, we will describe the technique that is utilized at Children's Mercy Hospital in Kansas City, Missouri (USA) and express our thoughts on the current debate regarding laparoscopic versus open inguinal hernia repair.

Technique

We begin by positioning the infant/child supine on the operative table with the head near the anesthesiologist. The patient is prepped from the xiphoid to the perineum. This allows for manual decompression of the scrotal or labial pneumoperitoneum at the conclusion of the case and drainage of a hydrocele if present. The operative surgeon stands to the patient's right while the assistant holds the camera from the left side. Pneumoperitoneum is established via a 5 mm cannula placed through the umbilicus using an open technique. Diagnostic laparoscopy is then performed. This approach allows for the visualization of both internal rings, provides a visual diagnosis and treatment, and eliminates wrong-side surgery. A 3 mm instrument is then introduced via a stab incision at the mid-axillary line and lateral to the umbilicus on either the patient's right or left, depending on surgeon preference (*Figure 1*, panel A). This instrument is manipulated by the surgeon's dominant

Figure 1 (A) This photograph shows the 5 mm cannula in the umbilicus through which the telescope has been introduced. It also shows a 2.7 mm grasping instrument (arrow) placed through the stab incision in the patient's left upper abdomen. The surgeon is performing a repair of an indirect right inguinal hernia with an extracorporeal technique, and his hand is introducing the awl into the abdominal cavity at the level of the internal ring; (B) 0.25% bupivacaine has been instilled through a 25-gauge needle to elevate the peritoneum away from the vas deferens and testicular vessels in the preperitoneal plane; (C) the non-absorbable braided suture has already been passed around the medial aspect of the right internal ring and is now being threaded through the eyelet in the awl, which has been placed around the lateral aspect of the right internal ring; (D) this laparoscopic view shows the indirect right inguinal hernia has been closed in this patient.

right hand. A second instrument can be inserted on the contralateral side of the abdomen if necessary to allow for the camera operator to assist in the operation. Cosmetically these stab incisions have recently been shown to approach invisibility to the observer (15). It has been shown experimentally that peritoneal injury intentionally created at the site of closure results in a more durable repair (16). Therefore, prior to ligation, we cauterize and mechanically disrupt the peritoneum around the internal opening that is not adjacent to the vessels and vas deferens. At the level of the internal ring, a 25-gauge needle is passed through the abdominal wall at approximately the 2 or 10 o'clock position, depending on the side of the hernia, to hydrodissect the peritoneum off the vas and vessels in the pre-peritoneal plane. The needle is advanced to just lateral to these structures and fluid is instilled under the peritoneal

tissue to lift it off the underlying structures (*Figure 1B*). The fluid utilized depends on the patient's weight. As we are limited on the volume of 0.25% bupivacaine in patients under 10 kg, we will usually dissect with isotonic injectable saline. In patients over 10 kg, we often use 0.25% bupivacaine which is also used for postoperative anesthesia at the inguinal, umbilical, and instrument sites. Typically no more than 3–5 mL of fluid is needed to elevate the vas and vessels from the peritoneal tissue.

We then close the internal ring percutaneously. Several variations of this technique, which involves placing a suture circumferentially around the internal ring, have been described (6–10). A 2 mm stab incision through the skin and fascia overlying the internal ring is created in order to place a circumferential suture around the internal ring. A non-absorbable braided suture is passed through the eyelet

Figure 2 This schematic drawing depicts the zygomatic arch awl which is used at our hospital for laparoscopic repair of indirect inguinal hernias.

of a zygomatic arch awl (*Figure 2*). The awl is used to pass the non-absorbable suture through the stab incision into the hydrodissected pre-peritoneal space around the medial portion of the internal ring, traversing over the vas and vessels, and finally piercing the peritoneum. The awl is then removed after the non-absorbable suture is pulled into the abdominal cavity with the 3 mm instrument. The awl is next introduced on the contralateral side of the internal ring and pierces the peritoneum just past the first suture. This overlap in sutures ensures that the entire circumference of the peritoneum at the internal opening is incorporated in the closure. The operative surgeon passes the tail end of the first suture through the eye of the awl (*Figure 1C*). The instrument is withdrawn and a single suture thereby circumferentially encircles the internal ring.

The assistant then compresses the scrotum or labia to try to evacuate any fluid or air. The assistant should also grasp the testis and ensure its proper location within the scrotum. The operating surgeon then ties the suture around the internal opening of the hernia sac, and ligates the hernia (*Figure 1D*). This is visualized with the laparoscope. The process is then repeated on the contralateral side, if necessary. The instruments are removed under visualization. The stab incisions are injected with bupivacaine (if not already done) and then closed with tissue adhesive or steri-strips. The umbilical site is anesthetized with bupivacaine

and then closed with absorbable sutures and a sterile dressing is applied. If a hydrocele was present prior to the operation and has not been evacuated, then it is aspirated with a large gauge needle and syringe before the patient is awakened from anesthesia.

Discussion

As previously mentioned, a number of surgeons (Schier, Yeung, Shalaby, Esposito, Ponsky, Montupet, among others) have described different techniques for the laparoscopic repair of an indirect inguinal hernia in an infant or child (1-11,16-22). The two approaches can be grouped as either extraperitoneal or intraperitoneal. The early reports were primarily intraperitoneal in nature. Montupet is credited with performing the first intracorporeal laparoscopic hernia (LH) repair in 1993 (14). His technique was initially published in the French literature (16). In 1999, Montupet and Esposito modified this technique by sectioning the periorificial peritoneum circumferentially, distal to the internal ring, before closing it with a purse string suture (18,19). In 1998, Felix Schier described a technique in girls in which “N” shaped sutures were placed on the periorificial peritoneum (21). In 2002, he described a three-center experience in 933 patients (22).

The extracorporeal approach has been described as well. CK Yeung in Hong Kong has been utilizing this approach for a number of years and has had excellent results (personal communication). Shalaby and Ponsky have also described their experience with this approach (8,12). These techniques are quite similar with the technique described in this article. Instead of using an awl, Shalaby uses a Reverdin needle. Interestingly, the use of cauterization to promote injury around the internal ring in the extracorporeal approach is conceptually similar to the sectioning of the peri-orificial peritoneum described by Montupet and Esposito with the intracorporeal approach (16,18). Our technique is a modification of the technique originally described by Patkowski *et al.* (10). The modifications include the use of hydrodissection to elevate the peritoneum away from the abdominal wall and core structures, and the use of an awl for passage of the non-absorbable braided suture. A third modification is the use of intentional peritoneal injury prior to closure of the hernia sac to help improve its durability.

At this time, there is no consensus as to the best technique for the laparoscopic repair. Benefits of the laparoscopic approach appear to be its consistency in that

the view seen by the surgeon is the same whether or not the hernia defect is small or large, whether or not incarceration has occurred and been reduced or not, whether it is a very premature infant or an older teenager. Thus, each surgeon is able to apply his/her technique to almost all patients presenting with an indirect inguinal hernia. In contrast, an understanding of the anatomy of the inguinal canal can be difficult, especially for young surgeons, and it can often be more difficult to manage an incarcerated inguinal hernia or a hernia in a very premature infant using the inguinal crease approach.

There have been several meta-analyses trying to compare the laparoscopic and open approaches for an indirect inguinal hernia (12,13,23,24). In one meta-analysis, 53 studies that described outcomes of the open hernia repair (OH) and the LH repair were reviewed (13). Several variables were evaluated including operative time, recurrence rate, and other complications. In that review, for an infant/child undergoing a unilateral hernia repair, there was no significant difference between the two approaches in regards to operative time. However, for bilateral disease, the laparoscopic approach was faster ($P=0.01$). There was also no significant difference between the two techniques regarding recurrence rate, but the rate of other complications (wound infection, hydrocele formation, iatrogenic cryptorchidism, testicular atrophy, among others) was higher for the open approach. In addition, in the laparoscopic series, it was noted that there was a contralateral patent processes vaginalis varying between 20–66% for those patients who were being operated on for a unilateral hernia. Finally, in this review, it was identified that an advantage of laparoscopy is the ability to treat rare hernias (femoral, etc.) if they are found.

One of the areas in which laparoscopy may have a definite advantage over the open operation is the patient with an incarcerated hernia (25). The bowel may be able to be reduced through external manual pressure on the inguinal canal and it should be easy to visualize the condition of the bowel at the time of reduction with the laparoscope. Also, it appears that repair of large indirect inguinal hernias in very small or premature babies may be advantageous using the laparoscopic approach (26). These operations can be difficult no matter which approach is utilized, but if a recurrence develops after the laparoscopic repair in these small infants, then it can be repaired at a later date using the open approach.

Three disadvantages of the laparoscopic approach include the fact that this technique utilizes a transabdominal

approach, whereas the open operation is extraperitoneal. Second, the small incisions that are utilized for the laparoscopic approach will be visualized above the underwear line as opposed to the traditional inguinal crease incision. However, in two recent studies, these incisions were barely visible 2 years after their operation (15,27). Finally, in those infants who present with incarceration and bowel distention, the bowel distention can be a hindrance to good (or even adequate) visualization to allow the surgeon to perform the LH repair.

As a general statement, at this time, the data are not convincing regarding whether or not one approach is more favorable versus the other. As more and more surgeons become facile with the laparoscopic approach, it may become the procedure of choice for inguinal hernia repair in infants and children. More data and prospective randomized trials are needed before we know for sure whether one approach is more advantageous than the other.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Yip KF, Tam PK, Li MK. Laparoscopic flip-flap hernioplasty: an innovative technique for pediatric hernia surgery. *Surg Endosc* 2004;18:1126-9.
2. Hassan ME, Mustafawi AR. Laparoscopic flip-flap technique versus conventional inguinal hernia repair in children. *JSLs* 2007;11:90-3.
3. Becmeur F, Philippe P, Lemandat-Schultz A, et al. A continuous series of 96 laparoscopic inguinal hernia repairs in children by a new technique. *Surg Endosc* 2004;18:1738-41.
4. Boo YJ, Han HJ, Ji WB, et al. Laparoscopic hernia sac transection and intracorporeal ligation show very low recurrence rate in pediatric inguinal hernia. *J Laparoendosc Adv Surg Tech A* 2012;22:720-3.
5. Zallen G, Glick PL. Laparoscopic inversion and ligation inguinal hernia repair in girls. *J Laparoendosc Adv Surg Tech A* 2007;17:143-5.
6. Spurbeck WW, Prasad R, Lobe TE. Two-year experience

- with minimally invasive herniorrhaphy in children. *Surg Endosc* 2005;19:551-3.
7. Oue T, Kubota A, Okuyama H, et al. Laparoscopic percutaneous extraperitoneal closure (LPEC) method for the exploration and treatment of inguinal hernia in girls. *Pediatr Surg Int* 2005;21:964-8.
 8. Shalaby RY, Fawy M, Soliman SM, et al. A new simplified technique for needlescopic inguinal herniorrhaphy in children. *J Pediatr Surg* 2006;41:863-7.
 9. Ozgediz D, Roayaie K, Lee H, et al. Subcutaneous endoscopically assisted ligation (SEAL) of the internal ring for repair of inguinal hernias in children: report of a new technique and early results. *Surg Endosc* 2007;21:1327-31.
 10. Patkowski D, Czernik J, Chrzan R, et al. Percutaneous internal ring suturing: a simple minimally invasive technique for inguinal hernia repair in children. *J Laparoendosc Adv Surg Tech A* 2006;16:513-7.
 11. Ostlie DJ, Ponsky TA. Technical options of the laparoscopic pediatric inguinal hernia repair. *J Laparoendosc Adv Surg Tech A* 2014;24:194-8.
 12. Ponsky TA, Nalugo M, Ostlie DJ. Pediatric laparoscopic inguinal hernia repair: a review of the current evidence. *J Laparoendosc Adv Surg Tech A* 2014;24:183-7.
 13. Esposito C, St Peter SD, Escolino M, et al. Laparoscopic versus open inguinal hernia repair in pediatric patients: a systematic review. *J Laparoendosc Adv Surg Tech A* 2014;24:811-8.
 14. Esposito C, Escolino M, Farina A, et al. Two decades of history of laparoscopic pediatric inguinal hernia repair. *J Laparoendosc Adv Surg Tech A* 2014;24:669-70.
 15. St Peter SD, Acher CW, Shah SR, et al. Parental and Volunteer Perception of Pyloromyotomy Scars: Comparing Laparoscopic, Open, and Nonsurgical Volunteers. *J Laparoendosc Adv Surg Tech A* 2016;26:305-8.
 16. Blatnik JA, Harth KC, Krpata DM, et al. Stitch versus scar--evaluation of laparoscopic pediatric inguinal hernia repair: a pilot study in a rabbit model. *J Laparoendosc Adv Surg Tech A* 2012;22:848-51.
 17. Montupet P, Esposito C, Roblot-Maigret B, et al. Le traitement de la hernie inguinale congénitale chez l'enfant par coelioscopie: Notes techniques. *J Coelochir* 1996;26:35-9.
 18. Esposito C, Montupet P. Laparoscopic treatment of recurrent inguinal hernia in children. *Pediatr Surg Int* 1998;14:182-4.
 19. Montupet P, Esposito C. Laparoscopic treatment of congenital inguinal hernia in children. *J Pediatr Surg* 1999;34:420-3.
 20. Esposito C, Montinaro L, Alicchio F, et al. Technical standardization of laparoscopic herniorrhaphy in pediatric patients. *World J Surg* 2009;33:1846-50.
 21. Schier F. Laparoscopic herniorrhaphy in girls. *J Pediatr Surg* 1998;33:1495-7.
 22. Schier F, Montupet P, Esposito C. Laparoscopic inguinal herniorrhaphy in children: a three-center experience with 933 repairs. *J Pediatr Surg* 2002;37:395-7.
 23. Yang C, Zhang H, Pu J, et al. Laparoscopic vs open herniorrhaphy in the management of pediatric inguinal hernia: a systemic review and meta-analysis. *J Pediatr Surg* 2011;46:1824-34.
 24. Alzahem A. Laparoscopic versus open inguinal herniotomy in infants and children: a meta-analysis. *Pediatr Surg Int* 2011;27:605-12.
 25. Nah SA, Giacomello L, Eaton S, et al. Surgical repair of incarcerated inguinal hernia in children: laparoscopic or open? *Eur J Pediatr Surg* 2011;21:8-11.
 26. Esposito C, Turial S, Escolino M, et al. Laparoscopic inguinal hernia repair in premature babies weighing 3 kg or less. *Pediatr Surg Int* 2012;28:989-92.
 27. Gasior AC, Knott EM, Holcomb GW 3rd, et al. Patient and parental scar assessment after single incision versus standard 3-port laparoscopic appendectomy: long-term follow-up from a prospective randomized trial. *J Pediatr Surg* 2014;49:120-2; discussion 122.

Cite this article as: Juang D, Fraser JD, Holcomb GW 3rd. The laparoscopic approach for repair of indirect inguinal hernias in infants and children. *Transl Pediatr* 2016;5(4):222-226. doi: 10.21037/tp.2016.10.03