Parent-Reported Bullying and Child Weight Gain between Ages 6 and 15

Angelina R. Sutin, PhD,¹ Eric Robinson, PhD,² Michael Daly, PhD,^{3,4} and Antonio Terracciano, PhD¹

Abstract

Background: Childhood bullying has long-term negative mental and physical health correlates, including weight gain and symptoms of depression. The purpose of this research is to examine whether bullying in the first year of school is associated with greater weight gain by early adolescence and whether adolescent depressive symptoms mediate this association.

Methods: Data were drawn from the Longitudinal Study of Australian Children. Children ($N=3929$) were measured every 2 years; BMI and waist circumference were available from ages 4 to 15. Parents reported on bullying at age 6. Children reported on their depressive symptoms at ages 12–13.

Results: Participants who weighed in the obese category at age 4 had an over 50% increased risk of being bullied in school at age 6. Being bullied at age 6 was associated with excess weight gain between ages 6 and 15, defined as either BMI or waist circumference. Depressive symptoms at age 12 partially explained the association between bullying and increases in adiposity. None of the associations varied by gender.

Conclusions: Similar to other forms of peer victimization, bullying early in school is associated with greater weight gain through early adolescence; depressive symptom is one mechanism that contributes to this association.

Keywords: childhood bullying; depression; LSAC; mechanism; weight gain

Many children have had the experience of being bullied at school. The CDC defines bullying as any unwanted aggressive behavior by another youth or group of youths who are not siblings or current dating partners that involves an observed or perceived power imbalance and is repeated multiple times or is highly likely to be repeated (p. 7¹). The CDC estimates that ~20% of school-aged children experience this form of peer victimization each year.² There are well-documented correlates of bullying. Children, who have been bullied, for example, are more prone to depressive symptoms,³ have academic difficulties,⁴ and are more likely to experience intimate partner violence in adolescent relationships.⁵ These correlates persist well into adulthood. Individuals, who were bullied in childhood, for example, have a greater risk of depression, attain less education, and are less likely to have a romantic partner four decades later.⁶

There are characteristics that make some children more vulnerable to being bullied. Body weight, in particular, has been associated consistently with risk of being bullied at

school.^{7,8} This association may not be unidirectional; peer victimization, such as bullying, is also associated with subsequent weight gain. Children, who experience bullying at age 11, for example, are at greater risk of obesity 4 years later,⁹ and adolescents with obesity who are bullied gain more weight over 4 years than those who are not bullied.³ Similar to other outcomes, the risk of obesity associated with childhood peer victimization extends to young adulthood¹⁰ and into middle age.¹¹ Studies to date, however, have not been able to explain why experiencing bullying may contribute to weight gain.

Poor psychological health is another common correlate of having been bullied. Children, who were bullied, for example, experience more depressive symptoms in adolescence¹² and are at greater risk of a diagnosis of depression in adulthood.⁶ Symptoms of depression are likewise one pathway that is associated with greater weight gain over time.¹³ As such, children who experience bullying may be at greater risk of subsequent weight gain over time because they experience more symptoms of depressed

¹Florida State University College of Medicine, Tallahassee, FL.

²Institute of Psychology, Health and Society, University of Liverpool, Liverpool, England.

³Behavioural Science Centre, University of Stirling, Stirling, Scotland.

⁴UCD Geary Institute, University College Dublin, Dublin, Republic of Ireland.

mood. Less research, however, has focused on identifying mechanisms of this association, such as depressive symptoms, relative to testing whether bullying is associated with subsequent weight gain.

The current research addressed the longitudinal relationship between bullying at the start of elementary school and weight gain between childhood and adolescence and whether depressive symptoms were one potential mechanism that accounted for this association. To address this question, we used a large longitudinal sample of Australian children measured every 2 years from ages 4–5 to 14–15. We take a developmental approach to examine whether weight status at age 4, before beginning elementary school, was associated with risk of being bullied during the first year of school at age 6, whether children who experienced bullying at age 6 gained more weight from age 6 to 15 than children who did not experience it, and whether depressive symptoms mediated the association between bullying and weight gain. Because previous research has suggested that the association between bullying and weight gain may vary by both gender^{3,10} and BMI category,³ we tested whether gender or weight category moderated these associations. We examined this process over a significant part of childhood (ages 6–15) and across multiple measures of adiposity (BMI, waist circumference).

Methods

Participants and Procedure

Participants were drawn from the Longitudinal Study of Australian Children (LSAC). The Ethics Committee at the Australian Institute of Family Studies approved the protocol and data collection in LSAC¹⁴; analyses of LSAC data were approved by the IRB at Florida State University. Families were recruited and consented into LSAC when the study child was four; families were reinterviewed every 2 years, with the most recent assessment at ages 14–15. Detailed information about sampling and recruitment can be found here.¹⁵ We selected the assessment at age 6 as the baseline for bullying because that was the first assessment when all children had been in school for at least a year (at age 6, 17 children were homeschooled and the parents of these children did not respond to the item on bullying at school). Child weight, height, and waist circumference were measured at each of the six waves of data collection, from ages 4 to 15. A total of 3929 children had the relevant data at age 6 to be included in the analyses. For the longitudinal analysis, there were 17,127 repeated assessments of child adiposity with an average of 2.81 ($SD=1.38$; range 1–5) assessments per child. All assessments were ~2 years apart. Missing data for 1054 children were primarily due to attrition between Waves 1 and 2 ($n=519$ families did not participate in the Wave 2 assessment) and missing data on one or more of the study variables ($n=535$: $n=378$ were missing household income, $n=113$ were missing bullying [$n=96$ were missing; $n=17$ were not asked the question because the child was homeschooled], $n=32$ were missing

child BMI at age 4, and $n=41$ were missing child BMI at age 6; the ns add to more than 535 because some participants were missing data on more than one variable). Compared with participants included in the analyses ($n=3929$), those excluded because of missing data ($n=1054$) were more likely to be from an indigenous ethnic group, were less likely to speak English in the home, and had a lower reported family income (all $ps<0.01$). There were no differences by gender, BMI, or whether the child had been bullied at age 6 between those included in the analysis and those who did not have complete data.

Measures

Bullied. As part of a self-completed questionnaire, parents were asked whether their child had been bullied at school in the last year (“In the last 12 months, has [name of study child] been bullied at school? ‘at school’ includes travel to and from school”). Parents responded yes or no to this question. Parents were not provided with the scientific definition of bullying, but rather relied on their own understanding of what it means to be bullied when responding to this item. Although parents tend to underreport bullying compared with child reports,¹⁶ the correlates of parent and child-reported bullying tend to be similar.¹⁷

Adiposity. During each in-home interview, trained staff measured the study child’s weight and height. Weight was measured with a glass bathroom scale and recorded to the nearest 50 grams. Height was measured with a portable stadiometer and recorded to the nearest 0.1 cm. The CDC growth charts¹⁸ were used to calculate age and sex-specific child BMI z-scores at each assessment, with increases in BMI z-scores denoting weight gain; raw BMI scores were also analyzed. Waist circumference was measured with a nonstretch dressmaker’s tape horizontally around the navel and recorded to the nearest 0.1 cm.

Depressive symptoms. At ages 12–13, children ($n=2756$) reported on their own depressive symptoms using the Short Mood and Feelings Questionnaire, which was developed to use in epidemiological surveys and has been found to correlate strongly with more in-depth assessments.¹⁹ Children rated 13 items about their mood (*e.g.*, miserable or unhappy) in the last 2 weeks on a scale with 1 = *true*, 2 = *sometimes*, and 3 = *not true*. Items were recoded to a scale that ranged from 0 (*not true*) to 2 (*true*) and summed to create an index of depressive symptoms (range 0–26).

Analytic Strategy

Logistic regression was used to predict the risk of having been bullied at age 6 from BMI weight category (underweight, overweight, and obese compared with normal weight), controlling for child gender, family income, child indigenous status, and language other than English spoken at home. Hierarchical linear modeling (HLM)²⁰ was used to model the trajectory of adiposity (raw BMI, BMI z-score, and waist circumference) between ages 6 and 15. HLM is a

Table 1. Descriptive Statistics of Study Variables

Variable	Percent/mean (SD)
Child sex (female)	49
Indigenous (yes)	3
Language other than English spoken in home (yes)	16
Household Income (scaled in 1000 seconds of Australian \$)	1649.09 (1077.21)
Bullied at school in the last year (yes)	32
Depressive symptoms at age 12	4.01 (5.32)
Child BMI (age 4)	
Underweight	5
Normal weight	75
Overweight	15
Obesity	5
N = 3929.	

flexible approach used to quantify within-individual change or growth trajectories that can accommodate variations in spacing and number of observations since the repeated assessments are used to estimate each individual's trajectory (Level 1) and those individual parameters are the basis of group estimates (Level 2). The repeated measures of adiposity were included at Level 1; bullying and the covariates were included at Level 2. A significant effect of bullying on the slope of adiposity would indicate that bullying was associated with greater weight gain. In this multilevel model, an association with the slope accounts for the differences in average weight across participants; we also controlled for the child's BMI at age 4 to account for differences in BMI associated with risk of being bullied. An interaction between bullying and gender was ad-

ded at Level 2 to test whether the associations varied by child gender.

We used standard bootstrapping procedures²¹ to test whether depressive symptoms mediated the association between bullying at age 6 and weight gain between ages 6 and 15. The model specified depressive symptoms at age 12 as the mediator between bullied at age 6 and weight gain between ages 6 and 15, again controlling for child gender, family income, child indigenous status, and language other than English spoken at home. For the bootstrapping procedure, cases were selected at random, with replacement, from the original sample of *N*, and the model estimated. Parameter estimates were saved from each bootstrap sample; the indirect effect was considered significant if the confidence interval around the distribution of estimates did not include zero. We tested whether this process was moderated by either BMI category at baseline or child gender. The mediation analyses were run in SPSS version 21.

Results

Approximately 30% of parents reported that their child had been bullied at school in the last year (Table 1). Compared with children who measured in the normal weight BMI category, children who measured in the obese BMI category at age 4 were ~55% more likely to have been bullied at school in the last year at age 6 [OR = 1.55, 95% confidence interval (CI) = 1.15–2.08] and children in the overweight BMI category were almost 40% more likely to have been bullied (OR = 1.38, 95% CI = 1.14–1.66); the underweight category was unrelated to risk of bullying (OR = 1.17, 95% CI = 0.85–1.62). This association was not moderated by gender.

The HLM analysis revealed that children who were bullied at school at age 6 weighed more on average (intercept) and gained more weight from ages 6 to 15 (slope) than children who were not bullied (Table 2). For example, the effect of bullying on the intercept indicated that children

Table 2. Hierarchical Linear Modeling Analysis of the Association between Bullying and Increases in Adiposity between Ages 6 and 15

Predictor	Raw BMI		BMI z-score		Waist circumference ^a	
	Intercept	Slope	Intercept	Slope	Intercept	Slope
Gender (female)	0.39 (0.07)**	0.10 (0.01)**	0.00 (0.02)	0.02 (0.00)**	-1.14 (0.18)**	-0.40 (0.03)**
Family income (scaled in 1000 seconds)	-0.13 (0.03)**	-0.02 (.00)**	-0.03 (0.01)**	0.00 (0.00)	-0.31 (0.09)**	-0.07 (0.01)**
Indigenous status (yes)	0.41 (0.20)*	0.11 (0.03)**	0.05 (0.06)	0.03 (0.01)**	1.52 (0.54)**	0.14 (0.08)
Language other than English	0.26 (0.09)**	-0.01 (0.01)	0.07 (0.03)*	0.00 (0.00)	0.44 (0.25)	0.00 (0.04)
Child age 4 BMI	1.31 (0.02)**	0.06 (0.00)**	0.65 (0.01)**	-0.02 (0.00)**	2.99 (0.06)**	0.11 (0.01)**
Parent-reported bullying at age 6	0.22 (0.07)**	0.05 (0.01)**	0.06 (0.02)*	0.01 (0.00)**	0.68 (0.20)**	0.06 (0.03)*

N = 3929.

^aResults were similar when also controlling for child height.

p* < 0.05; *p* < 0.01.

who were bullied had a nearly one-quarter point higher BMI on average and increased 0.45 more BMI points over the 9-year follow-up than children who were not bullied. The results were similar when adiposity was defined as raw BMI, BMI z-score, or waist circumference. The results for waist circumference were similar when controlling for child height at each assessment. These associations also held in supplemental analyses that controlled for parent BMI [$\beta_{\text{bullying}}=0.03$ (0.01), $p<0.01$, and $\beta_{\text{bullying}}=0.01$ (0.00), $p<0.05$, respectively, for raw and z-score BMI], with the exception of waist circumference [$\beta_{\text{bullying}}=0.02$ (0.03), *ns*], which is likely due to the reduced sample size and thus reduced power. There was no interaction between the BMI weight categories and bullying on subsequent weight gain, which indicated that the risk of weight gain associated with bullying was similar regardless of baseline weight. The results were also not moderated by gender, which indicated that the effects were similar for girls and boys (all interaction coefficients *ns*).

Parent reports of bullying at age 6 were associated with more symptoms of depression, as reported by the child at age 12 ($\beta=0.085$, $p<0.001$). These symptoms partially

mediated the relationship between bullying and increases in adiposity between ages 6 and 15. The results were again similar when adiposity was defined as raw BMI, z-score BMI, or waist circumference (Table 3). Depressive symptoms accounted for a modest amount of the association between bullying and weight gain (12% for raw BMI, 11% for z-score BMI, and 27% for waist circumference). Neither sex nor baseline weight category moderated these associations.

Discussion

Bullying is a relatively common experience among school children who have significant long-term correlates. The current study indicates that the child's weight at age 4 is a risk factor for this form of peer victimization by age 6 and that these experiences are associated with excess weight gain between childhood and adolescence. The present study further indicates that symptoms of depression may be one mechanism in the pathway from bullying to weight gain.

Bullying in childhood is associated with a range of negative outcomes, from poor performance in school⁴ to

Table 3. Depressive Symptoms as a Mediator between Bullied and Weight Gain between Ages 6 and 14

	Point estimate	SE	Bias-corrected 95% CI	Effect ratio
Raw BMI				
Bullied → depressive symptoms (IV to mediator, path a)	0.989**	0.212		
Depressive symptoms → age 14 BMI (mediator to DV, path b)	0.009**	0.002		
Bullied → age 14 BMI (total effect, path c)	0.075**	0.022		
Bullied → age 14 BMI (direct effect, path c')	0.066**	0.022		
Bullied → age 14 BMI (indirect effect through mediator)	0.009*	0.003	0.004–0.016	0.120
BMI z-score				
Bullied → depressive symptoms (IV to mediator, path a)	1.001**	0.212		
Depressive symptoms → age 14 z-BMI (mediator to DV, path b)	0.011**	0.003		
Bullied → age 14 z-BMI (total effect, path c)	0.104**	0.037		
Bullied → age 14 z-BMI (direct effect, path c')	0.092*	0.037		
Bullied → age 14 z-BMI (indirect effect through mediator)	0.011**	0.004	0.004–0.021	0.106
Waist circumference (WC)				
Bullied → depressive symptoms (IV to mediator, path a)	0.904**	0.213		
Depressive symptoms → age 14 WC (mediator to DV, path b)	0.110**	0.030		
Bullied → age 14 WC (total effect, path c)	0.367	0.339		
Bullied → age 14 WC (direct effect, path c')	0.268	0.340		
Bullied → age 14 WC (indirect effect through mediator)	0.099*	0.038	0.042–0.197	0.270

N=2756. Models are adjusted for baseline BMI, sex, indigenous status (yes vs. no), household income, and language other than English spoken at home (yes vs. no). Indirect effect coefficients are from a bootstrap analysis with *n*=5000 bootstrap samples.

* $p<0.05$, ** $p<0.01$.

CI, confidence interval; SE, standard error.

difficulties in romantic relationships.⁵ These associations are long lasting, with evidence of negative correlates at least four decades later.⁶ Health status, including weight gain, is not immune to the negative effects of bullying. Bullying in childhood and adolescence has been associated with excess weight gain over time²² and risk of obesity.⁹ The present findings extend this work to cover a significant part of childhood, from age 6 to 15, and to an alternative measure of adiposity (*i.e.*, waist circumference).

These findings also fit with the growing evidence base that shows that several types of peer and family victimization are associated with greater weight gain among both children and adults. Girls, who are labeled fat at age 10, for example, have a greater risk of obesity over the next decade than girls who are not given this label²³ and children labeled as overweight by their parents are at greater risk of future weight gain.²⁴ In adulthood, individuals who experience weight discrimination are at greater risk of weight gain²⁵ and obesity²⁶ over time. The stigmatization may even be internally generated; individuals who perceive themselves as overweight, either accurately or inaccurately, are at greater risk of weight gain and obesity over time.^{27,28} This evidence suggests that social and psychological stigmatization are implicated in weight gain and that these associations start early and persist across the life span.

There are a number of pathways through which bullying may lead to weight gain. In the present research, we tested depressive symptoms as one mechanism of this association. Consistent with previous research,¹² children who experienced bullying at the beginning of elementary school were more likely to report symptoms of depression in early adolescence. Consistent with expectations, the association between bullying and subsequent weight gain was mediated, in part, by depressive symptoms. This indirect effect through depression symptoms, however, was relatively modest, which suggests that there are other mechanisms that contribute to this association.

There may, for example, be both behavioral and physiological pathways from bullying to weight gain. The experience of bullying is stressful and it is associated with overeating as a strategy to cope with the stress.²⁹ Indeed, several forms of peer victimization have been associated with disordered eating. Adolescent girls teased because of their weight are more likely to have disordered eating in young adulthood,³⁰ and both appearance and weight-based teasing have been associated with attempts to limit food intake (*i.e.*, dietary restraint) and binge eating and/or purging behaviors.³¹ Weight-loss attempts through these unhealthy strategies tend to result in subsequent long-term weight gain.³² There may also be physiological mechanisms that account for the relationship between bullying and weight gain. Individuals who experience peer victimization have a stronger cortisol response to stressors³³ and more chronic inflammation.³⁴ Such physiological responses may increase risk of subsequent weight gain.³⁵

Previous studies on bullying and later outcomes, including weight gain, have generally found that the asso-

ciations may be moderated by gender.^{3,10} It is of note then that we found the association between bullying and subsequent weight gain was similar for both males and females, perhaps due to the younger age (age 6 vs. adolescence) of the reported bullying. It may be the case that bullying in adolescence is more detrimental for girls than boys, whereas it is equally harmful when experienced early in childhood. It may also be the difference between parent-reported and child-reported bullying. Parent reports of bullying are valid and predictive of emotional and behavioral outcomes, but provide different information than reports from the children themselves.¹⁷ Reports of bullying from children may show stronger gender differences with weight gain than reports from parents.

Children with obesity at age 4 had an ~50% increased risk of being bullied at age 6. This association is similar to other research that has found early childhood obesity to be a risk factor for subsequent bullying^{7,22} and that children and adolescents with obesity are more vulnerable to being bullied than children who weigh in the normal weight category.⁸ Of note, however, the results for subsequent weight gain were similar regardless of weight status, which indicated that the weight gain associated with bullying occurred regardless of the child's early weight status.

The present research had several strengths, including a large sample, repeated measurements of adiposity for each participant, and a measure of waist circumference, as well as BMI. There are also some limitations that could be addressed in future research. For example, we had parent-reported bullying, but not reports of bullying by the child. Although parent and child-reported bullying have similar correlates, there tends to be only a modest association between the two.¹⁷ In addition, parents were not given the definition of bullying, but rather relied on their lay understanding of what it means to be bullied. As such, it would be worthwhile in future research to get both child and parent reports using a more detailed measure to get a more precise assessment of bullying. In addition, we only tested depressive symptoms as one mediator between bullying and weight gain. Future research could test other potential behavioral (*e.g.*, disordered eating) and physiological (*e.g.*, inflammation) mediators of this relationship. Finally, we did not have information on why the child was bullied. For example, although we speculate that children with obesity who are bullied are bullied because of heavier body weight, we were not able to examine whether the bullying was related to the child's weight or another characteristic. Similar to experiences with discrimination, different forms of bullying may be associated with different outcomes, including weight gain. It would be worthwhile in future research to distinguish between bullying based on weight versus other reasons for bullying. Despite these limitations, the present research indicates that being a victim of bullying is associated with long-term implications for childhood weight: children who are bullied early in their time at school are at greater risk of excess weight gain into adolescence.

Acknowledgment

This research was supported by a grant from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (1R15HD083947) to Angelina R. Sutin. Eric Robinson was part funded by the Medical Research Council (United Kingdom).

Author Disclosure Statement

No competing financial interests exist.

References

- Gladden RM, Vivolo-Kantor AM, Hamburger ME, et al. *Bullying Surveillance Among Youths: Uniform Definitions for Public Health and Recommended Data Elements*, Version 1.0. Centers for Disease Control and Prevention, Atlanta, GA, 2014.
- CDC. Youth risk behavior surveillance—United States, 2013. *MMWR Surveill Summ* 2014;63:1–168.
- Adams RE, Bukowski WM. Peer victimization as a predictor of depression and body mass index in obese and non-obese adolescents. *J Child Psychol Psychiatry* 2008;49:858–866.
- Glew GM, Fan MY, Katon W, et al. Bullying, psychosocial adjustment, and academic performance in elementary school. *Arch Pediatr Adolesc Med* 2005;159:1026–1031.
- Hertz MF, Everett Jones S, Barrios L, et al. Association between bullying victimization and health risk behaviors among high school students in the United States. *J Sch Health* 2015;85:833–842.
- Takizawa R, Maughan B, Arseneault L. Adult health outcomes of childhood bullying victimization: Evidence from a five-decade longitudinal British birth cohort. *Am J Psychiatry* 2014;171:777–784.
- Lumeng JC, Forrest P, Appugliese DP, et al. Weight status as a predictor of being bullied in third through sixth grades. *Pediatrics* 2010;125:e1301–e1307.
- van Geel M, Vedder P, Tanilon J. Are overweight and obese youths more often bullied by their peers? A meta-analysis on the correlation between weight status and bullying. *Int J Obes* 2014;38:1263–1267.
- Sweeting H, Wright C, Minnis H. Psychosocial correlates of adolescent obesity, ‘slimming down’ and ‘becoming obese’. *J Adolesc Health* 2005;37:409.
- Mamun AA, O’Callaghan MJ, Williams GM, et al. Adolescents bullying and young adults body mass index and obesity: A longitudinal study. *Int J Obes* 2013;37:1140–1146.
- Takizawa R, Danese A, Maughan B, et al. Bullying victimization in childhood predicts inflammation and obesity at mid-life: A five-decade birth cohort study. *Psychol Med* 2015;45:2705–2715.
- Zwierzynska K, Wolke D, Lereya TS. Peer victimization in childhood and internalizing problems in adolescence: A prospective longitudinal study. *J Abnorm Child Psychol* 2013;41:309–323.
- Sutin AR, Zonderman A. Depressive symptoms are associated with weight gain among women. *Psychol Med* 2012;42:2351–2360.
- Sanson A, Nicholson J, Ungerer J, et al. Introducing the longitudinal study of Australian children. *LSAC Discussion Paper* 2002;1:1–60.
- Soloff C, Lawrence D, Johnstone R. Sample design. *LSAC Technical Paper No 1* 2005.
- Holt MK, Kantor GK, Finkelhor D. Parent/child concordance about bullying involvement and family characteristics related to bullying and peer victimization. *J Sch Violence* 2009;8:42–63.
- Shakoor S, Jaffee SR, Andreou P, et al. Mothers and children as informants of bullying victimization: Results from an epidemiological cohort of children. *J Abnorm Child Psychol* 2011;39:379–387.
- Must A, Anderson SE. Body mass index in children and adolescents: Considerations for population-based applications. *Int J Obes* 2006;30:590–594.
- Angold A, Costello EJ, Messer SC, et al. Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *Int J Methods Psychiatr Res* 1995;5:237–249.
- Raudenbush SW, Bryk AS. *Hierarchical Linear Models: Applications and Data Analysis Methods*, 2nd ed. Sage: Thousand Oaks, CA, 2002.
- Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* 2008;40:879–891.
- Qualter P, Murphy SM, Abbott J, et al. Developmental associations between victimization and body mass index from 3 to 10 years in a population sample. *Aggress Behav* 2015;41:109–122.
- Hunger JM, Tomiyama AJ. Weight labeling and obesity: A longitudinal study of girls aged 10 to 19 years. *JAMA Pediatr* 2014;168:579–580.
- Robinson E, Sutin AR. Parent perception of weight status and weight gain across childhood. *Pediatrics* 2016;137:e20153957.
- Jackson SE, Beeken RJ, Wardle J. Perceived weight discrimination and changes in weight, waist circumference, and weight status. *Obesity* 2014;22:2485–2488.
- Sutin AR, Terracciano A. Perceived weight discrimination and obesity. *PLoS One* 2013;8:e70048.
- Sutin AR, Terracciano A. Body weight misperception in adolescence and incident obesity in young adulthood. *Psychol Sci* 2015;26:507–511.
- Robinson E, Hunger JM, Daly M. Perceived weight status and risk of weight gain across life in US and UK adults. *Int J Obes* 2015;39:1721–1726.
- Olvera N, Dempsey A, Gonzalez E, et al. Weight-related teasing, emotional eating, and weight control behaviors in Hispanic and African American girls. *Eat Behav* 2013;14:513–517.
- Quick VM, McWilliams R, Byrd-Bredbenner C. Fatty, fatty, two-by-four: Weight-teasing history and disturbed eating in young adult women. *Am J Public Health* 2013;103:508–515.
- Menzel JE, Schaefer LM, Burke NL, et al. Appearance-related teasing, body dissatisfaction, and disordered eating: A meta-analysis. *Body Image* 2010;7:261–270.
- Neumark-Sztainer D, Wall M, Story M, et al. Dieting and unhealthy weight control behaviors during adolescence: Associations with 10-year changes in body mass index. *J Adolesc Health* 2012;50:80–86.
- Knack JM, Jensen-Campbell LA, Baum A. Worse than sticks and stones? Bullying is associated with altered HPA axis functioning and poorer health. *Brain Cogn* 2011;77:183–190.
- Copeland WE, Wolke D, Lereya ST, et al. Childhood bullying involvement predicts low-grade systemic inflammation into adulthood. *Proc Natl Acad Sci U S A* 2014;111:7570–7575.
- Adam TC, Epel ES. Stress, eating and the reward system. *Physiol Behav* 2007;91:449–458.

Address correspondence to:

Angelina R. Sutin, PhD

Florida State University College of Medicine

1115 W. Call Street

Tallahassee, FL 32306

E-mail: angelina.sutin@med.fsu.edu