Published in final edited form as:

Inorg Chem Front. 2016 May 1; 3(5): 711–717.

Rapid Synthesis of Redox-Active Dodecaborane B₁₂(OR)₁₂ Clusters Under Ambient Conditions

Alex I. Wixtrom^a, Yanwu Shao^a, Dahee Jung^a, Charles W. Machan^b, Shaunt N. Kevork^a, Elaine A. Qian^a, Jonathan C. Axtell^a, Saeed I. Khan^a, Clifford P. Kubiak^b, and Alexander M. Spokoyny^{a,*}

^aDepartment of Chemistry and Biochemistry, University of California, Los Angeles 607 Charles E. Young Drive East, Los Angeles, California 90095-1569

^bDepartment of Chemistry and Biochemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093-0358

Abstract

We have developed a fast and efficient route to obtain perfunctionalized ether-linked alkyl and benzyl derivatives of the *closo*-[B₁₂(OH)₁₂]²⁻ icosahedral dodecaborate cluster via microwave-assisted synthesis. These icosahedral boron clusters exhibit three-dimensional delocalization of the cage-bonding electrons, tunable photophysical properties, and a high degree of stability in air in both solid and solution states. A series of *closo*-[B₁₂(OR)₁₂]²⁻, *hypocloso*-[B₁₂(OR)₁₂]¹⁻ and *hypercloso*-[B₁₂(OR)₁₂]⁰ clusters have been prepared with reaction times ranging from hours to several minutes. This method is superior to previously reported protocols since it dramatically decreases the reaction times required and eliminates the need for inert atmosphere conditions. The generality of the new microwave-based method has been further demonstrated through the synthesis of several new derivatives, which feature redox potentials up to 0.6 V more positive than previously known B₁₂(OR)₁₂ cluster compounds. We further show how this method can be applied to a one-pot synthesis of hybrid, vertex-differentiated species B₁₂(OR)₁₁(OR) that was formerly accessible only via multi-step reaction sequence.

Graphical abstract

*Corresponding author. spokoyne@chem.ucla.edu.

Full experimental details and spectroscopic characterization details are provided in the supporting information. X-ray crystallographic data for **11** and **13** is provided in the CIF format. NMR spectra, mass spectra, IR spectra, cyclic voltammetry, XPS spectra, and EPR spectra are also provided. See DOI: 10.1039/x0xx00000x

Introduction

The existence of an icosahedral dodecaborate [B₁₂H₁₂]²⁻ cluster was first predicted by Lipscomb and co-workers in 1954.¹ In a subsequent theoretical molecular orbital-based approach published in 1955², Longuet-Higgins and Roberts predicted that such a cluster would only be stable as dianionic [B₁₂H₁₂]²⁻. A 1959 study by Shapiro and Williams³ suggested the possible formation of a [B₁₂H₁₂]²⁻ icosahedron, and in 1960 this cluster was first successfully isolated and characterized as a triethylammonium salt by Pitochelli and Hawthorne, albeit in a relatively low yield.⁴ Subsequent pioneering studies by Hawthorne, Knoth, Muetterties, and others initiated a new era in the field of boron cluster chemistry.^{5–19} Specifically, these groups have shown that [B₁₂H₁₂]²⁻ can be prepared on a large scale in a high yield (>90%)¹⁹ and undergo facile functionalization chemistry that parallels some properties of classical organic molecules (*e.g.* benzene).^{20–22} This was an exciting discovery, since previously many boron hydride clusters were perceived as highly unstable species prone to fast degradation by heat, acids, and bases. Conversely, [B₁₂H₁₂]²⁻ salts were shown to be stable in acids and bases, and were thermally stable as high as 810 °C with no observable decomposition.¹² Knoth and co-workers were the first to demonstrate the persubstitution of [B₁₂H₁₂]²⁻, producing halogenated [B₁₂F₁₂]²⁻, [B₁₂Cl₁₂]²⁻, [B₁₂Br₁₂]²⁻ and [B₁₂I₁₂]²⁻ derivatives.^{16,23} In the past two decades, persubstitution of [B₁₂H₁₂]²⁻ was improved with new synthetic methods and extended towards other functional groups including [B₁₂Me₁₂]²⁻ and [B₁₂(OH)₁₂]²⁻.^{24–28} Among the perfunctionalized derivatives synthesized, *closo*-[B₁₂(OH)₁₂]²⁻ is particularly appealing, as it is capable of undergoing further functionalization by forming ether, ester, carbonate, and carbamate linkages.^{27,29–31}

While controlled oxidation of the parent [B₁₂H₁₂]²⁻ anion leads to an irreversible cluster degradation forming a B–B linked dimer,¹⁷ several perfunctionalized variants have been previously observed to undergo reversible redox behavior.³² For example, Rupich reported that [B₁₂X₁₂]²⁻ could undergo a single electron oxidation to form a stable radical [B₁₂X₁₂]^{1•-},³³ which was recently isolated and structurally confirmed as the oxidized radical [B₁₂Cl₁₂]^{1•-} species by Knapp and co-workers.³⁴ Hawthorne and co-workers reported that the perfunctionalized B₁₂(OCH₂Ph)₁₂ cluster can exist in three distinct redox states accessible via two sequential and quasi-reversible one-electron oxidation reactions of the parent dianionic *closo* species.³⁵ The same group later showed that other benzyl and alkyl functionalized B₁₂(OR)₁₂ clusters can be synthesized, and all of these species exhibit similarly reversible redox behavior.²⁹ Schleid and co-workers subsequently showed that the

parent $[\text{B}_{12}(\text{OH})_{12}]^{2-}$ cluster can undergo a one-electron oxidation to form a stable radical $[\text{B}_{12}(\text{OH})_{12}]^{1-}$ species.³⁶ Interestingly, the redox potential of the ether-linked $\text{B}_{12}(\text{OR})_{12}$ species can be rationally tuned as a function of the O-bound substituent, reminiscent of many metal-based redox-active inorganic complexes.^{29,32,37} Unfortunately, all of the reported synthetic routes towards $\text{B}_{12}(\text{OR})_{12}$ clusters currently require either extremely long reaction times (weeks) or highly specialized high-pressure equipment. Furthermore, in all cases strict inert atmospheric conditions are also required for their synthesis.^{27,29,35}

Herein we report a rapid, scalable, and robust synthetic route to a wide range of perfunctionalized $\text{B}_{12}(\text{OR})_{12}$ cluster derivatives utilizing a bench-top microwave reactor. This technology has emerged over the past several decades and has been successfully employed in a large number of synthetic schemes³⁸ which necessitate the reaction heating above the boiling point of the solvent. Our work shows that the microwave-based method enables synthesis of perfunctionalized ether-linked boron clusters within minutes and does not require the use of inert atmosphere and rigorously dried solvents. We further show the synthetic utility of our method in the preparation of previously unknown $\text{B}_{12}(\text{OR})_{12}$ derivatives featuring highly oxidizing redox potentials as well as vertex-differentiated molecular architectures. Synthesized cluster species were all isolated and characterized using solution-based NMR and IR spectroscopic tools and mass spectrometry. Electron delocalization of the radical state in these species was further evaluated by electron paramagnetic resonance (EPR) spectroscopy and elucidated by X-ray photoelectron spectroscopy (XPS).

Results and Discussion

The tetrabutylammonium (TBA) salt of *closo*- $[\text{B}_{12}(\text{OH})_{12}]^{2-}$ ($\text{TBA}_2[\mathbf{1}]$) was chosen for use with our microwave synthesis due to its enhanced solubility in organic solvents compared to alkali metal salts of **1** (Figure 1).²⁹ The synthesis of $\text{TBA}_2[\mathbf{1}]$ was adapted from previously reported protocols by Hawthorne and co-workers (see SI for details).^{29,39,40}

Oxygen-free, anhydrous conditions (oven-dried glassware, dried and distilled solvents, nitrogen atmosphere) were initially employed for microwave-assisted syntheses of alkyl- and benzyl-functionalized *closo*- $[\text{B}_{12}(\text{OH})_{12}]^{2-}$ [**1**] ether-linked derivatives, due to previously described high-pressure and reflux-based methods necessitating stringent air and moisture free conditions. To our surprise, we discovered that microwave reactions utilizing previously reported conditions (benzyl bromide and $\text{TBA}_2[\mathbf{1}]$ in the presence of *N,N*-diisopropylethylamine (DIEA, Hünig's base) in acetonitrile) are driven at a much higher rate, resulting in quantitative formation of a mixture of charged 1-/2- TBA salts of **2** within 15 minutes at 140 °C, as indicated by *in situ* ^{11}B NMR spectroscopy. Specifically, no parent ^{11}B NMR resonance at $\delta = 18$ corresponding to the $[\text{B}_{12}(\text{OH})_{12}]^{2-}$ starting material is observed, and a singlet at $\delta = 16$ can be seen instead. Concomitant presence of $[\mathbf{2}]^{1-}$ radical species in the product mixture can be deduced from the diagnostic pink color of the solution and its measured signature EPR signal (G-factor = 2.008121). Oxidation of the reaction mixture using $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ in 90/10 ethanol/acetonitrile followed by column chromatography on silica gel produces the pure neutral cluster $[\mathbf{2}]^0$ in 63% yield (Figure 2). Oxidation can be conveniently monitored by ^{11}B NMR, where the fully oxidized cluster $[\mathbf{2}]^0$

exhibits a downfield resonance shift at δ 41.8. Overall, this represents a significant reduction in reaction duration from the originally reported 6 days and 4 hours required for reflux and high-pressure reactor methods, respectively, while retaining similar yield.²⁹

Encouraged by these results, we then optimized the microwave reaction times with allyl bromide and bromoethane reagents independently, and in both cases observed that complete substitution can be accomplished within 15 – 30 minutes at 140 °C. These compounds were isolated in their fully oxidized neutral form in a similar fashion to **[2]⁰**, (compounds **[3]⁰** and **[4]⁰**, respectively; Figure 3). Notably, previously reported high-pressure reactor synthesis of these species required 3 and 12 hours, respectively, for alkylation to occur at all twelve vertices, suggesting that the microwave-based method can be generally applied to several classes of ether-linked B₁₂(OR)₁₂ clusters and is potentially superior to the previously developed methods.²⁹

The relatively short perfunctionalization reaction times made possible by this new microwave technique prompted us to investigate whether it would be possible to utilize ambient synthetic conditions. Specifically, we hypothesized that the fast rate of product formation would outcompete the rate of degradation stemming from the presence of adventitious air and moisture during the synthesis. We therefore tested the synthesis of **2** using as-received non-dried acetonitrile (see SI) with the reagents added to a reaction vessel open to air. The open-air synthesis of **2** proceeded with full conversion in 15 minutes as indicated by ¹¹B NMR spectroscopy on the crude mixture. Following the normal work-up procedure (see SI), **[2]⁰** was isolated in a 63% yield, suggesting that rigorous exclusion of air and moisture is not necessary for this transformation. This open-air synthesis method was successfully used for all subsequent ether-based cluster syntheses reported in this work. We decided to further explore the scope of this transformation by using longer-chain alkyl substituents. Hexyl chain substitution required increased reaction times compared to the shorter ethyl substituent, yet persubstitution was still achieved within two hours, as opposed to 8 hours when using a high-pressure reactor²⁹ (isolated as **[5]⁰**, Figure 3). This increased reaction time likely stems from the increase in the length and size of the alkyl reagent affecting the kinetics of the reaction.^{41,42} To further probe the limits of the microwave-based method we tested hexene- and undecene-based electrophiles, and even with the requisite increase in reaction time to 7 and 8 hours respectively, persubstitution proceeded to full conversion. Neutral **[6]⁰** and **[7]⁰** were isolated in 43% and 28% yields, respectively, after oxidation and normal purification procedures (Figure 3). These derivatives have not been synthesized prior to this report, and their preparation illustrates how one can dramatically increase the size of these ether-based dodecaborate clusters via a direct linkage of large substituents featuring terminal olefins onto [B₁₂(OH)₁₂]²⁻.

Based on the success of the long-chain olefin-containing moieties, we investigated the compatibility of this system with less stable reagents by using ethyl 4-bromobutyrate with TBA₂[**1**]. Perfunctionalization with ethyl butyrate has been challenging using prior synthetic methods, requiring multiple-step sequential additions of the alkyl halide and Hünig's base for 20 days while being handled under inert atmosphere conditions.⁴³ However, utilizing microwave-assisted synthesis, the same product (Figure 3, **[8]⁰**) was obtained via a single

1.5 hour reaction, followed by oxidation with $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ overnight and purification with column chromatography on SephadexTM and silica gel.

Benzyl-substituted ether-based clusters can feature a high degree of electrochemical tunability as a function of the substituents attached to the aromatic ring.³⁷ Our method allows for the efficient synthesis of clusters containing both electron-rich (**9**) and electron-withdrawing (**10**) benzyl derivatives in yields matching previous methods but with significantly reduced reaction times (Figure 3). We were further intrigued by the possibility of extending the accessible electrochemical window for this class of compounds by utilizing benzyl halide precursors containing highly electron-withdrawing substituents. The perfunctionalized cluster featuring a trifluoromethyl (CF_3) group attached to the *para* position of the benzyl moiety was prepared using our method in 30 minutes, and following oxidation the isolated neutral compound [**11**]⁰ was obtained in 66% yield (Figure 3). The oxidation potential of [**11**]¹⁻/[**11**]⁰⁻ ($E_{1/2} = 0.56 \text{ V vs Fc/Fc}^+$) measured by cyclic voltammetry (CV) is particularly notable since it is higher than any reported $\text{B}_{12}(\text{OR})_{12}$ cluster to date (previously $0.09 \text{ V vs Fc/Fc}^+$).³⁷ Plotting the Hammett constants of various benzyl substituents⁴² versus the redox potentials of [$\text{B}_{12}(\text{OR})_{12}$] clusters perfunctionalized with these groups³⁷ (Figure 4) indicates the oxidation potential of these clusters can be rationally extended beyond the previously reported electrochemical window.

For example, according to the trend suggested by this Hammett plot, a *para*-nitrobenzyl-substituted cluster should exhibit a higher 1-/0 oxidation potential than **11** (Figure 3 and Figure 4).⁴² The perfunctionalized cluster **12** featuring *para*-nitro (NO_2) substituent attached to the benzyl was successfully synthesized using our method in 30 minutes, however, oxidation of the reaction mixture containing [**12**]^{2-/1-} with $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ did not produce any *hypercloso*-neutral species [**12**]⁰. Instead, the radical cluster species [**12**]¹⁻ was isolated as the only product in 67% yield (Figure 3). This is not surprising, given the predicted oxidation potential for the [**12**]¹⁻/[**12**]⁰ redox couple is more positive than the oxidizing strength of $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$.⁴⁴ Attempts to use stronger chemical oxidants (e.g. ceric ammonium nitrate) resulted in cluster degradation. Furthermore, insufficient solubility of [**12**]¹⁻ as a TBA salt precluded us from obtaining CV measurements for this derivative. Nevertheless, convinced we could expand the electrochemical window for $\text{B}_{12}(\text{OR})_{12}$ species featuring benzyl-based substituents, we turned our attention to a potentially more soluble compound containing a 3,5-*bis*(trifluoromethyl)benzyl group instead.

Using the microwave method described above, cluster **13** was synthesized in 30 minutes. Interestingly, unlike all of the synthesized clusters reported thus far, the post-microwave reaction mixture was colorless, which is a characteristic feature of the pure dianionic state [**13**]²⁻ for these clusters. The lack of color persisted even after column chromatography purification on silica gel in air. The identity of the pure isolated TBA salt of [**13**]²⁻ (73% yield) was validated by full spectroscopic characterization and mass spectrometry (see SI). Oxidation of [**13**]²⁻ with $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ did not produce the neutral cluster, rather the pure [**13**]¹⁻ radical species was isolated in 78 % yield, resulting in 56% net yield for [**13**]¹⁻ species starting from $\text{TBA}_2[\text{1}]$. Use of stronger chemical oxidants such as ceric ammonium nitrate (CAN) degraded the boron cage, producing a diagnostic ¹¹B NMR resonance at $\delta 20$ characteristic of borates.⁴⁷ Nevertheless, we were able to observe neutral cluster [**13**]⁰

electrochemically via infrared spectroelectrochemistry (IR-SEC) and cyclic voltammetry (CV) in CH_2Cl_2 (Figure 5, B, C). CVs of $[\mathbf{13}]^{1-}$ as the TBA salt in CH_2Cl_2 showed two quasi-reversible redox features at -0.05 V and 0.68 V vs Fc/Fc^+ , corresponding to the $2-/1-$ and $1-/0$ transitions, respectively. IR-SEC experiments where the applied potential was increased to more positive potentials incrementally on the TBA salt of $[\mathbf{13}]^{2-}$ showed subtle changes in the IR stretching modes for all three oxidation states of $[\mathbf{13}]$ around $1130\text{--}1140\text{ cm}^{-1}$ and $1200\text{--}1220\text{ cm}^{-1}$. The shift in these IR bands assigned to the B–O bond⁴⁸ to higher wavenumbers from $[\mathbf{13}]^{2-}/[\mathbf{13}]^{1-}/[\mathbf{13}]^0$ is consistent with those observed in the same region for the analogous *para*- CF_3 compound $[\mathbf{11}]^0$ which can be isolated in its neutral form via direct synthesis (see SI). The high oxidation potential for the $[\mathbf{13}]^{1-}/[\mathbf{13}]^0$ redox couple observed from cyclic voltammetry ($E_{1/2} = 0.70$ V vs Fc/Fc^+) is notable since it is the highest observed $1-/0$ oxidation potential for the $\text{B}_{12}(\text{OR})_{12}$ class of clusters reported to date³⁷ and is ~ 130 mV higher than the *para*- CF_3 benzyl cluster **11**. ^{19}F NMR spectroscopy provides another diagnostic handle on the oxidation state of this compound (Figure 5, D). ^{11}B NMR spectra for these clusters typically show a singlet around $\delta -14$ to -16 for the $2-$ state, though the $1-$ state is silent due to the presence of the paramagnetic radical (confirmed by EPR, Figure 5D (inset)). However, with ^{19}F NMR spectroscopy, a shift from a singlet resonance in $[\mathbf{13}]^{2-}$ at $\delta -63.36$ to a broad singlet at $\delta -63.24$ for $[\mathbf{13}]^{1-}$ was observed. This broadening is consistent with the paramagnetic nature of $[\mathbf{13}]^{1-}$, where F atoms are located far enough from the unpaired electron-carrying B_{12} -based cluster core to be resolved by ^{19}F NMR spectroscopy.

The structural parameters of the boron clusters featuring persubstituted vertices can exhibit significant distortions in the solid state as a function of the substituent and the redox state as determined by X-ray crystallographic studies. Specifically, $[\mathbf{2}]^{2-}$ exhibits nearly identical B–B bond distances ($1.781(4) - 1.824(4)$ Å) and angles ($\text{B–B–B } 107.798^\circ < \alpha < 109.229^\circ$) as expected for a perfect icosahedron, yet as the cluster is oxidized to the electron-deficient $[\mathbf{2}]^{1-}$ the structure expands and distorts slightly, with further distortion observed in the neutral state.³⁵ Additionally, the B–O bond lengths decrease as the cluster is oxidized from $[\mathbf{2}]^{2-}$, which contains the longest average B–O distances, to the neutral $[\mathbf{2}]^0$ state with the shortest B–O distances.³⁵ This observed trend of B–B bond lengthening, B–O bond contraction, and B–B–B angle distortion within the core as a function of cluster oxidation state is supported qualitatively by the crystal structures for neutral $[\mathbf{11}]^0$ and radical $[\mathbf{13}]^{1-}$ (Figure 6), which show comparable changes to those observed between $[\mathbf{2}]^{1-}$ and $[\mathbf{2}]^0$. Selected bond lengths and angles for $[\mathbf{11}]^0$ and $[\mathbf{13}]^{1-}$ are shown in Figure 6.

X-ray photoelectron spectroscopy (XPS) has been widely used to study oxidation states in inorganic compounds.^{49–52} We therefore decided to utilize this technique to further elucidate the oxidation state and effect of functionalization for these boron clusters. Boron XPS spectra for several representative clusters synthesized in our study (Figure 7) indicate a clear trend observed in the shift of B–B bond peak energies depending on the redox state of the functionalized cluster. The observed geometric distortion of these boron cluster icosahedra with oxidation from $2-$ to $1-$ and further to neutral species results in an increased B–B binding energy (Figure 7A). The nature of the substituents also produces a clear trend in these measurements, as compound **13** also exhibits an increase in B–B binding energy as the

cluster increases in oxidation state from $[13]^{2-}$ to $[13]^{1-}$ (Figure 7B), yet both are higher in energy than that of neutral cluster $[11]^0$. However, despite the change in oxidation potential and binding energy observed from the XPS data from substituent effects, the nature of the electron radical delocalization throughout the boron-based core remains consistent. A single, broad symmetric EPR signal centered between 3450 and 3500 Gauss was observed for all cluster species isolated in the radical form with g values ranging between 2.0079 and 2.0081 depending on the substituent (see SI). Due to the 3D delocalization of the single electron across the 12 boron nuclei comprising the cluster, there exist a large number of possible hyperfine couplings.³⁶ Overlap of these hyperfine couplings ultimately gives rise to the single broad line observed in the EPR spectra.

In addition to previously mentioned benefits such as shortened reaction durations and the lack of stringent requirements for inert reaction conditions, microwave-assisted synthesis allows for one-pot, single-step reactions that would otherwise require more elaborate protocols. Monosubstitution of a benzyl ligand followed by persubstitution of the remaining eleven vertices has previously required a lengthy process involving several steps,⁵³ whereas we demonstrate a one-pot approach enabled by our microwave-based method. For example, mixed-substituent $B_{12}(OEt)_{11}(OBn)$ cluster (**14**, Figure 8) can be formed in a single step simply by adding a stoichiometric amount of the desired reagents into a single reaction vessel.

This reaction was completed in 30 minutes, producing three distinct species as a mixture which were oxidized with $FeCl_3 \cdot 6H_2O$ as previously described and subsequently isolated via column chromatography on silica: the perfunctionalized ethyl cluster $[4]^0$, a small amount (<5%) of di-substituted $[B_{12}(OEt)_{10}(OBn)_2]^0$ clusters, and the desired $[14]^0$ in 18% yield. This method represents a significantly faster route to produce $[B_{12}(OR)_{11}(OR')]$ mixed-substituent clusters.⁵⁴

Conclusions

A rapid microwave-assisted synthetic route to perfunctionalized ether-linked $B_{12}(OR)_{12}$ clusters is disclosed and the robust nature of the technique demonstrated by the synthesis and characterization of multiple derivatives of **1**. For previously synthesized compounds, reaction duration was significantly reduced and prior requirements for oxygen-free and anhydrous reaction conditions were eliminated. Our method also allows for a unique one-pot synthesis of mixed-substituent clusters with good selectivity under the same open-air conditions. The cluster species described here maintain the attractive properties of earlier derivatives, behaving as redox-active cores which show delocalization of electrons throughout the entire 3D boron cage, while the new functional groups provide a significant expansion to the available tuneable redox potential window for this class of clusters.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge Department of Chemistry and Biochemistry at UCLA for start-up funds. Portions of this material are based upon work supported by the National Science Foundation (CHE-1048804). E.A.Q. acknowledges the USPHS of the National Institutes of Health (NIH) for the predoctoral training fellowship through the UCLA Chemistry-Biology Interface Training Program under the National Research Service Award (T32GM008496). C.W.M. and C.P.K. acknowledge support for this work from the AFOSR through a Basic Research Initiative (BRI) grant (FA9550-12-1-0414). We would like to thank Mr. Kent Kirlikovali for assisting with EPR sample preparation and Mr. Rafal Dziedzic for assistance with DART mass spectrometry. We thank Dr Liban Saleh for commenting on the manuscript.

Notes and references

1. Eberhardt WH, C B Jr, Lipscomb WN. *J Chem Phys.* 1954; 22:989–1001.
2. Longuet-Higgins HC, de V Roberts M. *Proc R Soc A Math Phys Eng Sci.* 1955; 230:110–119.
3. Shapiro I, Williams RE. *J Am Chem Soc.* 1959; 81:4787–4790.
4. Pitochelli AR, Hawthorne FM. *J Am Chem Soc.* 1960; 82:3228–3229.
5. Moore EB, Lohr LL, Lipscomb WN. *J Chem Phys.* 1961; 35:1329.
6. Hoffmann R, Lipscomb WN. *J Chem Phys.* 1962; 36:2179.
7. Hoffmann R, Lipscomb WN. *J Chem Phys.* 1962; 37:520.
8. Muetterties EL, Merrifield RE, Miller HC, Knoth WH, Downing JR. *J Am Chem Soc.* 1962; 84:2506–2508.
9. Hoffmann R, Lipscomb WN. *J Chem Phys.* 1962; 37:2872.
10. Miller HC, Miller NE, Muetterties EL. *J Am Chem Soc.* 1963; 85:3885–3886.
11. Ellis IA, Gaines DF, Schaeffer R. *J Am Chem Soc.* 1963; 85:3885–3885.
12. Muetterties EL, Balthis JH, Chia YT, Knoth WH, Miller HC. *Inorg Chem.* 1964; 3:444–451.
13. Hertler WR. *Inorg Chem.* 1964; 3:1195–1196.
14. Hertler WR, Raasch MS. *J Am Chem Soc.* 1964; 86:3661–3668.
15. Miller HC, Miller NE, Muetterties EL. *Inorg Chem.* 1964; 3:1456–1463.
16. Knoth WH, Miller HC, Sauer JC, Balthis JH, Chia YT, Muetterties EL. *Inorg Chem.* 1964; 3:159–167.
17. Wiersema RJ, Middaugh RL. *Inorg Chem.* 1969; 8:2074–2079.
18. Wunderlich JA, Lipscomb WN. *J Am Chem Soc.* 1960; 82:4427–4428.
19. (a) Sivaev IB, Bregadze VI, Sjöberg S. *Collect Czechoslov Chem Commun.* 2002; 67:679–727.(b) Davidson M. *Contemporary boron chemistry*, Royal Society of Chemistry. 2000; 253(c) Hosmane, NS. *Boron Science: New Technologies and Applications*. Taylor & Francis; 2011. (d) Knapp, C. *Comprehensive Inorganic Chemistry II*. Elsevier; 2013. p. 651–679.(e) Olid D, Núñez R, Viñas C, Teixidor F. *Chem Soc Rev.* 2013; 42:3318. [PubMed: 23318646] (f) Geis V, Gutsche K, Knapp C, Scherer H, Uzun R. *Dalt Trans.* 2009:2687.
20. Grimes RN. *J Chem Ed.* 2004; 81:657.
21. Hawthorne MF. *J Chem Ed.* 2009; 86:1131.
22. Spokoiny AM. *Pure Appl Chem.* 2013; 85:903–919.
23. Knoth WH, Miller HC, England DC, Parshall GW, Muetterties EL. *J Am Chem Soc.* 1962; 84:1056–1057.
24. Gu W, Ozerov OV. *Inorg Chem.* 2011; 50:2726–2728. [PubMed: 21361302]
25. Peryshkov DV, Popov AA, Strauss SH. *J Am Chem Soc.* 2009; 131:18393–18403. [PubMed: 19954188]
26. Ivanov SV, Miller SM, Anderson OP, Solntsev KA, Strauss SH. *J Am Chem Soc.* 2003; 125:4694–4695. [PubMed: 12696872]
27. Hawthorne MF. *Pure Appl Chem.* 2003; 75:1157–1164.
28. Peymann T, Herzog A, Knobler CB, Hawthorne MF. *Angew Chem, Int Ed.* 1999; 38:1061–1064.
29. Farha OK, Julius RL, Lee MW, Huertas RE, Knobler CB, Hawthorne MF. *J Am Chem Soc.* 2005; 127:18243–18251. [PubMed: 16366578]

30. Maderna A, Knobler CB, Hawthorne MF. *Angew Chem, Int Ed.* 2001; 40:1661–1664.
31. Jalisatgi SS, Kulkarni VS, Tang B, Houston ZH, Lee MW, Hawthorne MF. *J Am Chem Soc.* 2011; 133:12382–12385. [PubMed: 21766843]
32. Kaim W, Hosmane NS, Zálaiš S, Maguire JA, Lipscomb WN. *Angew Chem, Int Ed.* 2009; 48:5082–5091.
33. Rupich MW. *J Electrochem Soc.* 1985; 132:119.
34. Boeré RT, Kacprzak S, Keßler M, Knapp C, Riebau R, Riedel S, Roemmele TL, Rühle M, Scherer H, Weber S. *Angew Chem, Int Ed.* 2011; 50:549–552.
35. Peymann T, Knobler CB, Khan SI, Hawthorne MF. *Angew Chem, Int Ed.* 2001; 40:1664–1667.
36. Van N, Tiritiris I, Winter RF, Sarkar B, Singh P, Duboc C, Muñoz-Castro A, Arratia-Pérez R, Kaim W, Schleid T. *Chem - Eur J.* 2010; 16:11242–11245. [PubMed: 20726020]
37. Lee MW, Farha OK, Hawthorne MF, Hansch CH. *Angew Chem, Int Ed.* 2007; 46:3018–3022.
38. (a) de la Hoz A, Diaz-Ortiz A, Moreno A. *Chem Soc Rev.* 2005; 34:164. [PubMed: 15672180] (b) Baig RBN, Varma RS. *Chem Soc Rev.* 2012; 41:1559–1584. [PubMed: 22076552] (c) Gawande MB, Shelke SN, Zboril R, Varma RS. *Acc Chem Res.* 2014; 47:1338–1348. [PubMed: 24666323] (d) de la Hoz, A.; Loupy, A. *Microwaves in Organic Synthesis*, 2 Volume Set. Wiley; 2013. (e) Kappe, CO.; Dallinger, D.; Murphree, SS. *Practical Microwave Synthesis for Organic Chemists*. Wiley-VCH Verlag GmbH & Co KGaA; Weinheim, Germany: 2008.
39. Peymann T, Knobler CB, Khan SI, Hawthorne MF. *J Am Chem Soc.* 2001; 123:2182–2185. [PubMed: 11456863]
40. Bayer MJ, Hawthorne MF. *Inorg Chem.* 2004; 43:2018–2020. [PubMed: 15018524]
41. Taft RW. *J Am Chem Soc.* 1953; 75:4231–4238.
42. Hansch C, Leo A, Taft RW. *Chem Rev.* 1991; 91:165–195.
43. Pushechnikov A, Jalisatgi SS, Hawthorne MF. *Chem Commun.* 2013; 49:3579–81.
44. Connelly NG, Geiger WE. *Chem Rev.* 1996; 96:877–910. [PubMed: 11848774]
45. Zavarine IS, Kubiak CP. *J Electroanal Chem.* 2001; 495:106–109.
46. Machan CW, Sampson MD, Chabolla SA, Dang T, Kubiak CP. *Organometallics.* 2014; 33:4550–4559.
47. Salentine CG. *Inorg Chem.* 1983; 22:3920–3924.
48. Leites LA. *Chem Rev.* 1992; 92:279–323.
49. Alov NV. *J Anal Chem.* 2005; 60:431–435.
50. Temesghen W, Sherwood P. *Anal Bioanal Chem.* 2002; 373:601–608. [PubMed: 12185571]
51. Zhu XP, Yukawa T, Hirai M, Suematsu H, Jiang W, Yatsui K, Nishiyama H, Inoue Y. *Appl Surf Sci.* 2006; 252:5776–5782.
52. Xing MY, Li WK, Wu YM, Zhang JL, Gong XQ. *J Phys Chem C.* 2011; 115:7858–7865.
53. Goswami LN, Houston ZH, Sarma SJ, Li H, Jalisatgi SS, Hawthorne MF. *J Org Chem.* 2012; 77:11333–11338. [PubMed: 23164043]
54. Recent examples of other mixed substituted dodecaborate clusters:(a) Bondarev O, Khan AA, Tu X, Sevryugina Y, Jalisatgi SS, Hawthorne MF. *J Am Chem Soc.* 2013; 135:13204–13211. [PubMed: 23919884] (b) Jenne C, Kirsch C. *Dalton Trans.* 2015; 44:13119–13124. [PubMed: 26107425] (c) Zhang Y, Liu J, Duttwyler S. *Eur J Inorg Chem.* 2015; 31:5158–5162.

Figure 1.

Synthetic route to produce functionalized ether-linked derivatives of the *hypercloso* boron clusters (**2–14**) via microwave-assisted synthesis. Precursor synthesis of [**1**] $^{2-}$ was adapted from previously described methods (see SI for details).^{29,39,40}

Figure 2.

Synthesis of **2** from TBA₂[**1**] using microwave-based method along with two previously reported methods indicating dramatic reduction in reaction time without compromising the isolated yield of [**2**]⁰.²⁹

Figure 3.

Synthesized $B_{12}(OR)_{12}$ clusters via microwave-based method. Yields are reported for the species isolated in a designated oxidation state as an average of two independent trials. For previously synthesized species, reported yield is given for comparison.²⁹ * Additional 8% of 1-/2- non-oxidized species collected.

Figure 4.

(a) Redox potential of $[B_{12}(OR)_{12}]^{2-/1-}$ and (b) $[B_{12}(OR)_{12}]^{1-/0}$ substituted with various benzyl substituents plotted vs Hammett constants.^{22,42} Previously characterized³⁷ (black) and new (red) $[B_{12}(OR)_{12}]^0$ clusters are shown.

Figure 5.

(a) Reversible redox activity of **13**; (b) Cyclic voltammogram (CV) demonstrating two independent, one-electron oxidation/reduction waves between 2⁻/1⁻, and 1⁻/0 states (1 mM **13** with 0.1 M TBAPF₆ in CH₂Cl₂; glassy carbon working electrode, Pt wire counter electrode, Ag/AgCl pseudoreference electrode behind a CoralPor tip; referenced to an internal ferrocene standard); (c) Infrared spectroelectrochemical (IRSEC)^{45,46} analysis of 0/1⁻/2⁻ states of **13**; (d) ¹⁹F NMR spectra of [13]¹⁻ and [13]²⁻ with EPR of [13]¹⁻ (inset); (e) UV-Vis spectra of [13]¹⁻ and [13]²⁻.

Figure 6.

(A–B) Solid state X-ray structures for $[13]^{1-}$ and $[11]^0$ shown with 50% thermal ellipsoid probabilities for the boron atoms (hydrogen atoms omitted for clarity). B atoms are blue, O – red, C – black, and F – green. Selected bond lengths and angles of the boron cores (substituents omitted for clarity) for $[11]^0$ and $[13]^{1-}$ are shown on the bottom. Overlay of the two cores ($[13]^{1-}$ in blue, $[11]^0$ in red) is depicted in the middle.

Figure 7.

(a) Boron XPS spectra for $TBA_2[1]^{2-}$, $TBA[12]^{1-}$, and $[11]^0$ showing an increase in B–B bond energy with increasing oxidation state; (b) Boron XPS spectra for $TBA[13]^{2-}$ and $TBA[13]^{1-}$ indicating the higher B–B bond energies of the $[13]^{2-}$ and $[13]^{1-}$ anions compared to the other substituted clusters.

Figure 8.

One-pot synthesis of vertex-differentiated *hypercloso* cluster [14]⁰ from an approximately 60:1:1 molar mixture of benzyl bromide:bromoethane:TBA₂[1]. Isolated yield of [14]⁰ was 18% (compound [4]⁰ was also formed as an additional product of the reaction). ¹¹B NMR spectra indicates loss of the icosahedral symmetry due to vertex differentiation, ¹H NMR integrations show 24H (CH₂) and 36H (CH₃) for the 11 ethyl groups and 2H (CH₂) and 5H (Ph) for the single benzyl moiety.