Cost-Effectiveness of HIV Preexposure Prophylaxis for People Who Inject Drugs in the United States

Ms. Cora L. Bernard, MS,

Department of Management Science and Engineering, Huang Engineering Center, 475 Via Ortega, Stanford University, Stanford, CA 94305-4026

Dr. Margaret L. Brandeau, PhD,

Department of Management Science and Engineering, Huang Engineering Center, 475 Via Ortega, Stanford University, Stanford, CA 94305-4026

Dr. Keith Humphreys, PhD,

Stanford School of Medicine, Department of Psychiatry, 401 N. Quarry Road, Room C-305, Stanford, CA 94305-5717

Dr. Eran Bendavid, MD, MS,

Center for Primary Care and Outcomes Research/Center for Health Policy, Stanford University, 117 Encina Commons, Room 204, Stanford, CA 94305-6019

Dr. Mark Holodniy, MD,

AIDS Research Center, VA Palo Alto Health Care System, 3801 Miranda Ave. (132), Palo Alto, California 94304-5107

Mr. Christopher Weyant, MS,

Requests for Single Reprints: Cora L. Bernard, MS, Department of Management Science and Engineering, Huang Engineering Center, 475 Via Ortega, Stanford University, Stanford, CA 94305-4026; clb210@stanford.edu. Current author addresses and author contributions are available at www.annals.org.

Disclaimer: The views expressed are those of the authors and do not necessarily reflect the opinion of the Department of Veterans Affairs or U.S. government.

Disclosures: Ms. Bernard reports grants from National Science Foundation and Stanford University PACCAR Inc. Graduate Fellowship during the conduct of the study. Dr. Brandeau reports grants from National Institute on Drug Abuse during the conduct of the study. Dr. Weyant reports grants from National Science Foundation during the conduct of the study. Dr. Owens reports grants from NIH during the conduct of the study. Dr. Goldhaber-Fiebert reports grants from NIH/NIA (K01 AG037593) during the conduct of the study. Authors not named here have disclosed no conflicts of interest. Disclosures can also be viewed at www.acponline.org/authors/icmje/ConflictsOfInterestForms.do?msNum=M15-2634.

Reproducible Research Statement: *Study protocol:* Not applicable. *Statistical code:* Additional detail can be made available where technical appendix is insufficient by contacting Cora Bernard (clb210@stanford.edu). *Data set:* Not applicable.

Author Contributions: Conception and design: E. Bendavid, C.L. Bernard, M.L. Brandeau, J.D. Goldhaber-Fiebert, K. Humphreys, D.K. Owens, C. Weyant.

Analysis and interpretation of the data: C.L. Bernard, M.L. Brandeau, J.D. Goldhaber-Fiebert, M. Holodniy, K. Humphreys, D.K. Owens, C. Weyant.

Drafting of the article: C.L. Bernard, J.D. Goldhaber-Fiebert, D.K. Owens.

Critical revision for important intellectual content: E. Bendavid, C.L. Bernard, M.L. Brandeau, J.D. Goldhaber-Fiebert, M. Holodniy, K. Humphreys, D.K. Owens.

Final approval of the article: E. Bendavid, C.L. Bernard, M.L. Brandeau, J.D. Goldhaber-Fiebert, M. Holodniy, K. Humphreys, D.K. Owens, C. Weyant.

Statistical expertise: C.L. Bernard, J.D. Goldhaber-Fiebert, D.K. Owens.

Obtaining of funding: M.L. Brandeau, J.D. Goldhaber-Fiebert, D.K. Owens.

Administrative, technical, or logistic support: D.K. Owens.

Collection and assembly of data: E. Bendavid, C.L. Bernard, D.K. Owens, C. Weyant.

Department of Management Science and Engineering, Huang Engineering Center, 475 Via Ortega, Stanford University, Stanford, CA 94305-4026

Dr. Douglas K. Owens, MD, MS, and

Center for Primary Care and Outcomes Research/Center for Health Policy , Stanford University, 117 Encina Commons, Stanford, CA 94305-6019

Dr. Jeremy D. Goldhaber-Fiebert, PhD

Goldhaber-Fiebert: Center for Primary Care and Outcomes Research/Center for Health Policy, Stanford University, 117 Encina Commons, Room 217, Stanford, CA 94305-6019

Abstract

Background—The total population health benefits and costs of HIV preexposure prophylaxis (PrEP) for people who inject drugs (PWID) in the United States are unclear.

Objective—To evaluate the cost-effectiveness and optimal delivery conditions of PrEP for PWID.

Design—Empirically calibrated dynamic compartmental model.

Data Sources—Published literature and expert opinion.

Target Population—Adult U.S. PWID.

Time Horizon—20 years and lifetime.

Intervention—PrEP alone, PrEP with frequent screening (PrEP+screen), and PrEP+screen with enhanced provision of antiretroviral therapy (ART) for individuals who become infected (PrEP+screen+ART). All scenarios are considered at 25% coverage.

Outcome Measures—Infections averted, deaths averted, change in HIV prevalence, discounted costs (in 2015 U.S. dollars), discounted quality-adjusted life-years (QALYs), and incremental cost-effectiveness ratios.

Results of Base-Case Analysis—PrEP+screen+ART dominates other strategies, averting 26 700 infections and reducing HIV prevalence among PWID by 14% compared with the status quo. Achieving these benefits costs \$253 000 per QALY gained. At current drug prices, total expenditures for PrEP+screen+ART could be as high as \$44 billion over 20 years.

Results of Sensitivity Analysis—Cost-effectiveness of the intervention is linear in the annual cost of PrEP and is dependent on PrEP drug adherence, individual transmission risks, and community HIV prevalence.

Limitation—Data on risk stratification and achievable PrEP efficacy levels for U.S. PWID are limited.

Conclusion—PrEP with frequent screening and prompt treatment for those who become infected can reduce HIV burden among PWID and provide health benefits for the entire U.S. population, but, at current drug prices, it remains an expensive intervention both in absolute terms and in cost per QALY gained.

Primary Funding Source—National Institute on Drug Abuse.

People who inject drugs (PWID) account for a disproportionate share of the U.S. HIV burden (1). Although PWID make up less than 1% of the U.S. adult population, approximately 10% of new HIV infections are attributable to injection drug use alone or in combination with male-to-male sexual contact (2–4). Focusing HIV interventions on PWID may thus have an out-sized public health benefit.

Many trials have shown that daily oral preexposure prophylaxis (PrEP) can prevent heterosexual and same-sex transmission of HIV (5–7). The Bangkok Tenofovir Study, a randomized trial of PrEP, reported a 49% reduction in HIV infection among PWID in Thailand (8). The Centers for Disease Control and Prevention (CDC) revised their clinical practice guidelines in 2014 to recommend that PrEP be considered for any adult who injected drugs within the past 6 months and, additionally, shared injection equipment, enrolled in drug-dependence treatment, or was at increased risk for sexual transmission (9).

However, the population health benefits and costs of implementing a national program remain unclear. Currently, PrEP, a daily antiretroviral oral pill, is expensive. The U.S. Food and Drug Administration approved the combination of 300 mg of tenofovir disoproxil fumarate (TDF) and 200 mg of emtricitabine (FTC) (Truvada, Gilead Sciences), which alone costs approximately \$10 000 annually (10–12). The CDC's recommended package of preventive care around PrEP, including HIV screening and assessment for adverse effects every 3 months and monitoring for toxicities every 6 months, adds to the cost. Although prior studies have explored the cost-effectiveness of PrEP for men who have sex with men (MSM) (10, 11), PWID differ in risk behaviors and HIV incidence. Therefore, we performed a model-based evaluation of the cost-effectiveness of expanding PrEP for PWID in the United States. We incorporated new clinical trial results with epidemiologic and economic data to determine the optimal conditions under which PrEP can be delivered to this high-risk population.

Methods

Overview

We developed an empirically calibrated dynamic compartmental model of the U.S. HIV epidemic (**Figure 1** of the **Supplement**, available at www.annals.org) to evaluate the benefits and costs of a national PrEP program for PWID. Our model captured sexual and injection transmission between PWID with and without PrEP, MSM, and all other U.S. adult heterosexuals (a generally low-risk group) between 2015 and 2035. Sexual mixing patterns between groups captured risk behavior overlaps, such as the PWID–MSM population (**Supplement**, section 4.1.2). The model included opioid agonist therapy (for example, methadone maintenance), HIV screening and awareness, and antiretroviral treatment (ART) both at current and enhanced levels. Table 1 and **Table 1** of the **Supplement** show model inputs and their sources. We calibrated the model to CDC estimates of HIV prevalence (4, 13–17), incidence (1, 17), and infection awareness (4, 13–15, 18) by risk group (**Supplement**, section 5). We adopted a societal perspective and computed total lifetime quality-adjusted life-years (QALYs) and health care costs, discounted at 3% annually (19, 20). We evaluated each scenario in terms of HIV infections averted and incremental cost-effectiveness ratios (ICERs).

Population

To capture the actively injecting population, the model follows U.S. adults aged 18 to 64 years stratified by risk group (21, 22) with underlying age mixing. We subdivided PWID by enrollment in opioid agonist therapy and subdivided all groups by HIV status and awareness of this status. To capture differences in infectivity, mortality, costs, and quality of life, we further stratified infected individuals by HIV stage and ART status (**Supplement**, section 1.1). Initial 2015 HIV prevalence in each risk group is consistent with CDC estimates (16, 17) and independent of age. Most individuals in the model are low-risk heterosexuals (1, 23): People who inject drugs make up 0.56% of the U.S. adult population (2, 3, 21), and MSM, 1.5% (11, 18). Consistent with national estimates, starting HIV prevalence is 9.8% in PWID (4), 11.4% in MSM (14), and 0.18% in the low-risk group (16, 23). We used calibration to determine the distribution of subgroups within each risk group (for example, the distribution of CD4 counts among PWID unaware of their HIV status) (**Supplement**, sections 1.2, 5).

Models

Reflecting the subpopulations described above, our dynamic compartmental model has 62 active compartments (**Supplement**, section 1). During a 20-year time horizon, a system of differential equations updates each compartment's population at monthly time steps (**Supplement**, section 4). Additional Markov models capture the life trajectories and health care costs of individuals maturing out of the model at age 65 years and those still active in the model at the end of the time horizon (**Supplement**, section 6). All models are programmed in Matlab R2013a (MathWorks).

Mortality

All individuals face background mortality rates plus additional rates reflecting risk status (11, 24, 25), HIV status, and treatments (**Supplement**, section 3.1.3) (18, 26–29).

Injecting Behavior and Treatment

Injecting behaviors increase health risks and HIV transmission. Opioid agonist therapy includes methadone and buprenorphine maintenance therapies, which reduce injections by 55% (30) and decrease mortality but do not affect risky sexual behavior (31, 32). We assumed that 25% of PWID receive opioid agonist therapy and assessed varying levels of enrollment in sensitivity analysis (**Supplement**, section 4.1.1) (4, 13, 31, 33, 34). The model is calibrated to U.S. HIV dynamics and hence reflects current levels of injection, including the effect of needle–syringe exchange programs.

Awareness of HIV Status and Subsequent Behavior

Infected individuals discover their status through HIV testing, with uptake of testing depending on risk group and, for PWID, on enrollment in opioid agonist therapy (35, 36). We assumed that detection is more likely in later, more symptomatic HIV stages (29). Recent U.S. guidelines recommend ART for all HIV-infected patients, regardless of CD4 count (37). There is no estimate for how this will affect future viral suppression rates for high-risk populations; consequently, we calibrated the model to reflect current ART

enrollment levels (18, 28, 33). To match current ART uptake data, after a positive HIV diagnosis, individuals initiate treatment and become virally suppressed with a probability dependent on their risk characteristics (28, 38). We also incorporated risk behavior change (for example, condom use) after an HIV diagnosis (11, 15, 33, 34).

HIV Transmission

Transmission occurs via sexual contact and via sharing injection equipment. Infection depends on 2 factors: frequency of risky contact based on mixing patterns and probability of infection given a serodiscordant contact (**Supplement**, section 2).

We assumed preferential sexual mixing (for example, PWID are more likely to have high-risk sexual partners) (**Supplement**, section 2.2.1) (4, 33, 34, 36, 39–42). After accounting for preferential mixing, we assumed proportional mixing (for example, no preference is given to choosing a partner of similar HIV status) (11, 33, 43, 44). We assumed that PWID aware of their HIV infection and PWID in opioid agonist therapy are less likely to share equipment (30, 32) but that, when they do, injection equipment sharing follows proportional mixing (13, 31, 33, 42, 43).

Sexual transmission between serodiscordant partners depends on male condom use (4, 14, 15, 33, 34, 42, 45), condom effectiveness (46), the infected partner's HIV stage (43, 47–49, 74) and ART status (48, 50), along with the uninfected partner's use of PrEP (8). We assumed a higher transmission risk for unprotected sexual acts that occur between MSM (47). Transmission through shared injection equipment depends on the infected partner's HIV stage (47, 48) and ART status (8, 50), along with the uninfected partner's use of PrEP (8). PrEP provides partial protection: a 49% reduction for both sexual and injection-based risk for HIV (8). We varied the efficacy of PrEP from 10% to 90% in sensitivity analysis to account for uncertainty in effectiveness and differential adherence patterns (5–9, 51).

HIV Infection and Treatment

An HIV-infected individual enters an acute phase and then progresses through asymptomatic (CD4 count, 0.50 to 1.20×10^9 cells/L) and symptomatic (0.20 to 0.50×10^9 cells/L) stages before developing AIDS ($< 0.20 \times 10^9$ cells/L). Without ART, the duration in each HIV stage is consistent with the literature (21, 29, 48, 49). ART suppresses HIV-1 viral load, which improves quality of life, decreases infectivity, and extends life expectancy (26–28). We assumed that ART reduces the chance of sexual transmission by 90% (43, 48, 50) and transmission from shared injection equipment by 60% (8, 43).

PrEP Scenarios

Per the CDC's guidelines, PrEP includes both a daily oral pill and an accompanying package of clinical care (for example, HIV screening every 3 months, toxicity monitoring every 6 months) (9). To assess how much of the benefit from a PrEP program accrues from each component, we evaluated 3 delivery scenarios: 1) PrEP in isolation with screening at the same rate as in the general PWID population; 2) PrEP with increased HIV screening and toxicity monitoring (PrEP+screen), which decreases the time between infection and diagnosis; and 3) PrEP+screen in which 50% of newly diagnosed PWID in the early stages

of HIV receive prompt, sustained ART (PrEP+screen+ART). (In contrast, only 10% of PWID not enrolled in PrEP+screen+ART receive ART in the early stages.) Enhanced ART delivery also yields indirect benefits because shortened time to diagnosis and treatment for a patient reduces HIV transmission.

Because of PWID's generally lower access to health care and retention in treatment (4, 38), all delivery scenarios assume that 25% of uninfected PWID would enroll in a PrEP program (9). PrEP is discontinued at the time of a positive HIV diagnosis. A PrEP program moderately increases HIV awareness for all PWID because uninfected and infected-but-unaware individuals are equally likely to be screened before enrollment in PrEP (9).

Economic Model

Depending on risk group and health status, different costs and QALYs accrue to individuals (**Supplement**, section 7.1). We derived costs, adjusted for inflation by using the Consumer Price Index (52) and expressed in 2015 U.S. dollars, from previous HIV models (11, 33, 43). PrEP program costs reflect the cost of TDF–FTC (Truvada) (12, 53) as well as of monitoring for PrEP+screen and PrEP+screen+ART (10, 11, 54). We assessed value by comparing the incremental discounted costs and QALYs of each PrEP delivery scenario to the next-best alternative (20).

Model Calibration

We obtained initial estimates and distributions for each model input from the published literature and then empirically calibrated the model to U.S. epidemiologic data, resulting in 182 best-fitting parameter sets (**Supplement**, section 5) (55, 56). We accounted for parameter uncertainty by repeating our analyses across these calibrated sets (57).

Role of the Funding Source

No funder had a role in the design, conduct, and analysis of the study or in the decision to submit the manuscript for publication.

Results

Figure 1 and Table 2 show that PrEP+screen+ART is more effective and less expensive per QALY gained than both PrEP alone and PrEP+screen, illustrating that prompt and sustained ART provision is crucial to delivering PrEP with maximal value. PrEP+screen+ART for 25% of uninfected PWID (approximately 280 000 individuals) averts 21 500 new HIV infections (–20.0%) and 5300 deaths from AIDS (–7.4%) within the PWID population over 20 years. For all risk groups, 26 700 HIV infections (–8.2%) and 6300 deaths from AIDS (–1.4%) are averted. We projected that PrEP+screen+ART could reduce HIV prevalence by 14% in the PWID population by 2035 (from 6.5% to 5.6%). These health effects translate to an increase of 173 000 QALYs. However, PrEP+screen+ART increases costs by \$44 billion over the period despite the averted costs from prevented HIV infections; this is approximately \$2.2 billion per year or 9% of the 2015 federal budget for domestic HIV/AIDS (58). Compared with the status quo, PrEP+screen+ART costs \$253 000/QALY gained.

Sensitivity Analysis

For the PrEP+screen+ART delivery scenario, we performed multiple one-way sensitivity analyses on all model parameters (**Supplement**, section 8.1), with key results presented below. Our findings are robust to parameter uncertainty, with 98% of instances in a probabilistic sensitivity analysis costing more than \$100 000 per QALY gained (**Figure 12** of the **Supplement**; **Supplement**, section 8.2).

Cost of PrEP

The drug costs of PrEP are based on the U.S. Food and Drug Administration–approved combination of TDF–FTC (Truvada) (10–12). Figure 1 illustrates that drug costs drive most of the costs accruing over the next 20 years in all PrEP scenarios and thus critically determine cost-effectiveness. (Section 3.5.2 of the **Supplement** discusses sensitivity around screening costs.) Truvada is available at substantially reduced cost abroad and even within the United States for qualifying uninsured patients (59); moreover, patents for Truvada will expire within 5 years, when a generic drug might become available (**Supplement**, section 3.5.1) (60). We evaluated the cost-effectiveness of PrEP+screen+ART over a range of drug costs (Table 3). At a 65% price reduction, the intervention costs approximately \$100 000 per QALY gained; nonetheless, it would cost \$17 billion to deliver over 20 years (60).

PrEP Efficacy

PrEP's efficacy in reducing HIV acquisition critically determines cost-effectiveness. Consistent with trial data, we assume a 49% (10% to 70%) reduction in sexual and injection transmission (8). PrEP trials in other populations estimate reductions in HIV acquisition via sexual transmission ranging from negligible to 92%, with efficacy depending strongly on adherence, which may be lower outside of study settings (5–9, 51). We evaluated PrEP with efficacy ranging from 10% to 90%, finding that the cost per QALY gained remains greater than \$150 000 (Figure 2 and **Table 2** of the **Supplement**). Nonetheless, cost-effectiveness improves substantially, especially at the lower end of the efficacy range, as adherence increases. This suggests an adherence threshold below which PrEP provides very little value. An important consideration is whether those who are less adherent may fill PrEP prescriptions less frequently, implying a correlation between lower efficacy and lower cost. In a 2-way sensitivity analysis, we find that low adherence levels, even if they result in low drug costs, provide low value, thus underscoring the conclusion that high value necessitates sufficient adherence (**Figure 2** of the **Supplement**).

Transmission

The magnitude of PrEP's health benefits is largely driven by an individual's risk for HIV infection without PrEP (**Figure 3** of the **Supplement**). A good proxy for an individual's HIV risk is community prevalence; given similar ART coverage levels, populations with higher concentrations of infection will have higher transmission rates. Although average HIV prevalence among U.S. PWID is around 10%, prevalence varies from 2% to 19% among Metropolitan Statistical Areas (4, 13), suggesting a similarly wide range of incidence and risk levels. Targeting PrEP to highest-prevalence (and therefore highest-incidence)

communities where other intervention resources are scarce will thus deliver greatest value, potentially costing less than \$100 000/QALY gained.

Coverage

The main analysis assumes 25% coverage as complete population coverage is unlikely in the near term (4, 38). **Figure 4** of the **Supplement** and **Table 3** of the **Supplement** show the results of alternate coverage scenarios. Though PrEP would be maximally beneficial in a scenario in which herd immunity could be achieved, within feasible levels of coverage and efficacy in the PWID population, PrEP+screen+ART delivers slightly less incremental benefit as coverage increases (8).

Discussion

The value of PrEP is maximal when PrEP is delivered with frequent HIV screening and enhanced ART provision for those who become infected. Over 20 years, enrolling 25% of uninfected PWID in PrEP+screen+ART would reduce HIV burden among PWID and provide health benefits for the entire U.S. population, averting 26 700 new HIV infections and thereby gaining 173 000 QALYs. Nonetheless, PrEP+screen+ART is an expensive intervention at current prices of TDF–FTC (Truvada), costing \$253 000 per QALY gained, well above generally accepted thresholds for cost-effectiveness. In comparison, needle–syringe exchange programs cost in the range of \$4500 to \$34 000 per QALY gained (61, 62). Our analysis also indicates that a PrEP program creates affordability challenges. At current drug costs, PrEP for 25% of PWID would cost an additional \$44 billion over 20 years. This is equivalent to annually spending 9% of the current federal budget for domestic HIV/AIDS on PrEP for PWID (58). We found cost-effectiveness and total budget outlays to be approximately linear in the cost of PrEP: Thus, if the release of a generic medication reduced the drug cost by 65%, then PrEP+screen+ART would cost approximately \$100 000 per QALY gained. Additionally, in settings with high HIV transmission risk, a PrEP program could cost less than \$100 000 per QALY gained, although budget impact in both cases would still be substantial. These large expenditures will create challenges for both public and private payers that cover PrEP.

Cost-effectiveness is only one of many considerations for policymakers, who must also evaluate the ethical dimensions of an HIV prevention program for a population with generally low access to health services (4). PrEP for PWID can deliver individual-level protection and population-level health benefits, both of which provide substantial value in an underserved and high-risk group. In fact, the benefit of all PrEP programs is highest for individuals at greatest risk for HIV. Although it may be difficult for a clinician to distinguish between a lower- and higher-risk PWID, our analysis suggests that transmission rates within a population are a good proxy for individual risk, and thus PrEP delivers greater value in PWID communities with high HIV prevalence (and thus high HIV incidence) (13). Because high-prevalence populations are often the most marginalized, targeting PrEP to localized epidemics can serve the dual purpose of maximizing its value while addressing pressing social inequities (63).

The Bangkok Tenofovir Study participants were in drug-dependence treatment programs and received modest reimbursement for their time, which raises the question of whether their reductions in HIV transmission would generalize to PWID who are not in treatment. As is now evident from prior PrEP trials, efficacy depends on adherence (51). Whether adherence in the United States would reach (or exceed) levels achieved in the trial has not yet been demonstrated. If adherence were lower, then PrEP would be less cost-effective than we have estimated. These questions highlight the importance of ongoing evaluation of PrEP programs for PWID as they are initiated.

Our study has several limitations. First, PrEP+screen+ART could be more cost-effective than estimated if delivered to individuals centrally located in injecting or sexual networks. It could be substantially less cost-effective if considered within a portfolio of cheaper interventions (**Supplement**, section 9). **Figure 14** of the **Supplement** illustrates the substantially higher ICER of PrEP relative to a needle-syringe exchange program.

Second, because we are evaluating the effect of a national program, our input parameters as well as our results reflect population-level averages. Where possible, we accounted for this in sensitivity analyses: For instance, we assessed the value of PrEP+screen+ART over a range of HIV-risk scenarios. The model examines PWID as one risk group, although PWID have differing levels of risk (**Supplement**, section 4.1). Two important dimensions along which PWID could be stratified are substance injected, which affects frequency of injecting and effectiveness of opioid agonist therapy, and sexual behavior (for example, PWID who are also MSM). Despite a lack of data to properly stratify the model in these ways, our findings do suggest that PrEP would deliver greater value for both of these higher-risk subpopulations. The potential value for PrEP in high-risk/high-prevalence communities underscores the importance of localized epidemiologic surveillance to inform both future models and policy.

Third, new ART initiation guidelines will likely increase the number of virally suppressed individuals in coming years (37). Without data on future changes, we calibrated the model to reflect current practice patterns. More individuals on ART, while increasing HIV prevalence because of the longer lifespan of treated individuals, would decrease the risk for HIV transmission, making other prevention programs seem less favorable. Thus, if anything, our estimate of a high cost per QALY gained for PrEP+screen+ART is biased favorably toward the intervention.

Fourth, a concern about PrEP is the facilitation of drug-resistant HIV. However, resistance did not occur in the Bangkok Tenofovir Study (8). Moreover, the CDC's frequent screening guidelines decrease the likelihood of drug-resistant mutations in practice (5, 8, 9, 64, 65).

Finally, although we address parameter uncertainty within the model, we cannot address structural uncertainties, such as the choice of risk categories or transmission pathways (66). This is a limitation of disease modeling and should be considered when our conclusions are being evaluated.

Our study assesses the total population health benefits and costs of PrEP for U.S. PWID within a dynamic transmission framework. Other studies have explored the cost-

effectiveness of a national PrEP program for MSM and heterosexuals, where efficacy estimates are higher than in the PWID population (5–8), and have found that the intervention is both expensive in absolute cost and most valuable in highest-risk subpopulations (10, 11). Similarly, we find that greater adherence and higher transmission risk increase the value of the intervention, but at current costs a substantial budgetary increase of \$44 billion would be required to deliver PrEP+screen+ART to 25% of PWID over 20 years. For similar cost, greater benefit can be delivered by combining PrEP and screening services with enhanced ART provision for those who do become infected, highlighting the importance of the CDC's recommendations regarding services to be administered with PrEP (9).

In conclusion, we find that PrEP programs provide optimal value when combined with frequent HIV screening and enhanced ART provision for those who become infected. PrEP +screen+ART for PWID in the United States can reduce HIV incidence and have health benefits for the entire U.S. population, but it is currently expensive both in absolute terms and in cost per QALY gained. The cost-effectiveness of PrEP+ screen+ART depends greatly on the efficacy and cost of PrEP, as well as transmission rates in the covered population. Thus, PrEP+screen+ART has increased value for PWID with high levels of adherence or for PWID most at risk for HIV.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Grant Support: By grant R01-DA15612 from the National Institute on Drug Abuse. Ms. Bernard was supported by a PACCAR Inc. Stanford Graduate Fellowship and a National Science Foundation Graduate Fellowship (DGE-114747). Christopher Weyant was supported by a Stanford Department of Management Science and Engineering Graduate Fellowship and a National Science Foundation Graduate Fellowship (DGE-114747). Drs. Owens, Humphreys, and Holodniy were supported by the Department of Veterans Affairs. Dr. Goldhaber-Fiebert was supported by a Career Development Award (K01AG037593-01A1) from the National Institute on Aging.

References

- Centers for Disease Control and Prevention. Estimated HIV incidence in the United States, 2007–2010. HIV Surveillance Supplemental Report 2012. 17(No. 4) [12 August 2014] Accessed at www.cdc.gov/hiv/pdf/statistics_hssr_vol_17_no_4.pdf.
- Friedman SR, Tempalski B, Cooper H, Perlis T, Keem M, Friedman R, et al. Estimating numbers of injecting drug users in metropolitan areas for structural analyses of community vulnerability and for assessing relative degrees of service provision for injecting drug users. *J Urban Health*. 2004; 81:377–400. [PMID: 15273263]. [PubMed: 15273263]
- Centers for Disease Control and Prevention (CDC). Integrated prevention services for HIV infection, viral hepatitis, sexually transmitted diseases, and tuberculosis for persons who use drugs illicitly: summary guidance from CDC and the U.S. Department of Health and Human Services. *MMWR Recomm Rep*. 2012; 61:1–40. [PMID: 23135062].
- Spiller MW, Broz D, Wejnert C, Nerlander L, Paz-Bailey G, Centers for Disease Control and Prevention (CDC). HIV infection and HIV-associated behaviors among persons who inject drugs—20 cities, United States, 2012. *MMWR Morb Mortal Wkly Rep*. 2015; 64:270–5. [PMID: 25789742]. [PubMed: 25789742]
- Grant RM, Lama JR, Anderson PL, McMahan V, Liu AY, Vargas L, et al. iPrEx Study Team. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med*. 2010; 363:2587–99. [PMID: 21091279]. DOI: 10.1056/NEJMoa1011205 [PubMed: 21091279]

6. Murnane PM, Celum C, Mugo N, Campbell JD, Donnell D, Bukusi E, et al. Partners PrEP Study Team. Efficacy of preexposure prophylaxis for HIV-1 prevention among high-risk heterosexuals: subgroup analyses from a randomized trial. *AIDS*. 2013; 27:2155–60. [PMID: 24384592]. DOI: 10.1097/QAD.0b013e3283629037 [PubMed: 24384592]
7. Baeten JM, Donnell D, Ndase P, Mugo NR, Campbell JD, Wangisi J, et al. Partners PrEP Study Team. Antiretroviral prophylaxis for HIV prevention in heterosexual men and women. *N Engl J Med*. 2012; 367:399–410. [PMID: 22784037]. DOI: 10.1056/NEJMoa1108524 [PubMed: 22784037]
8. Choopanya K, Martin M, Suntharasamai P, Sangkum U, Mock PA, Leethochawalit M, et al. Bangkok Tenofovir Study Group. Antiretroviral prophylaxis for HIV infection in injecting drug users in Bangkok, Thailand (the Bangkok Tenofovir Study): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet*. 2013; 381:2083–90. [PMID: 23769234]. DOI: 10.1016/S0140-6736(13)61127-7 [PubMed: 23769234]
9. Centers for Disease Control and Prevention. U.S. Public Health Service. [12 June 2014] Preexposure prophylaxis for the prevention of HIV infection in the United States—2014: A Clinical Practice Guideline. 2014. Accessed at www.cdc.gov/hiv/pdf/prepguidelines2014.pdf
10. Paltiel AD, Freedberg KA, Scott CA, Schackman BR, Losina E, Wang B, et al. HIV preexposure prophylaxis in the United States: impact on lifetime infection risk, clinical outcomes, and cost-effectiveness. *Clin Infect Dis*. 2009; 48:806–15. [PMID: 19193111]. DOI: 10.1086/597095 [PubMed: 19193111]
11. Juusola JL, Brandeau ML, Owens DK, Bendavid E. The cost-effectiveness of preexposure prophylaxis for HIV prevention in the United States in men who have sex with men. *Ann Intern Med*. 2012; 156:541–50. [PMID: 22508731]. DOI: 10.7326/0003-4819-156-8-201204170-00001 [PubMed: 22508731]
12. U.S. Department of Veterans Affairs. [1 April 2015] National Acquisition Center (CCST). Accessed at www1.va.gov/nac/
13. Broz D, Wejnert C, Pham HT, DiNenno E, Heffelfinger JD, Crib-bin M, et al. National HIV Behavioral Surveillance System Study Group. HIV infection and risk, prevention, and testing behaviors among injecting drug users — National HIV Behavioral Surveillance System, 20 U.S. cities, 2009. *MMWR Surveill Summ*. 2014; 63:1–51. [PMID: 24990587].
14. Centers for Disease Control and Prevention (CDC). Prevalence and awareness of HIV infection among men who have sex with men—21 cities, United States, 2008. *MMWR Morb Mortal Wkly Rep*. 2010; 59:1201–7. [PMID: 20864920]. [PubMed: 20864920]
15. Centers for Disease Control and Prevention (CDC). HIV testing and risk behaviors among gay, bisexual, and other men who have sex with men - United States. *MMWR Morb Mortal Wkly Rep*. 2013; 62:958–62. [PMID: 24280915]. [PubMed: 24280915]
16. Centers for Disease Control and Prevention. HIV Surveillance Report, 2013. Feb.2015 25 [July 15, 2015] Accessed at www.cdc.gov/hiv/library/reports/surveillance/.
17. Hall HI, Green TA, Wolitski RJ, Holtgrave DR, Rhodes P, Lehman JS, et al. Estimated future HIV prevalence, incidence, and potential infections averted in the United States: a multiple scenario analysis. *J Acquir Immune Defic Syndr*. 2010; 55:271–6. [PMID: 20634702]. DOI: 10.1097/QAI.0b013e3181e8f90c [PubMed: 20634702]
18. Centers for Disease Control and Prevention. [June 29, 2015] HIV/AIDS: Infographics and Posters. Accessed at www.cdc.gov/hiv/library/infographics.html?s_cid=bb-dhap-hiv-ngmhaad-001-ngmhaad
19. Weinstein MC, O'Brien B, Hornberger J, Jackson J, Johannesson M, McCabe C, et al. ISPOR Task Force on Good Research Practices—Modeling Studies. Principles of good practice for decision analytic modeling in health-care evaluation: report of the ISPOR Task Force on Good Research Practices—Modeling Studies. *Value Health*. 2003; 6:9–17. [PMID: 12535234]. [PubMed: 12535234]
20. Gold M. Panel on cost-effectiveness in health and medicine. *Med Care*. 1996; 34:DS197–9. [PMID: 8969326]. [PubMed: 8969326]
21. Tempalski B, Pouget ER, Cleland CM, Brady JE, Cooper HL, Hall HI, et al. Trends in the population prevalence of people who inject drugs in US metropolitan areas 1992–2007. *PLoS One*. 2013; 8:e64789. [PMID: 23755143]. doi: 10.1371/journal.pone.0064789 [PubMed: 23755143]

22. Centers for Disease Control and Prevention (CDC). HIV infection and HIV-associated behaviors among injecting drug users - 20 cities, United States, 2009. *MMWR Morb Mortal Wkly Rep*. 2012; 61:133–8. [PMID: 22377843]. [PubMed: 22377843]
23. Central Intelligence Agency. [12 August 2014] The World Factbook: United States. Accessed at www.cia.gov/library/publications/the-world-factbook/geos/us.html
24. Degenhardt L, Hall W, Warner-Smith M. Using cohort studies to estimate mortality among injecting drug users that is not attributable to AIDS. *Sex Transm Infect*. 2006; 82(Suppl 3):iii56–63. [PMID: 16735295]. [PubMed: 16735295]
25. Arias E. United States life tables, 2010. *Natl Vital Stat Rep*. 2014; 63:1–63. [PMID: 25383611].
26. Samji H, Cescon A, Hogg RS, Modur SP, Althoff KN, Buchacz K, et al. North American AIDS Cohort Collaboration on Research and Design (NA-ACCORD) of IeDEA. Closing the gap: increases in life expectancy among treated HIV-positive individuals in the United States and Canada. *PLoS One*. 2013; 8:e81355. [PMID: 24367482]. doi: 10.1371/journal.pone.0081355 [PubMed: 24367482]
27. Wada N, Jacobson LP, Cohen M, French A, Phair J, Muñoz A. Cause-specific mortality among HIV-infected individuals, by CD4(+) cell count at HAART initiation, compared with HIV-uninfected individuals. *AIDS*. 2014; 28:257–65. [PMID: 24105030]. DOI: 10.1097/QAD.000000000000078 [PubMed: 24105030]
28. Spire B, Lucas GM, Carrieri MP. Adherence to HIV treatment among IDUs and the role of opioid substitution treatment (OST). *Int J Drug Policy*. 2007; 18:262–70. [PMID: 17689374]. [PubMed: 17689374]
29. Sanders GD, Bayoumi AM, Sundaram V, Bilir SP, Neukermans CP, Rydzak CE, et al. Cost-effectiveness of screening for HIV in the era of highly active antiretroviral therapy. *N Engl J Med*. 2005; 352:570–85. [PMID: 15703422]. [PubMed: 15703422]
30. MacArthur GJ, Minozzi S, Martin N, Vickerman P, Deren S, Bruneau J, et al. Opiate substitution treatment and HIV transmission in people who inject drugs: systematic review and meta-analysis. *BMJ*. 2012; 345:e5945. [PMID: 23038795]. doi: 10.1136/bmj.e5945 [PubMed: 23038795]
31. Zaric GS, Barnett PG, Brandeau ML. HIV transmission and the cost-effectiveness of methadone maintenance. *Am J Public Health*. 2000; 90:1100–11. [PMID: 10897189]. [PubMed: 10897189]
32. Sullivan LE, Moore BA, Chawarski MC, Pantaloni MV, Barry D, O'Connor PG, et al. Buprenorphine/naloxone treatment in primary care is associated with decreased human immunodeficiency virus risk behaviors. *J Subst Abuse Treat*. 2008; 35:87–92. [PMID: 17933486]. [PubMed: 17933486]
33. Cipriano LE, Zaric GS, Holodniy M, Bendavid E, Owens DK, Brandeau ML. Cost effectiveness of screening strategies for early identification of HIV and HCV infection in injection drug users. *PLoS One*. 2012; 7:e45176. [PMID: 23028828]. doi: 10.1371/journal.pone.0045176 [PubMed: 23028828]
34. Kral AH, Lorvick J, Ciccarone D, Wenger L, Gee L, Martinez A, et al. HIV prevalence and risk behaviors among men who have sex with men and inject drugs in San Francisco. *J Urban Health*. 2005; 82:i43–50. [PMID: 15738321]. [PubMed: 15738321]
35. Pollack HA, D'Aunno T. HIV testing and counseling in the nation's outpatient substance abuse treatment system, 1995–2005. *J Subst Abuse Treat*. 2010; 38:307–16. [PMID: 20171038]. DOI: 10.1016/j.jsat.2009.12.004 [PubMed: 20171038]
36. Sanchez T, Finlayson T, Drake A, Behel S, Cribbin M, Dinunno E, et al. Centers for Disease Control and Prevention (CDC). Human immunodeficiency virus (HIV) risk, prevention, and testing behaviors—United States, National HIV Behavioral Surveillance System: men who have sex with men, November 2003–April 2005. *MMWR Surveill Summ*. 2006; 55:1–16. [PMID: 16826162].
37. National Institutes of Health. [15 July 2015] Starting antiretroviral treatment early improves outcomes for HIV-infected individuals [press release]. May 27, 2015 Accessed at www.nih.gov/news/health/may2015/niaid-27.htm
38. Westergaard RP, Ambrose BK, Mehta SH, Kirk GD. Provider and clinic-level correlates of deferring antiretroviral therapy for people who inject drugs: a survey of North American HIV

- providers. *J Int AIDS Soc.* 2012; 15:10. [PMID: 22360788]. doi: 10.1186/1758-2652-15-10 [PubMed: 22360788]
39. Zule WA, Bobashev GV, Wechsberg WM, Costenbader EC, Coomes CM. Behaviorally bisexual men and their risk behaviors with men and women. *J Urban Health.* 2009; 86(Suppl 1):48–62. [PMID: 19513854]. DOI: 10.1007/s11524-009-9366-3 [PubMed: 19513854]
 40. Montgomery JP, Mokotoff ED, Gentry AC, Blair JM. The extent of bisexual behaviour in HIV-infected men and implications for transmission to their female sex partners. *AIDS Care.* 2003; 15:829–37. [PMID: 14617504]. [PubMed: 14617504]
 41. Wejnert C, Le B, Rose CE, Oster AM, Smith AJ, Zhu J, Gabriela Paz-Bailey for the NHBS Study Group. HIV infection and awareness among men who have sex with men—20 cities, United States, 2008 and 2011. *PLoS One.* 2013; 8:e76878. [PMID: 24194848]. doi: 10.1371/journal.pone.0076878 [PubMed: 24194848]
 42. Kapadia F, Latka MH, Hudson SM, Golub ET, Campbell JV, Bailey S, et al. DUIT Study Team. Correlates of consistent condom use with main partners by partnership patterns among young adult male injection drug users from five US cities. *Drug Alcohol Depend.* 2007; 91(Suppl 1):S56–63. [PMID: 17329041]. [PubMed: 17329041]
 43. Long EF, Brandeau ML, Owens DK. The cost-effectiveness and population outcomes of expanded HIV screening and antiretroviral treatment in the United States. *Ann Intern Med.* 2010; 153:778–89. [PMID: 21173412]. DOI: 10.7326/0003-4819-153-12-201012210-00004 [PubMed: 21173412]
 44. Semaan S, Neumann MS, Hutchins K, D'Anna LH, Kamb ML, Project RESPECT Study Group. Brief counseling for reducing sexual risk and bacterial STIs among drug users—results from project RESPECT. *Drug Alcohol Depend.* 2010; 106:7–15. [PMID: 19720471]. DOI: 10.1016/j.drugalcdep.2009.07.015 [PubMed: 19720471]
 45. Sionean C, Le BC, Hageman K, Oster AM, Wejnert C, Hess KL, et al. NHBS study group. HIV Risk, prevention, and testing behaviors among heterosexuals at increased risk for HIV infection—National HIV Behavioral Surveillance System, 21 U.S. cities, 2010. *MMWR Surveill Summ.* 2014; 63:1–39. [PMID: 25522191].
 46. Davis KR, Weller SC. The effectiveness of condoms in reducing heterosexual transmission of HIV. *Fam Plann Perspect.* 1999; 31:272–9. [PMID: 10614517]. [PubMed: 10614517]
 47. Public Health Agency of Canada. [9 February 2015] HIV transmission risk: a summary of the evidence. February. 2013 Accessed at www.catie.ca/sites/default/files/HIV-TRANSMISSION-RISK-EN.pdf
 48. Hollingsworth TD, Anderson RM, Fraser C. HIV-1 transmission, by stage of infection. *J Infect Dis.* 2008; 198:687–93. [PMID: 18662132]. DOI: 10.1086/590501 [PubMed: 18662132]
 49. Pilcher CD, Tien HC, Eron JJ Jr, Vernazza PL, Leu SY, Stewart PW, et al. Quest Study. Brief but efficient: acute HIV infection and the sexual transmission of HIV. *J Infect Dis.* 2004; 189:1785–92. [PMID: 15122514]. [PubMed: 15122514]
 50. Castilla J, Del Romero J, Hernando V, Marincovich B, García S, Rodríguez C. Effectiveness of highly active antiretroviral therapy in reducing heterosexual transmission of HIV. *J Acquir Immune Defic Syndr.* 2005; 40:96–101. [PMID: 16123689]. [PubMed: 16123689]
 51. Marrazzo JM, Ramjee G, Richardson BA, Gomez K, Mgodini N, Nair G, et al. VOICE Study Team. Tenofovir-based preexposure prophylaxis for HIV infection among African women. *N Engl J Med.* 2015; 372:509–18. [PMID: 25651245]. DOI: 10.1056/NEJMoa1402269 [PubMed: 25651245]
 52. U.S. Bureau of Labor Statistics. Consumer Price Index. U.S. Department of Labor; Washington, DC: 2014.
 53. New York State Department of Health. [February 9, 2015] Pre-exposure prophylaxis (PrEP) to prevent HIV infection: questions and answers. Dec. 2012 Accessed at www.health.ny.gov/publications/0265/
 54. Paltiel AD, Weinstein MC, Kimmel AD, Seage GR 3rd, Losina E, Zhang H, et al. Expanded screening for HIV in the United States—an analysis of cost-effectiveness. *N Engl J Med.* 2005; 352:586–95. [PMID: 15703423]. [PubMed: 15703423]
 55. Goldhaber-Fiebert JD, Stout NK, Ortendahl J, Kuntz KM, Goldie SJ, Salomon JA. Modeling human papillomavirus and cervical cancer in the United States for analyses of screening and vaccination. *Popul Health Metr.* 2007; 5:11. [PMID: 17967185]. [PubMed: 17967185]

56. Karnon J, Vanni T. Calibrating models in economic evaluation: a comparison of alternative measures of goodness of fit, parameter search strategies and convergence criteria. *Pharmacoeconomics*. 2011; 29:51–62. [PMID: 21142278]. DOI: 10.2165/11584610-000000000-00000 [PubMed: 21142278]
57. Stinnett AA, Paltiel AD. Estimating CE ratios under second-order uncertainty: the mean ratio versus the ratio of means. *Med Decis Making*. 1997; 17:483–9. [PMID: 9343807]. [PubMed: 9343807]
58. Henry, J. [4 September 2015] Kaiser Family Foundation. U.S. federal funding for HIV/AIDS: the President's FY 2016 budget request. Apr. 2015 Accessed at <http://kff.org/global-health-policy/fact-sheet/u-s-federal-funding-for-hiv-aids-the-presidents-fy-2016-budget-request/>
59. Gilead Sciences, Inc. [1 April 2015] Gilead expands initiatives providing access to new HIV therapy Truvada in United States and abroad. Aug 18. 2004 Accessed at www.gilead.com/news/press-releases/2004/8/gilead-expands-initiatives-providing-access-to-new-hiv-therapy-truvada-in-united-states-and-abroad
60. Livoti C. HIV drugs to be uniquely shielded from generic pricing pressure in public payer scheme. *Financial Times*. Feb 28.2013 [1 April 2015] Accessed at www.ft.com/cms/s/2/286774b4-81be-11e2-b050-00144feabdc0.html#axzz43viyCqtY.
61. Holtgrave DR, Pinkerton SD, Jones TS, Lurie P, Vlahov D. Cost and cost-effectiveness of increasing access to sterile syringes and needles as an HIV prevention intervention in the United States. *J Acquir Immune Defic Syndr Hum Retrovirol*. 1998; 18(Suppl 1):S133–8. [PMID: 9663636]. [PubMed: 9663636]
62. Kwon JA, Anderson J, Kerr CC, Thein HH, Zhang L, Iversen J, et al. Estimating the cost-effectiveness of needle-syringe programs in Australia. *AIDS*. 2012; 26:2201–10. [PMID: 22914579]. DOI: 10.1097/QAD.0b013e3283578b5d [PubMed: 22914579]
63. Blankenship KM, Smoyer AB, Bray SJ, Mattocks K. Black-white disparities in HIV/AIDS: the role of drug policy and the corrections system. *J Health Care Poor Underserved*. 2005; 16:140–56. [PMID: 16327113]. [PubMed: 16327113]
64. Lehman DA, Baeten JM, McCoy CO, Weis JF, Peterson D, Mbari G, et al. Partners PrEP Study Team. Risk of drug resistance among persons acquiring HIV within a randomized clinical trial of single- or dual-agent preexposure prophylaxis. *J Infect Dis*. 2015; 211:1211–8. [PMID: 25587020]. DOI: 10.1093/infdis/jiu677 [PubMed: 25587020]
65. Supervie V, Barrett M, Kahn JS, Musuka G, Moeti TL, Busang L, et al. Modeling dynamic interactions between pre-exposure prophylaxis interventions & treatment programs: predicting HIV transmission & resistance. *Sci Rep*. 2011; 1:185. [PMID: 22355700]. doi: 10.1038/srep00185 [PubMed: 22355700]
66. Basu S, Andrews J. Complexity in mathematical models of public health policies: a guide for consumers of models. *PLoS Med*. 2013; 10:e1001540. [PMID: 24204214]. doi: 10.1371/journal.pmed.1001540 [PubMed: 24204214]
67. Alistar SS, Owens DK, Brandeau ML. Effectiveness and cost effectiveness of oral pre-exposure prophylaxis in a portfolio of prevention programs for injection drug users in mixed HIV epidemics. *PLoS One*. 2014; 9:e86584. [PMID: 24489747]. doi: 10.1371/journal.pone.0086584 [PubMed: 24489747]
69. Koblin BA, Chesney MA, Husnik MJ, Bozeman S, Celum CL, Buchbinder S, et al. EXPLORE Study Team. High-risk behaviors among men who have sex with men in 6 US cities: baseline data from the EXPLORE Study. *Am J Public Health*. 2003; 93:926–32. [PMID: 12773357]. [PubMed: 12773357]
70. Bellis DJ. Reduction of AIDS risk among 41 heroin addicted female street prostitutes: effects of free methadone maintenance. *J Addict Dis*. 1993; 12:7–23. [PMID: 8381030]. [PubMed: 8381030]
72. Barnett PG, Chow A, Joyce VR, Bayoumi AM, Griffin SC, Sun H, et al. Effect of management strategies and clinical status on costs of care for advanced HIV. *Am J Manag Care*. 2014; 20:e129–37. [PMID: 25326927]. [PubMed: 25326927]
73. Tengs TO, Lin TH. A meta-analysis of utility estimates for HIV/AIDS. *Med Decis Making*. 2002; 22:475–81. [PMID: 12458977]. [PubMed: 12458977]

74. Quinn TC, Wamer MJ, Sewankambo N, Serwadda D, Li C, Wabwire-Mangen F, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. *N Engl J. Med.* 2000; 345:921–9. [PMID: 10738050].

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. Main analysis: incremental costs incurred and QALYs gained

To assess delivery scenarios, we evaluated 1) PrEP alone with status quo population-level screening rates for people who inject drugs; 2) PrEP with HIV screening every 3 mo (PrEP +screen); and 3) PrEP+screen with prompt and sustained linkage to ART for individuals who do become infected (PrEP+screen+ART). All cases assume 25% coverage, 49% PrEP efficacy, and \$10 000 annual PrEP drug cost. The *x*-axis shows incremental cost in billions of U.S. dollars compared with the status quo of no PrEP; the *y*-axis shows incremental QALYs in thousands. The point labeled PrEP+screen shows that at current levels of ART initiation and adherence, screening provides a small QALY increase for a large cost increase. When screening is combined with increased ART, however, screening delivers synergistic benefit, resulting in an ICER of \$253 000/QALY gained. This scenario, PrEP+screen+ART, dominates the other 2 scenarios (i.e., is more effective and has a lower ICER). ART = antiretroviral therapy; ICER = incremental cost-effectiveness ratio; PrEP = preexposure prophylaxis; QALY = quality-adjusted life-year.

Figure 2. Sensitivity analysis: PrEP value as a function of drug efficacy and drug cost

We evaluate a program of PrEP with frequent HIV screening and enhanced ART provision (PrEP+screen+ART) with 25% coverage and \$10 000 annual PrEP drug cost varying PrEP efficacy from 10% to 90%. The *y*-axis shows the ICER corresponding to efficacy levels specified on the *x*-axis. At current drug costs, the cost per QALY gained remains greater than \$150 000 for all efficacy levels but substantially decreases as efficacy improves, with the largest jumps in cost-effectiveness coming at lower levels of efficacy. At 49% efficacy (leftmost vertical line), we see our base-case analysis with an ICER of \$253 000/QALY gained. The Bangkok Tenofvir Study estimates a 74% reduction (rightmost vertical line) in HIV acquisition for high adherers. This results in a more favorable ICER of \$193 000/QALY gained. At a 65% cost reduction PrEP delivers higher value and crosses the \$100 000/QALY gained threshold at reported efficacy levels, although low levels of efficacy still result in high ICERs. ART = antiretroviral therapy; ICER = incremental cost-effectiveness ratio; PrEP = preexposure prophylaxis; QALY = quality-adjusted life-year.

Table 1

Estimates for Key Model Parameters

Parameter [*]	Value	Range [†]	Reference
Demographics, <i>n</i>			
Population (age 18–64 y)			
Total	200 million	198–203 million	1, 23
PWID	1.1 million	0.8–1.4 million	2, 3, 21
MSM	3.0 million	2.0–4.5 million	11, 18
Initial HIV conditions, %			
HIV prevalence			
PWID	9.8	8.0–12.1	4
MSM	11.4	8.8–14.7	11, 14, 17
Low-risk	0.18	0.14–0.23	Calculated (16, 17, 23)
Aware of HIV status			
PWID	69.9	59.1–78.6	4, 13
MSM	75.3	64.3–84.4	14, 15, 18
Low-risk	84.7	76.3–89.9	18
ART enrollment given aware			
PWID	40.9	29.3–51.7	18, 28
MSM	53.1	41.5–64.5	18
Low-risk	73.7	58.4–82.6	18, 33
Annual mortality rate			
PWID	0.015	0.011–0.021	24, 25
Hazard ratio for opioid agonist therapy	0.39	0.12–0.96	24, 33
MSM	0.0044	0.0034–0.0053	11, 25
Low-risk	0.0015	0.0011–0.0021	25
HIV natural history (average disease duration)			
Acute HIV, <i>mo</i>	2.2	1.2–4.6	48, 49
Asymptomatic HIV, <i>y</i>	5.3	3.1–9.8	21, 29, 48
Symptomatic HIV, <i>y</i>	7.1	4.5–14.4	21, 29, 48
Symptomatic HIV receiving ART, <i>y</i>	20.0	12.6–37.4	26, 27, 48
AIDS, <i>y</i>	1.7	0.9–3.8	48
AIDS receiving ART, <i>y</i>	5.3	3.1–9.8	26, 27, 48
ART (transmission reduction if partner is receiving ART), %			
Injecting partner	59	14–82	8, 43
Sexual partner	90	68–99	43, 48, 50
PrEP			
Infection reduction (injection and sexual routes), %	48.9	10.0–89.1	8
Screening frequency [‡] , <i>mo</i>	3	1.2–4.9	9
Injecting behavior			
Risky injections annually, <i>n</i>	48	17–89	4, 13, 31, 42
Probability of transmission per injection, asymptomatic HIV, %	0.05	0.03–0.07	47, 48

Parameter*	Value	Range [†]	Reference
Relative risk for transmission per injection			
Symptomatic relative to asymptomatic	2.85	2.85–3.28	47, 48
AIDS relative to symptomatic	2.49	1.99–3.17	47, 48
Acute relative to AIDS	3.45	2.33–5.10	47 – 49
Decrease in risky injections due to awareness of HIV status	23.2	0.0–55.0	33, 34
Opioid agonist therapy, %			
Initial PWID enrollment	24.8	13.2–34.4	4, 13, 31, 33, 34
PWID who quit annually	32.1	18.1–43.6	31, 70
PWID who transition to lower-risk groups annually	3.6	1.9–5.4	31, 70
Decrease in risky injections	54.7	0.004–82.2	30
Sexual behavior			
Annual partners, <i>n</i>			
PWID	3.1	1.2–5.4	4, 34, 42, 44
MSM	4.1	2.4–6.3	11, 34, 39, 69
Low-risk	1.5	0.8–2.5	33, 43–45
Partnerships, %			
PWID with other PWID	45.6	27.0–65.8	33, 42
PWID with MSM	8.0	4.9–11.2	4, 34, 39
MSM with other MSM	92.9	73.6–99.9	36, 39–41
Monthly probability of transmission in heterosexual partnership, asymptomatic HIV	0.87	0.49–1.29	43, 47, 48, 74
Relative risk for transmission per sexual partnership			
Symptomatic relative to asymptomatic	2.85	2.85–3.28	47, 48
AIDS relative to symptomatic	2.49	1.99–3.17	47, 48
Acute relative to AIDS	3.45	2.33–5.10	47 – 49
MSM partnership relative to heterosexual asymptomatic	4.46	2.45–7.27	43, 47
Annual costs, 2015 U.S. \$			
Uninfected PWID	3000	630–7200	31, 33
Uninfected non-PWID	5000	3060–7440	33
Opioid agonist therapy	7000	4320–10 430	31, 33
PrEP drug	10 000	7150–13 320	10–12, 53
PrEP screening services	800	100–2240	11, 54
ART	15 000	9170–22 300	29, 43, 54, 72
Asymptomatic HIV, not receiving ART	4000	2460–5950	29, 31, 43, 54, 72
Symptomatic HIV, not receiving ART	7000	4270–10 330	29, 31, 43, 54, 72
Symptomatic HIV, receiving ART [§]	6500	4190–9280	29, 31, 43, 54, 72
AIDS, not receiving ART	20 040	10 380–33 240	29, 31, 43, 54, 72
AIDS, receiving ART [§]	10 000	4870–17 160	29, 31, 43, 54, 72
Annual discount rate, %	3		19, 20
Quality-of-life multipliers ^{//}			
Uninfected (non-PWID)	1.00		11, 43, 67
PWID	0.90	0.67–1.00	43, 67

Parameter [*]	Value	Range [†]	Reference
PWID receiving opioid agonist therapy	0.95	0.82–1.00	31, 33, 67
Asymptomatic HIV	0.94	0.76–1.00	43, 73
Symptomatic HIV	0.81	0.68–0.95	43, 73
AIDS	0.70	0.57–0.86	43, 73
ART multiplier	1.15	1.00–1.35	11, 29, 43, 67

ART = antiretroviral therapy; MSM = men who have sex with men; PrEP = preexposure prophylaxis; PWID = people who inject drugs.

^{*} Additional model parameters in **Table 1** of the **Supplement** (available at www.annals.org).

[†] Range refers to the 95% CI used in probabilistic sensitivity analysis.

[‡] Denotes a control variable characterizing a specified intervention and is not considered uncertain. The associated range was used in 1-way sensitivity analysis only.

[§] Disease costs during ART exclude cost of ART.

^{||} The quality weight for any given compartment is capped at the underlying risk-group baseline quality weight (that is, cannot exceed 1).

Table 2

Results of Main Cost-Effectiveness Analysis

Scenario [*]	PWID HIV Infections Averted, <i>n</i> [†]	Total Population HIV Infections Averted, <i>n</i> [†]	Change in PWID HIV Prevalence at 20 y, % [†]	Total Costs, U.S. \$ (Trillions)	Total QALYs (Billions)	Incremental Costs, U.S. \$ (Billions) [†]	Incremental QALYs (Thousands) [†]	ICER, \$ [†]
No PrEP	–	–	–	32.528	6.4340	–	–	–
PrEP	18 600	21 800	–12.9	32.568	6.4341	39.9	126	Dominated
PrEP+screen	19 800	23 700	–14.1	32.571	6.4341	42.8	132	Dominated
PrEP+screen+ART	21 500	26 700	–14.0	32.572	6.4342	43.8	173	253 000

ART = antiretroviral therapy; ICER = incremental cost-effectiveness ratio; PrEP = preexposure prophylaxis; PWID = people who inject drugs; QALY = quality-adjusted life-year.

^{*} Every PrEP scenario assumes 25% coverage, or approximately 280 000 PWID enrolled, and 49% reduction in HIV acquisition from PrEP. In PrEP+screen, PrEP is delivered with HIV screening every 3 mo. PrEP+screen+ART includes enhanced linkage to ART for individuals who do become infected. (**Supplement**, section 7.2, available at www.annals.org, provides details and intuition on prevalence changes for each intervention.) PrEP+screen+ART (weakly) dominates the other 2 scenarios because it is more effective and less expensive per QALY gained. Therefore, PrEP+screen+ART is compared directly with no PrEP, which is the next-best alternative on the cost-effectiveness frontier.

[†] Relative to no-PrEP scenario.

Table 3

Sensitivity Analysis: PrEP Value as a Function of Drug Cost

Scenario [*]	Total Costs of PrEP, U.S. \$ (Trillions)	Total QALYs (Billions)	Incremental Costs, U.S. \$ (Billions) [†]	Incremental QALYs (Thousands) [†]	ICER, \$ [†]
No PrEP	32.528	6.4340	–	–	–
PrEP at \$2000 (80% price reduction)	32.539	6.4342	11.2	173	65 000
PrEP at \$3500 (65% price reduction)	32.545	6.4342	17.3	173	100 000
PrEP at \$6000 (40% price reduction)	32.555	6.4342	27.5	173	159 000
PrEP at \$10 000 (current price)	32.572	6.4342	43.8	173	253 000
PrEP at \$14 000 (40% price increase)	32.588	6.4342	60.0	173	347 000

ICER = incremental cost-effectiveness ratio; PrEP = preexposure prophylaxis; QALY = quality-adjusted life-year.

^{*} Every PrEP scenario assumes 25% coverage in a PrEP program with HIV screening every 3 mo and enhanced linkage to antiretroviral treatment (ART) for individuals who do become infected (PrEP+screen+ART), where PrEP is 49% effective. Costs are annual drug costs.

[†] Relative to no-PrEP scenario.